



HANG TUAH UNIVERSITY
FACULTY OF DENTISTRY PRESENT
INTERNATIONAL SCIENTIFIC MEETING

PROCEEDING BOOK

Dentisphere 3

Dentistry Update & Scientific Atmosphere

26th-27th, August 2016

Shangri-La Hotel
Surabaya-Indonesia



*Current Concepts and Technology
in Improving Dental and Oral Health Care*

ISBN 978-602-14590-1-0

PROCEEDING BOOK INTERNATIONAL SCIENTIFIC MEETING

**3rd DENTISPHERE (DENTISTRY UPDATE & SCIENTIFIC ATMOSPHERE)
CURRENT CONCEPTS AND TECHNOLOGY IN IMPROVING DENTAL AND ORAL
HEALTH CARE**

REVIEWER :

PROF. FUMIAKI KAWANO, DDS, Ph.D, FACULTY OF DENTISTRY TOKUSHIMA UNIVERSITY, JAPAN
PROF JOONGKI-KOOK SCHOOL OF DENTISTRY CHOSUN UNIVERSITY, KOREA
DRG HENI SUSILOWATI M.KES Ph.D, FKG UGM
DR. DIAN MULAWARMANTI, DRG, M.S, FKG UHT
DR KRISTANTI PARISIHNI, DRG, M.KES FKG UHT
DR. NOENGKI PRAMESWARI, DRG, M.KES FKG UHT
DRG. MEINAR NUR ASHIRIN, Ph.D FKG UHT

EDITOR :

DRG DIAN WIDYA DAMAIYANTI, M.KES
DRG AGNI FEBRINA P , M.KES
CARISSA ENDIANASARI, S.ST
RIZA FATMA WARDANI, AMD.AK

SETTING/LAY OUT :

DRG. DIAN WIDYA DAMAIYANTI, M.KES
CARISSA ENDIANASARI, S.ST

COVER DESIGN :

MONICA VITA, SKG

PRINTED AND PUBLISHED BY:

FKG HANG TUAH SURABAYA.PRESS
JL. ARIF RAHMAN HAKIM NO.150 SURABAYA 60111
TELP. 031-5945864, FAX. 031-5946261
WEBSITE: www.hangtuah.ac.id
Cetakan : SURABAYA, 2016-06-29

ISBN 978-602-14590-1-0

CONTENTS

DEAN OF FACULTY DENTISTRY HANG TUAH UNIVERSITY

WELCOME NOTE

CHAIRMAN 3RD DENTISPHERE WELCOME NOTE

CONTENTS

MAIN LECTURER

ML.1	Oclusal Schemes in Complete Denture Prof Fumiaki Kawano	1
ML.2	Achieving Aesthetic and Excellence with Modern Composite Dr. Anthony Tay, BDS	2
ML.3	Porous Titanium for Bone Substitute Materials Assoc. Prof. Yoshihito Naito, DDS., PhD	3
ML.4	The role of dentist in mass disaster AKBP Drg. Ahmad Fauzi, MM, GDipForOdont	4
ML.5	Basic research for development of oral hygiene products Prof Joong Ki-Kook	5
ML.6	Dental Readiness in Military Dentistry Kol. Laut (K) Ridwan Purwanto, drg., MARS - Ladokgi	6
ML.7	Occlusion Update : A Whole Elephant Perspective Dr. Yue Weng Cheu, BDS., FRACDS.,MJDF, RCSEng	7
ML.8	Things about root canal dilacerations Marino Sutedjo., drg., SpKG	8
ML.9	Irrigation at The One-Third of The Apical Root HM Bernard O Iskandar, drg., SpKG	9
ML.10	Emulating Nature : Dental Photography and Clinical Connection OnnyEryanto, drg	10
ML.11	Restorative Challenges and Treatment Option for Primary Teeth Assoc. Prof . Nagarajan M.PS	11
ML.12	Biological Response Around Graft and Implant Ika Dewi Ana, drg.,PhD	12

ML.13	Current concepts of dental caries in children Udijanto Tedjosongko, drg., PhD	13
ML.14	Exploration of Marine Biota and Hyperbaric Oxygen Therapy in Dentistry Dr Dian Mulawarmanti, drg, M.S	14
ML.16	Timing of Orthodontic Treatment Dr. Retno Widayati., drg., SpOrt (K)	15

SHORT LECTURER/ORAL PRESENTATION

SL. 1.1	Effect of Piper betle L. Leaves Extract In The Formation of Dental Plaque: Literature Review Poetry Oktanauli, Radinda Myrna Andiani	17
SL. 1.2	Treatment of Temporomandibular Disorder Using Full Occlusal Splint Erna Fakhriyana, Harry Laksono	25
SL. 1.4	Impression Technique Using A Sectional Impression Tray in Scleroderma's Patient : A Case Report Elin Hertiana	30
SL. 1.5	Effect of Denture Disinfection with Microwave to Dimensional Change and Water Sorption PutriWelda Utami Ritonga, Vincent	41
SL. 2.1	OrthodonticTreatment with Removable Appliance Pricillia Priska Sianita	46
SL. 2.4	Orthodontic Treatment Disharmony Dento Maxillare (DDM) by Extraction 4 First Premolare Paulus Maulana Soesilo Soesanto	53
SL. 2.5	Complete Examination Of Temporo Mandibular Joint for Detection in Temporo Mandibular Joint Disorder Samson Peter Louis Alfredo	59
SL. 2.6	RADIOGRAPHY ROLE IN FORENSIC IDENTIFICATION ON DISASTER Emy Khoironi	66
SL. 2.8	Biologic Width Concept In Gingivectomy Surgery (Case Report) Desy Fidyawati	74
SL. 2.9	Effect of Smoking on Gingival Melanin Pigmentation (Case Report) Veronica Septnina Primasari	81

SL. 2.10	The Influence of Interproximal Interface Towards Periodontal Tissue Billy Martin	86
SL. 2.11	Tissue Movement for Better Results in Preprosthetic Reconstructive Surgery: Case Report Britaria Theressy, Agung Krismariono	96
SL. 2.14	Distribution of Candida Species in Oral Candidiasis on Injection Drug User Fatma Yasmin Mahdani, Adiastuti Endah Parmadiati, Hening Tuti Hendarti, Annete Juwita Yukuri	107
SL. 2.15	<i>Comprehensive Approach of Severe Early Childhood Caries in Child with Post-palatoplasty: A Case Report</i> Lusiana Beatrice, Meirina Gartika	113
SL. 2.17	The Artistic Value of Gummy Smile Treatment Steffi Purnomo, Poernomo Agoes Wibisono	122
SL. 2.18	Management of Post Stroke Complete Edentulous Patient Using Suction Effective Method Rizki Purnamasari Nugraheni, Harry Laksono	127
SL. 2.19	Preschool Caries With Pufa Index In Summersari Districts Jember Ristya Widi Endah Yani	132
SL. 2.20	The Use of Pekkton® on Telescopic Crowns in Complete Overdenture: a Clinical Case Tika Rahardjo, Utari Kresnoadi, Harry Laksono	137
SL. 2.21	TREATMENT OF PATIENTS AMELOGENESIS IMPERFECTA WITH FULL VENEER METAL PORCELAIN CROWN (CASE REPORT) Fransiska Nuning Kusmawati	145
SL. 2.22	Restoring Facial Harmony and Chewing Function of Post Maxillectomy Patients: Rehabilitation of Maxillofacial Patients Widaningsih, Benny Dwi Cahyo	151
SL. 2.24	Zirconia All-Ceramic Bridge For Aesthetic Restoration Meinar Nur Ashrin, Ghita Hadi Hollanda	157
SL. 2.26	<i>Sticophushermanii</i> Extract Affected The Expression of TLR-4 and TNF- α in Periodontitis Induced by <i>Porphyromonas gingivalis</i> Kristanti Parisihni, Eddy Bagus Wasito, Retno Indrawati	163
SL. 2.27	<i>Integrin A2β1</i> And <i>Bmp-2</i> Regulated In Bone Remodelling To Accelerate Orthodontic Tooth Movement By Giving <i>Stichopus Hermanii</i> Noengki Prameswari, Arya Brahmanta	171

SL. 2.28	THE EXPRESSION OF MACROPHAGE CELL ON WOUND HEALING PROCESS IN RATTUS NORVEGICUS USING CHITOSAN GEL WITH DIFFERENT MOLECULAR WEIGHT Sularsih	178
SL. 2.29	EFFECTS OF <i>Stichopus hermanii</i> ETHANOLIC EXTRACT ON TLR-2 AND IL-17 EXPRESSION IN RATS WITH ORAL CANDIDIASIS IMMUNOSUPPRESSED MODEL Dwi Andriani, Syamsulina Revianti, Kristanti Parisihni	185
SL. 2.30	TGF- β 1 Expression on Traumatic Ulcer Healing Process Treated with Water Extract Gold Sea Cucumber Dian W Damaiyanti	193

POSTER PRESENTATION

P 1.3	Combination Technique For Gingival Depigmentation (Laporan Kasus) Tomy Juliyanto, Agung Krismariono	203
P1.4	Efek Terapi Oksigen Hiperbarik Dikombinasi Dengan Pemberian Bubuk Teripang Emas (<i>Stichopus hermanii</i>) terhadap Kadar Gula Darah pada Tikus Wistar Diabet yang Diinduksi Bakteri <i>Porphyromonas gingivalis</i> Rafika Rusydia Darojati, Yoifah Rizka, Syamsulina Revianti	209
P 1.8	The Comparison of Osteoblast and Osteoclast in the Pressure area and Tension area on Tooth Movement Because of Hyperbaric Oxygen Therapy Rizta Riztia Budianti, Rizki Kartika Putra, Arya Brahmanta	217
P 1.9	ComparisonOf Color Changes In Thermoplastic Nylon Resin Denture Base Material Soaked In Black Tea Debby Saputera, April Yastuti Rosandita, Dewi Puspitasari	232
P 1.13	The Effect of Alkaline Peroxide and Celery Extract (<i>Apium Greveolens</i> .L) 75% Solution to Flexural Strength of Heat Cured Typed Acrylic Resin Dewi Puspitasari, Reni Hamyulida, Debby Saputera	240
P 1.15	The Relation Of Body Mass Index StatusWith Dental Caries And Permanent Teeth Eruption Overview On Elementary School Students In District Hss Grade 1, 2, And 3 Rizki Indah Permatasari, RosihanAdhani, BayuIndraSukmana	247
P 1.16	Fluoride Concentration On Mice Teeth After Application Naf Patch On Back Mice That Shaved Manually And Ellectrically Diyah Fatmasari, Alya Maqdani	252

P 1.20	Management Of Maxillary Flat Edentulous Ridge With Magnetic Retained Immediate Complete Denture Ratih Prasetyowati, Mefina Kuntjoro, Harry Laksono	258
P 1.21	How to Manage Single Denture Syndrome?(Case Report) Primanda Nur Rahmania, Harry Laksono, Utari Kresnoadi	263
P 1.23	Maxillary Bare Root Complete Overdenture with Mandibular Removable Partial Denture Olivia Puspitasari Surya, Eha Djulaeha, Agus Dahlan	267
P 1.24	Precision Attachment Removable Partial Denture Is The Best Choice For Unilateral Free End Edentulous Ridge (Case Report) Happy Indra Bakhti, Agus Dahlan, Rostiny	271
P 1.25	Changes Spectrum Of Sound Frequency Consonant 'S' After Crossbite 21 Corrected Ani Subekti, Rinaldi Budi Utomo	275
P 1.26	Magnetic Attachment Retained Complete Overdenture As Treatment For Flat Alveolar Ridge (Case Report) Karina Mundiratri, Eha Djulaeha, Agus Dahlan	281
P 1.27	The Use of Facebow Transfer with Free-plane Articulator Marchello Marvin, Rostiny,Sukaedi	286
P 1.28	Management of Patient with Dentoalveolar Compensation and Ridge Resorption in Prosthodontics Herautami Caezar YS, Kris Biantoro, Harr Laksono, Eha Djulaeha	290
P 1.29	Management of Edentulous Patient Using Biofunctional Prosthetic System (BPS) Atika Rahmadina, Harry Laksono, Eha Djulaeha	298
P 2.33	Oropharyngeal Candidiasis in Diabetes Mellitus Patient Using Oral Glucosamine Hastin Sofyana, Hening Tuti Hendarti	303
P 2.34	Management Of Herpangina In A Young Adult Patient Ade Puspa Sari, Desiana Radithia	312
P 2.35	Manifestation of Recurrent Oral Ulceration Associated to Reactivation Rheumatic Heart Disease Silfra Yunus Kende, Rindang Tanjungsari, Adiastruti Endah, Desiana Raditya, Diah Savitri Ernawati	319

P 2.37	The Effectiveness of Snake And Ladder Game Method on Small Dentist Cudres' Level of Knowledge and Students' Oral Hygiene Hestieyonini Hadynanawati, Kiswaluyo, Zahara Meilawaty, Ristya Widi Endah Yani	329
P 2.39	Indirect Porcelain Veneer To Fix Instantly Palatoversi Tooth (Case Report) Diana Soesilo	337
P 2.40	Prosthetic Rehabilitation of a Partially Edentulous Patient with Chronic Periodontitis Chaterina Diyah Nanik.K	344
P 2.41	Apex Resection On Post Endodontic Treatment Tooth With Periapical Cystic (Case Report) Fani Pangabdian	353
P 2.42	Oral Mucocele in Pediatric Patient : a Case Report Ayulistya Paramita, Ghita Hadi Hollanda	359
P 2.47	<i>Expression of Osteopontin And Osteoblasts After Given Alloplast With PRF Compare To XenografWithPRF OnBone Defect</i> Hansen Kurniawan, Iwan Ruhadi, Noer Ulfah	365
P 2.48	An Obturator Bottle Feeding Appliance For A Newborn Baby With Cleft Palate Dika Agung Bakhtiar, Agus Dahlan	371
P 2.51	Maxillary Attachment Retained Removable Partial Denture And Mandibular Magnetic Retained Overdenture : A Case Report Rangga Surya Fathrianto, Harry Laksono	376
P 2.55	Direct Class II Resin Composite Restoration on Maxillary Right Posterior Tooth Diani Prisinda, Prilanita Gian	381

SL 2.11

CASE REPORT

Tissue Movement for Better Results in Preprosthetic Reconstructive Surgery

Britaria Theressy*, Agung Krismariono**

*Resident of Periodontics, Faculty of Dentistry, Airlangga University Surabaya

**Department of Periodontics, Faculty of Dentistry, Airlangga University Surabaya

ABSTRACT

Background: Alveolar bony defects can occur due to advance periodontal disease. These defects often cause a significant problems in dental treatment and rehabilitation. Many techniques exist for hard tissue augmentation. These techniques are based on specific procedures to perform preprosthetic rehabilitation. This article presents case report of preprosthetic surgery with severe bone loss in two different techniques. **Purpose:** To determine the superior techniques of two methods in preprosthetic surgery with severe bone loss. **Case and case management:** A nonsmoker 40 years old male was first examined in author's department in August 2015 with severe anterior mandible bone loss. The diagnosis was chronic periodontitis. His treatment plan included tooth extraction of hopeless teeth, bone augmentation and implant. Nonsurgical treatment was performed on all teeth, preprosthetic surgery with severe bone loss performed in anterior mandible. The first technique, incision made circularly right on each tooth sulcus without any horizontal incision on the apical of interdental papillae. Whereas the second technique, horizontal and vertical incision on the apical of interdental papillae was performed. **Results:** Three months post surgery, clinically the alveolar ridge level has increased. Radiographic evaluation: there are better bone formation after the second surgery either increasing of bone level condition as well as bone density and periodontal space recovery. **Conclusion:** The surgical technique with incision right on each sulcus of tooth and incision horizontally and vertically on the apical of interdental papillae led to novel possibilities to regenerate alveolar bone.

Keywords: Alveolar bony defects, chronic periodontitis, preprosthetic surgery techniques

Correspondence: Britaria Theressy, Resident of Periodontics, Faculty of Dentistry, Airlangga University, Jalan Mayjen Prof. Dr. Moestopo no. 47 Surabaya, Phone (031) 5030255, 5020256, E-mail: Iyad1519@gmail.com

BACKGROUND

The hallmarks of periodontal disease are inflammation and alveolar bone loss. Gingival inflammation can resolve but alveolar bone loss leads to eventual tooth loss. Resorption of alveolar bone is a common sequel of tooth loss and presents a clinical problem, especially in the esthetic zone.¹ Advanced alveolar bone loss (>7 mm) may result in esthetically and functionally compromised dental prosthesis like removable and fixed partial dentures and ideal implant placement in prosthetically driven position.² The deficient alveolar ridges always present with numerous challenges to the clinician for achieving predictable augmentation.³⁻⁹ The end goal of the therapy is to provide a functional restoration that is in harmony with the adjacent natural dentition. Thus augmentation of bone is often necessary.¹ Advances in biologic understanding of different bone regenerating materials and continuous innovations in surgical techniques have led to increased predictability in reconstruction of alveolar ridge defect.¹⁰

Bone is a dynamic structure with a continuous remodeling to ensure renewal of form and function.¹ Although bone tissue exhibits a large regeneration potential and may restore its original structure and function completely, bony defects may often fail to heal with bone tissue.¹¹ In order to facilitate and/or promote regeneration of lost bone, a variety of techniques may be employed, including particulate grafting, membrane use, block grafting, and distraction osteogenesis, either alone or in combination.¹ The biologic mechanisms forming the basis of bone

augmentation include three basic processes: osteogenesis, osteoconduction, and osteoinduction.¹¹ Bone augmentation techniques may be used for the applications of extraction socket defect grafting, horizontal ridge augmentation, vertical ridge augmentation, and sinus augmentation.¹ Many of these reconstructive efforts are limited in the efficacy due to inadequate flap coverage and vascular perfusion. Procedures to prevent the collapse of the alveolar ridge are highly technique sensitive and require different surgical designs depending upon the size of the defect.¹² Several flap techniques have a general concept associated with alveolar ridge augmentation. This include generating blood supply, protected shape of bone growth and achieving tension free flap in wound closure.¹³ In this article we presents two different techniques of flap design in preprosthetic surgery with severe alveolar bone loss for optimizing dental implant placement.

CASE REPORT

A nonsmoker 40-years-old male patient was first examined in Author's department in August 2015 with mobility in central incisors of mandible since 6 months, due to trauma wanted a permanent replacement of his mobile teeth. Clinical examination [Figure 1], radiographs [Figure 2], and a thorough history were obtained from the patient. On clinical examination were found as follows: grade III mobility of #32, grade II mobility of #34, generalized probing pocket depth 6-8mm of teeth on the mandible, missing teeth of #36 and #46 and generalized malposition

of teeth. The mandible anterior ridge deficiency was noticed. To evaluate the bone loss, panoramic radiographs was performed. The condition of periodontal tissues were found to be as followed: lack of bone density and supraposition of #32 and angular bone loss of #34. A final diagnosis of Generalized Chronic Periodontitis with severe bone loss on the mandible was established.

We decided his treatment plan included dental health education, scaling and root planning, splinting, occlusal adjustment, tooth extraction

of hopeless 32, bone augmentation and dental implant. Finding that the available bone condition was inadequate for an implant, we decided to horizontally augment the implant recipient site which is on region 32 with a suitable substitute. Either on 34, we decided to horizontally augment the angular bone loss. The complete treatment plan was explained to the patient and signed informed consent was obtained. After completion of initial treatment, surgical procedure was carried out at a later appointment.



Figure 1. clinical examination on first examined




Figure 2. radiographic examination before treatment

CASE MANAGEMENT

Surgical Procedure was performed 1 week after nonsurgical procedure.

Procedure of surgical I, bone augmentation on #32

1. After local anesthesia, a full thickness flap was performed which is incision made right on sulcus of #33 [Figure 3,4]. An incision at the sulcus was used to preserve interdental papillae.
2. #31 was extracted [Figure 5], followed by curettage of granulation tissue in tooth socket post extraction.
3. Incision was performed using Gracey curette instrument. While at incised, smoothing and

scrapping was performed to get rid of existing granulation tissue on the root surface using Gracey.

4. Bovine bone grafts, product of tissue bank of Dr. Soetomo Hospital Surabaya, was applied after all soft tissues were thoroughly removed from the recipient site. Once properly positioned, grafts was covered by released mucosa and then sutured [Figure 6,7].
5. For oral administration, amoxicillin 500mg t.i.d 3 days, doxycycline 20mg b.i.d 30 days, mefenamic acid 500 mg t.i.d Bio ATP s.i.d, calcitrol s.i.d and hyaluronic acid gel, were initiated.



Figure 3. flap incision using Gracey curette on the sulcus of #33



Figure 4. flap incision on the sulcus



Figure 5. The socket post extraction of #32



Figure 6. Bone grafts applied into the socket post removal of granulation tissues

Procedure of surgical I of bone augmentation #34

Bone augmentation on #34 was performed 3 weeks after augmentation #32. The flap design and techniques of bone augmentation #34 was similar to

#32 augmentation. The incision of flap was made right on each sulcus on #33 and #34 using Gracey curette followed by smoothing and scrapping on the tooth surface. This flap design was used to preserve the interdental papillae [Figure 8,9].



Figure 7. sutured using non absorbable silk 5.0



Figure 8. The flap incision on #33 and #34



Figure 9. bone grafts applied after removal of granulation tissues

After 5 months of healing post first surgery, on clinical evaluation there is a concave shape on the labial side of region #32 [Figure 10,11]. This means still has bone defect on the labial side of region #32. On radiographic evaluation, the alveolar bone density on region #32 has not

increased. The alveolar bone on #34 has increased [Figure 12]. Based on these results, we decided to perform second surgery of alveolar bone augmentation on region #32 and #34 in order to achieve adequate ridge width and bone density to facilitate the placement of implant.



Figure 10. the result after 5 months post first surgery. Clinically there is concave shape on the labial side (arrow)



Figure 11. the result after 5 months post first surgery. Clinically there is concave shape on the labial side (arrow)

Procedure of surgery II, bone augmentation of region #32

1. After local anesthesia, a full thickness flap was performed which is incision made horizontally along alveolar ridge of region #32 [Figure 13] using

blade number 15C. The vertical incision was made 5mm apically from the margin gingiva of #31 [Figure 14]. After the flap was reflected there was still bone defect on region #32 which this

- made a concave shape on the labial side of the bone.
2. Bone grafts was applied after all soft tissues were thoroughly removed from the recipient site. Bone grafts was applied at the

concave shape on the labial side [figure 15,16]. Once properly positioned, grafts was covered by released mucosa and then sutured [Figure 17]



Figure 13. flap was performed with horizontal incision on along alveolar



Figure 14. vertical flap incision on 5 mm apically from the margin gingiva of



Figure 15. the process of applying bone graft into the concave shape on the labial side of bone defect



Figure 16. bone graft on properly position before sutured



Figure 17. the flap was sutured using non absorbable suture

Procedure of surgery II, bone3. Bone graft was applied after all soft augmentation on #34

1. After adequate local anesthesia, a full thickness flap was performed with incision made horizontally along #33, #34, #35 included 3 mm apically of interdental papillae [Figure 18].




Figure 18. flap design on augmentation II of #34. A full thickness flap was performed with incision made horizontally along #33, #34, #35 included 3 mm apically of interdental papillae

Figure 19. The process of applying bone graft on the defect



Figure 20. sutured using non absorbable suture

RESULTS

The results of 3 months after second surgery of region #32 and #34 were as followed:

1. On clinical evaluation, there was an increased of alveolar ridge

level on region #32 [Figure 21,22] and there was not any concave shape on the labial side on the region #32 [Figure 23,24]



Figure 21. The result 4 months after first surgery



Figure 22. The result 3 months after second surgery



Figure 23. The concave shape post first surgery



Figure 24. there is no concave shape after second surgery

2. On radiographic evaluation, there was an increased of bone density on region #32 and there was a significant alveolar bone and periodontal space recovery on #34 compare to the results of first surgery [Figure 25, 26, 27]



Figure 25. The radiographic examination before treatment



Figure 26. The radiographic evaluation 4 months post first surgery




Figure 27. The radiographic evaluation
3 months post second surgery

3. On CBCT evaluation, the width of alveolar bone (axial) is 5.98 mm, 5.90 (coronal). There is severe bone loss on the interdental of #31 and #33. Intidental space between #31 and #33 is 5.93 mm. The average of bone profile with the region of interest 17x35 is 906.4 HU




Figure 28. The width of alveolar bone is
5.98 mm (CBCT evaluation, axial)

Figure 29. The width of alveolar bone
is 5.90 mm (CBCT evaluation,
coronal).




Figure 30. There is severe bone loss on
the interdental of #31 and #33. Intidental
space between #31 and #33 is 5.93 mm.
The average of bone profile with the
region of interest 17x35 is 906.4 HU

Figure 31. The 3D profile of present
alveolar bone on region #32 and #34

DISCUSSION

The purpose of the present case report was to compare two different flap designs in preprosthetic surgery in the treatment of advanced alveolar bone defect. The treatment plan of this case after tooth loss is included dental implant. The clinician is obligated to perform augmentation procedures to reconstruct lost bone in order to achieve adequate ridge width to facilitate the placement of implant in a prosthetically driven positioned. Surgical reconstructive procedure in preprosthetic surgery have become more numerous and complex. Depending on the size and morphology of the defect, various augmentation procedure can be used including particulate grafting, membrane use, block grafting, and distraction osteogenesis, either alone or in combination. All the proven techniques must be followed to achieve good results including the flap management techniques. These include generating a blood supply, maintaining a stable, protected space of bone growth and achieving tension-free flap wound closure.

On the first surgery, the clinician was intended making a minimally invasive surgery in order to save more blood supply so there was not any vertical incision performed. This include one of general concept for flap management associated with alveolar ridge augmentation. The use of vertical incisions, although often required for surgical access, should be minimized. The absence of horizontal and vertical incision on the first surgery may either interfered the flap tension while at wound closure which led to unsatisfied results. On the second surgery, either horizontal and vertical incision were

performed. This flap design was intended to open more access for smoothing and scrapping the root surface and bone defects from granulation tissues, easier bone graft placement and also reduce the tension of flap. Wound closure depends on how to adapt wound edges with suturing techniques. Good sutured depends on the free flap tension which is increase flap elasticity. This permits complete closure without stress on the wound margin. V.

Soft tissue management is a critical aspect of bone augmentation procedures. Incisions, reflection and manipulation should be design to optimized blood supply and wound closure. The design and management of mucoperiosteal flap must consider to increased dimensions of the ridge after augmentations as well as esthetics and approximation of the wound margins. Various surgical techniques have been investigated with the aim of achieving simultaneous complete wound closure of preprosthetic surgery and perfectly normal appearance related to the tissue of the adjacent teeth. Selection of the appropriate surgical technique depends on different factors, some of which are related to the patients while others related to anatomical characteristics of the defect. The surgical procedure must be executed with the utmost of care in order to preserve the vascularity of the flap and to minimized tissue injury.¹³ In this case, the flap design performed on second surgery which is included vertical and horizontal incision is more superior than flap design on the first surgery. The results of the second surgery is more significant than on the first surgery. The formation of alveolar bone included bone density and recovery of

periodontal space is better on the second surgery. The surgical technique performed with horizontal and vertical flap design led to novel possibilities to regenerate alveolar bone.

CONCLUSION

The surgical technique performed with vertical and horizontal flap design led to novel possibilities to regenerate alveolar bone.

REFERENCES

1. McAllister BS, Haghghat K. Bone augmentation techniques. *J Periodontol* 2007;78:377-96.
2. Tolstunov L, Classification of the Alveolar Ridge Width: Implant-Driven Treatment Considerations for the Horizontally Deficient Alveolar Ridges. *Int J Oral Implantology Vol 40 Issue S1 July 2014*.
3. Dhadse PV. Soft tissue expansion before vertical ridge augmentation: Inflatable silicone balloons or self-filling osmotic tissue expanders? *Int J Indian Society of periodontology* 2014 Jul-Aug. 18(4). 433-40.
4. Pikos MA. Mandibular block autografts for alveolar ridge augmentation. *Atlas Oral Maxillofac Surg Clin North Am* 2005;13:91-107.
5. Peleg M, Sawatari Y, Marx RN, Santoro J, Cohen J, Bejarano P, et al. Use of corticocancellous allogenic bone blocks for augmentation of alveolar bone defects. *Int J Oral Maxillofac Implants* 2010;25:153-62.
6. Rothame D. Dimensional ridge alterations following socket preservation using a nanocrystalline hydroxyapatite paste. A histomorphometrical study in dogs. *Int J Oral Maxillofacial Surgery*. August 2008. Volume 37, Issue 8, Pages 741–747
7. Nyström E, Nilson H, Gunne J, Lundgren S. A 9-14 year follow-up of onlay bone grafting in the atrophic maxilla. *Int J Oral Maxillofac Surg* 2009;38:111-6.
8. Harsha MB. Use of Subepithelial connective tissue graft for roort coverage. *World Journal of Dentistry*. April-June 2011.2(2).159-62
9. Işiksal E, Hazar S, Akyalçin S. Smile esthetics: perception and comparison of treated and untreated smiles. *Am J Orthod Dentofacial Orthop* 2006; 129: 8–16.
10. Klokkevold PR. Localized bone augmentation and implant site development. In: Newman MG, Takei HH, Klokkevold PR, Carranza FA, editors. *Clinical Periodontology*, 10 ed. Philadelphia: Saunders; 2006. p. 1133-47.
11. Lang NP, Araujo M, Karring T. alveolar bone formation. In: Lindhe J, Karring T, Lang NP, editors. *Clinical Periodontology and Implant Dentistry*, 4th edn. Oxford: Blackwell Munksgaard; 2003. p. 866-92
12. Alan S. Herford, Todd C. Cooper, Carlo Maiorana, Marco Cicciù, , Vascularized Connective Tissue Flap for Bone Graft Coverage, *Journal of Oral Implantology*. 2011;37(2):279-285. - See more at: <http://www.joionline.org/doi/full/10.1563/AAID-JOI-D-09-00146.1#sthash.W8CLFAsV.dpuf>
13. Newman, Takei, Klokkevold, Carranza . *Clinical Periodontology*, 11th ed. 2012. P.673