

Leptin serum and disease activity in spondyloarthritis

by Yuliasih Yuliasih

Submission date: 07-Mar-2021 11:30PM (UTC-0800)

Submission ID: 1527220841

File name: Leptin_serum_and_disease_activity_in_spondyloarthritis.pdf (377.46K)

Word count: 4408

Character count: 23737



Leptin serum and disease activity in spondyloarthritis

Yuliasih Yuliasih ^{1*}, Luluch Mulyani ²

¹ Rheumatology Division, Department of Internal Medicine, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60131, INDONESIA

² Department of Internal Medicine, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60131, INDONESIA

*Corresponding author: yuli177@yahoo.com

Abstract

Background: Spondyloarthritis (SpA) is an active chronic inflammatory disease involving pro-inflammatory cytokines; one of which is leptin. This study aimed to determine correlation between leptin serum level and disease activity in SpA patients.

Method: A cross-sectional study involving patients diagnosed with SpA was conducted according to the 2009 ASAS criteria. Leptin serum level is measured by using ELISA whilst disease activity in SpA by ASDAS-CRP. The correlation was analyzed using Spearman correlation test.

Results: Fifty subjects with SpA aged 54.3±9.32 years old showed mean body mass index of 23.63±3.05 kg/m², CRP level of 0.76±0.91 mg/dL, ESR of 40.88±16.92 mm/hour, Schober test of 12.97±1.18 cm, and chest expansion test of 1.39±0.67 cm. Median leptin level and ASDAS-CRP score were 9.6 (0.92-51.57) ng/dL and 2.2 (1.3-3.86), respectively. No correlation was found between leptin level and disease activity (p = 0.174).

Conclusion: Leptin serum level cannot be used as a modality to assess disease activity in SpA.

Keywords: leptin, spondyloarthritis, ASDAS, disease activity, adipokines

Yuliasih Y, Mulyani L (2020) Leptin serum and disease activity in spondyloarthritis. Eurasia J Biosci 14: 1881-1886.

© 2020 Yuliasih and Mulyani

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

INTRODUCTION

Spondyloarthritis (SpA) is a chronic progressive low-grade inflammation disease which is often accompanied by metabolic syndrome consisting of obesity, dyslipidemia, and hypertension due to interference with adipokines regulation (Dougados et al. 2011, Genre et al. 2014). SpA may also cause the occurrence of secondary osteoarthritis (Ahmad et al. 2018). Furthermore, other studies presented that chronic arthritis is associated with other diseases including Lyme disease and risk factors for TB (Dharmajaya 2018, Kusmiati et al. 2016, Massi et al. 2017, Rotan et al. 2018).

Some researchers report that leptin is closely related to the activity of SpA disease (Elolemy et al. 2013, Kim et al. 2012, ParkMin-Chan et al. 2009). Leptin is an adipokines (proinflammatory cytokine) which is the main component underlying immunopathology that connects rheumatoid arthritis and obesity. This process is assumed to also have a role in the pathogenesis of SpA and obesity (Daïen et al. 2015). In addition, when immune system dysregulation occurs, especially in genetically susceptible individuals, bacterial dysbiosis can trigger autoimmune reactions, including spondyloarthritis (SpA) (Ciccio et al. 2015, 2016, Uotani et al. 2015).

Inflammation is a response to eliminate various pathogens and preserve host integrity (Putra et al. 2018, Umit et al., 2019). Some studies report an association between leptin level, inflammatory markers, and disease activity (ParkMin-Chan et al. 2009). An in vitro study on peripheral blood mononuclear cell (PBMC) culture shows that the leptin level in SpA patients are higher than controls and often formed in organs with active inflammation (Kim et al. 2012). In contrast, other studies found no difference or lower leptin level in SpA patients compared to control group (Derdemezis et al. 2010, Toussirot et al. 2013). Further studies show that the differences in the results of these studies may be due to differences in the population of patients studied, study designs (in vitro or in vivo), and duration of illness (Derdemezis et al. 2010, Kim et al. 2012, ParkMin-Chan et al. 2009).

In SpA, cytokines and acute phase proteins systemically increase by 2-3 times normal level (Dougados et al. 2011). Low-grade inflammation increases the production of leptin and triggers an inflammatory process that occurs continuously. Hyperleptinemia further triggers insulin resistance and

Received: March 2020

Accepted: May 2020

Printed: June 2020

32 obesity. Pro-inflammatory action increases when there is infection (Simanjuntak et al. 2018). Leptin acts as proinflammatory cytokines that activates endothelial cells and increases the accumulation of macrophages in adipose tissue which ultimately releases proinflammatory cytokines and aggravates inflammatory process (Castro et al. 2017).

18 Disease activity in SpA can be measured by the Ankylosing Spondylitis Disease Activity Score with CRP (ASDAS-CRP) score, in order to determine the presence of active inflammation in SpA. In 2009 The Assessment of Spondyloarthritis International Society (ASAS) introduced the ASDAS score system which can be used to measure disease activity categories more objectively (Markov Alexander 2019). In Indonesia, there has been no study linking between leptin and ASDAS-CRP scores. The results of this study are expected to shed a light about the controversy that arises between the roles of leptin in SpA. This study aimed to determine correlation between leptin serum level and disease activity in SpA patients (Shahsavari et al, 2013).

METHODS

Subjects

This cross-sectional study involved 50 subjects who were consecutively sampled. All subjects were outpatients in Dr. Soetomo General Hospital, Surabaya, diagnosed with SpA according to the 2009 Assessment of Spondyloarthritis International Society (ASAS) criteria. Patients with BMI >30kg/m², diabetes mellitus, liver cirrhosis, asthma, tuberculosis, and history of smoking were all excluded. Every patient who was enrolled in this study had voluntarily signed a consent form stating their willingness without force nor coercion to participate as study subjects. All study subjects were aware that they were involved in this cross-sectional study and their rights and confidentiality as patients were preserved with utmost respect. This study was conducted strictly according to the 2016 ICH-GCP and the Declaration of Helsinki and had been approved by Ethics Committee of Dr. Soetomo General Hospital, Surabaya.

Leptin Serum

Leptin serum level was measured by Enzyme Linked Immunosorbent Assay (ELISA), using Human Leptin (Quantikine® ELISA Human Leptin Immunoassay, R&D System, IncCatalog D1700, USA) as reagent. The minimum detectable level was 7.8 pg/ml. Blood sample was drawn in a serum separator tube for 30 minutes, then centrifuged for 15 minutes and stored in < -20 C fridge. 100 µL of each standard, control, and sample were added into tubes containing 100 µL of Assay Diluent RD1-19. Tubes were sealed with adhesives and incubated in room temperature for 2 hours before aspirated and washed 4 times. Aspiration and washing were repeated after adding 200 µL of Leptin Conjugate

Table 1. Subjects' characteristics

Characteristics	Results
Sex	
Male	13 (26%)
Female	37 (74%)
Age (year old)	54.3±9.32 ^a
Body Mass Index (kg/m ²)	23.6±3.05 ^a
CRP (mg/dL)	0.76±0.91 ^a
ESR (mm/hour)	40.88±16.92 ^a
Schober Test (cm)	12.97±1.18 ^a
Chest Expansion Test (cm)	1.39±0.67 ^a
ASDAS-LED (score)	3.38±0.81 ^a
ASDAS-CRP (score)	2.2(1.3-3.86) ^b

^a Mean±SD; ^b Median(min-max)

Table 2. Leptin Serum According to SpA type and BMI grade

Variable	Category	Leptin serum (ng/dL)
Type of SpA	Axial SpA	10.10 (0.92-51.57) ^b
	Peripheral SpA	16.72±1.38 ^a
BMI	Normal	6.03(0.92-48.21) ^b
	Overweight	10.2 (1.33-51.57) ^b
	Obese I	19.84±3.39 ^a

^a Mean±SD; ^b Median (min-max)

2 and one-hour incubation period. Furthermore, 200 µL of substrate solution was added, and tubes were stored in light-proof storage for 30-minute incubation period. After that, 50 µL of stop solution was added, and finally sample was run through microplate reader with 450 nm wavelength for 30 minutes.

Disease Activity

39 Severity of SpA is evaluated by Ankylosing Spondylitis Disease Activity Score (ASDAS). This scoring system evaluates back pain, duration of morning stiffness, peripheral joint pain or swelling, and global assessment of disease activity. The score is often combined with C-reactive protein (CRP) level or erythrocyte sedimentation rate (ESR). ASDAS-CRP is more recommended to evaluate SpA than its alternative, ASDAS-LED (Zochling 2011).

Statistical Analysis

Data collected from this study was analyzed in SPSS 17 (IBM Corp., Armonk, NY, USA). Subjects' demographics and characteristics were shown as descriptive data. Correlation of leptin serum level and ASDAS were analyzed by Spearman's correlation test, with significance level of 0.05.

RESULTS

Table 1 shows that the subjects are all middle-aged (54.3±9.32 years old) with female as majority (74%). Median ASDAS-CRP score was 2.2. In Table 2, out of 50 subjects, median serum leptin level was 9.6(0.92-51.57) ng/dL, with higher leptin level found in the peripheral SpA group of 16.72±1.38 ng/dL. Based on BMI grade, subjects in Obese I category had the highest mean leptin level (19.84±3.39 ng/dL). According to ASDAS-CRP as seen in Table 3, this study has no case categorized as inactive disease whilst the majority of the

Table 3. Leptin level and other various parameters to disease activity in SpA

Parameters	Moderate Disease n=17 (34%)	High Disease n=27 (54%)	Very High disease n=6 (12%)
ESR	33.5±13.5 ^a	43.22±16.31 ^a	51.67±21.97 ^a
Schober Test	12.76±0.79 ^a	13.12±1.19 ^a	12.5±1.48 ^a
Chest Expansion	1.32±0.53 ^a	1.37±0.76 ^a	1.66±0.6 ^a
CRP	0.3(0.1-2.3) ^b	0.5 (0.1-5.7) ^b	0.95 (0.5-1.3) ^b
Leptin	7.57 (1.33-36.7) ^b	9.04 (0.92-45.73) ^b	33.05(7.17-51.57) ^b

^a Mean±SD; ^b Median(min-max)




Fig. 1. Scatterplot for Leptin level and Disease activity

subjects fell into high disease category (54%). Leptin level appears to be the highest in subjects with very high disease.

Based on Kolmogorov-Smirnov analysis of disease activity to leptin level, the study appears to be having abnormal data distribution with p-value of 0.000. Furthermore, this study obtained insignificant p-value of 0.174 through Spearman test, hence no correlation between leptin level and disease activity in SpA (ASDAS-CRP score) as seen in **Fig. 1**. Statistically, the scatterplot showed that leptin level among all three categories of disease activity overlaps. Correlation test was performed for each category of disease activity to leptin level. The results showed no correlation between serum leptin level and ASDAS-CRP scores in patients with either moderate, high, or very high disease activity (p-value of 0.217, 0.278, and 0.178). Correlation coefficient (r-value) was not analyzed as there was no significant value found in the first place (Shahsavari et al, 2013).

DISCUSSION

There was no association between leptin serum level and disease activity using ASDAS-CRP score. Leptin may play a role at local sites of inflammation rather than systemically. This adipokine does not show any effect on the disease activity in SpA (Mei et al. 2016).

The role of leptin in the pathogenesis of SpA is still unclear. Past studies have shown contradictory results

about this matter. Leptin is produced by adipocytes. Adipose tissue synthesizes hormones, cytokines and immune factors, such as TNF-α, IL-1 and IL-6. Chronic inflammation triggers white adipose tissue (WAT) to secrete adipokines, which are inflammatory cytokines consisting of leptin, adiponectin, resistin, visfatin, etc. (Moon et al. 2013). Chronic inflammation can reduce the amount of body fat and reducing the production of leptin by adipocytes which are the main source of leptin (Sari et al. 2007). The reduction of adipose tissue mass leads to low production of leptin (Tousssirot et al. 2013). A study compared serum leptin level in SpA patients to a control group (Sari et al. 2007). The study reported that leptin level in SpA patients was significantly lower than that of controls. Furthermore, previous studies reported no difference of leptin levels between SpA patients and control group and no correlation between serum leptin levels and disease activity (Derdemezis et al. 2010, Tousssirot et al. 2013). A systematic review and a meta-analysis mentioned that serum leptin level was not associated with other inflammatory markers, such as ESR, CRP, TNF-α and also BASDAI (Mei et al. 2016).

In contrast, several other studies reported significant correlations between leptin level and disease activity SpA. Research in 2013 reported that serum leptin level increased significantly in SpA patients and was correlated to disease activity parameters (BASDAI, BASFI) (Elolemy et al. 2013). SpA patients with syndesmophytes have higher leptin level than those without (Kim et al. 2012). A prospective research also

studied leptin level and disease activity in SpA patients (ParkM-C et al. 2007). After 31 months, it was found that there was a decrease of the disease activity parameters (BASDAI, ESR, and CRP) and serum leptin level. These contradictory findings are possibly due to differences in the demographics of the study population, duration of disease, and treatment provided.

Leptin level in this study was showing similarity with a study in Egypt that found elevated leptin serum in axial SpA (7.6 ±3.3 ng/dL), peripheral SpA (10.9±1.9 ng/dL), and overall cases (9.1±3.9) ng/dL (Elolemy et al. 2013). Patients with peripheral SpA showed higher leptin level as the disease itself exhibits profound arthritis or Rheumatoid Arthritis (RA)-like manifestations. Another study reported that leptin serum level in RA patients was 19.8±2.7 ng/dL (Toussirot et al. 2013).

The American College of Rheumatology reported that SpA occurred three times higher in men than women whilst other researchers state there was no difference. This study had a majority of female patients with the female to male ratio of 3:1 possibly due to sampling method. Among 80% of SpA patients, the initial symptoms appeared under age of 30 and less than 5% starts older than 45 years old. In this study, the average age of the samples was 53.58±9.28 years old (Braun et al. 2008). The mean age reported in this study was higher than previous studies, possibly due to SpA patients coming to a health facility on later age when the disease had already advanced. From the study report, the average SpA patient was diagnosed after 8-10 years after the initial symptoms due to "insidious" symptoms. A study in Brazil showed that axial SpA was more prevalent than peripheral SpA (Sampaio-Barros et al. 2010). Another report mentioned the prevalence of axial SpA was 60% higher than peripheral SpA cases in America (Strand et al. 2013). In this study, there were 35 axial patients (70%) and 15 peripheral patients (30%). The numbers are in accordance with other studies.

In early stage of SpA, erosion occurs at proximal entheses and it gradually will heal on later stage and the formation of a spur at the distal attachment site will cause vertebral fusion (McGonagle et al. 2009). Spinal mobility can be assessed, among others, by chest expansion and the Schober test. Abnormal spinal mobility can indicate structural damage as well as the severity of SpA. In this study, the Schober test results showed similarity to previous studies which showed 13.3±1.2 cm and 12.9±1.6 cm, respectively (Rezvani et al. 2012, Tian et al. 2014). It is suspected that the progression of the disease between the races is slower, considering that subjects in this study are older than 40 years old. Furthermore, earlier studies reported mean chest expansion of 3.07±1.66 cm and 3.6±1.3 cm

(Grubisić et al. 2014, Rezvani et al. 2012). However, this study reported only 1.39±0.67 cm for chest expansion test, which is similar to previous studies.

Study in the Netherlands reported an average ASDAS-CRP value of 2.4±0.9 in active SpA patients (Rubio Vargas et al. 2016). People's Republic of China reported higher ASDAS-CRP score in patients with axial SpA (2.05±1.07) than those with peripheral SpA (1.86±1.07) because ASDAS-CRP evaluates axial disease activity better than peripheral one (Xue et al. 2012). This study gave a median value of ASDAS-CRP score of 2.2 (1.3-3.86). The majority SpA patients in this study were in high activity disease category, and none was categorized as having inactive disease. Most of SpA patients did not seek treatment until much later they exhibited major symptoms. Considering the socio-economy conditions, these patients were prescribed with conventional Disease Modifying Anti-Rheumatic Drugs (DMARDs) such as Methotrexate (MTX) and/or Sulfasalazine (SSZ) instead of using targeting therapy. Patient's compliance in medication might also contribute in the lack of inactive cases in this study.

ASDAS-CRP heavily relies on clinical symptoms which are subject to patients' perspective and understanding about their disease. To confirm a worsening in disease activity, physicians need to evaluate spinal mobility (Reveille et al. 2012). Any limitations in spinal mobility can be evaluated by performing Schober test, occiput-to-wall test and chest expansion. A study in Morocco showed an association between spinal mobility as measured by the Schober test, occiput-to-wall test and chest expansion with disease activity using BASDAI, BASMI, BASFI and BASRI (Ibn Yacoub et al. 2012). Assessing SpA activity is not simple and it requires a lot of parameters because SpA is a low-grade inflammation with activity and progression that are clinically difficult to assess.

CONCLUSION

There was no association between leptin serum level and disease activity using ASDAS-CRP score. Leptin serum level cannot be used as a modality to assess disease activity in SpA.

ACKNOWLEDGEMENT

Authors would like to thank Poernomo Boedi Setiawan, MD, as Head of Internal Medicine Department, Universitas Airlangga; Endang Retnowati, MD from the Department of Clinical Pathology, Universitas Airlangga; Cita Rosita, MD, PhD, as Head of Research and Development Division of Dr. Soetomo General Hospital; and Harsono, MD, as Director of Dr. Soetomo General Hospital.

REFERENCES

- Ahmad IW, Rahmawati LD, Wardhana TH (2018) Demographic Profile, Clinical and Analysis of Osteoarthritis Patients in Surabaya. *Biomolecular and Health Science Journal* 1(1): 34-39.
- Braun J, Baraliakos X, Listing J, Fritz C, Alten R, Burmester G, Krause A, Schewe S, Schneider M, Sørensen H, Zeidler H, Sieper J (2008) Persistent clinical efficacy and safety of anti-tumour necrosis factor alpha therapy with infliximab in patients with ankylosing spondylitis over 5 years: evidence for different types of response. *Annals of the rheumatic diseases* 67(3): 340-345. <https://doi.org/10.1136/ard.2007.075879>
- Castro AM, Macedo-de la Concha LE, Pantoja-Meléndez CA (2017) Low-grade inflammation and its relation to obesity and chronic degenerative diseases. *Revista Médica del Hospital General de México* 80(2): 101-105. <https://doi.org/10.1016/j.hgmx.2016.06.011>
- Ciccía F, Guggino G, Rizzo A, Saieva L, Peralta S, Giardina A, Cannizzaro A, Sireci G, De Leo G, Alessandro R, Triolo G (2015) Type 3 innate lymphoid cells producing IL-17 and IL-22 are expanded in the gut, in the peripheral blood, synovial fluid and bone marrow of patients with ankylosing spondylitis. *Annals of the Rheumatic Diseases* 74(9): 1739-1747. <https://doi.org/10.1136/annrheumdis-2014-206323>
- Ciccía F, Rizzo A, Triolo G (2016) Subclinical gut inflammation in ankylosing spondylitis. *Current Opinion in Rheumatology* 28(1): 89-96. <https://doi.org/10.1097/BOR.0000000000000239>
- Daïen CI, Sellam J (2015) Obesity and inflammatory arthritis: impact on occurrence, disease characteristics and therapeutic response. *RMD open* 1(1): e000012.
- Derdemezis CS, Filippatos TD, Voulgari P V, Tselepis AD, Drosos AA, Kiortsis DN (2010) Leptin and adiponectin levels in patients with ankylosing spondylitis. The effect of infliximab treatment. *Clin Exp Rheumatol* 28(6): 880-883.
- Dharmajaya R (2018) Tuberculous spondylitis in Haji Adam Malik hospital, Medan. In: IOP Conference Series Earth and Environmental Science Vol. 125. <https://doi.org/10.1088/1755-1315/125/1/012095>
- Dougados M, Baeten D (2011) Spondyloarthritis. *The Lancet* 377(9783): 2127-2137.
- Eloemy GG, Ganeb SS, Ghanima ATA, Abdelgwad ER (2013) Influence of adipocytokines and IL-6 on ankylosing spondylitis disease activity and functional status. *The Egyptian Rheumatologist* 35(2): 65-70.
- Genre F, López-Mejías R, Miranda-Filloo JA, Ubilla B, Carnero-López B, Blanco R, Pina T, González-Juanatey C, Llorca J, González-Gay MA (2014) Adipokines, biomarkers of endothelial activation, and metabolic syndrome in patients with ankylosing spondylitis. *BioMed research international* 2014.
- Grubisić F, Grazio S, Balenović A, Nemčić T, Kusić Z (2014) Osteoporosis, spinal mobility and chest expansion index in patients with ankylosing spondylitis. *Collegium antropologicum* 38(1): 63-68.
- Ibn Yacoub Y, Amine B, Laatiris A, Hajjaj-Hassouni N (2012) Gender and disease features in Moroccan patients with ankylosing spondylitis. *Clinical rheumatology* 31(2): 293-297. <https://doi.org/10.1007/s10067-011-1819-x>
- Kim K-J, Kim J-Y, Park S-J, Yoon H, Yoon C-H, Kim W-U, Cho C-S (2012) Serum leptin levels are associated with the presence of syndesmophytes in male patients with ankylosing spondylitis. *Clinical rheumatology* 31(8): 1231-1238.
- Kusmiati T, Narendrani HP (2016) POTT'S Disease. *Jurnal Respirasi* 2(3): 99-106.
- Massi MN, Biatko KT, Handayani I, Pratama MY, Septriani S, Nurdin GM, Ali MB (2017) Evaluation of rapid GeneXpert MTB/RIF method using DNA tissue specimens of vertebral bones in patients with suspected spondylitis TB. *Journal of Orthopaedics* 14(1): 189-191. <https://doi.org/10.1016/j.jor.2016.12.003>
- McGonagle D, Benjamin M (2009) Entheses, Enthesitis and Enthesopathy. *Reports on the Rheumatic Diseases Series 6. Arthritis Research UK* 4.
- Mei Y, Wang P, Chen L, Li Z (2016) Plasma/Serum Leptin Levels in Patients with Ankylosing Spondylitis: A Systematic Review and Meta-analysis. *Archives of Medical Research* 47(2): 111-117. <https://doi.org/10.1016/j.arcmed.2016.03.001>
- Moon H-S, Dalamaga M, Kim S-Y, Polyzos SA, Hamnvik O-P, Magkos F, Paruthi J, Mantzoros CS (2013) Leptin's role in lipodystrophic and nonlipodystrophic insulin-resistant and diabetic individuals. *Endocrine reviews* 34(3): 377-412. <https://doi.org/10.1210/er.2012-1053>
- Park M-C, Lee S-W, Choi S-T, Park Y-B, Lee S-K (2007) Serum leptin levels correlate with interleukin-6 levels and disease activity in patients with ankylosing spondylitis. *Scandinavian journal of rheumatology* 36(2): 101-106. <https://doi.org/10.1080/03009740600991760>

- Park Min-Chan, Chung S-J, Park Y-B, Lee S-K (2009) Pro-inflammatory effect of leptin on peripheral blood mononuclear cells of patients with ankylosing spondylitis. *Joint bone spine* 76(2): 170–175. <https://doi.org/10.1016/j.jbspin.2008.04.018>
- Putra A, Ridwan FB, Putridewi AI, Kustiyah AR, Wirastuti K, Sadyah NAC, Rosdiana I, Munir D (2018) The role of $\text{tnf-}\alpha$ induced mscs on suppressive inflammation by increasing $\text{tgf-}\beta$ and il-10 . *Open Access Macedonian Journal of Medical Sciences* 6(10): 1779-1783. <https://doi.org/10.3889/oamjms.2018.404>
- Reveille JD, Witter JP, Weisman MH (2012) Prevalence of axial spondylarthritis in the United States: estimates from a cross-sectional survey. *Arthritis care & research* 64(6): 905-910. <https://doi.org/10.1002/acr.21621>
- Rezvani A, Ergin O, Karacan I, Oncu M (2012) Validity and reliability of the metric measurements in the assessment of lumbar spine motion in patients with ankylosing spondylitis. *Spine* 37(19): E1189-96. <https://doi.org/10.1097/BRS.0b013e31825ef954>
- Rotan H, Ginting Y, Loesnihari R, Kembaren T, Marpaung B (2018) Correlation between chronic arthritis patients confirmed with questionnaire and serologic test of Lyme disease. In: *IOP Conference Series Earth and Environmental Science* Vol. 125. Institute of Physics Publishing <https://doi.org/10.1088/1755-1315/125/1/012043>
- Rubio Vargas R, van den Berg R, van Lunteren M, Ez-Zaitouni Z, Bakker PAC, Dagfinrud H, Ramonda R, Landewé R, Molenaar E, van Gaalen FA, van der Heijde D (2016) Does body mass index (BMI) influence the Ankylosing Spondylitis Disease Activity Score in axial spondyloarthritis?: Data from the SPACE cohort. *RMD open* 2(1): e000283. <https://doi.org/10.1136/mdopen-2016-000283>
- Sampaio-Barros PD, Bortoluzzo AB, Conde RA, Costallat LTL, Samara AM, Bértolo MB (2010) Undifferentiated spondyloarthritis: a longterm followup. *The Journal of rheumatology* 37(6): 1195-1199. <https://doi.org/10.3899/jrheum.090625>
- Sari I, Demir T, Kozaci LD, Akar S, Kavak T, Birlik M, Onen F, Akkoc N (2007) Body composition, insulin, and leptin levels in patients with ankylosing spondylitis. *Clinical rheumatology* 26(9): 1427-1432. <https://doi.org/10.1007/s10067-006-0509-6>
- Shahsavari F, Ghanadi K, Jafarzadeh M, Nasiri B, Forutani SH. (2013). Opium addiction decreases T3 uptake amount. *Life Science Journal*. 10(5s): 608-610.
- Shahsavari F, Jafarzadeh M, Azaragooon A, Hedayati M, Asadifar B. (2013). Small ubiquitin-like modifier 4 M55V polymorphism is not associated with diabetic nephropathy in Iranian type 2 diabetes patients. *Indian Journal of Human Genetics*. 19(2),179-183.
- Simanjuntak TP, Hatta M, Rauf S, Yusuf I, Tahir M (2018) Forkhead box P3 messenger-RNA expression after Curcuma longa extract intervention in early pregnant mice with toxoplasmosis. *Research Journal of Immunology* 11(1): 1-6. <https://doi.org/10.3923/rji.2018.1.6>
- Strand V, Rao SA, Shillington AC, Cifaldi MA, McGuire M, Ruderman EM (2013) Prevalence of axial spondyloarthritis in United States rheumatology practices: Assessment of SpondyloArthritis International Society criteria versus rheumatology expert clinical diagnosis. *Arthritis care & research* 65(8): 1299-1306. <https://doi.org/10.1002/acr.21994>
- Tian G, Liang J-N, Wang Z-Y, Zhou D (2014) Emerging role of leptin in rheumatoid arthritis. *Clinical and experimental immunology* 177(3): 557-570. <https://doi.org/10.1111/cei.12372>
- Toussiroit É, Grandclément É, Gaugler B, Michel F, Saas P, Dumoulin G (2013) Serum adipokines and adipose tissue distribution in rheumatoid arthritis and ankylosing spondylitis. A comparative study. *Frontiers in immunology* 4: 453.
- Umit EG, Baysal M, Bas V, Goze H, Asoglu V, Kirkizlar O, Demir AM (2019) Value of Extracellular High Mobility Group Box 1 (HMGB1) in the Clinical Context of Immune Thrombocytopenia. *Journal of Clinical and Experimental Investigations*, 10(2): em00724. <https://doi.org/10.5799/jcei/5833>
- Uotani T, Miftahussurur M, Yamaoka Y (2015) Effect of bacterial and host factors on Helicobacter pylori eradication therapy. *Expert opinion on therapeutic targets* 19(12): 1637-1650.
- Xue Y, Jiang L, Cheng Q, Chen H, Yu Y, Lin Y, Yang X, Kong N, Zhu X, Xu X, Wan W, Zou H (2012) Adipokines in psoriatic arthritis patients: the correlations with osteoclast precursors and bone erosions. *PLoS one* 7(10): e46740. <https://doi.org/10.1371/journal.pone.0046740>
- Zochling J (2011) Measures of symptoms and disease status in ankylosing spondylitis: Ankylosing Spondylitis Disease Activity Score (ASDAS), Ankylosing Spondylitis Quality of Life Scale (ASQoL), Bath Ankylosing Spondylitis Disease Activity Index (BASDAI), Bath Ankylosing S. *Arthritis care & research* 63 Suppl 1: S47-58. <https://doi.org/10.1002/acr.20575>

Leptin serum and disease activity in spondyloarthritis

ORIGINALITY REPORT

19%

SIMILARITY INDEX

14%

INTERNET SOURCES

15%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1	repository.unair.ac.id Internet Source	3%
2	hdl.handle.net Internet Source	1%
3	Hamed Mohammadi, Nadia Sharafkandi, Maryam Hemmatzadeh, Gholamreza Azizi et al. "The role of innate lymphoid cells in health and disease", Journal of Cellular Physiology, 2018 Publication	1%
4	worldwidescience.org Internet Source	1%
5	www.hindawi.com Internet Source	1%
6	"MYOPATHIES AND SOFT TISSUE RHEUMATISM INCLUDING FIBROMYALGIA", APLAR Journal of Rheumatology, 8/2006 Publication	1%
7	china.iopscience.iop.org Internet Source	1%

8	link.springer.com Internet Source	1%
9	Bora Nam, Bon San Koo, Tae-Han Lee, Ji-Hui Shin, Jin-Ju Kim, Seunghun Lee, Kyung Bin Joo, Tae-Hwan Kim. "Low BASDAI score alone is not a good predictor of anti-tumor necrosis factor treatment efficacy in ankylosing spondylitis: a retrospective cohort study", BMC Musculoskeletal Disorders, 2021 Publication	1%
10	sysrevpharm.org Internet Source	1%
11	www.e-sciencecentral.org Internet Source	1%
12	A.M. Castro, L.E. Macedo-de la Concha, C.A. Pantoja-Meléndez. "Low-grade inflammation and its relation to obesity and chronic degenerative diseases", Revista Médica del Hospital General de México, 2017 Publication	<1%
13	Juergen Dunst, Axel Becker, Christine Lautenschläger, Silke Markau, Heike Becker, Kersten Fischer, Gabriele Haensgen. "Anemia and Elevated Systemic Levels of Vascular Endothelial Growth Factor (VEGF)", Strahlentherapie und Onkologie, 2002 Publication	<1%

14 Yong-jun Mei, Peng Wang, Lin-jie Chen, Zhi-jun Li. "Plasma/Serum Leptin Levels in Patients with Ankylosing Spondylitis: A Systematic Review and Meta-analysis", Archives of Medical Research, 2016
Publication <1%

15 repository.unej.ac.id
Internet Source <1%

16 www.ema.europa.eu
Internet Source <1%

17 brage.bibsys.no
Internet Source <1%

18 Matteo Colina, Letizia Penolazzi, Martina di Ciano, Elisabetta Lambertini et al. "Osteoclasts from peripheral blood mononuclear cells culture of ankylosing spondylitis subjects are resistant to apoptosis", Biomedicine & Preventive Nutrition, 2013
Publication <1%

19 Nadia Hamjane, Fatiha Benyahya, Naima Ghailani Nourouti, Mohcine Bennani Mechita, Amina Barakat. "Cardiovascular diseases and metabolic abnormalities associated with obesity: What is the role of inflammatory responses? A systematic review", Microvascular Research, 2020 <1%

20

Sousan Kolahi, Alireza Khabbazi, Nastaran Kazemi, Aida Malek Mahdavi. "Does vitamin D deficiency contribute to higher disease activity in patients with spondyloarthritis?", Immunology Letters, 2019

Publication

<1%

21

www.spandidos-publications.com

Internet Source

<1%

22

Gehan Gamal Elolemy, Sahar Saad Ganab, Ahmed Taha Abou Ghanima, Eman Ramadan Abdelgwad. "Influence of adipocytokines and IL-6 on ankylosing spondylitis disease activity and functional status", The Egyptian Rheumatologist, 2013

Publication

<1%

23

José Agustín Tapia-Hernández, Francisco Rodríguez-Felix, Josué Elías Juárez-Onofre, Saúl Ruiz-Cruz et al. "Zein-polysaccharide nanoparticles as matrices for antioxidant compounds: A strategy for prevention of chronic degenerative diseases", Food Research International, 2018

Publication

<1%

24

Martin H. Lundqvist, Kristina Almby, Niclas Abrahamsson, Jan W. Eriksson. "Is the Brain a Key Player in Glucose Regulation and

<1%

Development of Type 2 Diabetes?", *Frontiers in Physiology*, 2019

Publication

25

Metabolic Syndrome, 2016.

Publication

<1%

26

chimie-biologie.ubm.ro

Internet Source

<1%

27

etheses.whiterose.ac.uk

Internet Source

<1%

28

Akash V Ajmera. "Suboptimal Response to Ferrous Sulfate in Iron-Deficient Patients Taking Omeprazole :", *American Journal of Therapeutics*, 12/2010

Publication

<1%

29

Faten Abdulhady Zakareia. "Correlation of peripheral arterial blood flow with plasma chemerin and VEGF in diabetic peripheral vascular disease", *Biomarkers in Medicine*, 02/2012

Publication

<1%

30

Jing Wang, Jinyu Su, Yuan Yuan, Xiaxia Jin, Bo Shen, Guoguang Lu. "The role of lymphocyte-monocyte ratio on axial spondyloarthritis diagnosis and sacroiliitis staging ", *Research Square*, 2020

Publication

<1%

31 Ming-Yue Zhang, Ahmed Ali Dini, Xiao-Ke Yang, Lian-Ju Li, Guo-Cui Wu, Rui-Xue Leng, Hai-Feng Pan, Dong-Qing Ye. "Association between serum/plasma adiponectin levels and immune-mediated diseases: a meta-analysis", Archives of Dermatological Research, 2017

Publication

32 Tigor Peniel Simanjunta, Mochammad Hatta, Syahrul Rauf, Irawan Yusuf, Mardiah Tahir. "Forkhead Box P3 Messenger-RNA Expression after Curcuma longa Extract Intervention in Early Pregnant Mice with Toxoplasmosis", Research Journal of Immunology, 2018

Publication

33 aplar2019.com

Internet Source

34 journals.sagepub.com

Internet Source

35 Tian, Guo, Jia-Ning Liang, Zhuo-Yun Wang, and Dian Zhou. "Emerging role of leptin in rheumatoid arthritis", Clinical & Experimental Immunology, 2014.

Publication

36 D'Aprile, Paola. "Clinical Assessment of Spondyloarthritis", MRI of Rheumatic Spine, 2014.

Publication

37

Hendra Gunawan, Ihdinal Mukti, Sony Wibisono Mudjanarko, Awalia, Lita Diah Rahmawati, Agung Pranoto. "Correlation of radiographic damage and central obesity in ankylosing spondylitis: A cross sectional study", Obesity Medicine, 2019

Publication

<1%

38

Iona MacDonald, Shan-Chi Liu, Chien-Chung Huang, Shu-Jui Kuo, Chun-Hao Tsai, Chih-Hsin Tang. "Associations between Adipokines in Arthritic Disease and Implications for Obesity", International Journal of Molecular Sciences, 2019

Publication

<1%

39

pure.uva.nl
Internet Source

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

Leptin serum and disease activity in spondyloarthritis

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
