

Serum Interleukin-2 Level Associated with TregTh17 Ratio in Active Systemic Lupus Erythematosus

by Riarita Subamayanti

Submission date: 08-Mar-2021 03:34PM (UTC+0800)

Submission ID: 1527223554

File name: d_with_TregTh17_Ratio_in_Active_Systemic_Lupus_Erythematosus.pdf (511.38K)

Word count: 3710

Character count: 19467

Serum Interleukin-2 Level Associated with Treg/Th17 Ratio in Active Systemic Lupus Erythematosus

RIARITA SUBAMAYANTI¹, YULIASIH^{1*}, LITA DIAH RAHMAWATI²

¹Department of Internal Medicine, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60131, Indonesia

²Rheumatology Division, Department of Internal Medicine, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60131, Indonesia

*Corresponding Author

Email ID: yuli177@yahoo.com

Received: 28.03.20, Revised: 30.04.20, Accepted: 22.05.20

ABSTRACT

Introduction: Systemic lupus erythematosus (SLE) is resulted from loss of immunological tolerance. Interleukin-2 (IL-2) plays an important role in loss of tolerance in SLE through its influence on Treg and Th17 balance. This study was conducted to determine the correlation between serum IL-2 level with Treg/Th17 ratio in active SLE.

Methods: This cross-sectional analytic study included thirty newly diagnosed SLE patients. IL-2 level was measured using ELISA, whereas Treg and Th17 expressions were measured using the flow cytometry method, then the ratio was calculated. The Systemic Lupus Activity Measure (SLAM) scores were used to assessed SLE disease activity.

Results: All patients were female with mean age of 31±10 years. Hematological conditions and arthritis were the most prominent clinical manifestations. All patients were in active SLE conditions with a high mean SLAM score of 29±4. Serum IL-2 level was lower in SLE patients compared to healthy subjects (median 7.12 pg/mL vs 49.31 pg/mL, respectively). Treg/Th17 ratio was also lower in SLE patients compared to healthy subjects (median 0.405 vs 0.830, respectively). There was a significant correlation between serum IL-2 level with Treg/Th17 ratio ($r = 0.463$; $P = 0.01$).

Conclusion: Serum IL-2 level is associated with Treg/Th17 ratio in active SLE.

Keywords: SLE, IL-2, Treg/Th17 ratio, disease activity, SLAM.

INTRODUCTION

Systemic Lupus Erythematosus (SLE) is a chronic autoimmune systemic disease that involves diverse pathogenesis and various illnesses, from mild to life-threatening [1,2]. Lupus symptoms that appear at any time are very potential to disrupt daily activities and cause many other problems. To achieve optimal health status and high quality of life, SLE patients must be proactive in managing the disease [3]. All changes in conditions must be experienced by patients with SLE both in environmental aspects, such as social support, physical aspects and emotional aspects affecting changes in their life quality [4].

It is known that the pathogenesis of SLE is based on interactions of genetic predisposition, hormonal factors, and environmental triggers. The previous study presented the marker of SLE disease activity might be Midkine serum level and have a role in the pathogenesis of SLE [5]. However, to date, the specific cellular and humoral mechanisms underlying immune dysfunction in SLE have not been fully understood. The main factor in the occurrence of autoimmune

in SLE is the loss of tolerance [6]. SLE pathogenesis is characterized by the absence of self-tolerance, with autoreactive T-cells and the activation of autoreactive B-cell, resulting in pathogenic autoantibodies and tissue damage. Self-antigen tolerance is retained by T cell subpopulations, referred to as Regulatory T (Tregs), which help to control immunological balance, and tolerance loss is an underlying autoimmune pathology [7,8]. Decreased production of IL-2 in SLE is a factor in immune dysregulation that underlies the loss of tolerance in SLE. IL-2 is important in maintaining self-tolerance and inhibit autoimmunity. Low IL-2 contributes to a decrease in the number of Treg cells, which are essential to the regulation of autoreactive T cells expansion [9]. The decrease of IL-2 levels in SLE have been reported in various studies and are thought to play a major role in loss of tolerance[9,10] through their influence on Treg and Th17 balance [11,12].

The reduction in IL-2 development also results in a disruption in cell death induced by activation

induced cell death (AICD) that is necessary to downregulate T cell clone growth [13]. In order to produce the CD8 + T-Cell Effector functions and CD8 + T cell memory development, IL-2 is also necessary. The failure of the CD8+T cytotoxic feature increases the risk for SLE patients of intracellular infection [9]. Increasing the number of Treg cells by IL-2 can be a specific therapeutic strategy to restore self-tolerance to SLE, but functional improvement and abnormal immune responses must be carefully analyzed in SLE patients given the complexity of the pathogenesis and etiology of SLE disease.

Data from various studies on IL-2 in SLE have shown opposing results about the role of IL-2 in SLE. Multiple studies reported a relationship between IL-2 levels and SLE disease activity [14–16]. Whereas, another study reported that IL-2 levels were not associated with SLE disease activity [17]. Various studies on the therapy of low-dose IL-2 in SLE have also been carried out. Low-dose IL-2 therapy increases the population of Treg cells and decreases the number of TFH and Th17 cells, followed the decreased of disease activity in SLE patients [18,19].

Considering the importance of IL-2 in SLE and the limited research data on the role of IL-2 in the mechanism and activity of SLE disease in Indonesia, this study was conducted to determine the correlation between serum IL-2 levels and Treg/Th17 ratio in active SLE patients. Serum IL-2 level was measured using the ELISA method, while Treg and Th17 counts were measured using flow cytometry technique. The Systemic Lupus Activity Measure (SLAM) score was used to assessed SLE disease activity because it is the easiest and most efficient, does not require a high cost, reliable, valid, responsive to changes in disease activity over time, with sensitivity and specificity equivalent to SLEDAI. To avoid bias in the results of the study, the study subjects were naive SLE patients who had never received prior intervention. This study aimed to regarding the relationship between Treg/Th17 and various cytokines with disease activity in SLE patients at Dr. Soetomo Hospital, Surabaya.

METHODS

This was an observational cross-sectional analysis study. The study was conducted at dr. Soetomo General Hospital, Surabaya. The population studied were all patients with SLE who were treated as an inpatient at dr. Soetomo General Hospital, Surabaya. A consecutive sampling method was used to select the participant for this study.

Criteria for inclusion in this study were that the subjects agreed to participate in the study, patient with SLE based on ACR criteria, active SLE based on SLAM score, and aged 16-60 years. Patients

that were or have been receiving steroid and or immunosuppressant therapies, patients with infection, tuberculosis, morbus hansen, HIV/AIDS, malignancy, asthma, acute coronary syndrome, mixed connective tissue disease and smoking were excluded.

Serum IL-2 level was measured using ELISA, while Treg and Th17 counts were measured using flow cytometry. SLAM scores was used the assessed SLE disease activity.

Data were analyzed with analytic statistics and presented in tables, graphics and diagrams. Correlation between serum IL-2 level and Treg/Th17 ratio was analyzed using Spearman's correlation test. Statistical analysis was performed using the SPSS software version 23.0, and a p value <0.05 was considered statistically significant. The Ethics Committee Health Research of Faculty of Medicine, Universitas Airlangga has approved this study with 0346/KEPK/VI/2018.

RESULTS

Thirty SLE patients who were treated at the inpatient department of the internal medicine department of the dr. Soetomo General Hospital, Surabaya, Indonesia who met the inclusion and exclusion criteria were enrolled in this study, consisting of thirty (100%) females. General characteristics of study subjects are shown in Table 1. Data such as age, hemoglobin, WBC, lymphocyte, ESR, C3 and SLAM score were normally distributed, whereas data such as thrombocyte, CRP and C4 were not normally distributed. Test of normality were performed on all data using Shapiro-Wilk test.

The diagnosis of SLE in this study was assessed using the ACR 1997 criteria, fulfilling at least 4 of 11 symptoms and signs. Table 2 shows the distribution of clinical and laboratory manifestations of research subjects based on ACR 1997 criteria.

Serum IL-2 levels, Treg percentage, Th17 percentage and Treg/Th17 ratio were measured on all study samples. The normality test using Saphiro-Wilk showed abnormal distribution of IL-2, Treg percentage and Treg/Th17 ratio, whereas Th17 percentage showed normal distribution. The results of serum IL-2 levels, Treg percentage, Th17 percentage and Treg/Th17 ratio are shown in Table 3. For comparison, measurements were made on healthy subjects with the same methods and examination tools. In healthy subjects the median of serum IL-2 level was 49.31 pg/mL; the median of Treg percentage was 11.75%; the average of Th17 percentage was $13.01 \pm 0.91\%$; and the Treg/Th17 ratio was 0.830. Figure 1 shows there was a significant correlation ($r_s = 0.463$; $P = 0.010$) between IL-2 and Treg/Th17 ratio.

Table 1: General characteristics of the patients

Indicator	Results (n=30)				
	Frequency (%)	Mean ± SD	Median	Min	Max
Age		31 ± 10			
16-20 year	4 (13%)				
20-30 year	12 (40%)				
30-40 year	8 (27%)				
>40 year	6 (20%)				
Hemoglobin (g/dL)		7.1 ± 2.3			
< 12.0	29 (96.7%)				
> 12.0	1 (3.3%)				
WBC (cell/mm ³)		8056 ± 3442			
< 4000	1 (3.3%)				
> 4000	29 (96.7%)				
Lymphocyte (cell/mm ³)		968 ± 428			
< 1500	3 (10%)				
> 1500	27 (90%)				
Thrombocyte (cell/mm ³)			190500	2000	744000
< 100000	12 (40%)				
> 100000	18 (60%)				
ESR (mm/hour)		62 ± 38			
CRP (mg/dL)			7.84	0.10	290.77
C3 (mg/dL)		28.88 ± 12.98			
C4 (mg/dL)			15.25	5.00	54.30
SLAM		29 ± 4			

Table 2: Distribution of clinical and laboratory manifestations based on ACR 1997 criteria

ACR 1997 criteria	Total	Percentage (%)
Malar rash	4	13.3
Discoid rash	8	26.7
Photosensitivity	13	43.3
Oral Ulcer	10	33.3
Arthritis	23	76.7
Serositis	14	46.7
Kidney Disorder	8	26.7
Neurological Disorders	6	20.0
Hematological Disorders		
Hemolytic Anemia	4	13.3
Leukopenia < 4000/mm ³	1	3.3
Lymphopenia < 1500/mm ³	27	90.0
Thrombocytopenia < 100.000/mm ³	12	40.0
Immunology Disorder:		
Anti ds-DNA negative (≤ 92.6 WHO unit/mL)	18	60
Anti ds-DNA positive (> 92.6 WHO unit/mL)	12	40
ANA test:		
Negative (< 20 unit)	7	23.3
Positive (≥ 20 unit)	22	73.3

Table 3: Serum IL-2 levels, Treg and Th17 percentage, Treg/Th17 ratio

Indicator	Results (n=30)		
	Results	Minimum	Maximum
Treg (%)	11.92	1.39	41.20
Th17 (%)	30.09 ± 12.60	5.62	56.90
Treg/Th17 ratio	0.405	0.100	0.918




Fig. 1: Correlation between serum IL-2 level and Treg/Th17 ratio in active SLE using Spearman test, r showed the correlation coefficient.

DISCUSSION

There was a significant correlation between serum IL-2 levels and Treg/Th17 ratio in active SLE. The decline in development of IL-2 affects the balance of Treg/Th17 and induces a transition to T17. In SLE, there is a disturbance of Treg and Th17 balance in the form of a shift towards Th17. Furthermore, there is a decrease in the Treg/Th17 ratio. The moderate size of correlation coefficient might be due to other cytokines that play a role in Treg/Th17 balance such as TGF- β , TNF- α , IL-1 β , IL-6, IL-10, IL-21, and IL-23 [2,9].

From the total of thirty patients involved in this study, all patients (100%) were female. Other studies also reported female dominance in SLE [20–22]. These studies support the theory that SLE affects women more than men, presumably due to the influence of the hormones estrogen and prolactin. However, another report showed case in male who had six signs and symptoms, such as malar rash, arthritis, photosensitivity, anemia, immunologic disorder, and antinuclear antibody [23].

The mean age of the study subjects were almost the same when compared to some previous studies [16,20–22]. These studies support the theory that SLE is more common in reproductive age, presumably due to the influence of the hormones estrogen and prolactin.

The most common clinical manifestations based on the ACR 1997 criteria were arthritis, obtained

in 76.7% of the study subjects. Several other studies also reported arthritis as the most clinical manifestation [20,24,25]. Arthritis can be found as an initial manifestation at the time of diagnosis or as an accompanying manifestation of flare. Some flares are caused by bacterial infections [26,27]. These studies support the theory that musculoskeletal manifestations are prominent and often found in active SLE.

This study used the SLAM score to assess disease activity and obtained an average SLAM score of 29.3. The score <20 indicates the disease is inactive, while the score \geq indicates the disease is active. In this study, all study subjects had SLAM scores >20 so it was concluded that all subjects in this study were in active disease condition. All subjects in this study were naive patients who had not received corticosteroid therapy or other immunosuppressants. Various studies reported that Asian races are associated with higher disease activity [28–30]. A study in Yogyakarta found that the average SLAM score of hospitalized SLE patients was by 16.7 [30]. Whereas, research conducted in 3 countries with a predominantly white population found that the average SLAM score were lower at 4.1 in America, 6.3 in the UK, and 7.3 in Canada [31]. These studies support the theory that SLE disease activity in Asian populations is higher than that of white populations. The study by reported a lower SLAM score than this study, it might be related to the effect of therapy.

Indeed, Treg can affect individual cells' ability to generate cytokines of effectors [32]. For Treg and Th17 cell growth, IL-2 is a cytokine that is essential. The differentiation of Treg effector cells is influenced by IL-2, so in low IL-2 conditions there is a decrease in the amount of Treg. IL-2 plays a role in the expansion and conversion of CD4 + Foxp3 T cells into CD4 + Foxp3 + or Treg T cells. In the pathogenesis of SLE, besides affecting Treg, IL-2 also affects Th17. IL-2 is known to inhibit IL-17A expression and differentiation of Th17 cells. IL-2 inhibits differentiation of Th17 through activation of STAT5, decreased ROR γ t, and production of Ets-1 [33-35].

Some limitation of this study that could influence the results, among others, are that this study was a cross-sectional research so that the serum IL-2 level, Treg and Th17 percentage, Treg/Th17 ratio measurement was only done once. Moreover, serum IL-2 levels, Treg and Th17 percentages, Treg/Th17 ratio do not have a universal reference value that applies universally.

CONCLUSION

There was a significant correlation between serum IL-2 levels and Treg/Th17 ratio in active SLE. In active SLE, there is a decrease in IL-2 levels which does not fully affect the Treg/Th17 balance. It is thought that there are other cytokines that affect the balance. Further research need to be done with a comparative study between healthy and SLE naive patients with a larger sample size and balanced proportions between the two groups. A pre- and post-research is needed to assess changes in IL-2 levels, Treg percentage, Th17 percentage, Treg/Th17 ratio, and SLE disease activity after administration of steroid therapy and immunosuppressants. Various other cytokines and chemokines that might be involved in the Treg/Th17 balance in SLE disease need to be studied and then analyzed for their strength and effects with each other.

29

CONFLICT OF INTEREST

The author stated that there is no conflict of interest.

REFERENCES

1. Carter EE, Barr SG, Clarke AE. The global burden of SLE: Prevalence, health disparities and socioeconomic impact, *Nature Reviews Rheumatology*, 2016; 12(10): 605-20.
2. Crispin J, Tsokos G: T cells and systemic lupus erythematosus. W: Lahita R, Tsokos G, Buyon J, Koike T, red. *Systemic Lupus Erythematosus*, 5th edn. San Diego: Elsevier; 2010, s. 129-42.
3. Sari NPWP. Precipitating Factors and Preventive Behavior towards the Exposures of Systemic

- Lupus Erythematosus, *Jurnal Ners*, 2016; 11(2): 212-9.
4. Yanih I. Quality of Life in Patient with Systemic Lupus Erythematosus (SLE), *Jurnal Berkala Epidemiologi*, 2016; 4(1): 1-12.
5. Marpaung B, Ginting AR, Sjah OM. Serum Midkine levels in systemic lupus erythematosus, *Rawal Medical Journal*, 2019; 44(2): 414-7.
6. Azevedo PC, Murphy G, Isenberg DA: Pathology of systemic lupus erythematosus: the challenges ahead. W: *Systemic Lupus Erythematosus*. Springer; 2014, s. 1-16.
7. Nelson BH. IL-2, Regulatory T Cells, and Tolerance, *The Journal of Immunology*, 2004; 172(7): 3983-8.
8. Malek TR, Bayer AL. Tolerance, not immunity, crucially depends on IL-2, *Nature Reviews Immunology*, 2004; 4(9): 665-74.
9. Lieberman LA, Tsokos GC. The IL-2 Defect in Systemic Lupus Erythematosus Disease Has an Expansive Effect on Host Immunity, *Journal of Biomedicine and Biotechnology*, 2010; 2010.
10. Malek TR, Castro I. Interleukin-2 Receptor Signaling: At the Interface between Tolerance and Immunity, *Immunity*, 2010; 33(2): 153-65.
11. Bachmann MF, Oxenius A. Interleukin 2: From immunostimulation to immunoregulation and back again, *EMBO Reports*, 2007; 8(12): 1142-8.
12. Malek TR. The Biology of IL-2, Annual review of stem cell development, 2008(Annu. Rev. Immunol. 2008. 26:453-79 First): 453-79.
13. Xu L, Zhang L, Yi Y, Kang HK, Datta SK. Human lupus T cells resist inactivation and escape death by upregulating COX-2, *Nature Medicine*, 2004; 10(4): 411-5.
14. Bermas BL, Petri M, Goldman D i wsp. T helper cell dysfunction in systemic lupus erythematosus (SLE): Relation to disease activity, *Journal of Clinical Immunology*, 1994; 14(3): 169-77.
15. Viillard JF, Pellegrin JL, Ranchin V i wsp. Th1 (IL-2, interferon-gamma (IFN- γ)) and Th2 (IL-10, IL-4) cytokine production by peripheral blood mononuclear cells (PBMC) from patients with systemic lupus erythematosus (SLE), *Clinical and Experimental Immunology*, 1999; 115(1): 189-95.
16. Sedighi S, Aghaei M, Musavi S, Nomali M. Relationship between serum level of interleukin-2 in patients with systemic lupus erythematosus and disease activity in comparison with control group, *Journal of Clinical and Diagnostic Research*, 2014; 8(7): 18-20.
17. Ortega C, Garcia-cozar F, Pena J, Sanchez-guijo P, Santamaria M. Relationship of with Disease Activity in SLE Patients IL-4, b.d.: 257-60.
18. Mizui M, Koga T, Lieberman LA i wsp. IL-2 Protects Lupus-Prone Mice from Multiple End-Organ Damage by Limiting CD4 - CD8 - IL-17-Producing T Cells, *The Journal of Immunology*, 2014; 193(5): 2168-77.
19. He J, Zhang X, Wei Y i wsp. Low-dose interleukin-2 treatment selectively modulates CD4+ T cell subsets in patients with systemic

- lupus erythematosus, *Nature Medicine*, 2016; 22(9): 991–3.
20. Zian Z, Maamar M, Aouni M El i wsp. Immunological and Clinical Characteristics of Systemic Lupus Erythematosus: A Series from Morocco, *BioMed Research International*, 2018; 2018.
 21. Li M, Zhang W, Leng X i wsp. Chinese SLE Treatment and Research group (CSTAR) registry: I Major clinical characteristics of Chinese patients with systemic lupus erythematosus, *Lupus*, 2013; 22(11): 1192–9.
 22. AlSaleh J, Jassim V, ElSayed M, Saleh N, Harb D. Clinical and immunological manifestations in 151 SLE patients living in Dubai, *Lupus*, 2008; 17(1): 62–6.
 23. Sibarani H, Zubir Z. Systemic lupus erythematosus in a male patient, *IOP Conf Series: Earth and Environmental Science*, 2018; 125(1).
 24. Fawzy SM, Gheita TA, El-Nabarawy E, El-Demellawy HH, Shaker OG. Serum BAFF level and its correlations with various disease parameters in patients with systemic sclerosis and systemic lupus erythematosus, *Egyptian Rheumatologist*, 2011; 33(1): 45–51.
 25. Tydén H, Lood C, Gullstrand B i wsp. Increased serum levels of S100A8/A9 and S100A12 are associated with cardiovascular disease in patients with inactive systemic lupus erythematosus, *Rheumatology (United Kingdom)*, 2013; 52(11): 2048–55.
 26. Patrick J, Marpaung B, Ginting Y. Differences of serum procalcitonin levels between bacterial infection and flare in systemic lupus erythematosus patients, *IOP Conference Series Earth and Environmental Science*, 2018; 125(1).
 27. Miftahussurur M, Yamaoka Y. Helicobacter pylori virulence genes and host genetic polymorphisms as risk factors for peptic ulcer disease, *Expert review of gastroenterology & hepatology*, 2015; 9(12): 1535.
 28. Boers A, Li Q, Wong M, Miller M, Littlejohn G. Differences in SLE disease activity between patients of Caucasian and South-East Asian/Chinese background in an Australian hospital, *APLAR Journal of Rheumatology*, 2006; 9(1): 43–8.
 29. Golder V, Connelly K, Staples M, Morand E, Hoi A. Association of Asian ethnicity with disease activity in SLE: An observational study from the Monash Lupus Clinic, *Lupus*, 2013; 22(13): 1425–30.
 30. Eritomatosus L, Les S, Paramaiswari A, Kertia N, Achadiyono DW. Ayu Paramaiswari, 2014; 4(1): 15–22.
 31. Panopalis P, Petri M, Manzi S i wsp. The systemic lupus erythematosus tri-nation study: Cumulative indirect costs, *Arthritis Care and Research*, 2007; 57(1): 64–70.
 32. Wammes LJ, Hamid F, Wiria AE i wsp. Regulatory T cells in human geohelminth infection suppress immune responses to BCG and Plasmodium falciparum, *European Journal of Immunology*, 2010; 40(2): 437–42.
 33. Harada T, Kyttaris V, Li Y, Juang YT, Wang Y, Tsokos GC. Increased expression of STAT3 in SLE T cells contributes to enhanced chemokine-mediated cell migration, *Autoimmunity*, 2007; 40(1): 1–8.
 34. Laurence A, Tato CM, Davidson TS i wsp. Interleukin-2 Signaling via STAT5 Constrains T Helper 17 Cell Generation, *Immunity*, 2007; 26(3): 371–81.
 35. Moisan J, Grenningloh R, Bettelli E, Oukka M, Ho IC. Ets-1 is a negative regulator of Th17 differentiation, *Journal of Experimental Medicine*, 2007; 204(12): 2825–35.

Serum Interleukin-2 Level Associated with TregTh17 Ratio in Active Systemic Lupus Erythematosus

ORIGINALITY REPORT

22%

SIMILARITY INDEX

15%

INTERNET SOURCES

20%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Masoud Keikha, Mohammad Ali-Hassanzadeh, Mohsen Karbalaei. "Association of Helicobacter pylori vacA genotypes and peptic ulcer in Iranian population: a systematic review and meta-analysis", Research Square, 2020
Publication 1%

- 2** "Original Research", European Heart Journal Supplements, 2017
Publication 1%

- 3** Claire Bombardier, Dafna D. Gladman, Murray B. Urowitz, Dominique Caron et al. "Derivation of the sledai. A disease activity index for lupus patients", Arthritis & Rheumatism, 1992
Publication 1%

- 4** Lee Ann Garrett-Sinha, Alyssa Kearly, Anne B. Satterthwaite. "The Role of the Transcription Factor Ets1 in Lupus and Other Autoimmune Diseases", Critical Reviews in Immunology, 2016
Publication 1%

5

Yorleny Vicioso, Hermann Gram, Rose Beck, Abhishek Asthana, Keman Zhang, Derek P. Wong, John Letterio, Reshmi Parameswaran. "Combination Therapy for Treating Advanced Drug-Resistant Acute Lymphoblastic Leukemia", *Cancer Immunology Research*, 2019

Publication

1 %

6

S. Phatak, S. Chaurasia, S. K. Mishra, R. Gupta, V. Agrawal, A. Aggarwal, R. Misra. "Urinary B cell activating factor (BAFF) and a proliferation-inducing ligand (APRIL): potential biomarkers of active lupus nephritis", *Clinical & Experimental Immunology*, 2017

Publication

1 %

7

www.em-consulte.com

Internet Source

1 %

8

Miao Shao, Jing He, Ruijun Zhang, Xia Zhang, Yue Yang, Chun Li, Xiaoyun Liu, Xiaolin Sun, Zhanguo Li. "Interleukin-2 Deficiency Associated with Renal Impairment in Systemic Lupus Erythematosus", *Journal of Interferon & Cytokine Research*, 2019

Publication

1 %

9

N/A. "Current World Literature", *Current Opinion in Rheumatology*, 09/2008

Publication

1 %

10	onlinelibrary.wiley.com Internet Source	1 %
11	Vaishali R. Moulton, Abel Suarez-Fueyo, Esra Meidan, Hao Li, Masayuki Mizui, George C. Tsokos. "Pathogenesis of Human Systemic Lupus Erythematosus: A Cellular Perspective", Trends in Molecular Medicine, 2017 Publication	1 %
12	doaj.org Internet Source	1 %
13	edoc.hu-berlin.de Internet Source	1 %
14	jxym.amegroups.com Internet Source	1 %
15	arthritis.pure.elsevier.com Internet Source	<1 %
16	sinta3.ristekdikti.go.id Internet Source	<1 %
17	April Jorge, Victor M. Castro, April Barnado, Vivian Gainer et al. "Identifying lupus patients in electronic health records: Development and validation of machine learning algorithms and application of rule-based algorithms", Seminars in Arthritis and Rheumatism, 2019 Publication	<1 %

18

Miao Miao, Xian Xiao, Jiayi Tian, Yunzhi Zhufeng et al. "Therapeutic Potential of Targeting Tfr/Tfh Cell Balance by Low-Dose-IL-2 in Active SLE: A Post-Hoc Analysis from a Double-Blind RCT Study", Research Square, 2020

Publication

<1 %

19

jkms.kams.or.kr

Internet Source

<1 %

20

lup.lub.lu.se

Internet Source

<1 %

21

Artono, Bakti Surarto, Nyilo Purnami, Fransiska Hutahaen, M. Reza Mahardhika. "The Association of IL-1 Alpha Level and TNF Alpha Expressions on Bone Destruction in Chronic Suppurative Otitis Media and Cholesteatoma", Indian Journal of Otolaryngology and Head & Neck Surgery, 2019

Publication

<1 %

22

H Sibarani, Z Zubir. "Systemic lupus erythematosus in a male patient", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

23

J. Tyrrell-Price. "The effect of interleukin-10 and of interleukin-12 on the in vitro

<1 %

production of anti-double-stranded DNA antibodies from patients with systemic lupus erythematosus", Clinical & Experimental Immunology, 4/2001

Publication

24

www.pubmedcentral.nih.gov

Internet Source

<1 %

25

cancerres.aacrjournals.org

Internet Source

<1 %

26

emedicine.medscape.com

Internet Source

<1 %

27

www.karger.com

Internet Source

<1 %

28

S K Katkam, L Rajasekhar, V K Kutala. " The influence of functional polymorphic positions of molecules on risk for South Indian systemic lupus erythematosus patients ", Lupus, 2018

Publication

<1 %

29

www.intechopen.com

Internet Source

<1 %

30

Kelly C Cushing, Xiaomeng Du, Yanhua Chen, LC Stetson et al. "Inflammatory Bowel Disease Risk Variants Are Associated with an Increased Risk of Skin Cancer", Cold Spring Harbor Laboratory, 2021

Publication

<1 %

- | | | |
|----|---|------|
| 31 | aricjournal.biomedcentral.com
Internet Source | <1 % |
| 32 | docs.google.com
Internet Source | <1 % |
| 33 | gradworks.proquest.com
Internet Source | <1 % |
| 34 | repository.wima.ac.id
Internet Source | <1 % |
| 35 | www.jrheum.org
Internet Source | <1 % |
| 36 | Jing He, Ruijun Zhang, Miao Shao, Xiaozhen Zhao et al. "Efficacy and safety of low-dose IL-2 in the treatment of systemic lupus erythematosus: a randomised, double-blind, placebo-controlled trial", <i>Annals of the Rheumatic Diseases</i> , 2020
Publication | <1 % |
| 37 | José C. Crispín. "T-Cells and Systemic Lupus Erythematosus", <i>Systemic Lupus Erythematosus</i> , 2011
Publication | <1 % |
| 38 | Linda A. Lieberman, George C. Tsokos. "The IL-2 Defect in Systemic Lupus Erythematosus Disease Has an Expansive Effect on Host Immunity", <i>Journal of Biomedicine and Biotechnology</i> , 2010 | <1 % |

39

f1000researchdata.s3.amazonaws.com

Internet Source

<1 %

40

www.jehp.net

Internet Source

<1 %

41

www.onlinelibrary.wiley.com

Internet Source

<1 %

42

Abel Suárez-Fueyo, Sean J. Bradley, David Klatzmann, George C. Tsokos. "T cells and autoimmune kidney disease", *Nature Reviews Nephrology*, 2017

Publication

<1 %

43

Hideyuki Murata, Ryutaro Matsumura, Akio Koyama, Takao Sugiyama et al. "T cell receptor repertoire of T cells in the kidneys of patients with lupus nephritis", *Arthritis & Rheumatism*, 2002

Publication

<1 %

44

Rahul Sharma, Gilbert R Kinsey. "Regulatory T cells in Acute and Chronic Kidney Diseases", *American Journal of Physiology - Renal Physiology*, 2017

Publication

<1 %

45

Teresa A Lillis, Vanessa Tirone, Nisarg Gandhi, Stacy Weinberg et al. "Sleep Disturbance and Depression Symptoms Mediate the Relationship between Pain and Cognitive

<1 %

Dysfunction in Lupus Patients", Arthritis Care & Research, 2018

Publication

46	archive.org Internet Source	<1 %
47	cwww.intechopen.com Internet Source	<1 %
48	d-nb.info Internet Source	<1 %
49	journal-cdn.frontiersin.org Internet Source	<1 %
50	repository.um.edu.my Internet Source	<1 %
51	worldwidescience.org Internet Source	<1 %
52	www.medicaljournals.se Internet Source	<1 %
53	"Moderated Poster Presentations", International Journal of Urology, 8/2006 Publication	<1 %
54	Christian Lood. "Protein synthesis of the pro-inflammatory S100A8/A9 complex in plasmacytoid dendritic cells and cell surface S100A8/A9 on leukocyte subpopulations in systemic lupus erythematosus", Arthritis Research & Therapy, 2011	<1 %

55

J Patrick, B Marpaung, Y Ginting. "Differences of serum procalcitonin levels between bacterial infection and flare in systemic lupus erythematosus patients", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

56

Joo, Young Bin, and Sang-Cheol Bae. "Assessment of clinical manifestations, disease activity and organ damage in 996 Korean patients with systemic lupus erythematosus: comparison with other Asian populations", International Journal of Rheumatic Diseases, 2014.

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Serum Interleukin-2 Level Associated with TregTh17 Ratio in Active Systemic Lupus Erythematosus

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
