

DAFTAR PUSTAKA

- Abbasinia, M. (2016) 'The Effect of a Designed Respiratory Care Program on the Incidence of Ventilator-Associated Pneumonia: A Clinical Trial', *Journal of Caring Sciences*, 5(2), pp. 161–167. doi: 10.15171/jcs.2016.017.
- Adini, S. (2018) 'the Comparison of the Effect of Honey and Chlorhexidine in Preventing Ventilator Associated Pneumonia in Patients on Mechanical Ventilaton', *Belitung Nursing Journal*, 4(2), pp. 161–167. doi: 10.33546/bnj.355.
- Akl, E. (2019) 'Cochrane Handbook for Systematic Reviews of Interventions', *Cochrane Handbook for Systematic Reviews of Interventions*, (October). doi: 10.1002/9781119536604.
- Alecrim, R. X. (2019) 'Strategies for preventing ventilator-associated pneumonia: an integrative review', *Revista brasileira de enfermagem*, 72(2), pp. 521–530. doi: 10.1590/0034-7167-2018-0473.
- Ataee, R. A. (2016) 'To: The use of 2% chlorhexidine gel and toothbrushing for oral hygiene of patients receiving mechanical ventilation: Effects on ventilator-associated pneumonia', *Revista Brasileira de Terapia Intensiva*, 26(4), pp. 438–440. doi: 10.5935/0103-507X.20140069.
- Atient, P. (2016) 'E DUCATION ON PATIENT- VENTILATOR SYNCHRONY, CLINICIANS' KNOWLEDGE LEVEL, AND DURATION OF MECHANICAL VENTILATION', 25(6).
- Berry, A. M. (2011) 'Effects of three approaches to standardized oral hygiene to reduce bacterial colonization and ventilator associated pneumonia in mechanically ventilated patients: A randomised control trial', *International Journal of Nursing Studies*. Elsevier Ltd, 48(6), pp. 681–688. doi: 10.1016/j.ijnurstu.2010.11.004.
- Branch-Elliman, W., Wright, S. B. and Howell, M. D. (2015) 'Determining the ideal strategy for ventilator-associated pneumonia prevention: Cost-benefit analysis', *American Journal of Respiratory and Critical Care Medicine*, 192(1), pp. 57–63. doi: 10.1164/rccm.201412-2316OC.
- de Camargo, L., da Silva, S. N. and Chambrone, L. (2019) 'Efficacy of toothbrushing procedures performed in intensive care units in reducing the risk of ventilator-associated pneumonia: A systematic review', *Journal of Periodontal Research*, 54(6), pp. 601–611. doi: 10.1111/jre.12668.
- Chacko, R. (2017) 'Oral decontamination techniques and ventilator-associated pneumonia', *British Journal of Nursing*, 26(11), pp. 594–599. doi: 10.12968/bjon.2017.26.11.594.
- Chang, I. and Schibler, A. (2016) 'Ventilator Associated Pneumonia in Children', *Paediatric Respiratory Reviews*. Elsevier Ltd, 20, pp. 10–16. doi: 10.1016/j.prrv.2015.09.005.
- Chen, Y. (2016) 'Prospective observational study to compare oral topical metronidazole versus 0.2% chlorhexidine gluconate to prevent nosocomial pneumonia', *American Journal of Infection Control*. Elsevier Inc., 44(10), pp. 1116–1122. doi: 10.1016/j.ajic.2016.03.054.
- Chulay, M. and Gallagher, J. J. (2017) 'AACN Practice Alert Prevention of Ventilator-Associated Pneumonia in Adults', *American Association of Critical-*

- Care Nurses*, 37(3), pp. 22–6. Available at: <https://doi.org/10.4037/ccn2017460>.
- Da Collina, G. A. (2017) ‘Oral hygiene in intensive care unit patients with photodynamic therapy: Study protocol for randomised controlled trial’, *Trials*, 18(1). doi: 10.1186/s13063-017-2133-y.
- Cooper, V. B. and Haut, C. (2013) ‘Preventing ventilator-associated pneumonia in children: An evidence-based protocol’, *Critical Care Nurse*, 33(3), pp. 21–29. doi: 10.4037/ccn2013204.
- Deschepper, M. (2018) ‘Effects of chlorhexidine gluconate oral care on hospital mortality: a hospital-wide, observational cohort study’, *Intensive Care Medicine*. Springer Berlin Heidelberg, 44(7), pp. 1017–1026. doi: 10.1007/s00134-018-5171-3.
- Enwere, E. N. (2016) ‘Impact of chlorhexidine mouthwash prophylaxis on probable ventilator-associated pneumonia in a surgical intensive care unit’, *International Journal of Critical Illness and Injury Science*, 6(1), pp. 3–8. doi: 10.4103/2229-5151.177368.
- Fernanda, C. (2017) ‘Impact of oral hygiene involving toothbrushing versus chlorhexidine in the prevention of ventilator-associated pneumonia: A randomized study’, *BMC Infectious Diseases*. BMC Infectious Diseases, 17(1). doi: 10.1186/s12879-017-2188-0.
- Garrido-Benedicto, P. (2017) ‘Effect of daily cleaning with chlorhexidine on the incidence of contamination of blood cultures in the critical patient’, *Enfermería Intensiva (English ed.)*. Sociedad Española de Enfermería Intensiva y Unidades Coronarias (SEEIUC), 28(3), pp. 97–104. doi: 10.1016/j.enfie.2017.03.001.
- Gomaa, M. M., Wahba, Y. and El-Bayoumi, M. A. (2017) ‘Pre versus post application of a 0.12% chlorhexidine based oral hygiene protocol in an Egyptian pediatric intensive care unit: Practice and effects’, *The Egyptian Journal of Critical Care Medicine*. The Egyptian College of Critical Care Physicians, 5(3), pp. 87–91. doi: 10.1016/j.ejccm.2017.11.002.
- Haghighi, A. (2017) ‘The impact of oral care on oral health status and prevention of ventilator-associated pneumonia in critically ill patients’, *Australian Critical Care*. Australian College of Critical Care Nurses Ltd, 30(2), pp. 69–73. doi: 10.1016/j.aucc.2016.07.002.
- Hanifi, N. (2017) ‘The effect of ozonated water and chlorhexidine gluconate on prevention of ventilator-associated pneumonia: A double-blind, randomized, clinical trial’, *Iranian Red Crescent Medical Journal*, 19(10), pp. 0–7. doi: 10.5812/ircmj.60576.
- Hassan, Z. M. and Wahsheh, M. A. (2017) ‘Knowledge level of nurses in Jordan on ventilator-associated pneumonia and preventive measures’, *Nursing in Critical Care*, 22(3), pp. 125–132. doi: 10.1111/nicc.12273.
- Hillier, B. (2013) ‘Preventing ventilator-associated pneumonia through oral care, product selection, and application method: A literature review’, *AACN Advanced Critical Care*, 24(1), pp. 38–58. doi: 10.1097/NCI.0b013e31827df8ad.
- Khan, M. (2017) ‘Oral Care Effect on Intubated Patient with 0.2 per cent Chlorhexidine Gluconate and Tooth Brushing in Intensive Care Unit’, *Journal of Academy of Advanced Dental Research*, p. 222941121772909. doi: 10.1177/2229411217729099.

- Klarin, B. (2018) 'Can probiotics be an alternative to chlorhexidine for oral care in the mechanically ventilated patient? A multicentre, prospective, randomised controlled open trial', *Critical Care*. *Critical Care*, 22(1), pp. 1–10. doi: 10.1186/s13054-018-2209-4.
- Kusahara, D. M. (2012) 'Oral care and oropharyngeal and tracheal colonization by Gram-negative pathogens in children', *Nursing in Critical Care*, 17(3), pp. 115–122. doi: 10.1111/j.1478-5153.2012.00494.x.
- Kusahara, D. M., Peterlini, M. A. S. and Pedreira, M. L. G. (2012) 'Oral care with 0.12% chlorhexidine for the prevention of ventilator-associated pneumonia in critically ill children: Randomised, controlled and double blind trial', *International Journal of Nursing Studies*, 49(11), pp. 1354–1363. doi: 10.1016/j.ijnurstu.2012.06.005.
- Lambert, M.-L. (2013) 'Prevention of ventilator-associated pneumonia in intensive care units: an international online survey', *Antimicrobial Resistance and Infection Control*, 2(1), p. 9. doi: 10.1186/2047-2994-2-9.
- Lee, S. (2019) 'Chlorhexidine-related mortality rate in critically ill subjects in intensive care units: A systematic review and meta-analysis', *Respiratory Care*, 64(3), pp. 337–349. doi: 10.4187/respcare.06434.
- Lev, A., Aied, A. S. and Arshed, S. (2015) 'The effect of different oral hygiene treatments on the occurrence of ventilator associated pneumonia (VAP) in ventilated patients', *Journal of Infection Prevention*, 16(2), pp. 76–81. doi: 10.1177/1757177414560252.
- Lin, Y. J. (2015) 'Reduced occurrence of ventilator-associated pneumonia after cardiac surgery using preoperative 0.2% chlorhexidine oral rinse: Results from a single-centre single-blinded randomized trial', *Journal of Hospital Infection*, 91(4), pp. 362–366. doi: 10.1016/j.jhin.2015.08.018.
- McCue, M. K. and Palmer, G. A. (2019) 'Use of Chlorhexidine to Prevent Ventilator-Associated Pneumonia in a Long-term Care Setting: A Retrospective Medical Record Review', *Journal of Nursing Care Quality*, 34(3), pp. 263–268. doi: 10.1097/NCQ.0000000000000367.
- Meidani, M. (2017) 'Oropharyngeal Irrigation to Prevent Ventilator-Associated-Pneumonia: Comparing Potassium Permanganate with Chlorhexidine Abstract', *International Journal of Preventive Medicine*, 8, pp. 2018–2021. doi: 10.4103/ijpvm.IJPVM.
- Mogyoródi, B. (2016) 'Ventilator-associated pneumonia and the importance of education of ICU nurses on prevention - Preliminary results', *Interventional Medicine and Applied Science*, 8(4), pp. 147–151. doi: 10.1556/1646.8.2016.4.9.
- Nasiriani, K. (2016) 'The effect of brushing with a soft toothbrush and distilled water on the incidence of ventilator-associated pneumonia in the intensive care unit', *Tanaffos*, 15(2), pp. 101–107.
- Okgün Alcan, A., Demir Korkmaz, F. and Uyar, M. (2016) 'Prevention of ventilator-associated pneumonia: Use of the care bundle approach', *American Journal of Infection Control*. Elsevier Inc., 44(10), pp. e173–e176. doi: 10.1016/j.ajic.2016.04.237.
- Ory, J. (2017) 'Comparative study of 2 oral care protocols in intensive care units', *American Journal of Infection Control*. Elsevier Inc., 45(3), pp. 245–250. doi: 10.1016/j.ajic.2016.09.006.

- Osman, S. (2020) 'The incidence of ventilator-associated pneumonia (VAP) in a tertiary-care center: Comparison between pre- and post-VAP prevention bundle', *Journal of Infection and Public Health*. King Saud Bin Abdulaziz University for Health Sciences, 13(4), pp. 552–557. doi: 10.1016/j.jiph.2019.09.015.
- Panchabhai, T. S. (2009) 'Oropharyngeal cleansing with 0.2% chlorhexidine for prevention of nosocomial pneumonia in critically ill patients: An open-label randomized trial with 0.01% potassium permanganate as control', *Chest*, 135(5), pp. 1150–1156. doi: 10.1378/chest.08-1321.
- Roberts, N. and Moule, P. (2011) 'Chlorhexidine and tooth-brushing as prevention strategies in reducing ventilator-associated pneumonia rates.', *Nursing in critical care*, 16(6), pp. 295–302. doi: 10.1111/j.1478-5153.2011.00465.x.
- Saensom, D. (2016) 'Oral health and ventilator-associated pneumonia among critically ill patients: a prospective study', *Oral Diseases*, 22(7), pp. 709–714. doi: 10.1111/odi.12535.
- Sharif-Abdullah, S. S. B. (2016) 'The effect of chlorhexidine in reducing oral colonisation in geriatric patients: A randomised controlled trial', *Singapore Medical Journal*, 57(5), pp. 262–266. doi: 10.11622/smedj.2016091.
- Tuon, F. F. (2017) 'Prospective, randomised, controlled study evaluating early modification of oral microbiota following admission to the intensive care unit and oral hygiene with chlorhexidine', *Journal of Global Antimicrobial Resistance*. Taibah University, 8, pp. 159–163. doi: 10.1016/j.jgar.2016.12.007.
- Veitz-Keenan, A. and Ferraiolo, D. M. (2017) 'Oral care with chlorhexidine seems effective for reducing the incidence of ventilator-associated pneumonia', *Evidence-Based Dentistry*. Nature Publishing Group, 18(4), pp. 113–114. doi: 10.1038/sj.ebd.6401272.
- Villar, C. C. (2016) 'Effectiveness of intraoral chlorhexidine protocols in the prevention of ventilator-associated pneumonia: Meta-analysis and systematic review', *Respiratory Care*, 61(9), pp. 1245–1259. doi: 10.4187/respcare.04610.
- Wang, C. H. (2017) 'Normal saline instillation before suctioning: A meta-analysis of randomized controlled trials', *Australian Critical Care*. Australian College of Critical Care Nurses Ltd, 30(5), pp. 260–265. doi: 10.1016/j.aucc.2016.11.001.
- Zand, F. (2017) 'The effects of oral rinse with 0.2% and 2% chlorhexidine on oropharyngeal colonization and ventilator associated pneumonia in adults' intensive care units', *Journal of Critical Care*. Elsevier Inc., 40, pp. 318–322. doi: 10.1016/j.jcrc.2017.02.029.
- Zuckerman, L. M. (2016) 'Oral Chlorhexidine Use to Prevent Ventilator-Associated Pneumonia in Adults: Review of the Current Literature', *Dimensions of Critical Care Nursing*, 35(1), pp. 25–36. doi: 10.1097/DCC.0000000000000154.

DAFTAR LAMPIRAN

Lampiran 1