

## DAFTAR PUSTAKA

- Anton, S. D. *et al.* (2014) ‘Guidelines for Americans How Much Walking Is Needed To Improve Cardiorespira-’, (October 2014), pp. 37–41. doi: 10.1080/02701367.2011.10599766.
- Bouaziz, W. *et al.* (2018) ‘Effects of Interval Aerobic Training Program with Recovery bouts on cardiorespiratory and endurance fitness in seniors’, *Scandinavian Journal of Medicine and Science in Sports*, 28(11), pp. 2284–2292. doi: 10.1111/sms.13257.
- Cadore, E. L. *et al.* (2018) ‘Cardiorespiratory adaptations in elderly men following different concurrent training regimes’, 22(4), pp. 483–490.
- Cegielski, J. *et al.* (2017) ‘A 4-week, lifestyle-integrated, home-based exercise training programme elicits improvements in physical function and lean mass in older men and women: a pilot study’, (0), pp. 1–12. doi: 10.12688/f1000research.11894.1.
- Coetsee, C. and Terblanche, E. (2017) ‘The effect of three different exercise training modalities on cognitive and physical function in a healthy older population’, *European Review of Aging and Physical Activity*. *European Review of Aging and Physical Activity*, 14(1), pp. 1–11. doi: 10.1186/s11556-017-0183-5.
- DeBolt, L. S. and McCubbin, J. A. (2004) ‘The Effects of Home-Based Resistance Exercise on Balance, Power, and Mobility in Adults with Multiple Sclerosis’, *Archives of Physical Medicine and Rehabilitation*, 85(2), pp. 290–297. doi: 10.1016/j.apmr.2003.06.003.
- Department of Health (2011) ‘Start Active, Stay Active: A report on physical activity from the four home countries’ Chief Medical Officers’, *Report*, p. 62.
- Esmail, A. *et al.* (2020) ‘Effects of Dance/Movement Training vs. Aerobic Exercise Training on cognition, physical fitness and quality of life in older adults: A randomized controlled trial’, *Journal of Bodywork and Movement Therapies*, 24(1), pp. 212–220. doi: 10.1016/j.jbmt.2019.05.004.
- Fernández-García, Á. I. *et al.* (2020) ‘How to Improve the Functional Capacity of Frail and Pre-Frail Elderly People? Health, Nutritional Status and Exercise Intervention. The EXERNET-Elder 3.0 Project’, *Sustainability*, 12(15), p. 6246. doi: 10.3390/su12156246.
- Ferreira, C. B. *et al.* (2018) ‘Effects of a 12-Week Exercise Training Program on Physical Function in Institutionalized Frail Elderly’, *Journal of Aging Research*, 2018. doi: 10.1155/2018/7218102.
- Gallè, F. *et al.* (2016) ‘Improving physical fitness and health status perception in community-dwelling older adults through a structured program for physical activity promotion in the city of Naples , Italy : A randomized controlled

- trial', pp. 5–8. doi: 10.1111/ggi.12879.
- Glynn, N. W. *et al.* (2020) 'Impact of Baseline Fatigue on a Physical Activity Intervention to Prevent Mobility Disability', *Journal of the American Geriatrics Society*, 68(3), pp. 619–624. doi: 10.1111/jgs.16274.
- Hong, J. *et al.* (2017) 'Effects of home-based tele-exercise on sarcopenia among community-dwelling elderly adults: Body composition and functional fitness', *Experimental Gerontology*. Elsevier Inc., 87, pp. 33–39. doi: 10.1016/j.exger.2016.11.002.
- Hong, S. Y. and Jun, S. Y. (2017) 'Community Capacity Building Exercise Maintenance Program for Frail Elderly Women', *Asian Nursing Research*. Korean Society of Nursing Science, 11(3), pp. 166–173. doi: 10.1016/j.anr.2017.06.001.
- Hurst, C., Weston, K. L. and Weston, M. (2018) 'The effect of 12 weeks of combined upper- and lower-body high- intensity interval training on muscular and cardiorespiratory fitness in older adults', *Aging Clinical and Experimental Research*. Springer International Publishing, 0(0), p. 0. doi: 10.1007/s40520-018-1015-9.
- Kim, D. Il *et al.* (2018) 'Six weeks of combined aerobic and resistance exercise using outdoor exercise machines improves fitness, insulin resistance, and chemerin in the Korean elderly: A pilot randomized controlled trial', *Archives of Gerontology and Geriatrics*. Elsevier, 75(November 2017), pp. 59–64. doi: 10.1016/j.archger.2017.11.006.
- Kucharski, D. *et al.* (2019) 'Moderate-to-high intensity exercise with person-centered guidance influences fatigue in older adults with rheumatoid arthritis', *Rheumatology International*. Springer Berlin Heidelberg, 39(9), pp. 1585–1594. doi: 10.1007/s00296-019-04384-8.
- Kusmana, D. (2001) *Senam Kesehatan Jantung*. Jakarta: FKUI.
- Lee, T. E. (2018) 'Effects of regular exercise on functional fitness of community elderly', *MATEC Web of Conferences*, 201, pp. 10–14. doi: 10.1051/mateconf/201820106001.
- Leirós-Rodríguez, R. *et al.* (2018) 'Comparisons of the Health Benefits of Strength Training, Aqua-Fitness, and Aerobic Exercise for the Elderly', *Rehabilitation Research and Practice*, 2018, pp. 1–8. doi: 10.1155/2018/5230971.
- Lin, W. Q. *et al.* (2015) 'Factors associated with fatigue among men aged 45 and older: A cross-sectional study', *International Journal of Environmental Research and Public Health*, 12(9), pp. 10897–10909. doi: 10.3390/ijerph120910897.
- Liu, J. Y. W. *et al.* (2017) 'An individualized exercise programme with and without behavioural change enhancement strategies for managing fatigue among frail older people: A quasi-experimental pilot study', *Clinical Rehabilitation*,

31(4), pp. 1–11. doi: 10.1177/0269215516649226.

Maruya, K. *et al.* (2016) ‘Effect of a simple and adherent home exercise program on the physical function of community dwelling adults sixty years of age and older with pre-sarcopenia or sarcopenia’, *Journal of Physical Therapy Science*, 28(11), pp. 3183–3188. doi: 10.1589/jpts.28.3183.

McLellan, A. G. *et al.* (2018) ‘Can older adults improve the identification of moderate intensity using walking cadence?’, *Aging Clinical and Experimental Research*. Springer International Publishing, 30(1), pp. 89–92. doi: 10.1007/s40520-017-0746-3.

Meng, N. H. *et al.* (2020) ‘Effects of concurrent aerobic and resistance exercise in frail and pre-frail older adults: A randomized trial of supervised versus home-based programs’, *Medicine*, 99(29), p. e21187. doi: 10.1097/MD.00000000000021187.

Mun, S. *et al.* (2016) ‘Interrelationships among common symptoms in the elderly and their effects on health-related quality of life: A cross-sectional study in rural Korea’, *Health and Quality of Life Outcomes*. Health and Quality of Life Outcomes, 14(1), pp. 1–10. doi: 10.1186/s12955-016-0549-9.

Nelson, M. E. *et al.* (2004) ‘The Effects of Multidimensional Home-Based Exercise on Functional Performance in Elderly People’, *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 59(2), pp. M154–M160. doi: 10.1093/gerona/59.2.m154.

Nicklas, B. J. *et al.* (2019) ‘Effects of Caloric Restriction on Cardiorespiratory Fitness, Fatigue, and Disability Responses to Aerobic Exercise in Older Adults with Obesity: A Randomized Controlled Trial’, *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*, 74(7), pp. 1084–1090. doi: 10.1093/gerona/gly159.

Outayanik, B. *et al.* (2017) ‘Effects of a physical activity intervention program on nutritional status and health-related physical fitness in thai older adults: Pilot study’, *Asian Journal of Sports Medicine*, 8(1). doi: 10.5812/asjms.37508.

Pedersen, M. T. *et al.* (2017) ‘Effect of team sports and resistance training on physical function, quality of life, and motivation in older adults’, *Scandinavian Journal of Medicine and Science in Sports*, 27(8), pp. 852–864. doi: 10.1111/sms.12823.

Prasetyo, Y. *et al.* (2017) ‘Effects of Aerobic Exercise and Weight Training-Aerobic towards Physical Fitness of Elderly Hajj Candidates’, 7(1), pp. 106–117.

Rofika, A. and Yuniastuti, A. (2018) ‘Effectiveness of Tera Gymnastics and Healthy Heart Gymnastics on Blood Pressure Among Elderly With Hypertension in Sinomwidodo Village, Tambakromo Subdistrict, Pati District’, *Public Health Perspective Journal*, 3(1), pp. 7–13. Available at:

<http://journal.unnes.ac.id/sju/index.php/phpj>.

- Sukartini, T. and Nursalam (2009) 'Manfaat senam tera terhadap kebugaran lansia', 8(3), pp. 153–158.
- Tanaka, E. H. *et al.* (2016) 'The effect of supervised and home based exercises on balance in elderly subjects: a randomized controlled trial to prevent falls', *Revista Brasileira de Geriatria e Gerontologia*, 19(3), pp. 383–397. doi: 10.1590/1809-98232016019.150027.
- Timmons, J. *et al.* (2018) 'Comparison of time-matched aerobic, resistance or concurrent exercise training in older adults', (0), pp. 0–1. doi: 10.1111/sms.13254.
- Vellas, B. and Fleck, F. (2014) 'Most people can stay fit and healthy into old age', *Bulletin of the World Health Organization*, 92(9), pp. 628–629. doi: 10.2471/BLT.14.030914.
- WHO (2018) *Aging and Health*. Available at: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>.
- WHO (2019) *Physical Activity and Older Adults*. Available at: [https://www.who.int/dietphysicalactivity/factsheet\\_olderadults/en/](https://www.who.int/dietphysicalactivity/factsheet_olderadults/en/) (Accessed: 13 September 2019).