

**DAFTAR PUSTAKA**

- Adil, M., Adil, S., Syed, K., Aziz, T. and Badshah, A. (2016) ‘Comparison of Inter Premolar, Molar Widths and Arch Depth Among Different Malocclusions’, *Pakistan Oral & Dental Journal*, 36(2), p. 2016
- ADIL, M., ADIL, S., SYED, K., AZIZ, T. and BADSHAH, A. (2016) ‘Comparison of Inter Premolar, Molar Widths and Arch Depth Among Different Malocclusions’, *Pakistan Oral & Dental Journal*, 36(2), p. 2016
- Aju Wahju Ardani, I. G., Kannayyah, D. and Triwardhani, A. (2019) ‘Correlation of maxillary and mandibular arch form and tooth size ratio in ethnic Javanese malocclusion patient’, *Journal of International Oral Health*. DOI: 10.4103/jioh.jioh\_8\_19.
- Alabdullah, M., Saltaji, H., Abou-Hamed, H. and Youssef, M. (2015) ‘Association between facial growth pattern and facial muscle activity: A prospective cross-sectional study’, *International orthodontics / Collège européen d'orthodontie*. Elsevier Masson SAS, 13(2), pp. 181–194. DOI: 10.1016/j.ortho.2015.03.011.
- Alhammadi, M. S., Halboub, E., Fayed, M. S., Labib, A. and El-Saaidi, C. (2018) ‘Global distribution of malocclusion traits: A systematic review’, *Dental Press Journal of Orthodontics*, 23(6), pp. e1–e10. DOI: 10.1590/2177-6709.23.6.40.e1-10.onl.
- Alhammadi, M. S. (2019) ‘Dentoalveolar compensation in different anteroposterior and vertical skeletal malocclusions’, *Journal of Clinical and Experimental Dentistry*. DOI: 10.4317/jced.56020.
- Ardani, I. G. A. W., Rahmawati, D., Narmada, I. B., Nugraha, A. P., Nadia, S., Taftazani, H. and Kusumawardani, M. K. (2020) ‘Surface Electromyography Unveil the Relationship between Masticatory Muscle Tone and Malocclusion Class I & II in Javanese Ethnic Patient’, *Journal of International Dental and Medical Research*, 13(4), pp. 1447–1454
- Ardani, I. G. A. W., Sanjaya, M. L. and Sjamsudin, J. (2018) ‘Cephalometric characteristic of skeletal Class II malocclusion in Javanese Population at Universitas Airlangga Dental Hospital’, *Contemporary Clinical Dentistry*. DOI: 10.4103/ccd.ccd\_465\_18.
- Ardani, I. G. A. W., Willyanti, I. and Narmada, I. B. (2018) ‘Correlation between vertical components and skeletal class ii malocclusion in ethnic javanese’, *Clinical, Cosmetic and Investigational Dentistry*, 10, pp. 297–302. DOI: 10.2147/CCIDE.S188414.
- Bae, J., Son, W. S., Kim, S. S., Park, S. B. and Kim, Y. Il (2017) ‘Comparison of masticatory efficiency according to Angle’s classification of malocclusion’,

- Korean Journal of Orthodontics*, 47(3), pp. 151–157. DOI: 10.4041/kjod.2017.47.3.151.
- Becht, M. P., Mah, J., Martin, C., Razmus, T., Gunel, E. and Ngan, P. (2014) ‘Évaluation De La Morphologie Du Muscle Masséter Dans Différents Types De Malocclusions En Utilisant La Tomographie Volumétrique À Faisceau Conique’, *International Orthodontics*. Elsevier Masson SAS, 12(1), pp. 32–48. DOI: 10.1016/j.ortho.2013.12.003.
- Biondi, K., Lorusso, P., Fastuca, R., Mangano, A., Zecca, P. A., Bosco, M., Caprioglio, A. and Levrini, L. (2016) ‘Evaluation of masseter muscle in different vertical skeletal patterns in growing patients’, *European Journal of Paediatric Dentistry*, 17(1), pp. 47–52
- Bourzgui, F., Khribchi, A., Rachdy, Z., Housbane, S. and Othmani, M. B. (2016) ‘Evaluation of arch forms depending on the angle classification’, *International Orthodontics*. Elsevier Masson SAS, 14(4), pp. 528–536. DOI: 10.1016/j.ortho.2016.10.017.
- Cassidy, K. M., Harris, E. F., Tolley, E. A. and Keim, R. G. (1998) ‘Genetic influence on dental arch form in orthodontic patients’, *Angle Orthodontist*, pp. 445–454. DOI: 10.1043/0003-3219(1998)068<0445:GIODAF>2.3.CO;2.
- Claro, C. A. de A., Abrão, J. and Reis, S. A. B. (2010) ‘Association between overbite and craniofacial growth pattern’, *Brazilian Oral Research*, 24(4), pp. 425–432. DOI: 10.1590/S1806-83242010000400009.
- Custodio, W., Gomes, S. G. F., Faot, F., Garcia, R. C. M. R. and Del Bel Cury, A. A. (2011) ‘Occlusal force, electromyographic activity of masticatory muscles and mandibular flexure of subjects with different facial types’, *Journal of Applied Oral Science*, 19(4), pp. 343–349. DOI: 10.1590/S1678-77572011005000008.
- Dmytrenko, M. I. (2018) ‘Analysis of electromyographic indexes of temporal and masticatory muscles in patients with distal occlusion complicated by dental crowding’, *Wiadomosci lekarskie (Warsaw, Poland : 1960)*, 71(2), pp. 295–298
- Enlow DH, H. M. (1996) *Essentials of facial growth. 2nd edition*, WB Saunders Company
- Enomoto, A., Watahiki, J., Yamaguchi, T., Irie, T., Tachikawa, T. and Maki, K. (2010) ‘Effects of mastication on mandibular growth evaluated by microcomputed tomography’, *European Journal of Orthodontics*, 32(1), pp. 66–70. DOI: 10.1093/ejo/cjp060.
- Eyquem, A. P., Kuzminsky, S. C., Aguilera, J., Astudillo, W. and Toro-Ibacache, V. (2019) ‘Normal and altered masticatory load impact on the range of

- craniofacial shape variation: An analysis of pre-Hispanic and modern populations of the American Southern Cone’, *PLoS ONE*, 14(12), pp. 1–20. DOI: 10.1371/journal.pone.0225369.
- Graber, L., Vanarsdall, R., Vig, K. and Huang, G. (2017) *Orthodontics: Current Principles and Techniques: Sixth Edition*. Sixth Edit. St. Louis (Missouri): Elsevier
- Gupta, A., Singh, K., Tandon, R., Chauhan, A., Kulshrestha, R. and Trivedi, H. (2017) ‘Functional Electromyographic Activities of Facial Muscles in Different Growth Patterns’, *SM Dentistry Journal*, 3(3), pp. 1–5. DOI: 10.36876/smd.1017s.
- Hall, J. E. (2016) *Guyton and Hall Textbook of Medical Physiology*. 13th edn. Elsevier
- Hichijo, N., Kawai, N., Mori, H., Sano, R., Ohnuki, Y., Okumura, S., Langenbach, G. E. J. and Tanaka, E. (2014) ‘Effects of the masticatory demand on the rat mandibular development’, *Journal of Oral Rehabilitation*, 41(8), pp. 581–587. DOI: 10.1111/joor.12171.
- Hwang, S., Jeong, S., Choi, Y. J., Chung, C. J., Lee, H. S. and Kim, K. H. (2018) ‘Three-dimensional evaluation of dentofacial transverse widths of adults with various vertical facial patterns’, *American Journal of Orthodontics and Dentofacial Orthopedics*. American Association of Orthodontists, 153(5), pp. 692–700. DOI: 10.1016/j.ajodo.2017.08.026.
- Isola, G., Anastasi, G. P., Matarese, G., Williams, R. C., Cutroneo, G., Bracco, P. and Piancino, M. G. (2018) ‘Functional and molecular outcomes of the human masticatory muscles’, *Oral Diseases*, 24(8), pp. 1428–1441. DOI: 10.1111/odi.12806.
- Jaradat, M. (2018) ‘An Overview of Class III Malocclusion (Prevalence, Etiology and Management)’, *Journal of Advances in Medicine and Medical Research*, 25(7), pp. 1–13. DOI: 10.9734/jammr/2018/39927.
- Lione, R., Franchi, L., Noviello, A., Bollero, P., Fanucci, E. and Cozza, P. (2013) ‘Three-dimensional evaluation of masseter muscle in different vertical facial patterns: A cross-sectional study in growing children’, *Ultrasonic Imaging*, 35(4), pp. 307–317. DOI: 10.1177/0161734613502468.
- Mageet, A. O. (2016) ‘Classification of Skeletal and Dental Malocclusion: Revisited’, *Stomatology Edu Journal*, 3(2), pp. 205–211. DOI: 10.25241/2016.3(2).11.
- Mitchell, L., Littlewood, S. J., Nelson-moon, Z. L. and Dyer, F. (2013) *Introduction to Orthodontics*. Fourth Edi. Oxford: Oxford University Press
- Moyers, R. E. (1988) *Handbook Of Orthodontics*. 4th edn

- Nishi, S. E., Basri, R., Alam, M. K., Komatsu, S., Komori, A., Sugita, Y. and Maeda, H. (2017) 'Evaluation of masticatory muscles function in different malocclusion cases using surface electromyography', *Journal of Hard Tissue Biology*, 26(1), pp. 23–28. DOI: 10.2485/jhtb.26.23.
- De Novaes Benedicto, E., Kairalla, S. A., Oliveira, G. M. S., Junior, L. R. M., Rosário, H. D. and Paranhos, L. R. (2016) 'Determination of vertical characteristics with different cephalometric measurements', *European Journal of Dentistry*, 10(1), pp. 116–120. DOI: 10.4103/1305-7456.175694.
- Ödman, A., Bresin, A. and Kiliaridis, S. (2019) 'The effect of retraining hypofunctional jaw muscles on the transverse skull dimensions of adult rats', *Acta Odontologica Scandinavica*. Taylor & Francis, 77(3), pp. 184–188. DOI: 10.1080/00016357.2018.1531437.
- Olmez, S. and Dogan, S. (2011) 'Comparison of the arch forms and dimensions in various malocclusions of the Turkish population', *Open Journal of Stomatology*, 01(04), pp. 158–164. DOI: 10.4236/ojst.2011.14023.
- Omar, H., Alhajrasi, M., Felemban, N. and Hassan, A. (2018) 'Dental arch dimensions, form and tooth size ratio among a Saudi sample', *Saudi Medical Journal*, 39(1), pp. 86–91. DOI: 10.15537/smj.2018.1.21035.
- Patel, D., Mehta, F., Patel, N., Mehta, N., Trivedi, I. and Mehta, A. (2015) 'Evaluation of arch width among Class I normal occlusion, Class II Division 1, Class II Division 2, and Class III malocclusion in Indian population', *Contemporary Clinical Dentistry*, 6(June), pp. S202–S209. DOI: 10.4103/0976-237X.166842.
- Prasad, M., Kannampallil, S. T., Talapaneni, A. K., George, S. A. and Shetty, S. K. (2013) 'Evaluation of arch width variations among different skeletal patterns in South Indian population', *Journal of Natural Science, Biology and Medicine*, 4(1), pp. 94–102. DOI: 10.4103/0976-9668.107267.
- Prates, L. da S., Gois, M., Berwig, L. C., Blanco-Dutra, A. P., Busanello-Stella, A. R. and Silva, A. M. T. da (2016) 'Avaliação clínica e eletromiográfica da mastigação nos diferentes padrões de crescimento facial', *Revista CEFAC*, 18(1), pp. 104–112. DOI: 10.1590/1982-021620161817015.
- Premkumar, S. (2011) *Textbook of Craniofacial Growth, Textbook of Craniofacial Growth*
- Premkumar, S. (2015) *Textbook of Orthodontics*. New Delhi: Elsevier
- Proffit, W. R., Fields, H. W., Larson, B. E. and Sarver, D. M. (2019) *CONTEMPORARY ORTHODONTICS*. 6th edn. Elsevier

- Quiudini, P. R., Pozza, D. H., Pinto, A. dos S., de Arruda, M. F. and Guimarães, A. S. (2017) ‘Differences in bite force between dolichofacial and brachyfacial individuals: Side of mastication, gender, weight and height’, *Journal of Prosthodontic Research*. Japan Prosthodontic Society, 61(3), pp. 283–289. DOI: 10.1016/j.jpor.2016.10.003.
- Raty Utari, T. and Kurnia Putri, M. (2019) ‘Orthodontic Treatment Needs in Adolescents Aged 13-15 Years Using Orthodontic Treatment Needs Indicators’, *Journal of Indonesian Dental Association*, 2(2), p. 49. DOI: 10.32793/jida.v2i2.402.
- Ravosa, M. J., Ning, J., Costley, D. B., Daniel, A. N., Stock, S. R. and Stack, M. S. (2010) ‘Masticatory biomechanics and masseter fiber-type plasticity’, *Journal of Musculoskeletal Neuronal Interactions*, 10(1), pp. 46–55
- Sanders, D. A., Rigali, P. H., Neace, W. P., Uribe, F. and Nanda, R. (2010) ‘Skeletal and dental asymmetries in Class II subdivision malocclusions using cone-beam computed tomography’, *American Journal of Orthodontics and Dentofacial Orthopedics*. American Association of Orthodontists, 138(5), pp. 542.e1-542.e20. DOI: 10.1016/j.ajodo.2010.02.027.
- Sandhu, S. S., Utreja, A., Prabhakar, S., Sandhu, N. and Kashyap, R. (2013) ‘A Study of Electromyographic Activity of Masseter and Temporalis Muscles and Maximum Bite Force in Patients with Various Malocclusions’, *Journal of Indian Orthodontic Society*, 47(2), pp. 53–61. DOI: 10.1177/0974909820130201.
- Shirai, M., Kawai, N., Hichijo, N., Watanabe, M., Mori, H., Mitsui, S. N., Yasue, A. and Tanaka, E. (2018) ‘Effects of gum chewing exercise on maximum bite force according to facial morphology’, *Clinical and Experimental Dental Research*, 4(2), pp. 48–51. DOI: 10.1002/cre2.102.
- Singh, G. (2015) *Textbook of Orthodontics*. 3rd edn. New Delhi: Jaypee Brothers Medical Publishers (P) Ltd
- Soyoye, O. A., Otuyemi, O. D., Kolawole, K. A. and Ayoola, O. O. (2018) ‘Relationship between masseter muscle thickness and maxillofacial morphology in pre-orthodontic treatment patients’, *International Orthodontics*. Elsevier Masson SAS, 16(4), pp. 698–711. DOI: 10.1016/j.ortho.2018.09.015.
- Spassov, A., Toro-Ibacache, V., Krautwald, M., Brinkmeier, H. and Kupczik, K. (2017) ‘Congenital muscle dystrophy and diet consistency affect mouse skull shape differently’, *Journal of Anatomy*, 231(5), pp. 736–748. DOI: 10.1111/joa.12664.
- van Spronsen, P. H. (2010) ‘Long-Face Craniofacial Morphology: Cause or Effect of Weak Masticatory Musculature?’, *Seminars in Orthodontics*. Elsevier

- Inc., 16(2), pp. 99–117. DOI: 10.1053/j.sodo.2010.02.001.
- Staley, R. N. and Reske, N. T. (2011) *Essentials of Orthodontics Diagnosis and Treatment*. 1st edn. Wiley-Blackwell
- Swetah, V. and Pandian, S. (2016) ‘Review Article Maxillary and Mandibular Arch Forms’, 40(19), pp. 80–82
- Toniolo, L., Cancellara, P., Maccatrazzo, L., Patruno, M., Mascarello, F. and Reggiani, C. (2008) ‘Masticatory myosin unveiled: First determination of contractile parameters of muscle fibers from carnivore jaw muscles’, *American Journal of Physiology - Cell Physiology*, 295(6), pp. 1535–1543. DOI: 10.1152/ajpcell.00093.2008.
- Yelampalli, M. R. and Rachala, M. R. (2012) ‘Timely management of developing class III malocclusion’, *Journal of Indian Society of Pedodontics and Preventive Dentistry*, 30(1), pp. 78–84. DOI: 10.4103/0970-4388.95590.
- Zhiyi, S., Min, G. and Yanqi, Y. (2018) ‘International Journal of Dentistry and Oral Science ( IJDOS ) ISSN : 2377-8075 The Association between Mastication , Malocclusion , and Craniofacial Morphology Malocclusion and Masticatory Function Long-term effect of Mastication on Malocclusion and Crani’, pp. 6–11