

# THE INDIAN VETERINARY JOURNAL

SINCE - 1924

**Journal of the  
INDIAN VETERINARY ASSOCIATION**  
ESTD - 1922      Regd. No. Sl. No. 96/1967 (CHENNAI)



No. 11, Pasumpon Muthuramalinga Thevar Salai (Chamiers Road),  
Nandanam, Chennai - 600 035, Tamil Nadu, India  
Tel. : +91 44 2435 1006  
Email : [ivj83@yahoo.com](mailto:ivj83@yahoo.com)  
ONLINE : [www.ivj.org.in](http://www.ivj.org.in)

**THE INDIAN VETERINARY JOURNAL**  
(Official Organ of the Indian Veterinary Association)

**EDITORIAL COMMITTEE**

**Dr A.V. KRISHNAN, Chief Editor**

B.V.Sc., M.V.Sc (Path.)

Retd. Additional Director of A.H., TN

**Dr S. SUKUMAR, Managing Editor**

B.V.Sc., M.V.Sc (Vet. Micro), Ph.D. (Biotech)

Retd. Professor of Biotechnology, TANUVAS, TN

**Dr V. Titus George, Editor**

B.V.Sc., M.V.Sc., Ph.D. (Patho)

Retd. Prof. of Pathology, TANUVAS

**Dr I. Ponnu Pandian, Editor**

B.V.Sc., PGDAEM

**Dr K. Venukopalan, Editor**

B.V.Sc., M.V.Sc., Ph.D. (Poul.)

Retd. Prof. of Poultry Science, TANUVAS

**EDITORIAL BOARD**

**CHAIRMAN**

**Dr. Chirantan Kadian**

B.V.Sc & A.H.

**President, Indian Veterinary Association**

**MEMBERS**

**Prof. Dr C. Balachandran**

M.V.Sc, Ph.D., PGDAJ, PGDEVP, DICVP

FAO Fellow, FIAVP, FNAVS, FASAW

Vice-Chancellor

Tamilnadu Veterinary and Animal Sciences University

Madhavaram Milk Colony, Chennai - 600 051.

**Dr. P.K. Panwar**

BVSc & AH, MVSc,

Department of Animal Husbandry

Uttar Pradesh

**Dr. Subhash Chander Ahlawat**

BVSc & AH.

Department of Animal Husbandry

Haryana

**MANAGING COMMITTEE OF  
INDIAN VETERINARY ASSOCIATION**

**President :**

Dr. Chirantan Kadian

**Secretary General :**

Dr. R.S. Patel

**Treasurer :**

Dr. I. Ponnu Pandian

**Joint Secretary :**

Dr. M.K. Srikanth

**Vice Presidents :**

Dr. L. Chandrasekaran, HQ, Tamilnadu

Dr. Arun Sirkeck, Himachal Pradesh

Dr. S.G. Yalagod, Karnataka

Dr. Ilang, Nagaland

Dr. Ramji Lal Meena, Rajasthan

Dr. Sudhir Kumar, Uttar Pradesh

**Zonal Secretaries :**

Dr. Amit Nain, Punjab

Dr. R. Vasantha Rayalu, Andhra Pradesh

Dr. Bhuban Ch. Sarmah, Assam

Dr. Umesh Balaji Sontakke, Maharashtra

Dr. Pankaj Kumar, Jharkhand

**Managing Editor, IVJ :**

Dr. S. Sukumar, Tamil Nadu

# THE INDIAN VETERINARY JOURNAL

(Official organ of the Indian Veterinary Association)

Vol. 96

October 2019

No. 10

## CONTENTS

### GENERAL ARTICLES :

**Performance of “*Tanuvus Namakkal Gold Japanese Quail*” Fed Diet with Different Levels of Energy and Lysine**

K. Shibi Thomas, R. Amutha, M.R. Purushothaman, P.N. Richard Jagatheesan and S. Ezhil Valavan ... 09

**Supplementation of *Phaleria macrocarpa* Leaf Extract Significantly Reduced *E.Coli* Infection in Broiler**

Diyantoro and Siti Eliana Rochmi ... 12

**Drug-Drug Interactions and Prescribing Pattern in Veterinary Practice in Iraq**

Sarhan R. Sarhan, Hayfaa A. Moheisen, Elaph K. Kamel and Zahraa J. Mohammad ... 14

**A Case of Cranioschisis with Meningocele in a Large White Yorkshire Piglet**

E.Rachel Jemimah, S. Meenakshi Sundaram, A. Shanmuga sundaram, T. Muthuramalingam and R. Venkataramanan ... 18

**Antisperm Antibody in Repeat Breeder Friesian Holstein Cows at KPSP Setia Kawan Nongkojajar, Tutur District, Pasuruan, Indonesia**

FadilaZikri Amanda, Sri PantjaMadyawati, IsnainiFadilah and Erma Safitri ... 20

**Economics of Inclusion of Turmeric (*Curcuma longa*) and Ginger (*Zingiber officinale*) in Broiler Feeds**

S.A. Jadhav, S.M. Bhalerao, A.V. Khanvilkar, S.D. Changan and L.A. Pangaonkar ... 23

**Molecular Identification of Extended Spectrum Beta-Lactamase (ESBL) Producing**

***Escherichia coli* Isolated From Dairy Cows in East Java Province, Indonesia**

Akyun Rozaqi Syah Putra, Mustofa Helmi Effendi, Setiawan Koesdarto and Wiwiek Tyasningsih ... 26

**Acupuncture Could Increase Spermatogonic Cells in Albino Rats Exposed to Heat Stroke**

Muhammad Thohawi Elziyad Purnama, Imas Hapsari Rahmaningtyas, Ainun Septia Putri, Angela Swasti Ivana Lee, Fachrun Nisa Tatimma and Herlina Masyitoh ... 30

***Phaleria macrocarpa* (Scheff.) Boerl. Pulp Extract Increases the Sperm Characteristics in *Rattus norvegicus***

Amaq Fadholly, Arif N. M. Ansori, Annise Proboningrat, Muhammad K. J. Kusala, Naimah Putri, Soeharsono and Budi Utomo ... 32

**Red Dragon Fruit Peel (*Hylocereus polyrhizus*) in Rabbit Nutrition**

Ragil Angga Prastiya, Moh. Anam Al-Arif, Fachrun Nisa' Tatimma and Bodhi Agustono ... 36

**Eyelid Occlusion induced Form Deprivation Myopia (FDM) on Axial Length and Morphological Changes of the Sclera in Rabbits**

Rini Kusumawardhany, GatutSuhendro, Tjahjono D. Gondhowiardjo, Nurwasis, AtinaYustisia Lestari and LitaRakhmaYustinasari ... 38

**RUNX2 and SOX9 Expression on Chondrocyte Hypertrophy Formation in Post-Menopausal Osteoarthritis Mechanism (An Experimentation on Rat Model)**

Bimo Sasono, Heri Suroto, Dwikora N. Utomo, Aulani Am, Hari Basuki N, Damayanti Tinduh and Fedik A. Rantam ... 41

**Concentration of Microcystin in an Intensive Vannamei Farm at Banyuwangi, East Java Indonesia**

EndangDewi Masithah, Fitri Annisha and Agoes Soegianto ... 45

## CLINICAL AND FIELD ARTICLES :

<b>The Effect of Feeding High Level of Protein on Reproductive Performance of Bali Starling</b> Mas'ud Hariadi, Budi Utomo, Herry A. Hermadi, Rezha S.W. Hadi, Alfian Zulfahmi and Erma Safitri	... 48
<b>Management Very Large Urolith in Pomerian Dog - A Case Report</b> E. Erwin, E. Etriwati and D. Noviana	... 51
<b>Fusion Protein of Aminoacid Mutations in Newcastle Disease Isolated from Swan Goose Caused Resistance to Infection</b> Naimah Putri, Rahaju Ernawati, Suwarno, Jola Rahmahani and Fedik Abdul Rantam	... 53
<b>Post Coital Paraphimosis in a Chippiparai Dog</b> S.Kokila, M.Bharathidasan, A.Ganesan, S.Dharmaceelan, A.R.Ninu and D.Vishnugurubaran	... 56
<b>Prevalence of Ticks in Sheep and Goats in Perambalur District of Tamilnadu</b> P. Suresh Kumar	... 58
<b>The Antioxidant Activity of Pinus merkusii Ethanol Extract as Hepatoprotector in Mice</b> A.Herdiansyah, Rimayanti and S.A. Sudjarwo	... 60
<b>Medical Management of Cellulitis of Limb in a Mule</b> Aneesh. A and Nitin Sheoran	... 62
<b>Haemorrhagic Colitis Due to <i>Buxtonella Sulcata</i> and <i>E.Coli</i> in a Cross Bred Cow</b> S. Saravanan, T.Anna, K.K. Ponnu Swamy, P. A. Enbavelan, R.C. Sundararajan and R. Ramprabhu	... 65
<b>Surgical Management of Peribulbar Abscess in Indian Star Tortoise (<i>Geocheloneelegans</i>) – A Case Report</b> Vinny Anna Varghese, Shashi KantMahajan, VandanaSangwan and Kuldip Gupta	... 67
<b>Effective Clearance of Vermic Toxins by Haemodialyzer with Small Surface Area</b> M.Chandrasekar, M.Arthi, M.G.Jayathangaraj, Stella Esther and S.Nethaji	... 69
<b>Surgical Management of Uterine Rupture in a Bitch</b> G. Shanmuga Priya and M.G. Mohamed Ali	... 71
<b>Cestode Infection in a Free Ranging Indian Rock Python (<i>Python molurus</i>)</b> M.S. Sreelakshmi, Mammen J. Abraham, Ajith Jacob George and N.Divakaran Nair	... 73
<b>Management of Atopic Dermatitis in Dog</b> Steven Taufic Leo and LitaRakhmaYustinasari	... 74
<b>Intussusception and its Surgical Management Under General Anaestheisa in a Cow – A Case Report</b> S.Vigneshwaran, S. Senthilkumar, S. Dharmaceelan, S.Kathirvel, A. Kumaresan and K.Jayakumar	... 76
<b>Meloxicam Toxicity in Labrador Dog Due to Dispensing Error and its Reversal by Misoprostol</b> K. Kannan, M. Saravanan, P.K. Ram Kumar, T. Arul Kumar, S. Senthil Kumar and N. Premalatha	... 78
<b>Cost Effectiveness of Different Diet Fed to “Tanuvas Namakkal Gold Japanese Quail”</b> K. Shibi Thomas, R. Amutha, M.R. Purushothaman, P.N. Richard Jagatheesan and S. Ezhil Valavan	... 81
<b><i>Hymenolepis lanceolata</i> in a Duck from Cauvery Delta Region</b> M.K. Vijayarathi and A. Meenakshisundaram	... 84
<b>Enhancement of Biohydrogen Production from Cattle Rumen Fluid by Optimization of Micronutrients</b> Merli Maman, S. Meignanalakshmi, K.Vijayarani, H. Gopi and S. Meenakshi Sundaram	... 85

## Author and Subject Index

89 & 90

## RUNX2 and SOX9 Expression on Chondrocyte Hypertrophy Formation in Post-Menopausal Osteoarthritis Mechanism (An Experimentation on Rat Model)

Bimo Sasono, Heri Suroto, Dwikora N. Utomo, Aulani Am, Hari Basuki N, Damayanti Tinduh and Fedik A. Rantam<sup>1</sup>

Stem Cells Research and Development Center, Laboratory of Virology and Immunology, Microbiology Department, Faculty of Veterinary Medicine Airlangga University, Indonesia.

(Received : January, 2019 04/19 Accepted : May, 2019)

### Abstract

In osteoarthritis post-menopause, decreasing estrogen influenced formation of chondrocyte hypertrophy. This study would explain role of RUNX2 and SOX9 to chondrocyte hypertrophy formation in decreasing estrogen condition. Fifty four rats were divided into six groups. Level of 17  $\beta$  estradiol and TGF- $\beta$  were examined by ELISA. SOX9 and RUNX2 used immunohistochemistry. Histopathology for chondrocyte hypertrophy. Level of 17  $\beta$  estradiol and SOX9 on treatment group decreased significantly, TGF- $\beta$  and RUNX2 increased significantly. Chondrocyte hypertrophy increased significantly in treatment group. The decreasing of 17  $\beta$  estradiol level caused increasing hypertrophy Chondrocyte through increasing level of TGF- $\beta$  and RUNX2.

**Key words:** Chondrocyte hypertrophy, Estrogen, RUNX2, TGF- $\beta$ .

Osteoarthritis is a degenerative disease that causes joint cartilage damages and joint movement inhibition. The formation of chondrocyte hypertrophy indicates "Point of No Return" leading to osteoarthritis. The impact of estrogen towards Type-2 collagen remains unclear. Chondroprogenitor cells found in the cartilage may become a factor that explains the effect of decreasing estrogen on the level of Type-2 collagen and answers the question why chondrocyte hypertrophy happens on osteoarthritis post menopause (Goldring *et al.*, 2012).

The prevalence of osteoarthritis increases on post-menopausal women. Decreasing estrogen

level causes changes on micro-environment, namely the increasing TGF- $\beta$  level that triggers differentiation on chondroprogenitor cells into chondrocyte hypertrophy by increasing of RUNX2 (Jayasuria *et al.*, 2016), which initiates cellular changes towards osteoarthritis. The increasing number of chondrocyte hypertrophy enhances the level of MMP-13 so that the level of Type-2 collagen will decrease in cartilage (Roman Blast, *et al.*, 2009). Role of RUNX2 and SOX9 on decreasing estrogen level and the number of Hypertrophic Chondrocyte, and the effects on TGF- $\beta$  level on chondroprogenitor cells and their hypertrophy is to be studied (Cheng and Genever., 2010).

### Materials and Methods

The rats (*Rattus norvegicus*) from Veterinary Medical Faculty of Gajah Mada University used for the estimation. Monoclonal antibody for RUNX2 using Mouse anti-Rat RUNX2 Monoclonal Antibody (Santacruz Biotechnology, USA). Monoclonal antibody for SOX9 was estimated using Mouse anti-Rat SOX9 Monoclonal Antibody (LSBIO, USA), MMP-13 was estimated by Rabbit anti-Rat MMP-13 Polyclonal Antibody (BIOSS, USA). Collagen type 2 was estimated using Rabbit anti-Rat Collagen type 2 Polyclonal Antibody (BIOSS, USA). Levels of 17  $\beta$  estradiol were estimated using Rat 17  $\beta$  Estradiol Elisa Kit (Bioassay Technology Laboratory Shanghai, CHINA) and TGF- $\beta$  estimated using Rat Transforming Growth Factor  $\beta$  Elisa Kit (Bioassay Technology Laboratory Shanghai, CHINA).

Blood samples of 2cc were collected with EDTA in the control group (K) and treatment

<sup>1</sup>Corresponding author : Email : fedik-a-r@fkh.unair.ac.id

group (P), and centrifuged at 3000 rpm/minute for 10 minutes to separate blood serum for estimation of 17  $\beta$  estradiol and TGF- $\beta$  level, at department of Science and Technology.

54 rats were divided into three groups (K1 to K3) and three treatment groups (P1 to P3) consisting of 9 rats each. Cartilage samples were taken from knee joint and calcified in 2% nitric acid for 3 weeks consecutively and they were examined at 4, 8 and 12<sup>th</sup> weeks.

Sandwich ELISA test was conducted to measure the levels of 17 $\beta$  estradiol the preparation of all reagents, standard solution (50  $\mu$ L contains 6 ng/L), and standard samples by considering the room temperature; 40  $\mu$ L standard solution was mixed by 10  $\mu$ L anti-estradiol antibody solution and added into 50  $\mu$ L streptavidin HRP. The solution was incubated at 37°C for an hour. Plates were cleaned in 350  $\mu$ L wash solution for five times. 50  $\mu$ L solution A and 50  $\mu$ L solution B were mixed and incubated at 37°C for 10 minutes until its color changed and then 50  $\mu$ L stop solution was added. OD value was read using microplate reader set at  $\lambda$  450 nm and measured. TGF examination was conducted based on the following procedures: preparing the preparations, samples, and standard solution (120  $\mu$ L concentration 2400 ng/L). 50  $\mu$ L aquadest was added into 50  $\mu$ L standard solution until the concentration reached 75 ng/L. 50  $\mu$ L streptavidin-HRP was added into 50  $\mu$ L standard solution (composed of 40  $\mu$ L sample and 10  $\mu$ L TGF- $\beta$  antibody) and incubated at 37 °C for an hour. Plates were cleansed in 350  $\mu$ L buffer solution. 50  $\mu$ L substrate A and 50  $\mu$ L substrate B were added and incubated at 37 °C in dark room condition. 50  $\mu$ L stop solution was added until the color changed from blue to yellow. OD value was measured using microplate recorder set at  $\lambda$  450 nm 30 minutes after stop solution added.

Immunohistochemistry examination comprised of dehydration phase, clearing phase, and embedding phase. Dehydration phase is reducing water molecules by using alcohol with concentrations of 70%, 80%, 90%, and finally 100% (absolute) for an hour consecutively. Clearing phase refers to cleaning the alcohol and chemical traces using xylol 98 % for an

hour with three repetitions. Embedding phase is the process of creating tissue block by dipping cartilage tissue samples into paraffin to create paraffin block. Tissue samples were stored at 56-58°C temperature for 2 hours with three repetitions. The tissues in paraffin block, were cut using microtom with 4  $\mu$ m thickness. The slice were put into water bath at 50°C temperature and stuck to the object glass. The preparations were heated to thaw the paraffin before coloration. Immunohistochemistry coloration technique applied in this experiment was biotin streptavidin technique based on the following procedures: first Betazoid Statrec Universal staining kit (SCYTECH, CANADA) 1:100 in substrate for 3 minutes then the object glass was rinsed in aquadest, followed by Hematoxylin Meyers coloration for 30 second and the object glass was rinsed in water. Dehydration process was conducted by using ethanol 50 %, 70%, 80%, 90%, and 100% followed by xylol for 3 x 3 minutes. Then, mounting was conducted to close the tissue with glass decks and glued. Using the microscope, expression of RUNX2 and SOX9 were examined (Nikon H600L, Digital camera DS Fi2,300 megapixel)

Cartilage tissue block paraffins were made and then cut with a microtom with a thickness of 4  $\mu$ m, then stained using hematoxylin and eosin stain. Preparations were read using a Nikon (Eclipse Ci microscope, 16.25megapixel Camera DS-Ri2) to see changes in the number of chondrocyte hypertrophy with 400x magnification.

The data was analyzed statistically by Anova test. The levels of SOX9 and RUNX2 were analyzed with Brown Forsythe using SPSS 23 for windows with significance level at ( $p < 0,05$ ).

## Results and Discussion

This study showed the level of 17  $\beta$  estradiol significantly decreased on the treatment group compared with the control group, especially on P3 (12-week treatment) caused by ovariectomy in 4 weeks (Table I). 17  $\beta$  estradiol was mainly produced in the ovaries so ovariectomy treatment would reduce estradiol level and cause similar symptoms as menopause.

The level of TGF- $\beta$  significantly increased

**Table I.** Levels of 17 estradiol, TGF- $\beta$ , SOX9, RUNX2 on different groups

	17 estradiol	TGF-	SOX9	RUNX2
K1	9.888 $\pm$ 1.6	143.3 $\pm$ 56.21	5.28 $\pm$ 3.5	8.25 $\pm$ 1.74
K2	8.441 $\pm$ 0.8	235.8 $\pm$ 81.24	3.84 $\pm$ 2	11.01 $\pm$ 1.96
K3	8.125 $\pm$ 1.39	339.5 $\pm$ 87.61	4.55 $\pm$ 2.2	11.79 $\pm$ 1.77
P1	7.966 $\pm$ 1.1	339.8 $\pm$ 46.88	2.31 $\pm$ 1.6	12.03 $\pm$ 2.75
P2	7.201 $\pm$ 1.16	382.8 $\pm$ 90.64	1.9 $\pm$ 1.1	14.19 $\pm$ 2.31
P3	6.848 $\pm$ 0.92	394.7 $\pm$ 75.6	1.4 $\pm$ 1.1	16.27 $\pm$ 3.89

**Table II.** Chondrocyte Hypertrophy Number

Group	n	Mean $\pm$ SD
K1	9	17.78 $\pm$ 3.99 <sup>ab</sup>
K2	9	14.00 $\pm$ 5.15 <sup>a</sup>
K3	9	18.33 $\pm$ 6.86 <sup>ab</sup>
P1	9	27.22 $\pm$ 7.71 <sup>c</sup>
P2	8	25.75 $\pm$ 11.68 <sup>bc</sup>
P3	8	41.50 $\pm$ 10.89 <sup>d</sup>

\*Different superscripts in the same column indicated a significant difference ( $p < 0.05$ ).

on the treatment groups (P) compared to control group (K) and seen in the 4th, 8th, and 12th week. Matsuda, *et al.*, (2001) Found that Estrogen level correlated with TGF- $\beta$  level on human cells. The study showed there was an increase of TGF- $\beta$  level caused by the decreasing level of estradiol. There was cross-reaction between estradiol and TGF- $\beta$ .

Ito, *et al.*, (2009) found that estrogen receptor  $\alpha$  affected TGF- $\beta$  by inhibiting Smad proteins of cytoplasm. Cheng, *et al.*, (2016) found the effect of estrogen on cells through estrogen receptor  $\beta$  that inhibited type-2 collagen and sclerostin expression, but he didn't explain how did estrogen affect the collagen.

Table I shows the concentration of estradiol was decreasing significantly (Kruskal-Wallis test) on treatment group compared control group ( $p < 0.05$ ). The level of TGF- $\beta$  in blood was increasing significantly on the treatment group (P1, P2, P3) ( $p \leq 0.05$ ). This result indicates that the decreasing estradiol level increase the level of blood TGF- $\beta$ . Level of SOX9 decreased significantly on the treatment group (P1, P2, P3) compared to the control group ( $p < 0.05$ ) and level of RUNX2 increased significantly in treatment

group compared to control group ( $p < 0.05$ ). This result shows that decreasing level of estrogen was increasing TGF- $\beta$  and RUNX2.

Table II shows the number of chondrocyte hypertrophy cells significantly ( $p < 0.05$ ) increased in the treatment groups compared to the control groups.

The effect of increasing TGF- $\beta$  on chondroprogenitor cells is SOX9 and RUNX2 expression on the nuclei. RUNX2 and SOX9 are competing one another so that the differentiation depends on which factor is more dominant. If RUNX2 is dominant, the differentiation would be oriented to osteogenesis. But, if SOX9 is dominant, the differentiation would be orientated to chondrogenesis. The increasing of chondrocyte hypertrophy cells is commonly caused by RUNX2 expression. As an intracrine, RUNX2 stimulates differentiation towards chondrocyte hypertrophy. RUNX2 competes with SOX9 which functions as RUNX2 inhibiting factor (Cheng and Genever, *loc. cit*). In this research, the number of chondrocyte hypertrophy cells increased when RUNX2 level was increasing and SOX9 level was decreasing. It showed that the expression of RUNX2 influenced differentia-

tion of chondroprogenitor cell into hypertrophic chondrocyte.

The increasing TGF- $\beta$  level in blood may affect various extracellular matrix conditions and stimulate chondroprogenitor cells in cartilage. The effect of TGF- $\beta$  on chondroprogenitor cells would change micro-environment caused by decreasing estrogen level then stimulating differentiation towards chondrocyte hypertrophy (Dexheimer *et al.*, 2016). Chondrocyte hypertrophy expresses several proteins, namely Alkali Phosphatase (ALP) Venous Endothelial Growth Factor (VEGF), Osteopontin, Matrix Metalloprotease 13 (MMP-13), and type-10 collagen. The role of MMP-13 as collagenase enzyme is clearly proven to degrading type-2 collagen. Degradation of type-2 collagen in cartilage caused by processes above will reduce the strength of cartilage because type 2 collagen is important to the extracellular matrix (ECM). Many opinions mentioned that mechanical, inflammation, pathological apoptosis and aging process might be the factors of hypertrophic chondrocytes formation. However, estrogen, as one of osteoarthritis factors, needs to be accounted considering the increased prevalence of osteoarthritis in post-menopausal period.

### Summary

It can be concluded that decreasing level of 17  $\beta$  estradiol caused increasing amount of chondrocyte hypertrophy through increasing level of TGF- $\beta$  and RUNX2.

### Acknowledgement

We would like to thank the Rector of Universitas Airlangga and Dean of Veterinary Medicine Faculty, Universitas Airlangga.

### References

- Cheng A., and Genever P.G. (2010) Sox determines Runx2 transactivity by directing intracellular degradation. *J. of Bone and Mineral Res.*, **25**(12) : 1680-1689.
- Cheng Q., Tang W., Sheu T.J., Gan J., Li H., and Zhang X. (2016) Circulating TGF- $\beta$  1 levels are negatively correlated with sclerostin levels in postmenopausal women. *Clinica Chimica Acta*, **455** : 87-92.
- Dexheimer V., Gabbler J., Boman K., Sims T., and Omlor G. (2016) Differential expression of TGF- $\beta$  superfamily members and role of versus endochondral chondrocyte differentiation. Nature Publishing Group, June, pp 1-14.
- Goldring MB. (2012) Chondrogenesis, chondrocyte differentiation, and articular cartilage metabolism in health and osteoarthritis p269–85.
- Ito I, Hanyu A, Wayama M, Goto N, Katsuno Y, and Kawasaki S, (2009). Estrogen inhibits transforming growth factor signaling by promoting Smad2/3 degradation. *J Biol Chem*. **285**(19): 14747–55.
- Jayasuriya CT, Chen Y, Liu W, and Chen Q. (2016) The influence of tissue microenvironment on stem cell-based cartilage repair. *Ann N Y Acad Sci*. **1383**(1) : 21–33.
- Matsuda T, Yamamoto T, Muraguchi A, and Saatcioglu F. (2001) Cross-talk between Transforming Growth Factor and Estrogen Receptor Signaling through Smad3. *J Biol Chem*; **24**(5): 42908–42914.
- Roman-Blas JA, Castañeda S, Largo R, and Herrero-Beaumont G. (2009) Osteoarthritis associated with estrogen deficiency. *Arthritis Res Ther*, **11**(5): 241.