

DAFTAR PUSTAKA

- A, Sentkowska., K, Pyrzynska. 2017. *Flavonoid Content and Antioxidant Properties of Different Black Tea Infusions. Journal of Nutrition and Health Sciences* Volume 4.
- Arief, D. Z., Rohdiana, D., & Somantri, M. (2012). **Analisis Polifenol Total dan Aktivitas Penangkapan Radikal Bebas DPPH (1,1-Diphlnyl, 2-Picrylhidrazl) Teh Putih (*Camellia sinensis* LO Kuntze) Berdasarkan Suhu dan Lama Penyeduhannya.** *Artikel Penelitian.*
- Astawan, M., 2008. *Khasiat warna-warni makanan.* Gramedia Pustaka Utama.
- Badan, P.O.M. and MODAL, P., 2000. Republik Indonesia. *Cara Uji Klinik yang Baik.* Jakarta: Badan POM.
- Budavari, S. 1996. *Merck Index Whitehouse Station.* NJ Merck.
- Badan Standardisasi Nasional. 2016. **SNI 1902-2016 Teh Hitam.** Jakarta. 17 hal.
- Chang, M, Y., Lin Y, Y ., Chang, Y, C., Huang, W, Y., Lin, W, S., Chen, C, Y., Huang, S, L., Lin, Y, S. 2020. *Effects of Infusion and Storage on Antioxidant Activity and Total Phenolic Content of Black Tea. Journal Of Applied Sciences Volume* 10, p. 2685.
- Crozier, A., Clifford, M.N. and Del Rio, D., 2012. *Bioavailability of dietary monomeric and polymeric flavan-3-ols. Bioavailability and function of flavonoids: oxidative stress and disease, 30,* pp.45-78.
- Davies, H.S., Pudney, P.D., Georgiades, P., Waigh, T.A., Hodson, N.W., Ridley, C.E., Blanch, E.W. and Thornton, D.J., 2014. *Reorganisation of the salivary mucin network by dietary components: insights from green tea polyphenols. PloS one, 9(9),* p.e108372.
- Demir, E., Serdar, G. and Sökmen, M., 2015. *Comparison of some extraction methods for isolation of catechins and caffeine from Turkish green tea. International Journal of Secondary Metabolite, 2(2),* pp.16-25.
- Fadda, A., Serra, M., Molinu, M.G., Azara, E., Barberis, A. and Sanna, D., 2014. *Reaction time and DPPH concentration influence antioxidant activity and kinetic parameters of bioactive molecules and plant extracts in the reaction with the DPPH radical. Journal of Food composition and analysis, 35(2),* pp.112-119.

- Farooq, S. and Sehgal, A., 2018. *Antioxidant activity of different forms of green tea: loose leaf, bagged and matcha*. *Current Research in Nutrition and Food Science Journal*, 6(1), pp.35-40.
- Gandjar, I.G. and Rohman, A., 2012. **Analisis obat secara spektrofotometri dan kromatografi**. *Pustaka Pelajar, Yogyakarta*, pp.71-75.
- Graham, H. N. 1992. *Green tea composition, consumption, and polyphenol chemistry*. *Preventive medicine*, 21(3), 334-350.
- Handa, S.S., 2008. *An overview of extraction techniques for medicinal and aromatic plants*. *Extraction technologies for medicinal and aromatic plants*, 1.
- Hamilton-Miller, J.M.T., 2001. *Anti-cariogenic properties of tea (Camellia sinensis)*. *Journal of medical microbiology*, 50(4), pp.299-302.
- Hara, Y. 2001. *Methods of Extracting Polyphenolic Constituents of Tea Leaves*. In *green tea: Health Benefits and Applications* (pp. 20–23). New York: Marcel Dekker.
- Harmita, H., 2012. **Petunjuk pelaksanaan validasi metode dan Cara Perhitungannya**. *Pharmaceutical Sciences and Research (PSR)*, 1(3), pp.117-135.
- Huerta, C., Lanes, S.F. and Rodríguez, L.A.G., 2005. *Respiratory medications and the risk of cardiac arrhythmias*. *Epidemiology*, pp.360-366.
- Hartoyo, A., 2003. **Teh dan khasiatnya bagi kesehatan**. *Kanisius, Yogyakarta*.
- Hilal, Y. and Engelhardt, U., 2007. *Characterisation of white tea—Comparison to green and black tea*. *Journal für Verbraucherschutz und Lebensmittelsicherheit*, 2(4), pp.414-421.
- Indarti, D., 2015. **Outlook teh komoditas pertanian subsektor perkebunan**. Pusat Data dan Sistem Informasi Pertanian, p.14.
- Indonesia, D.K.R., 2014. **Farmakope Indonesia Edisi V**. *Jakarta: Departemen Kesehatan Republik Indonesia*.
- Jang, S. I., Jun, M. H., Lillehoj, H. S., Dalloul, R. A., Kong, I. K., Kim, S., & Min, W. 2007. *Anticoccidial effect of green tea-based diets against Eimeria maxima*. *Veterinary Parasitology*, 144(1-2), 172-175.
- Jigisha, A., Nishant, R., Navin, K., & Pankaj, G. 2012. *Green tea: a magical herb with miraculous outcomes*. *Int. Res. J. Pharm*, 3(5), 139-148.
- Juneja, L. R., Kapoor, M. P., Okubo, T., & Rao, T. 2013. *Green tea polyphenols: nutraceuticals of modern life*. CRC Press.

- Khalaf, N. A., Ashok, K., Shakya, A. K., Al-Othman, A., El-Agbar, Z., Farah, H. 2008. *Antioxidant Activity of Some Common Plants. Journal Of Turkey Biology* Volume 32 p. 51-55.
- Khan, N. and Mukhtar, H., 2013. *Tea and health: studies in humans. Current pharmaceutical design*, 19(34), pp.6141-6147.
- Kedare, S.B. and Singh, R.P., 2011. *Genesis and development of DPPH method of antioxidant assay. Journal of food science and technology*, 48(4), pp.412-422.
- Keith, L., Ko, W. H., and Sato, D. M. 2006. *Identification guide for diseases of tea (Camellia sinensis)*.
- Ku, C.S., Jang, J.P. and Mun, S.P., 2007. *Exploitation of polyphenol-rich pine barks for potent antioxidant activity. Journal of Wood Science*, 53(6), pp.524-528.
- Lee, C.H., Yang, L., Xu, J.Z., Yeung, S.Y.V., Huang, Y. and Chen, Z.Y., 2005. *Relative antioxidant activity of soybean isoflavones and their glycosides. Food Chemistry*, 90(4), pp.735-741.
- Leung, L. K., Su, Y., Chen, R., Zhang, Z., Huang, Y., & Chen, Z. Y. (2001). Theaflavins in black tea and catechins in green tea are equally effective antioxidants. *The Journal of nutrition*, 131(9), 2248-2251.
- Lung, J.K.S. and Destiani, D.P., 2017. **Uji Aktivitas Antioksidan Vitamin A, C, E dengan Metode DPPH.** *Farmaka*, 15(1), pp.53-62.
- Lutterodt, H., Luther, M., Slavin, M., Yin, J.J., Parry, J., Gao, J.M. and Yu, L.L., 2010. *Fatty acid profile, thymoquinone content, oxidative stability, and antioxidant properties of cold-pressed black cumin seed oils. LWT-Food Science and Technology*, 43(9), pp.1409-1413.
- Maesaroh, K., Kurnia, D., & Al Anshori, J. 2018. **Perbandingan Metode Uji Aktivitas Antioksidan DPPH, FRAP dan FIC Terhadap Asam Askorbat, Asam Galat dan Kuersetin.** *Chimica et Natura Acta*, 6(2), 93-100.
- Maruyama, K., Kihara-Negishi, F., Ohkura, N., Nakamura, Y., Nasui, M. and Saito, M., 2017. *Simultaneous determination of catechins and caffeine in green tea-based beverages and foods for specified health uses. Food and Nutrition Sciences*, 8(03), p.316.
- Martono, B., Falah, S. and Nurlaela, E., 2016. *Antioxidant Activities of GMB 7 Variety of Tea at Different Altitude.*
- Martono, Y., Riyanto, S., Rohman, A. and Martono, S., 2016, July. *Improvement method of fast and isocratic RP-HPLC analysis of major diterpene glycoside*

- from Stevia rebaudiana leaves*. In *AIP Conference Proceedings* (Vol. 1755, No. 1, p. 080001). AIP Publishing.
- Marwanto. 2014. **Rekayasa alat pemereras air jeruk siam dengan sistem ulir**. Sambas: Poltesa.
- Milašiene, R., Rawicka, K., Kornyšova, O., Ligor, M., Maruška, A. and Buszewska, B., 2007. *Evaluation of antioxidactivity of green and black tea (Camellia sinensis) and rooibos (Aspalathus linearis) tea extracts by means of hplc with reaction detector*. *Ars Separatoria Acta*, pp.27-33.
- Mishra, K., Ojha, H., & Chaudhury, N. K. 2012. *Estimation of antiradical properties of antioxidants using DPPH Å assay: A critical review and results*. *Food Chemistry*, 130(4), 1036–1043. <https://doi.org/10.1016/j.foodchem.2011.07.127>.
- Miura, Y., Chiba, T., Tomita, I., Koizumi, H., Miura, S., Umegaki, K., ... & Ikeda, M. 2001. *Tea catechins prevent the development of atherosclerosis in apoprotein E-deficient mice*. *The Journal of nutrition*, 131(1), 27-32.
- Molyneux, P. 2004. *The Use of the Stable Free Radical Diphenylpicryl-hydrazyl (DPPH) for Estimating Antioxidant Activity*. *Songklanakarinn Journal of Science and Technology*, 26(December 2003), 211–219. <https://doi.org/10.1287/isre.6.2.144>
- Mulja, M., Suharman, 1995. **Analisis instrumental**, 188.
- Nugraha, A., Sumarwan, U. and Simanjuntak, M., 2017. **Faktor Determinan Preferensi dan Perilaku**
- Pękal, A., Droźdz, P., Biesaga, M., Pyrzynska, K. 2011. *Evaluation of the antioxidant properties of fruit and flavoured black teas*. *The Journal Of European Nutrition* Volume 50 p.681–688.
- Pękal, A., Biesaga, M. and Pyrzynska, K., 2011. *Interaction of quercetin with copper ions: complexation, oxidation and reactivity towards radicals*. *Biometals*, 24(1), pp.41-49.
- Rohdiana, D., 2015. **Teh: proses, karakteristik dan komponen fungsionalnya**. *Food Review Indonesia*, 10(8), pp.34-37.
- Sahala, A. and Soegihardjo, C.J., 2012. **Uji Aktivitas Antioksidan Dan Penetapan Kadar Fenolat Total Fraksi Air Daun Ketapang (Terminalia Catappa L.) Dengan Metode Dpph (2,2-diphenyl-1-picrylhydrazyl) Dan Metode Folin-ciocalteu**. *Jurnal Farmasi Sains dan Komunitas (Journal of Pharmaceutical Sciences and Community)*, 9(2).

- Sánchez-Moreno, C., 2002. *Methods used to evaluate the free radical scavenging activity in foods and biological systems*. *Food science and technology international*, 8(3), pp.121-137.
- Satiadarma, K., Mulja, M., Tjahjono, D.H. and Kartasasmita, R.E., 2004. **Asas pengembangan prosedur analisis**. *Cetakan Pertama*. Surabaya: Airlangga University Press. Hal, 46(49), pp.89-90.
- Setyamidjaja, D. 2000. **Teh Budidaya & Pengolahan Pascapanen**. Kanisius.
- Shah, S., Gani, A., Ahmad, M., Shah, A., Gani, A., Masoodi, F, A. 2015. *In vitro antioxidant and antiproliferative activity of microwave-extracted green tea and black tea (Camellia sinensis): a comparative study*. *Journal Of Nutrafoods*.
- Shankar, S., Ganapathy, S., and Srivastava, R. K. 2007. *Green tea polyphenols: biology and therapeutic implications in cancer*. *Front Biosci*, 12(4881), 99.
- Sharangi, A. B. 2009. *Medicinal and therapeutic potentialities of tea (Camellia sinensis L.)—A review*. *Food Research International*, 42(5-6), 529-535.
- Shekhar, T.C. and Anju, G., 2014. *Antioxidant activity by DPPH radical scavenging method of Ageratum conyzoides Linn. leaves*. *Am J Ethnomed*, 1(4), pp.244-249.
- Shimamura, T., Sumikura, Y., Yamazaki, T., Tada, A., Kashiwagi, T., Ishikawa, H., ... Ukeda, H. 2014. *Applicability of the DPPH Assay for Evaluating the Antioxidant Capacity of Food Additives – Inter-laboratory Evaluation Study –*. *Analytical Sciences*, 30, 717–721.
- Skoog DA, Holler FJ, Nieman TA. 2007. *Principles of Instrumental Analysis 6 th Edition*. Philadelphia (US): Harcourt Brace College Publishers.
- Song, Y. A., Park, Y. L., Kim, K. Y., Chung, C. Y., Lee, G. H., Cho, D. H., ... & Kim, N. 2011. *Black tea extract prevents lipopolysaccharide-induced NF- κ B signaling and attenuates dextran sulfate sodium-induced experimental colitis*. *BMC complementary and alternative medicine*, 11(1), 91.
- Srinivasan, P., Sabitha, K. E., & Shyamaladevi, C. S. 2007. *Attenuation of 4-Nitroquinoline 1-oxide induced in vitro lipid peroxidation by green tea polyphenols*. *Life sciences*, 80(12), 1080-1086.
- Sri Rahayu, D., Kusriani, D. and Fachriyah, E., 2009. *Penentuan Aktivitas Antioksidan dari Ekstrak Etanol Daun Ketapang (Terminalia catappa L) dengan Metode 1, 1-Difenil-2-Pikrilhidrazil (DPPH)*.
- Sugihartini, N., Fudholi, A., Pramono, S., & Sismindari, S. 2014. **Validasi Metode Analisa Penetapan Kadar Epigallocatekin Galat Dengan Klt**

- Densitometri.** *Jurnal Pharmacia*, 4(1).
- Sukmawati, S. 2018. **Optimasi Dan Validasi Metode Analisis Dalam Penentuan Kandungan Total Flavonoid Pada Ekstrak Daun Gedi Hijau (*Abelmoscus manihot* L.) yang Diukur Menggunakan Spektrofotometer Uv-Vis.** *PHARMACON*, 7(3).
- Turkmen, N., Velioglu, Y, S., Sari, F., Polat, G. 2007. ***Effect of Extraction Conditions on Measured Total Polyphenol Contents and Antioxidant and Antibacterial Activities of Black Tea.*** *Journal Of Molecuas Volume* 12, p. 484-496
- USP, 2018. USP 41/NF 36. Rockville, MD: *United States Pharmacopeia Convention, Inc.*.
- Visht, S. and Chaturvedi, S., 2012. ***Isolation of natural products.*** *Current Pharma Research*, 2(3), p.584.
- Waris, R., Najib, A. and Pratiwi, E.D., 2016. ***Radical scavenging activity of leaf extract of edible Hibiscus (Abelmoschus manihot (L.) Medik) using 1, 1-Diphenyl-2-Picryl Hydrazil (DPPH).*** *International Journal of PharmTech Research*, 9(6), pp.343-347.
- Widowati, W., Herlina, T., Ratnawati, H., Constantia, G., Deva, I, D, G, S., Maesaroh, M. 2015. ***Antioxidant Potential of Black, Greenand Oolong Tea Methanol Extracts.*** *The Journal of Biology, Medicine, & Natural Product Chemistry Volume 4 Number 2. P. 35-39.*
- Yamaguchi, T., Takamura, H., Matoba, T. and Terao, J., 1998. ***HPLC method for evaluation of the free radical-scavenging activity of foods by using 1, 1-diphenyl-2-picrylhydrazyl.*** *Bioscience, biotechnology, and biochemistry*, 62(6), pp.1201-1204.
- Yang, C.S., Wang, X., Lu, G. and Picinich, S.C., 2009. ***Cancer prevention by tea: animal studies, molecular mechanisms and human relevance.*** *Nature Reviews Cancer*, 9(6), p.429.
- Yang, J., Liu R, H. 2012. ***The phenolic profiles and antioxidant activity in different types of tea.*** USA. *International Journal of Food Science and Technology*.
- Yuwono, M., dan Indrayanto, G. 2005. ***Validation of chromatographic methods of analysis.*** *Profiles of Drug Substances, Excipients and Related Methodology*, 32, 243-259.
- Zhou, K., and Yu, L. 2004. ***Effects of extraction solvent on wheat bran antioxidant activity estimation.*** *LWT-Food Science and Technology*, 37(7), 717-721.

Zuo, Y., Chen, H., and Deng, Y. 2002. *Simultaneous determination of catechins, caffeine and gallic acids in green, Oolong, black and pu-erh teas using HPLC with a photodiode array detector*. *Talanta*, 57(2), 307-316.