


eISSN: 2251-7308

pISSN: 2228-5881

Advanced Pharmaceutical Bulletin



Official Journal of
Faculty of Pharmacy
&

Drug Applied Research Center
Tabriz University of Medical Sciences

APB

Quarterly

<http://apb.tbzmed.ac.ir>
apb@tbzmed.ac.ir

Advanced Pharmaceutical Bulletin

(/)

Title

Title

Search

ISSN: 2228-5881 (<https://portal.issn.org/resource/ISSN/2228-5881>) eISSN: 2251-7308 (<https://portal.issn.org/resource/ISSN/2251-7308>)

Editors



Hadi Valizadeh (PharmD, PhD)

Founder and Editor-in-chief

Professor of Pharmaceutics

Faculty of Pharmacy,

Tabriz University of Medical Sciences, Tabriz, IRAN

Office Phone: +98(41)33348801

Email: valizadeh@tbzmed.ac.ir (<mailto:valizadeh@tbzmed.ac.ir>)

Google scholar (<http://scholar.google.com/citations?user=j1p6wZ4AAAAJ&hl=en>)



Hossein Babaei (PhD)

Director-in-Charge

Professor of Pharmacology

Faculty of Pharmacy,

Tabriz University of Medical Sciences

Tabriz, IRAN

Office Phone: +98(41)33363311

Fax: +98 (41) 33363231

E-mail: babaeih@tbzmed.ac.ir (<mailto:babaeih@tbzmed.ac.ir>)

(<https://scholar.google.com/citations?user=LpGJ520AAAAJ&hl=en>) Google Scholar (<https://scholar.google.com/citations?user=LpGJ520AAAAJ&hl=en>)

Associate Editors



Parvin Zakeri-Milani, Ph.D

Faculty of Pharmacy Tabriz University of Medical Sciences
Tabriz, IRAN
Phone: +98(41)33341315
E-mail: pzakeri@tbzmed.ac.ir (mailto:pzakeri@tbzmed.ac.ir)
Google scholar (<https://scholar.google.com/citations?user=zChcMm07hoC&hl=en>)



Hojjatollah Nozad Charoudeh

Associate Professor of Stem Cell Biology
Faculty of Medicine, Tabriz University of Medical Sciences
Tabriz, Iran,
Email: nozadh@tbzmed.ac.ir
Tel: +98(41)33342086
Google scholar (<https://scholar.google.com/citations?user=pgLFTBAAAAAJ&hl=en>)



Afshin Gharekhani

Department of Clinical Pharmacy,
Faculty of Pharmacy,
Tabriz University of Medical Sciences
Tabriz, IRAN
Email: Gharekhania@tbzmed.ac.ir (mailto:Gharekhania@tbzmed.ac.ir)
Google scholar (<https://scholar.google.com/citations?user=4aUxAV4AAAAJ&hl=en>)



Jafar Ezzati Nazhad Dolatabadi (PhD)

Assistant Professor of Pharmaceutical Nanotechnology
Tabriz University of Medical Sciences
Tabriz, IRAN
Email: ezzatij@tbzmed.ac.ir (mailto:ezzati@tbzmed.ac.ir)
Google scholar (<https://scholar.google.com/citations?user=Hdi1uF8AAAAJ&hl=en>)



Saurabh Bhatia (PhD)

Associate Professor,
Amity Institute of Pharmacy (AIP), Amity university,
Manesar, Gurgaon, INDIA
Tel.: 0091-9991634366
Email: sbhatia@ggn.amity.edu (mailto:sbhatia@ggn.amity.edu)
Google scholar (https://scholar.google.com/citations?user=C__7sY4AAAAJ&hl=en), linkedin (<https://www.linkedin.com/in/dr-saurabh-bhatia-65750b1a/>)



Hamed Hamishehkar, Ph.D

Professor of Pharmaceutics
Drug applied research center,
Tabriz University of Medical Sciences, Tabriz, IRAN
Tel: +98(41)-33378163
Email: hamishehkar.hamed@gmail.com (mailto:hamishehkar.hamed@gmail.com)
Google scholar (https://scholar.google.com/citations?hl=en&user=le6cgp4AAAAJ&view_op=list_works&sortby=pubdate)

International Editorial and Advisory Board



Raimar Loebenber, PhD

Professor, Pharmaceutical Sciences
Faculty of Pharmacy and Pharmaceutical Sciences, University of Alberta
3-142J Katz Group Centre, Edmonton, AB, Canada
Email: raimar.loebenberg@ualberta.ca (mailto:rloebenberg@pharmacy.ualberta.ca)
Tel: +1 (780) 492-1255
Google scholar, (<http://scholar.google.com/citations?user=wP6VnFMAAA&hl=en>) LinkedIn (<http://www.linkedin.com/pub/raimar-loebenberg/5b/5bb/a66>)



Gabriele Caviglioli, PhD

Professor at Università degli Studi di Genova
via Brigata Salerno, 13, 16147 Genova, Italy
Email: caviglioli@difar.unige.it (mailto:caviglioli@difar.unige.it)
LindedIn (<https://www.linkedin.com/in/gabriele-caviglioli-aa125116/>)



Hamid Omidian, PhD

Associate Professor, Pharmaceutical Sciences
Nova Southeastern University
3200 S University Drive
Fort Lauderdale, FL 33328, USA
Email: omidian@nova.edu (mailto:omidian@nova.edu)
Tel: 954-262-1334
Google scholar (<http://scholar.google.com/citations?hl=en&user=vWT5lowAAAAJ>)



Mohammad Barzegar Jalali, Ph.D

Professor of Pharmaceutics
Faculty of Pharmacy,
Tabriz University of Medical Sciences, Tabriz, IRAN
Office Phone: +98(41)33376148
Fax: +98 (41)33344798
E-mail: barzegar_jalali@yahoo.com (mailto:barzegar_jalali@yahoo.com)
Google scholar (<https://scholar.google.com/citations?user=ozaHfhkAAAAJ&hl=en>)



Miguel de la Guardia, PhD

Professor of Analytical Chemistry
Department of Analytical Chemistry
University of Valencia,
Valencia, SPAIN
Scopus ID: 7006639927
Email: miguel.delaguardia@uv.es (email: miguel.delaguardia@uv.es)
Google scholar (https://scholar.google.com/citations?hl=en&user=3oOU98IAAAA&view_op=list_works&sortBy=pubdate)



Ali Nokhodchi, PhD

Professor of Pharmaceutics and Drug Delivery
School of Life Sciences (<http://www.sussex.ac.uk/profiles/355690>)
University of Sussex
Brighton, UK BN1 9QG
email: A.Nokhodchi@sussex.ac.uk (<http://apb.tbzmed.ac.ir/AboutJournal/EditorialBoard/tabid/76/ct/Edit/mid/398/A.Nokhodchi@sussex.ac.uk>)
Google scholar (<http://scholar.google.com/citations?user=LiWqf28AAAAJ&hl=en>)



Yadollah Omid (PharmD, PhD)

Professor of Pharmaceutical Sciences
College of Pharmacy, Nova Southeastern University
Fort Lauderdale, FL, USA
Email: yomidi@yahoo.com (mailto:yomidi@yahoo.com)
Google Scholar (<http://scholar.google.com/citations?user=xBc5Y1YAAAA&hl=en>)



Mehrdad Hamidi, Ph.D

Professor of Pharmaceutics
Faculty of Pharmacy, Zanjan University of Medical Sciences
Zanjan, IRAN
Email: hamidim@zums.ac.ir (mailto:hamidim@zums.ac.ir)
Google scholar (<http://scholar.google.com/citations?user=S5gjK9sAAAAJ&hl=en>)



Gabriel Hancu, PhD

Associate professor (<http://www.umftgm.ro/facultatea-de-farmacie/departamente-si-discipline/departamentul-de-stiinte-farmaceutice-de-specialitate-dsfs/chimie-farmaceutica/colectivul-disciplinei.html>), Department of Pharmaceutical Chemistry,
Faculty of Pharmacy, University of Medicine and Pharmacy Târgu Mures (<http://www.umftgm.ro/facultatea-de-farmacie/departamente-si-discipline/departamentul-de-stiinte-farmaceutice-de-specialitate-dsfs/chimie-farmaceutica/colectivul-disciplinei.html>)
Gh. Marinescu 38, 540139 Târgu Mures, Romania
Tel: 0040-265-215551
E-mail: gabriel.hancu@umftgm.ro (<https://apb.tbzmed.ac.ir/page/gabriel.hancu@umftgm.ro>)
Google scholar (<http://scholar.google.com/citations?user=eRfui44AAAAJ&hl=en>)



Khaled Nabih Zaki Rashed, PhD

National Research Centre (NRC),
Pharmacognosy Department,
Pharmaceutical and Drug Industries Research Division,
Dokki, Giza, Egypt
E-mail: khalednabih2015@yahoo.co.uk (mailto:khalednabih2015@yahoo.co.uk), khaledrashed_352@ymail.com (mailto:khaledrashed_352@ymail.com)
Google scholar (<http://scholar.google.com/citations?user=oFPg8YYAAAAJ&hl=en>)



Kasturi Pawar, PhD

R&D Manager, Bausch Health Inc
Rohnert Park, California, USA
Email: kasturi.pawar@bauschheath.com (mailto:kasturi.pawar@bauschheath.com).
linkedIn (<https://www.linkedin.com/in/kasturi-p-34614314>)

[Submit Your Paper \(/Login\)](#)

[Instructions for Authors \(/InstructionsforAuthors\)](#)

[Aims and Scope \(/AimsandScope\)](#)

[Editorial Board \(/EditorialBoard\)](#)

[Editorial Policies \(/EditorialPolicies\)](#)

[Publication Ethics \(/Ethics\)](#)

[About APB & Contacts \(/ContactsandEditorialOffice\)](#)

[Articles ahead of print \(/Inpress\)](#)

[Archive \(/Archive\)](#)

Journal Info:

Freq.: Quarterly

eISSN: 2251-7308 (https://road.issn.org/issn_search?afs:query=2251-7308#.Vw7aFpwwiUk)

pISSN: 2228-5881 (https://road.issn.org/issn_search?afs:query=2228-5881#.Vw7aFpwwiUk)

Permission (<https://creativecommons.org/licenses/>): CC BY



([//creativecommons.org/licenses/](https://creativecommons.org/licenses/))

Platinum Open Access

(no submission or publication fees)

Powered by



Abstracting and Indexing

PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/journals/2246/>), Pubmed (<https://pubmed.ncbi.nlm.nih.gov/?term=%22Adv%20Pharm%20Bull%22%5BJournal%5D>),

Emerging Sources Citation Index (<http://mjl.clarivate.com/cgi-bin/jrnlst/jlresults.cgi?PC=MASTER&ISSN=2228-5881>)

Scopus (<https://www.scopus.com/sourceid/21100232413?origin=sbrowse>), Google Scholar (<https://scholar.google.com/citations?user=WWDFCCgAAAAJ&hl=enhttps://scholar.google.com/citations?user=WWDFCCgAAAAJ&hl=en>),

EMBASE/Excerpta Medica (<https://www.elsevier.com/solutions/embase-biomedical-research/embase-coverage-and-content>)

EBSCOhost (<https://www.ebscohost.com/titleLists/a9h-journals.htm>); CAS (https://cassi.cas.org/publication.jsp?P=LglBQf5Q2NQyz133K_lI3zLPXfcr-WXf_a0s1c_1-2Y3O7R7eJ0GCdLPXfcr-WXfimsBlkq8XcUjhmk0WtYxmzLPXfcr-WXfSHEqQVQf5dZsXLBusJR_0w); DOAJ (https://doaj.org/toc/2251-7308?source=%7B%22query%22%3A%7B%22filtered%22%3A%7B%22filter%22%3A%7B%22bool%22%3A%7B%22must%22%3A%5B%7B%22terms%22%3A%7B%22index.issn.exact%22%3A%5B%22228-5881%22%2C%222251-7308%22%5D%7D%7D%2C%7B%22term%22%3A%7B%22_type%22%3A%22article%22%7D%7D%5D%7D%7D%2C%22query%22%3A%7B%22match_all%22%3A%7B%7D%7D%7D%2C%22size%22%3A100%2C%22_source%22%3A%7E)

EBSCOhost (<https://www.ebscohost.com/titleLists/a9h-journals.htm>); CAS (https://cassi.cas.org/publication.jsp?P=LglBQf5Q2NQyz133K_lI3zLPXfcr-WXf_a0s1c_1-2Y3O7R7eJ0GCdLPXfcr-WXfimsBlkq8XcUjhmk0WtYxmzLPXfcr-WXfSHEqQVQf5dZsXLBusJR_0w); DOAJ (https://doaj.org/toc/2251-7308?source=%7B%22query%22%3A%7B%22filtered%22%3A%7B%22filter%22%3A%7B%22bool%22%3A%7B%22must%22%3A%5B%7B%22terms%22%3A%7B%22index.issn.exact%22%3A%5B%22228-5881%22%2C%222251-7308%22%5D%7D%7D%2C%7B%22term%22%3A%7B%22_type%22%3A%22article%22%7D%7D%5D%7D%7D%2C%22query%22%3A%7B%22match_all%22%3A%7B%7D%7D%7D%2C%22size%22%3A100%2C%22_source%22%3A%7E)

EBSCOhost (<https://www.ebscohost.com/titleLists/a9h-journals.htm>); CAS (https://cassi.cas.org/publication.jsp?P=LglBQf5Q2NQyz133K_lI3zLPXfcr-WXf_a0s1c_1-2Y3O7R7eJ0GCdLPXfcr-WXfimsBlkq8XcUjhmk0WtYxmzLPXfcr-WXfSHEqQVQf5dZsXLBusJR_0w); DOAJ (https://doaj.org/toc/2251-7308?source=%7B%22query%22%3A%7B%22filtered%22%3A%7B%22filter%22%3A%7B%22bool%22%3A%7B%22must%22%3A%5B%7B%22terms%22%3A%7B%22index.issn.exact%22%3A%5B%22228-5881%22%2C%222251-7308%22%5D%7D%7D%2C%7B%22term%22%3A%7B%22_type%22%3A%22article%22%7D%7D%5D%7D%7D%2C%22query%22%3A%7B%22match_all%22%3A%7B%7D%7D%7D%2C%22size%22%3A100%2C%22_source%22%3A%7E)

EBSCOhost (<https://www.ebscohost.com/titleLists/a9h-journals.htm>); CAS (https://cassi.cas.org/publication.jsp?P=LglBQf5Q2NQyz133K_lI3zLPXfcr-WXf_a0s1c_1-2Y3O7R7eJ0GCdLPXfcr-WXfimsBlkq8XcUjhmk0WtYxmzLPXfcr-WXfSHEqQVQf5dZsXLBusJR_0w); DOAJ (https://doaj.org/toc/2251-7308?source=%7B%22query%22%3A%7B%22filtered%22%3A%7B%22filter%22%3A%7B%22bool%22%3A%7B%22must%22%3A%5B%7B%22terms%22%3A%7B%22index.issn.exact%22%3A%5B%22228-5881%22%2C%222251-7308%22%5D%7D%7D%2C%7B%22term%22%3A%7B%22_type%22%3A%22article%22%7D%7D%5D%7D%7D%2C%22query%22%3A%7B%22match_all%22%3A%7B%7D%7D%7D%2C%22size%22%3A100%2C%22_source%22%3A%7E)

EBSCOhost (<https://www.ebscohost.com/titleLists/a9h-journals.htm>); CAS (https://cassi.cas.org/publication.jsp?P=LglBQf5Q2NQyz133K_lI3zLPXfcr-WXf_a0s1c_1-2Y3O7R7eJ0GCdLPXfcr-WXfimsBlkq8XcUjhmk0WtYxmzLPXfcr-WXfSHEqQVQf5dZsXLBusJR_0w); DOAJ (https://doaj.org/toc/2251-7308?source=%7B%22query%22%3A%7B%22filtered%22%3A%7B%22filter%22%3A%7B%22bool%22%3A%7B%22must%22%3A%5B%7B%22terms%22%3A%7B%22index.issn.exact%22%3A%5B%22228-5881%22%2C%222251-7308%22%5D%7D%7D%2C%7B%22term%22%3A%7B%22_type%22%3A%22article%22%7D%7D%5D%7D%7D%2C%22query%22%3A%7B%22match_all%22%3A%7B%7D%7D%7D%2C%22size%22%3A100%2C%22_source%22%3A%7E)

Advanced Pharmaceutical Bulletin

Q1 Pharmacology, Toxicology and Pharmaceutics... best quartile

SJR 2019 0.57



powered by scimagojr.com

(<https://www.scimagojr.com/journalsearch.php?q=21100232413&tip=sid&exact=no>)



Commentary

Comments on "Dissolution Enhancement of Atorvastatin Calcium by Cocrystallization"


(/Inpress/apb-28083)
Shahram Emami  (<https://orcid.org/0000-0002-5182-1728>), Ali Shayanfar*  (<https://orcid.org/0000-0001-7507-9057>)

Adv Pharm Bull. doi: 10.34172/apb.2021.067 (<https://doi.org/10.34172/apb.2021.067>) | Published on: 2020 May 26

 PDF (/Files/Inpress/apb-28083.pdf)

Review Article

Breast Cancer: a Global Concern, Diagnostic and Therapeutic Perspectives, Mechanistic Targets in Drug Development

(/Inpress/apb-28663)
Gul-e-Saba Chaudhry*  (<https://orcid.org/0000-0002-4555-8572>), Rehmat Jan, Abdah Akim, Muhammad Naveed Zafar, Yeong Yik Sung, Tengku Sifzizul Tengku Muhammad

Adv Pharm Bull. doi: 10.34172/apb.2021.068 (<https://doi.org/10.34172/apb.2021.068>) | Published on: 2020 Oct 14

 PDF (/Files/Inpress/apb-28663.pdf)

Review Article

Ligament/Tendon Culture under Hypoxic Conditions: A Systematic Review


(/Inpress/apb-30198)
Sholahuddin Rhatomy*  (<https://orcid.org/0000-0002-5512-6706>), Dwikora Novembri Utomo, Cita Rosita Sigit Prakoeswa, Fedik Abdul Rantam, Heri Suroto, Ferdiansyah Mahyudin

Adv Pharm Bull. doi: 10.34172/apb.2021.069 (<https://doi.org/10.34172/apb.2021.069>) | Published on: 2020 Oct 20

 PDF (/Files/Inpress/apb-30198.pdf)

Review Article

Medicinal Plants in the Treatment of Hypertension: A Review

(/Inpress/apb-28754)
Raha Kamyab, Hossein Namdar, Mohammadali Torbati*  (<https://orcid.org/0000-0001-5040-9769>), Morteza Ghojzadeh, Mostafa Araj-Khodaei, Seyed Mohammad Bagher Fazliljou*

Adv Pharm Bull. doi: 10.34172/apb.2021.090 (<https://doi.org/10.34172/apb.2021.090>) | Published on: 2020 Nov 01


 PDF (/Files/Inpress/apb-28754.pdf)

Mini Review

DR-5 and DLL-4 mAb Functionalized SLNs of Gamma-Secretase Inhibitors- An Approach for TNBC Treatment

(/Inpress/apb-28814)
Mamta Kumari  (<https://orcid.org/0000-0002-9917-2077>), Praveen T. Krishnamurthy*  (<https://orcid.org/0000-0003-2666-0878>), Sai kiran S. S. Pinduprolu, Piyong

Sola
Adv Pharm Bull. doi: 10.34172/apb.2021.070 (<https://doi.org/10.34172/apb.2021.070>) | Published on: 2020 Oct 19

 PDF (/Files/Inpress/apb-28814.pdf)

Research Article

Application of Carbonic Nanosheets Based on Urea Precursors as Dispersive Solid Phase Extraction Adsorbent for Extraction of Methamphetamine from Urine Samples

(/Inpress/apb-29006)
Arezou Taghvimi  (<https://orcid.org/0000-0003-0773-5547>), Siavoush Dastmalchi, Yousef Javadzadeh*  (<https://orcid.org/0000-0001-7283-3560>)

Adv Pharm Bull. doi: 10.34172/apb.2021.071 (<https://doi.org/10.34172/apb.2021.071>) | Published on: 2020 Oct 20

 PDF (/Files/Inpress/apb-29006.pdf)

Research Article

A Novel Approach for Development of Intraocular Biodegradable Ranibizumab Implant: A Solution for Stability of Protein Activity

(/Inpress/apb-29068)
Fatemeh Darigar Pakdel  (<https://orcid.org/0000-0003-3863-1599>), Ahmad Mirshahi, Payam Zahedi, Kazem Mohammad, Farkhondeh Hemmati, Javad Dadgar Pakdel, Mohammad Hossein Nicknam*  (<https://orcid.org/0000-0001-9132-1675>), Farid Abedin Dorkoosh*  (<https://orcid.org/0000-0002-0417-7300>)

Adv Pharm Bull. doi: 10.34172/apb.2021.072 (<https://doi.org/10.34172/apb.2021.072>) | Published on: 2020 Oct 20

 PDF (/Files/Inpress/apb-29068.pdf)

How to cite this article: Rhatomy Sh, Utomo DN, Prakoeswa CRS, Rantam FA, Suroto H, Mahyudin F. Ligament/Tendon Culture under Hypoxic Conditions: A Systematic Review. *Advanced Pharmaceutical Bulletin*, doi:[10.34172/apb.2021.069](https://doi.org/10.34172/apb.2021.069)

Ligament/Tendon Culture under Hypoxic Conditions: A Systematic Review

Author's details:

Sholahuddin Rhatomy¹ <https://orcid.org/0000-0002-5512-6706>, Dwikora Novembri Utomo², Cita Rosita Sigit Prakoeswa³, Fedik Abdul Rantam^{4,5}, Heri Suroto², and Ferdiansyah Mahyudin²

1. Doctoral Program, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia
2. Department of Orthopaedic and Traumatology, Dr. Soetomo General Hospital, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia
3. Department of Dermatology and Venereology, Dr. Soetomo General Hospital, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia
4. Virology and Immunology Laboratory, Microbiology Department, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya Indonesia
5. Stem Cell Research and Development Center, Universitas Airlangga, Surabaya Indonesia
- 6.

Corresponding author:

Name : Sholahuddin Rhatomy, MD, Adult Reconstructive Surgeon

Sholahuddin Rhatomy, MD (Sport and Adult Reconstructive Surgeon)

Department of Orthopaedics and Traumatology, Faculty of Medicine, Universitas Airlangga, Indonesia

Mayjen Prof. Dr. Moestopo Street, number 47, Pacar Kembang, Tambaksari, Surabaya, East Java, Indonesia 60132

Tel : (+6231)5020251

Email: doktergustomrhatomy@yahoo.com

ORCID ID : <http://orcid.org//0000-0002-5512-6706>

Ligament/tendon culture under hypoxic conditions: a systematic review

Abstract

The hypoxic environment is a substantial factor in maintenance, proliferation, and differentiation of the cell cultures. Low oxygen is known as a potent chondrogenesis stimulus in stem cells that is important for clinical application and engineering of functional cartilage. Hypoxia can potentially induce angiogenesis process by secretion of cytokines. This systematic review goal is to discover the effect of hypoxic condition on tendon/ligament culture and the best oxygen level of hypoxia for in vitro and in vivo studies. We included 21 articles. A comprehensive review of this database confirms that the hypoxic condition is a substantial factor in the maintenance, proliferation, and differentiation of ligament/tendon cultures. Cell proliferation in the severe hypoxic (oxygen concentration of 1%) group at 24 h postcultivation was considered significant, but cell proliferation was markedly inhibited in the severe hypoxic group after 48 h.

Keywords: Hypoxic condition, ligament, tendon, culture

Introduction

Hypoxia is a physiological condition in the environment of various stem cells. In a previous study, the oxygen level for most cell cultures was maintained at approximately 20% in vitro.¹ The native microenvironment in the body has a low oxygen pressure; on average, this pressure is approximately 12% in arterial blood and 3% in tissues, which may vary depending on the cell location.¹

Hypoxia has been used in tissue cultivation. Some studies have reported a significant increase in the proliferation of cells cultured under low oxygen pressure.^{2,3} Comparisons of growth in normal and low oxygen levels during preconditioning showed that embryonic stem cells grew more efficiently with low oxygen. Among the spectrum of adult stem cells, Grayson et al. examined human bone marrow-derived mesenchymal stem cells (MSCs) and showed that low oxygen levels resulted in a 30-fold increase in cell growth compared to normoxic conditions.¹

Tendon or ligament tears are frequent injuries that may disturb the patient's quality of life. The healing process is often inadequate, leading to scar tissue formation. Various treatments for these injuries have been developed, including physiotherapy, pain medications, steroid injections, and surgical procedures. However, these therapeutic methods have not been satisfactory for patients. Stem cells have become a new therapeutic modality because of their ability to induce cell rejuvenation, stemness potential, and immunomodulatory activity. A systematic review by Ahmad et al. revealed that stem cell treatment has been applied to heal injured tendons in animals and humans. Among the various sources of stem cells, mesenchymal stem cells (MSCs) are highly useful for healing tendons and ligaments and preventing scar formation.^{4,5}

Some studies have examined various levels of hypoxic conditions and their effects.⁷ However, limited evidence is available regarding the use of hypoxic conditions for ligament/tendon cultures. An in-depth review of in vitro and in vivo studies is needed to evaluate the effectiveness of hypoxic conditions for ligament/tendon cultures and to translate their use into clinical applications. Thus, this review aimed to determine the effect of hypoxic conditions and to identify the optimal oxygen level for use in in vivo and in vitro tendon/ligament cultures.

Methods

Review protocol

Our review aimed to elucidate the effect of hypoxic levels on tendon/ligament cultures as well as the optimal hypoxic condition (oxygen concentration) for in vitro and in vivo studies.

Literature search and study selection

In May 2020, a database search was carried out in the Scopus, PubMed (Medline), Web of Science, Science Direct, Google Scholar, and Cochrane Library databases to identify all studies published in English that described the outcomes of ligament/tendon cultures under hypoxic conditions. A comprehensive search of the literature was performed based on the Animal Research: Reporting of In Vivo Experiments (ARRIVE) guidelines and the Checklist for Reporting In vitro Studies (CRIS Guidelines).

The search keywords included “ligament”, “tendon”, “culture”, “condition”, “precondition”, and “hypoxic” alone and combined the Boolean operators “AND” or “OR”.

Two authors (D.N.U. and S.R.) independently scanned for eligibility based on the title and abstract after duplicate and review articles were removed from the collected articles. These two authors also performed an additional search based on references to the included articles. The final search result was included in this review. The included articles were read by the authors to match the inclusion and exclusion criteria. Any discordance between authors was resolved through discussion.

Outcome measures

The following outcome measures were analysed: 1) the most common hypoxic condition (oxygen concentration) used and 2) the main outcomes.

Eligibility criteria

The following inclusion criteria were used: 1) peer-reviewed level 1 to 4 studies, 2) studies published in English, 3) studies of ligament or tendon cultures, 4) a description of the hypoxic condition (oxygen concentration), and 5) an obvious description of the outcomes. Studies that were not published in English, duplicate articles, review studies, and irrelevant articles were excluded.

Assessment of methodological quality and risk of bias

The methodological quality of the included studies was assessed using ARRIVE and CRIS guidelines.^{9,10} Internal validity was assessed using the Systematic Review Center for Laboratory Animal Experimentation Risk of Bias tool (SYRCLE RoB tool).¹¹ Two authors (D.N.U and S.R.) independently performed all assessments.

Data Extraction and synthesis

Two authors (D.N.U. and S.R.) independently analysed and recorded the data from the included studies. The following data were extracted: study design, types of animals used for in vivo

study, cell types for in vitro study, process for establishing animal or cell models, hypoxic condition, stem cell isolation procedures, interventions, comparisons, follow-up duration, primary outcomes of in vivo and in vitro studies and their results, any significant deviation from the control, and other outcomes.

We also recorded the blinding method of data collection of the included studies when available. We collectively reported the types of animals used in the study, interventions, and duration of follow-up. We reported any adverse reactions, quantitative outcome measurements, and biomechanical tests for in vivo studies. For in vitro studies, we reported cell proliferation, viability and migration; tenogenic differentiation; and immunomodulatory properties. We could not perform a meta-analysis because of the substantial heterogeneity of the data (i.e., different intervention levels of hypoxic conditions and different measurements of the outcomes).

Results

Study selection

In total, 9540 articles were collected from the databases; 8789 articles were excluded based on the title or duplication. Title and abstract screening was conducted for 70 articles. Sixteen articles were excluded because they did not match the inclusion criteria. After enrolling 54 articles for full-text assessment, 25 were excluded due to insufficient description of hypoxic conditions ($n = 21$), short communications/reviews ($n = 2$), and multiple reports of the same experiment ($n = 2$). Ultimately, 29 articles were enrolled in this systematic review. The article selection process is provided in Figure 1.

Characteristics of selected studies

Of the 29 studies, 24 (82.7%) were published between 2010 and April 2020. Human periodontal ligament-derived stem cells (86%) were the main cell type used among these studies. All included studies were conducted in vitro, but 2 studies also reported the results of an in vivo experiment.^{7,12}

Stem cell source

Of the 29 studies, 27 (93%) used human stem cells, and only 2 studies used stem cells from an animal (rat). Moreover, 21 studies (72%) used the human periodontal ligament, whereas the remaining studies used the Achilles tendon or periodontal ligament from rats, human hamstring, or human anterior cruciate ligament.

Hypoxic conditions (oxygen concentration)

Fourteen studies performed a comparison with normoxic conditions (20% or 21% oxygen), three used 1% oxygen,^{20,24,28} five used 2% oxygen,^{8,22,25,26,29} two used 3% oxygen,^{12,30} and four studies compared various hypoxia levels, including 1%, 2%, and 5% oxygen;³⁷ 1% oxygen and a reoxygenation group (from 1% to 20% oxygen);^{8,15} and 5% and 1% oxygen.³¹ Overall, 15 studies evaluated the effects of certain oxygen concentrations: 3 used 5% oxygen,^{13,14,32} 1 used 3% oxygen,³⁴ 5 used 2% oxygen,^{7,19,21,27,35} 2 used 1% oxygen,^{20,36} 3 used less than 1% oxygen,¹⁶⁻¹⁸ and another used less than 0.1% oxygen.³³

Main Outcomes

Seven studies (24%) reported increased cell proliferation.^{6-8,13,25,36,37} Four studies showed that hypoxic conditions increased alkaline phosphatase activity.^{8,13,19,37} Six studies revealed a significant increase in vascular endothelial growth factor (VEGF) expression.^{8,14,15,18,19,33} Eight studies reported a substantial increase in the expression of cytokines, such as interleukin (IL)-1 β ,^{15,18,23} IL-6,^{15,18,20,23} IL-8,¹⁸

IL-10,¹² IL-37,¹² and tumour necrosis factor- α (TNF- α),²³ in cells cultured under hypoxic conditions. Ito et al. observed no effects by hypoxia on the secretion of IL-1 β , IL-6, IL-8, IL-17A, macrophage migration inhibitory factor, monocyte chemoattractant protein-1, TNF- α , and macrophage colony-stimulating factor in the culture media.³² Moreover, other studies revealed a significant increase in collagen type I²⁸ and III^{20,28} and glycosaminoglycan (GAG).²⁸

Six studies reported an increase in osteogenic differentiation.^{7,19,22,26,29,36} Hypoxic conditions also increased osteoclastogenesis,^{17,27} improved the maintenance of pluripotent stem cells,²⁵ increased the expression of several mediators of apoptosis, and promoted tenocyte apoptosis.²⁰

Discussion

This study was a systematic review of the available literature describing ligament/tendon cultures under hypoxic conditions. In this review, we assessed the characteristics of the selected studies, stem cell sources, hypoxic conditions (i.e., oxygen concentration), and the main study outcomes.

The oxygen level of the tissue culture in the in vitro study was maintained at approximately 20%. Hypoxia plays an important role in marker secretion and especially in haematologic features, such as gene expression in haematopoietic stem cells (HSCs) and hypoxia-inducible factor-dependent expression in supporting cells. This suggests that hypoxia may improve HSC growth and maintenance during ex vivo culture.³⁸ Hypoxia promotes MSC cultivation by improving cell expansion, osteogenesis increment, and bone and cartilage cell proliferation. Furthermore, MSC preculture under hypoxic conditions improved their regenerative potential.¹

Matsuda et al. first studied ligament/tendon cultures using human periodontal ligament exposed to 5% oxygen,¹³ after which research on ligament or tendon cultures under hypoxic conditions developed over the last decade. Therefore, most studies (82.7%, 24 studies) were published from 2010-2020.

Human derived periodontal ligament (HDPL) stem cells (86%) were predominantly used in the articles included in this review. Periodontitis disturbs the chewing mechanism between teeth and is highly prevalent worldwide. Research on periodontal regeneration was pioneered in 2004, and other stem cells have been reported to have the capability to differentiate into periodontal tissue under proper conditions for induction.³⁹

All included studies were conducted in vitro, but 2 studies also included in vivo experiments.^{7,12} Stem cell growth and function are used to determine the optimal in vivo microenvironment conditions, including the oxygen level. Natural cellular microenvironments seem to contain lower oxygen levels, with a mean oxygen level of approximately 12% in arterial blood and 3% in the tissues as well as considerable variation depending on the location. During early pregnancy, the surface of the uterine cavity has approximately 2% oxygen. After utero-placental circulation develops, the placental oxygen level increases to approximately 8%.^{1,40}

No uniformity was found between the studies regarding the optimum oxygen concentration/hypoxic conditions for ligament/tendon cultures, but 11 studies (37%) used 2% oxygen and 9 studies (31%) used 1% oxygen. Zhang et al.³⁷ compared the effects of oxygen concentrations of 1%, 2%, 5%, and 21% and concluded that there was a significant increase in cell proliferation in the severe hypoxia group (1% oxygen concentration) at 24 h post-cultivation ($p < 0.05$); moreover, after 48 h, cell proliferation was markedly restrained ($p < 0.05$). The level of

alkaline phosphatase activity in the severe hypoxia group was also greatly reduced ($p < 0.05$) after 24 h.

Lastly, this review aimed to evaluate the main outcomes of ligament/tendon cultures under hypoxic conditions. The examined studies did not analyse similar parameters or outcomes; thus, a meta-analysis of the data was not feasible.

Seven included studies (24%) reported increases in cell proliferation.^{6,8,13,25,36,37} This result is consistent with that of many other studies that observed a significant increase in the proliferation of embryonic stem cells or adult stem cells exposed to low oxygen tension.¹

Eight studies (27.5%) reported an increase in the expression of cytokines, such as IL-1 β ,^{15,18,23} IL-6,^{15,18,20,23} IL-8,¹⁸ IL-10,¹² IL-37,¹² and TNF- α .²³ However, other studies examined different cytokines. Ito et al. observed no effects on the secretion of IL-1 β , IL-6, IL-8, IL-17A, macrophage migration inhibitory factor, monocyte chemoattractant protein-1, TNF- α , and macrophage colony-stimulating factor in the culture media.³² The production of the pro-inflammatory cytokine IL-8 was suppressed in MSCs cultured under hypoxic conditions, whereas the levels of the anti-inflammatory cytokines IL-17A and GM-CSF increased following short-term hypoxia.⁴¹

Other studies identified a significant increase in the production of collagen type I²⁸ and III^{20,28} and GAG.²⁸ These findings show promise for tissue engineering and offer the option to regenerate the tendon tissue by producing a stronger tendon construct for injury repair, thicker collagen fibre density, denser tissue architecture, and more normal restoration of the tendon-bone interface.

Six studies described a significant increase in VEGF expression.^{8,14,15,18,19,33} When stem cells were placed in low-oxygen conditions, most studies examining MSCs revealed an increase in growth factor secretion, particularly VEGF. Angiogenesis may be induced by stem cells through cell differentiation that contributes to angiogenesis directly or indirectly via hypoxia-stimulated cytokine production.^{1,14}

This review has some limitations, including the use of different oxygen concentrations and the predominant use of stem cells derived from the HDPL. Further investigations are needed to determine which oxygen concentrations are effective in achieving the best outcome. Research comparing different oxygen concentrations is needed to identify the optimal conditions for tissue culture.

Conclusion

Primarily, this systematic review found that a low oxygen concentration is an important factor contributing to the preservation, proliferation, differentiation, and function of ligament/tendon cultures. In a notable study, a significant increase in cell proliferation was observed in the severe hypoxia group (1% oxygen concentration) at 24 h post-cultivation, but cell proliferation was markedly reduced at 48 h post-cultivation. Areas for future research might include optimising the preconditions in terms of the timing and oxygen concentrations that are most effective for achieving each outcome.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

Conflicts of interest

None of the authors have conflicts of interest related to the study to disclose.

Funding

We did not receive funding from any source.

Authors' contributions

Sholahuddin Rhatomy and Dwikora Novembri Utomo designed and conducted the experiments.

Sholahuddin Rhatomy, Dwikora Novembri Utomo, Heri Suroto, Cita Rosita Sigit Prakoeswa, Fedik Abdul Rantam and Ferdiansyah Mahyudin searched the database.

Sholahuddin Rhatomy, Dwikora Novembri Utomo, Heri Suroto and Ferdiansyah collected and analyzed data.

Sholahuddin Rhatomy and Dwikora Novembri Utomo wrote the draft; all authors read, reviewed and approved the final manuscript.

Guarantor

Sholahuddin Rhatomy

References

1. Abdollahi H, Harris LJ, Zhang P, et al. The role of hypoxia in stem cell differentiation and therapeutics. *J Surg Res*. 2011;165(1):112-117. doi:10.1016/j.jss.2009.09.057
2. Hu X, Wei L, Taylor TM, Wei J, Zhou X, Wang JA, et al. Hypoxic preconditioning enhances bone marrow mesenchymal stem cell migration via Kv2.1 channel and FAK activation. *Am J Physiol - Cell Physiol*. 2011;301(2):362-72. doi: 10.1152/ajpcell.00013.2010.
3. Iida K, Takeda-Kawaguchi T, Tezuka Y, Kunisada T, Shibata T, Tezuka KI. Hypoxia enhances colony formation and proliferation but inhibits differentiation of human dental pulp cells. *Arch Oral Biol*. 2010;55(9):648-54. <https://doi.org/10.1016/j.archoralbio.2010.06.005>
4. Ahmad Z, Wardale J, Brooks R, Henson F, Noorani A, Rushton N. Exploring the application of stem cells in tendon repair and regeneration. *Arthroscopy*. 2012;28(7):1018-1029. doi:10.1016/j.arthro.2011.12.009
5. Rhatomy S, Prasetyo TE, Setyawan R, Soekarno NR, Romaniyanto FNU, Sedjati AP, et al. Prospect of stem cells conditioned medium (secretome) in ligament and tendon healing: A systematic review. *Stem Cells Transl Med*. 2020 ; 9 (8):895-902. <https://doi.org/10.1002/sctm.19-0388>
6. He Y, Jian CX, Zhang HY, Zhou Y, Wu X, Zhang G, et al. Hypoxia enhances periodontal ligament stem cell proliferation via the MAPK signaling pathway. *Genet Mol Res*. 2016;15(4):1-10. DOI <http://dx.doi.org/10.4238/gmr15048965>
7. Zhang QB, Zhang ZQ, Fang SL, Liu YR, Jiang G, Li KF. Effects of hypoxia on proliferation and osteogenic differentiation of periodontal ligament stem cells: an in

- vitro and in vivo study. *Genet Mol Res*. 2014;13(4):10204-10214. Published 2014 Dec 4. doi:10.4238/2014.December.4.15
8. Amemiya, H., Matsuzaka, K., Kokubu, E., Ohta, S. and Inoue, T. (2008), Cellular responses of rat periodontal ligament cells under hypoxia and re-oxygenation conditions *in vitro*. *Journal of Periodontal Research*, 43: 322-327. doi:10.1111/j.1600-0765.2007.01032.x
 9. Kilkenny C, Browne W, Cuthill IC, Emerson M, et al. Animal research: Reporting in vivo experiments: The ARRIVE guidelines. *Br J Pharmacol*. 2010;160(7):1577–9. <https://doi.org/10.1111/j.1476-5381.2010.00872.x>
 10. Krithikadatta J, Gopikrishna V, Datta M. CRIS guidelines (Checklist for Reporting In- vitro Studies): A concept note on the need for standardized guidelines for improving quality and transparency in reporting in vitro studies in experimental dental research. *J Conserv Dent*. 2014;17(4):301–4. <https://dx.doi.org/10.4103%2F0972-0707.136338>
 11. Hooijmans, C.R., Rovers, M.M., de Vries, R.B. *et al*. SYRCLE's risk of bias tool for animal studies. *BMC Med Res Methodol* 14, 43 (2014) : 1-9. <https://doi.org/10.1186/1471-2288-14-43>
 12. Giacoppo S, Thangavelu SR, Diomede F, Bramanti P, Conti P, Trubiani O, et al. Anti-inflammatory effects of hypoxia-preconditioned human periodontal ligament cell secretome in an experimental model of multiple sclerosis: a key role of IL-37. *Faseb J*. 2017;31(12):5592–608. doi: 10.1096/fj.201700524R
 13. Naoki Matsuda, Naoko Morita, Kazuko Matsuda and Masami Watanabe. Proliferation and Differentiation of Human Osteoblastic Cells Associated with Differential Activation of MAP Kinases in Response to Epidermal Growth Factor, Hypoxia, and Mechanical Stress *In Vitro*. *Biochem Biophys Res Commun*. 1998;249(2):350-354. <https://doi.org/10.1006/bbrc.1998.9151>
 14. Petersen W, Pufe T, Zantop T, Tillmann B, Mentlein R. Hypoxia and PDGF have a synergistic effect that increases the expression of the angiogenic peptide vascular endothelial growth factor in Achilles tendon fibroblasts. *Arch Orthop Trauma Surg*. 2003;123(9):485-488. doi:10.1007/s00402-003-0493-0
 15. Motohira H, Hayashi J, Tatsumi J, Tajima M, Sakagami H, Shin K. Hypoxia and Reoxygenation Augment Bone-Resorbing Factor Production From Human Periodontal Ligament Cells. *J Periodontol*. 2007;78(9):1803–9. doi: 10.1902/jop.2007.060519
 16. Kitase Y, Yokozeki M, Fujihara S, Izawa T, Kuroda S, Tanimoto K, et al. Analysis of gene expression profiles in human periodontal ligament cells under hypoxia: The protective effect of CC chemokine ligand 2 to oxygen shortage. *Arch Oral Biol*. 2009;54(7):618–24. doi:10.1016/j.archoralbio.2009.03.010
 17. Park HJ, Baek KH, Lee HL, Kwon A, Hwang HR, Qadir AS, et al. Hypoxia inducible factor-1 α directly induces the expression of receptor activator of nuclear factor- κ B ligand in periodontal ligament fibroblasts. *Mol Cells*. 2011;31(6):573–8. <https://doi.org/10.1007/s10059-011-1055-x>
 18. Chae HS, Park HJ, Hwang HR, Kwon A, Lim WH, Yi WJ, et al. The effect of antioxidants on the production of pro-inflammatory cytokines and orthodontic tooth movement. *Mol Cells*. 2011;32(2):189–96.

- <https://doi.org/10.1007/s10059-011-0071-1>
19. Wu Y, Cao H, Yang Y, et al. Effects of vascular endothelial cells on osteogenic differentiation of noncontact co-cultured periodontal ligament stem cells under hypoxia. *J Periodontol Res*. 2013;48(1):52-65. doi:10.1111/j.1600-0765.2012.01503.x
 20. Millar NL, Reilly JH, Kerr SC, Campbell AL, Little KJ, Leach WJ, et al. Hypoxia: A critical regulator of early human tendinopathy. *Ann Rheum Dis*. 2012;71(2):302–10. doi:10.1136/ard.2011.154229
 21. Kato R, Ishihara Y, Kawanabe N, Sumiyoshi K, Yoshikawa Y, Nakamura M, et al. Gap- junction-mediated communication in human periodontal ligament cells. *J Dent Res*. 2013;92(7):635–40. DOI: 10.1177/0022034513489992
 22. Zhao L, Wu Y, Tan L, Xu Z, Wang J, Zhao Z, et al. Coculture With Endothelial Cells Enhances Osteogenic Differentiation of Periodontal Ligament Stem Cells via Cyclooxygenase-2/Prostaglandin E 2/Vascular Endothelial Growth Factor Signaling Under Hypoxia. *J Periodontol*. 2013;84(12):1847–57. doi: 10.1902/jop.2013.120548
 23. Jian C, Li C, Ren Y, He Y, Li Y, Feng X, et al. Hypoxia Augments Lipopolysaccharide-Induced Cytokine Expression in Periodontal Ligament Cells. *Inflammation*. 2014;37(5):1413–23. DOI: 10.1007/s10753-014-9865-6
 24. Gözl L, Memmert S, Rath-Deschner B, Jäger A, Appel T, Baumgarten G, et al. LPS from *P. gingivalis* and hypoxia increases oxidative stress in periodontal ligament fibroblasts and contributes to periodontitis. *Mediators Inflamm*. 2014:1-13. <http://dx.doi.org/10.1155/2014/986264>
 25. Zhou Y, Fan W, Xiao Y. The effect of hypoxia on the stemness and differentiation capacity of PDLC and DPC. *Biomed Res Int*. 2014:1-7. <http://dx.doi.org/10.1155/2014/890675>
 26. Li L, Han MX, Li S, Xu Y, Wang L. Hypoxia regulates the proliferation and osteogenic differentiation of human periodontal ligament cells under cyclic tensile stress via mitogen-activated protein kinase pathways. *J Periodontol*. 2014;85(3):498-508. doi:10.1902/jop.2013.130048
 27. Li ML, Yi J, Yang Y, et al. Compression and hypoxia play independent roles while having combinative effects in the osteoclastogenesis induced by periodontal ligament cells. *Angle Orthod*. 2016;86(1):66-73. doi:10.2319/121414.1
 28. Kowalski TJ, Leong NL, Dar A, Wu L, Kabir N, Khan AZ, et al. Hypoxic culture conditions induce increased metabolic rate and collagen gene expression in ACL-derived cells. *J Orthop Res*. 2016;34(6):985–94. DOI 10.1002/jor.23116
 29. Chen D, Wu L, Liu L, Gong Q, Zheng J, Peng C, et al. Comparison of HIF1A-AS1 and HIF1A-AS2 in regulating HIF-1 α and the osteogenic differentiation of PDLCs under hypoxia. *Int J Mol Med*. 2017;40(5):1529–36. DOI: 10.3892/ijmm.2017.3138
 30. Murabayashi D, Mochizuki M, Tamaki Y, Nakahara T. Practical methods for handling human periodontal ligament stem cells in serum-free and serum-containing culture conditions under hypoxia: implications for regenerative medicine. *Hum Cell*. 2017;30(3):169-180. doi:10.1007/s13577-017-0161-2

31. Xiao Z, Han Y, Zhang Y, Zhang X. Hypoxia-regulated human periodontal ligament cells via Wnt/ β -catenin signaling pathway. *Medicine (Baltimore)*. 2017;96(16):e6562 :1-9. doi:10.1097/MD.00000000000006562
32. Ito H, Kifune T, Ishiyama M, et al. Effect of hypoxia on the expression of CCAAT/enhancer-binding protein β and receptor activator of nuclear factor kappa-B ligand in periodontal ligament cells. *J Oral Sci*. 2018;60(4):544-551. doi:10.2334/josnusd.17-0436
33. Kifune T, Ito H, Ishiyama M, Iwasa S, Takei H, et al. Hypoxia-induced Upregulation of Angiogenic Factors in Immortalized Human Periodontal Ligament Fibroblasts. *J Oral Sci*. 2018;60(4):519–25. doi.org 10.2334 josnusd.17-0441
34. Liu Z, Guo L, Li R, Xu Q, Yang J, Chen J, et al. Transforming growth factor- β 1 and hypoxia inducible factor-1 α synergistically inhibit the osteogenesis of periodontal ligament stem cells. *Int Immunopharmacol*. 2019;75:105834 : 1-8. <https://doi.org/10.1016/j.intimp.2019.105834>
35. Yan K, Wu C, Ye Y, Li L, Wang X, He W, et al. A20 inhibits osteoclastogenesis via TRAF6-dependent autophagy in human periodontal ligament cells under hypoxia. *Cell Prolif*. 2020;53(3) : 1-10. DOI: 10.1111/cpr.12778
36. Mao H, Yang A, Zhao Y, Lei L, Li H. Succinate Supplement Elicited “pseudohypoxia” Condition to Promote Proliferation, Migration, and Osteogenesis of Periodontal Ligament Cells. *Stem Cells Int*. 2020;2020.10.3892/etm.2013.1349 : 1-14. <http://doi.org/10.1155/2020/2016809>
37. Zhang HY, Liu R, Xing YJ, Xu P, Li Y, Li CJ. Effects of hypoxia on the proliferation, mineralization and ultrastructure of human periodontal ligament fibroblasts in vitro. *Exp Ther Med*. 2013;6(6):1553–9. <https://dx.doi.org/10.3892%2Fetm.2013.1349>
38. Eliasson P, Jönsson JI. The hematopoietic stem cell niche: Low in oxygen but a nice place to be. *J Cell Physiol*. 2010;222(1):17–22. <https://doi.org/10.1002/jcp.21908>
39. Seo BM, Miura M, Gronthos S, Bartold PM, Batouli S, Brahim J, et al. Investigation of multipotent postnatal stem cells from human periodontal ligament. *Lancet*. 2004;364(9429):149–55. [https://doi.org/10.1016/s0140-6736\(04\)16627-0](https://doi.org/10.1016/s0140-6736(04)16627-0)
40. Genbacev O, Zhou Y, Ludlow JW, Fisher SJ. Regulation of human placental development by oxygen tension. *Science*. 1997;277(5332):1669-1672. doi:10.1126/science.277.5332.1669
41. Antebi B, Rodriguez LA, Walker KP, Asher AM, Kamucheka RM, Alvarado L, et al. Short-term physiological hypoxia potentiates the therapeutic function of mesenchymal stem cells. *Stem Cell Res Ther*. 2018;9(1):1–15. <https://doi.org/10.1186/s13287-018-1007-x>