

**DAFTAR PUSTAKA**

- Adjepong, M. *et al.* (2016) 'The role of antioxidant micronutrients in the rate of recovery of burn patients: a systematic review', *Burns & Trauma*. *Burns & Trauma*, 4(1). doi: 10.1186/s41038-016-0044-x.
- Alamdari, N. *et al.* (2008) 'Temporal changes in the involvement of pyruvate dehydrogenase complex in muscle lactate accumulation during lipopolysaccharide infusion in rats', *Journal of Physiology*, 586, pp. 1767-1775.
- Attaluri, P. *et al.* (2018) 'Thiamine Deficiency: An Important Consideration in Critically Ill Patients', *American Journal of the Medical Sciences*. Elsevier Inc., 356(4), pp. 382–390. doi: 10.1016/j.amjms.2018.06.015.
- Boekstegers, P. *et al.* (1994) 'Skeletal muscle partial pressure of oxygen in patients with sepsis', *Critical Care Medicine*, 22, pp. 640-650.
- Campbell, K. *et al.* (2016) 'The Management of Nausea and Vomiting of Pregnancy', *Journal of Obstetrics and Gynaecology Canada*. Elsevier Inc, 38(12), pp. 1127–1137. doi: 10.1016/j.jogc.2016.08.009.
- COON, W. W. and WOLFMAN, E. F. (1961) 'Thiamine deficiency in surgical patients.', *Annals of surgery*, 154. doi: 10.1097/00000658-196112000-00031.
- Corcoran, TB. *et al.* (2009) 'Prevalence of vitamin deficiencies on admission: relationship to hospital mortality in critically ill patients', *Anaesth Intensive Care*, 37, pp. 254-260.
- Costa, N. A. *et al.* (2014) 'Serum thiamine concentration and oxidative stress as predictors of mortality in patients with septic shock', *Journal of Critical Care*, 29, pp. 249-252.
- Crook, M. A. and Sriram, K. (2014) 'Thiamine deficiency: The importance of recognition and prompt management', *Nutrition*. Elsevier Inc., 30(7–8), pp. 953–954. doi: 10.1016/j.nut.2014.03.003.
- Cruikshank, AM. *et al.* (1988) 'Thiamine deficiency in the critically ill', *Intensive Care Med*, 14(4), pp. 384-387.
- Donnino, M. W. *et al.* (2010) 'Thiamine deficiency in critically ill patients with sepsis', *Journal of Critical Care*. Elsevier Inc., 25(4), pp. 576–581. doi: 10.1016/j.jcrc.2010.03.003.
- Ferguson, T. I. *et al.* (2014) 'A review of stability issues associated with vitamins in parenteral nutrition', *e-SPEN Journal*. Elsevier Ltd, 9(2), pp. e49–e53.

doi: 10.1016/j.clnme.2014.01.001.

- Filho, R.R. *et al.* (2016) 'Blood lactate levels cutoff and mortality prediction in sepsis - time for a reappraisal? A retrospective cohort study', *SHOCK*, 46(5), pp. 480-485.
- Gattinoni, L. *et al.* (2015) 'Sepsis: needs for defining severity', *Intensive Care Med*, 41(3), pp. 551-552.
- Kaukonen, K.M. *et al.* (2014) 'Mortality related to severe sepsis and septic shock among critically ill patients in Australia and New Zealand' *JAMA*, 311(13), pp. 1308-1316.
- Kalantari, A. and Rezaie, S. R. (2019) 'Challenging the one-hour sepsis bundle', *Western Journal of Emergency Medicine*, 20(2), pp. 185–190. doi: 10.5811/westjem.2018.11.39290.
- Khan, P. and Divatia, J. V. (2010) 'Severe sepsis bundles', *Indian Journal of Critical Care Medicine*, 14(1), pp. 8–13. doi: 10.4103/0972-5229.63028.
- Leite, H. P. *et al.* (2017) 'Effect of blood thiamine concentrations on mortality: Influence of nutritional status', *Nutrition*. Elsevier Inc, 48, pp. 105-110. doi: 10.1016/j.nut.2017.11.020.
- Leppäniemi, A. *et al.* (2015) 'Management of abdominal sepsis - A paradigm shift?', *Anaesthesiology Intensive Therapy*, 47(4), pp. 400–408. doi: 10.5603/AIT.a2015.0026.
- Lie, K.C. *et al.* (2018) 'Utility of SOFA score, management and outcomes of sepsis in Southeast Asia: a multinational multicenter prospective observational study', *Journal of Intensive Care*, 6(9), pp. 1-8.
- Maiorana, A. *et al.* (2014) 'Acute thiamine deficiency and refeeding syndrome: Similar findings but different pathogenesis', *Nutrition*, 30(7–8), pp. 948–952. doi: 10.1016/j.nut.2014.02.019.
- Manzanares, W. and Hardy, G. (2011) 'Thiamine supplementation in the critically ill', *Curr Opin Clin Nutr Metab Care*, 14:610-617.
- Meena, S. and Gupta, A. (2014) 'Dorsal wrist ganglion: Current review of literature', *Journal of Clinical Orthopaedics and Trauma*. Elsevier Ltd, 5(2), pp. 59–64. doi: 10.1016/j.jcot.2014.01.006.
- Michael W. Donnino, M.D.1, 2, Lars W. Andersen, M.D.1, 3, Maureen Chase, M.D., M.P.H.1, Katherine M. Berg, M.D.2, Mark Tidswell, M.D.4, Tyler Giberson, B.S.1, Richard Wolfe, M.D.1, Ari Moskowitz, M.D.5, Howard Smithline, M.D.6, Long Ngo, Ph.D.5, and Micha, M. . (2017) 'Randomized, Double-Blind, Placebo-Controlled Trial of Thiamine as a Metabolic Resuscitator in Septic Shock: A Pilot Study', *Physiology & behavior*,

- 176(3), pp. 139–148. doi: 10.1016/j.physbeh.2017.03.040.
- Moskowitz, A. *et al.* (2018) 'Ascorbic acid, corticosteroids, and thiamine in sepsis: a review of the biologic rationale and the present state of clinical evaluation', *Critical Care*. *Critical Care*, 22(1), pp. 1–7. doi: 10.1186/s13054-018-2217-4.
- Ozasa, H. *et al.* (2006) 'Clinical examination of the water-soluble vitamin levels in blood during peripheral parenteral nutrition', *Kurume Medical Journal*, 53(3–4), pp. 79–87. doi: 10.2739/kurumemedj.53.79.
- Pertkiewicz, M. *et al.* (2009) 'Basics in clinical nutrition: Composition of nutritional admixtures and formulas for parenteral nutrition', *e-SPEN*, 4(4), pp. 161–163. doi: 10.1016/j.eclnm.2009.01.003.
- Ratih, E.P. *et al.* (2017) 'Prognostic accuracy of the sofa score, sirs kriteria, and qsofa score for in-hospital mortality among adults with suspected infection admitted to the intensive care unit', *JAMA*, 317, pp. 290-300.
- Rhodes, A. *et al.* (2017) *Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016, Intensive Care Medicine*. Springer Berlin Heidelberg. doi: 10.1007/s00134-017-4683-6.
- Ross, J. T., Matthay, M. A. and Harris, H. W. (2018) 'Secondary peritonitis: Principles of diagnosis and intervention', *BMJ (Online)*, 361. doi: 10.1136/bmj.k1407.
- Sartelli, M. *et al.* (2014) 'Current concept of abdominal sepsis: WSES position paper', *World Journal of Emergency Surgery*, 9(1), pp. 1–16. doi: 10.1186/1749-7922-9-22.
- Seear, M. *et al.* (1992) 'Thiamine, riboflavine, and pyridoxine deficiencies in a population of critically ill children', *J Pediatr*, 121(4), pp. 533-538.
- Shin, T. G. *et al.* (2019) 'Early Vitamin C and Thiamine Administration to Patients with Septic Shock in Emergency Departments: Propensity Score-Based Analysis of a Before-and-After Cohort Study', *Journal of Clinical Medicine*, 8(1), p. 102. doi: 10.3390/jcm8010102.
- Singer, P. *et al.* (2009) 'ESPEN Guidelines on Parenteral Nutrition: Intensive care', *Clinical Nutrition*. Elsevier Ltd, 28(4), pp. 387–400. doi: 10.1016/j.clnu.2009.04.024.
- Siregar, E.S. *et al.* (2016) 'Lactate clearance: predictor for mortality and therapeutic response on severe sepsis patient', *Bali Medical Journal*, 5(2), pp. 292-294.
- Sriram, K. and Lonchyna, V. A. (2009) 'Micronutrient supplementation in adult nutrition therapy: Practical considerations', *Journal of Parenteral and*

*Enteral Nutrition*, 33(5), pp. 548–562. doi: 10.1177/0148607108328470.

Veličković, J. *et al.* (2019) ‘The association of early postoperative lactate levels with morbidity after elective major abdominal surgery’, *Bosnian Journal of Basic Medical Sciences*, 19(1), pp. 72–80. doi: 10.17305/BJBMS.2018.3186.

Weimann, A. *et al.* (2017) ‘ESPEN guideline: Clinical nutrition in surgery’, *Clinical Nutrition*. Elsevier Ltd, 36(3), pp. 623–650. doi: 10.1016/j.clnu.2017.02.013.

Wischmeyer, P. E. (2018) ‘Nutrition Therapy in Sepsis’, *Critical Care Clinics*. Elsevier Inc, 34(1), pp. 107–125. doi: 10.1016/j.ccc.2017.08.008.