

## DAFTAR PUSTAKA

- ACSM. (2013). *ACSM's guidelines for exercise testing and prescription* (M. P. Bayles & A. M. Swank Eds.): Lippincott Williams & Wilkins.
- ACSM. (2016). *American College of Sport Medicine's Guidelines for Exercise Testing and Prescription* (L. S. Pescatello, R. Arena, D. Riebe, & P. D. Thompson Eds. 9 ed.). Philadelphia: Lippincott Williams & Wilkins.
- Akiskal, H. S. (2009a). Mood Disorders: Clinical Features. In B. J. Sadock, V. A. Sadock, & P. Ruiz (Eds.), *Kaplan & Saddock's Comprehensive Textbook of Psychiatry* (9 ed., Vol. 1, pp. 1693-1733). Philadelphia: Lippincott Williams & Wilkins.
- Akiskal, H. S. (2009b). Mood Disorders: Historical Introduction and Conceptual Overview. In B. J. Sadock, V. A. Sadock, & P. Ruiz (Eds.), *Kaplan & Saddock's Comprehensive Textbook of Psychiatry* (9 ed., Vol. 1, pp. 1629-1645). Philadelphia: Lippincott Williams & Wilkins.
- Amir, N. (2005). *Depresi: Aspek Neurobiologi, Diagnosis dan Tatalaksana*. Jakarta: Balai Penerbit FKUI.
- Berlim, M. T., den Eynde Van, F., dan Daskalakis, Z. J. (2013). High-frequency repetitive transcranial magnetic stimulation accelerates and enhances the clinical response to antidepressants in major depression: a meta-analysis of randomized, double-blind, and sham-controlled trials.
- Blair, M., Coleman, K., Jesso, S., Jodoin, V. D., Smolewska, K., Warriner, E., Finger, E., dan Pasternak, S. H. (2016). Depressive symptoms negatively impact Montreal Cognitive Assessment performance: a memory clinic experience. *Canadian Journal of Neurological Sciences*, 43(4), 513-517.
- Bretlau, L., Lunde, M., Lindberg, L., Uden, M., Dissing, S., dan Bech, P. (2008). Repetitive transcranial magnetic stimulation (rTMS) in combination with escitalopram in patients with treatment-resistant major depression. A double-blind, randomised, sham-controlled trial. *Pharmacopsychiatry*, 41(02), 41-47.
- Bruce, M. L., Seeman, T. E., Merrill, S. S., dan Blazer, D. G. (1994). The impact of depressive symptomatology on physical disability: MacArthur Studies

- of Successful Aging. *American Journal of Public Health*, 84(11), 1796-1799.
- Channon, S., dan Green, P. (1999). Executive function in depression: the role of performance strategies in aiding depressed and non-depressed participants. *Journal of Neurology, Neurosurgery & Psychiatry*, 66(2), 162-171.
- Demitrack, M. A., dan Brock, D. G. (2014). A Review of Current Clinical Practice in the Treatment of Major Depression. In A. Rotenberg, C. Horvath, & A. Pascual-Leone (Eds.), *Springer Protocols: Transcranial Magnetic Stimulation* (pp. 293-311). New York: Humana Press.
- Dimeo, F., Bauer, M., Varahram, I., Proest, G., dan Halter, U. (2001). Benefits from aerobic exercise in patients with major depression: a pilot study. *British journal of sports medicine*, 35(2), 114-117.
- Drevets, W. C. (2000). Neuroimaging studies of mood disorders. *Biological psychiatry*, 48(8), 813-829.
- Dwivedi, Y., Rizavi, H. S., Conley, R. R., Roberts, R. C., Tamminga, C. A., dan Pandey, G. N. (2003). Altered gene expression of brain-derived neurotrophic factor and receptor tyrosine kinase B in postmortem brain of suicide subjects. *Archives of general psychiatry*, 60(8), 804-815.
- Elliott, R. (1998). The neuropsychological profile in unipolar depression. *Trends in cognitive sciences*, 2(11), 447-454.
- Erdos, J., Inaki, I.-G., dan Ibarгойen-Roteta, N. (2017). *Repetitive transcranial magnetic stimulation for treatment resistant major depression: Rapid assessment of other technologies using the HTA Core Model for Rapid Relative Effective Assessment*. Vienna: Ludwig Boltzmann Institut.
- Erickson, K. I., Voss, M. W., Prakash, R. S., Basak, C., Szabo, A., Chaddock, L., Kim, J. S., Heo, S., Alves, H., dan White, S. M. (2011). Exercise training increases size of hippocampus and improves memory. *Proceedings of the National Academy of Sciences*, 108(7), 3017-3022.
- Fitzgerald, P. B., dan Daskalakis, Z. J. (2013). *Repetitive transcranial magnetic stimulation treatment for depressive disorders: A practical guide*: Springer Science & Business Media.

- Freitas, S., Prieto, G., Simões, M. R., dan Santana, I. (2014). Psychometric properties of the Montreal Cognitive Assessment (MoCA): an analysis using the Rasch model. *The Clinical Neuropsychologist*, 28(1), 65-83.
- Fujimura, H., Altar, C. A., Chen, R., Nakamura, T., Nakahashi, T., Kambayashi, J.-i., Sun, B., dan Tandon, N. N. (2002). Brain-derived neurotrophic factor is stored in human platelets and released by agonist stimulation. *Thrombosis and haemostasis*, 87(04), 728-734.
- Garcia-Toro, M., Pascual-Leone, A., Romera, M., Gonzalez, A., Mico, J., Ibarra, O., Arnillas, H., Capllonch, I., Mayol, A., dan Tormos, J. (2001). Prefrontal repetitive transcranial magnetic stimulation as add on treatment in depression. *Journal of Neurology, Neurosurgery & Psychiatry*, 71(4), 546-548.
- Gomez-Pinilla, F., Vaynman, S., dan Ying, Z. (2008). Brain-derived neurotrophic factor functions as a metabotrophin to mediate the effects of exercise on cognition. *European Journal of Neuroscience*, 28(11), 2278-2287.
- Hasler, G. (2010). pathophysiology of depression: do we have any solid evidence of interest to clinicians? *Official Journal of the World Psychiatric Association*, 9, 155-161.
- Hilberg, T., Menzel, K., Gläser, D., Zimmermann, S., dan Gabriel, H. H. W. (2008). Exercise intensity: platelet function and platelet-leukocyte conjugate formation in untrained subjects. *Thrombosis research*, 122(1), 77-84.
- Huang, M.-l., Luo, B.-y., Hu, J.-b., Wang, S.-S., Zhou, W.-h., Wei, N., Hu, S.-h., dan Xu, Y. (2012). Repetitive transcranial magnetic stimulation in combination with citalopram in young patients with first-episode major depressive disorder: a double-blind, randomized, sham-controlled trial. *Australian & New Zealand Journal of Psychiatry*, 46(3), 257-264.
- Husein, N., Lumempouw, S., dan Ramli, Y. (2010). Montreal cognitive assessment versi indonesia (moca-ina) untuk skrining gangguan fungsi kognitif. *Neurona (Majalah Kedokteran Neuro Sains Perhimpunan Dokter Spesialis Saraf Indonesia)*, 27(4).

- Istrianana, E., Kurnia, A., Weijers, A., Hidayat, T., Pinxten, L., de Jong, C., dan Schellekens, A. (2013). Excellent reliability of the Hamilton Depression Rating Scale (HDRS-21) in Indonesia after training. *Asia-Pacific Psychiatry*, 5(3), 141-146.
- Jaeger, J., Berns, S., Uzelac, S., dan Davis-Conway, S. (2006). Neurocognitive deficits and disability in major depressive disorder. *Psychiatry research*, 145(1), 39-48.
- Kamijo, K., Hayashi, Y., Sakai, T., Yahiro, T., Tanaka, K., dan Nishihira, Y. (2009). Acute effects of aerobic exercise on cognitive function in older adults. *Journals of Gerontology: Series B*, 64(3), 356-363.
- Khatri, P., Blumenthal, J. A., Babyak, M. A., Craighead, W. E., Herman, S., Baldewicz, T., Madden, D. J., Doraiswamy, M., Waugh, R., dan Krishnan, K. R. (2001). Effects of exercise training on cognitive functioning among depressed older men and women. *Journal of aging and physical activity*, 9(1), 43-57.
- Kubesch, S., Bretschneider, V., Freudenmann, R., Weidenhammer, N., Lehmann, M., Spitzer, M., dan Gron, G. (2003). Aerobic endurance exercise improves executive functions in depressed patients. *Journal of Clinical Psychiatry*, 64(9), 1005-1012.
- Kusumawardhani, A., Amir, N., Maramis, M. M., Redayani, P., Desmiarti, dan Khamelia. (2013). *Panduan Gangguan Depresi Mayor*. Jakarta: PDSKJI.
- Laske, C., Banschbach, S., Stransky, E., Bosch, S., Straten, G., Machann, J., Fritsche, A., Hipp, A., Niess, A., dan Eschweiler, G. W. (2010). Exercise-induced normalization of decreased BDNF serum concentration in elderly women with remitted major depression. *International Journal of Neuropsychopharmacology*, 13(5), 595-602.
- Levkovitz, Y., Caftori, R., Avital, A., dan Richter-Levin, G. (2002). The SSRI drug Fluoxetine, but not the noradrenergic tricyclic drug Desipramine, improves memory performance during acute major depression. *Brain research bulletin*, 58(4), 345-350.

- Loh, A. Z., Tan, J. S., Zhang, M. W., dan Ho, R. C. (2017). The global prevalence of anxiety and depressive symptoms among caregivers of stroke survivors. *Journal of the American Medical Directors Association*, 18(2), 111-116.
- Maslim, R. (2013). *Buku Saku Diagnosis Gangguan Jiwa: Rujukan Ringkas dari PPDGJ-III dan DSM-5*. Jakarta: Bagian Ilmu Kedokteran Jiwa FK Unika Atmajaya.
- McClintock, S. M., Reti, I. M., Carpenter, L. L., McDonald, W. M., Dubin, M., Taylor, S. F., Cook, I. A., O'Reardon, J., Husain, M. M., dan Wall, C. (2018). Consensus recommendations for the clinical application of repetitive transcranial magnetic stimulation (rTMS) in the treatment of depression. *The Journal of clinical psychiatry*, 79(1).
- McIntyre, R. S., Cha, D. S., Soczynska, J. K., Woldeyohannes, H. O., Gallagher, L. A., Kudlow, P., Alsuwaidan, M., dan Baskaran, A. (2013). Cognitive deficits and functional outcomes in major depressive disorder: determinants, substrates, and treatment interventions. *Depression and anxiety*, 30(6), 515-527.
- McMorris, T. C., Jo. (2018). Neurobiological Changes as an Explanation of Benefits of Exercise. In H. Budde & M. Wegner (Eds.), *The Exercise Effect of Mental Health: Neurobiological Mechanisms* (pp. 30-66). New York: Taylor & Francis.
- Mikkelsen, K., Stojanovska, L., Polenakovic, M., Bosevski, M., dan Apostolopoulos, V. (2017). Exercise and mental health. *Maturitas*, 106, 48-56.
- Moirand, R., Galvao, F., Lecompte, M., Poulet, E., Haesebaert, F., dan Brunelin, J. (2018). Usefulness of the Montreal Cognitive Assessment (MoCA) to monitor cognitive impairments in depressed patients receiving electroconvulsive therapy. *Psychiatry research*, 259, 476-481.
- Moreau, D. (2018). Physical Exercise and Cognitive Enhancement. In H. Budde & M. Wegner (Eds.), *The Exercise Effect of Mental Health: Neurobiological Mechanisms* (pp. 171-187). New York: Taylor & Francis.

- Moser, D., Jorge, R., Manes, F., Paradiso, S., Benjamin, M., dan Robinson, R. (2002). Improved executive functioning following repetitive transcranial magnetic stimulation. *Neurology*, *58*(8), 1288-1290.
- Najib, U., dan Horvath, C. (2014). Transcranial Magnetic Stimulation (TMS) Safety Considerations and Recommendations. In A. Rotenberg, J. C. Horvath, & A. Pascual-Leone (Eds.), *Transcranial Magnetic Stimulation* (pp. 15-30). New York: Springer.
- Nasreddine, Z. S., Phillips, N. A., Bédirian, V., Charbonneau, S., Whitehead, V., Collin, I., Cummings, J. L., dan Chertkow, H. (2005). The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*, *53*(4), 695-699.
- Neurosoft. (2014). Neuro MS/D Transcranial Magnetic Stimulation. Ivanovo: Neurosoft.
- NICE. (2015). NICE Interventional Procedure Guidance: Repetitive Transcranial Magnetic Stimulation for Depression: National Institute for Health and Care Excellence.
- O'Connor, M. G., Jerskey, B. A., Robertson, E. M., Brenninkmeyer, C., Ozdemir, E., dan Leone, A. P. (2005). The effects of repetitive transcranial magnetic stimulation (rTMS) on procedural memory and dysphoric mood in patients with major depressive disorder. *Cognitive and behavioral neurology*, *18*(4), 223-227.
- Oberman, L. (2014). Repetitive Transcranial Magnetic Stimulation (rTMS) Protocols. In A. Rotenberg, C. Horvath, & A. Pascual-Leone (Eds.), *Springer Protocols: Transcranial Magnetic Stimulation* (pp. 129-139). New York: Humana Press.
- Ogborn, D. I., dan Gardiner, P. F. (2010). Effects of exercise and muscle type on BDNF, NT-4/5, and TrKB expression in skeletal muscle. *Muscle & Nerve: Official Journal of the American Association of Electrodiagnostic Medicine*, *41*(3), 385-391.
- Rasmussen, P., Brassard, P., Adser, H., Pedersen, M. V., Leick, L., Hart, E., Secher, N. H., Pedersen, B. K., dan Pilegaard, H. (2009). Evidence for a

- release of brain-derived neurotrophic factor from the brain during exercise. *Experimental physiology*, 94(10), 1062-1069.
- Richter-Levin, G., dan Segal, M. (1996). Serotonin, Aging and Cognitive Functions of the Hippocampus. *Reviews in the Neurosciences*, 7(2), 103-114.
- Rossi, S., Hallett, M., Rossini, P. M., Pascual-Leone, A., dan Group, S. o. T. C. (2009). Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clinical neurophysiology*, 120(12), 2008-2039.
- Rossini, D., Magri, L., Lucca, A., Giordani, S., Smeraldi, E., dan Zanardi, R. (2005). Does rTMS hasten the response to escitalopram, sertraline, or venlafaxine in patients with major depressive disorder? A double-blind, randomized, sham-controlled trial. *The Journal of clinical psychiatry*, 66(12), 1569-1575.
- Rush, A. J., dan Nierenberg, A. A. (2009). Mood Disorders: Treatment of Depression. In B. J. Sadock, V. A. Sadock, & P. Ruiz (Eds.), *Kaplan & Saddock's Comprehensive Textbook of Psychiatry* (9 ed., Vol. 1, pp. 1734-1743). Philadelphia: Lippincott Williams & Wilkins.
- Savitz, J. B., dan Drevets, W. C. (2009). Brain Circuit in Major Depressive Disorder. In B. J. Sadock, V. A. Sadock, & P. Ruiz (Eds.), *Kaplan & Saddock's Comprehensive Textbook of Psychiatry* (9 ed., Vol. 1, pp. 1675-1685). Philadelphia: Lippincott Williams & Wilkins.
- Sehatzadeh, S., Tu, H. A., Palimaka, S., Yap, B., O'Reilly, D., dan Bowen, J. (2016). Repetitive transcranial magnetic stimulation for treatment-resistant depression: a systematic review and meta-analysis of randomized controlled trials. *Ontario health technology assessment series*, 16(5), 1.
- Sheline, Y. I. (2003). Neuroimaging studies of mood disorder effects on the brain. *Biological psychiatry*, 54(3), 338-352.
- Spampinato, C., Aguglia, E., Concerto, C., Pennisi, M., Lanza, G., Bella, R., Cantone, M., Pennisi, G., Kavasidis, I., dan Giordano, D. (2013). Transcranial magnetic stimulation in the assessment of motor cortex excitability and treatment of drug-resistant major depression. *IEEE*

- Transactions on Neural Systems and Rehabilitation Engineering*, 21(3), 391-403.
- Springer, B. A., Marin, R., Cyhan, T., Roberts, H., dan Gill, N. W. (2007). Normative values for the unipedal stance test with eyes open and closed. *Journal of geriatric physical therapy*, 30(1), 8-15.
- Srisurapanont, M., Eurviriyankul, K., Suttajit, S., dan Varnado, P. (2017). Internal consistency and concurrent validity of the Montreal Cognitive Assessment in individuals with major depressive disorder. *Psychiatry research*, 253, 333-337.
- Stahl, S. M. (2013a). Antidepressants *Stahl's essential psychopharmacology: neuroscientific basis and practical applications* (pp. 284-369). Cambridge: Cambridge university press.
- Stahl, S. M. (2013b). Mood disorder *Stahl's essential psychopharmacology: neuroscientific basis and practical applications* (pp. 237-283). Cambridge: Cambridge university press.
- Stahl, S. M. (2013c). Psychosis and Schizophrenia *Stahl's essential psychopharmacology: neuroscientific basis and practical applications* (pp. 79-128). Cambridge: Cambridge university press.
- Stanton, R., dan Reaburn, P. (2014). Exercise and the treatment of depression: a review of the exercise program variables. *Journal of Science and Medicine in Sport*, 17(2), 177-182.
- Su, T.-P., Huang, C.-C., dan Wei, I.-H. (2005). Add-on rTMS for medication-resistant depression: a randomized, double-blind, sham-controlled trial in Chinese patients. *The Journal of clinical psychiatry*, 66(7), 930-937.
- Tam, N. D. (2013). Improvement of processing speed in executive function immediately following an increase in cardiovascular activity. *Cardiovascular psychiatry and neurology*, 2013.
- Thase, M. E. (2009). Mood Disorders: Neurobiology. In B. J. Sadock, V. A. Sadock, & P. Ruiz (Eds.), *Kaplan & Saddock's Comprehensive Textbook of Psychiatry* (9 ed., Vol. 1, pp. 1664-1674). Philadelphia: Lippincott Williams & Wilkins.



- Tsai, J.-C., Chen, C.-W., Chu, H., Yang, H.-L., Chung, M.-H., Liao, Y.-M., dan Chou, K.-R. (2016). Comparing the sensitivity, specificity, and predictive values of the Montreal Cognitive Assessment and Mini-Mental State Examination when screening people for mild cognitive impairment and dementia in Chinese population. *Archives of psychiatric nursing*, 30(4), 486-491.
- Vasques, P. E., Moraes, H., Silveira, H., Deslandes, A. C., dan Laks, J. (2011). Acute exercise improves cognition in the depressed elderly: the effect of dual-tasks. *Clinics*, 66(9), 1553-1557.
- Wang, H.-Y., Crupi, D., Liu, J., Stucky, A., Cruciatu, G., Di Rocco, A., Friedman, E., Quartarone, A., dan Ghilardi, M. F. (2011). Repetitive transcranial magnetic stimulation enhances BDNF–TrkB signaling in both brain and lymphocyte. *Journal of Neuroscience*, 31(30), 11044-11054.
- Wassermann, E. M., dan Zimmermann, T. (2012). Transcranial magnetic brain stimulation: therapeutic promises and scientific gaps. *Pharmacology & therapeutics*, 133(1), 98-107.
- WHO. (2008). *The Global Burden of Disease: 2004 Update*: WHO.
- WHO. (2017) Depression and Other Common Mental Disorders: Global Health Estimates. Geneva: WHO.