

DAFTAR PUSTAKA

- ACSM. (2013). *ACSM's guidelines for exercise testing and prescription* (M. P. Bayles & A. M. Swank Eds.): Lippincott Williams & Wilkins.
- Akiskal, H. S. (2009). Mood Disorders: Clinical Features. In B. J. Sadock, V. A. Sadock, & P. Ruiz (Eds.), *Kaplan & Saddock's Comprehensive Textbook of Psychiatry* (9 ed., Vol. 1, pp. 1693-1733). Philadelphia: Lippincott Williams & Wilkins.
- Barge-Schaapveld, D. Q., Nicolson, N. A., Berkhof, J., & Marten, W. d. (1999). Quality of life in depression: daily life determinants and variability. *Psychiatry research*, 88(3), 173-189.
- Beusterien, K. M., Steinwald, B., & Ware Jr, J. E. (1996). Usefulness of the SF-36 Health Survey in measuring health outcomes in the depressed elderly. *Journal of geriatric psychiatry and neurology*, 9(1), 13-21.
- Demitrack, M. A., & Brock, D. G. (2014). A Review of Current Clinical Practice in the Treatment of Major Depression. In A. Rotenberg, C. Horvath, & A. Pascual-Leone (Eds.), *Springer Protocols: Transcranial Magnetic Stimulation* (pp. 293-311). New York: Humana Press.
- Dimeo, F., Bauer, M., Varahram, I., Proest, G., & Halter, U. (2001). Benefits from aerobic exercise in patients with major depression: a pilot study. *British journal of sports medicine*, 35(2), 114-117.
- Erdos, J., Inaki, I.-G., & Ibargoyen-Roteta, N. (2017). *Repetitive transcranial magnetic stimulation for treatment resistant major depression: Rapid assessment of other technologies using the HTA Core Model for Rapid Relative Effective Assessment*. Vienna: Ludwig Boltzmann Institut.
- Erickson, K. I., Voss, M. W., Prakash, R. S., Basak, C., Szabo, A., Chaddock, L., . . . White, S. M. (2011). Exercise training increases size of hippocampus and improves memory. *Proceedings of the National Academy of Sciences*, 108(7), 3017-3022.
- Fitzgerald, P. B., & Daskalakis, Z. J. (2013). *Repetitive transcranial magnetic stimulation treatment for depressive disorders: A practical guide*: Springer Science & Business Media.
- Gomez-Pinilla, F., Vaynman, S., & Ying, Z. (2008). Brain-derived neurotrophic factor functions as a metabotrophin to mediate the effects of exercise on cognition. *European Journal of Neuroscience*, 28(11), 2278-2287.
- Istriana, E., Kurnia, A., Weijers, A., Hidayat, T., Pinxten, L., de Jong, C., & Schellekens, A. (2013). Excellent reliability of the Hamilton Depression Rating Scale (HDRS-21) in Indonesia after training. *Asia-Pacific Psychiatry*, 5(3), 141-146.
- Kamijo, K., Hayashi, Y., Sakai, T., Yahiro, T., Tanaka, K., & Nishihira, Y. (2009). Acute effects of aerobic exercise on cognitive function in older adults. *Journals of Gerontology: Series B*, 64(3), 356-363.
- Kusumawardhani, A., Amir, N., Maramis, M. M., Redayani, P., Desmiarti, & Khamelia. (2013). *Panduan Gangguan Depresi Mayor*. Jakarta: PDSKJI.
- Laske, C., Banschbach, S., Stransky, E., Bosch, S., Straten, G., Machann, J., . . . Eschweiler, G. W. (2010). Exercise-induced normalization of decreased

- BDNF serum concentration in elderly women with remitted major depression. *International Journal of Neuropsychopharmacology*, 13(5), 595-602.
- Loh, A. Z., Tan, J. S., Zhang, M. W., & Ho, R. C. (2017). The global prevalence of anxiety and depressive symptoms among caregivers of stroke survivors. *Journal of the American Medical Directors Association*, 18(2), 111-116.
- Moreau, D. (2018). Physical Exercise and Cognitive Enhancement. In H. Budde & M. Wegner (Eds.), *The Exercise Effect of Mental Health: Neurobiological Mechanisms* (pp. 171-187). New York: Taylor & Francis.
- Najib, U., & Horvath, C. (2014). Transcranial Magnetic Stimulation (TMS) Safety Considerations and Recommendations. In A. Rotenberg, J. C. Horvath, & A. Pascual-Leone (Eds.), *Transcranial Magnetic Stimulation* (pp. 15-30). New York: Springer.
- Neurosoft. (2014). Neuro MS/D Transcranial Magnetic Stimulation. Ivanovo: Neurosoft.
- Oberman, L. (2014). Repetitive Transcranial Magnetic Stimulation (rTMS) Protocols. In A. Rotenberg, C. Horvath, & A. Pascual-Leone (Eds.), *Springer Protocols: Transcranial Magnetic Stimulation* (pp. 129-139). New York: Humana Press.
- Rossi, S., Hallett, M., Rossini, P. M., Pascual-Leone, A., & Group, S. o. T. C. (2009). Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clinical neurophysiology*, 120(12), 2008-2039.
- Rush, A. J., & Nierenberg, A. A. (2009). Mood Disorders: Treatment of Depression. In B. J. Sadock, V. A. Sadock, & P. Ruiz (Eds.), *Kaplan & Suddock's Comprehensive Textbook of Psychiatry* (9 ed., Vol. 1, pp. 1734-1743). Philadelphia: Lippincott Williams & Wilkins.
- Schuch, F., Vasconcelos-Moreno, M., Borowsky, C., Zimmermann, A., Rocha, N., & Fleck, M. (2015). Exercise and severe major depression: effect on symptom severity and quality of life at discharge in an inpatient cohort. *Journal of psychiatric research*, 61, 25-32.
- Solvason, H. B., Husain, M., Fitzgerald, P. B., Rosenquist, P., McCall, W. V., Kimball, J., . . . Lisanby, S. H. (2014). Improvement in quality of life with left prefrontal transcranial magnetic stimulation in patients with pharmacoresistant major depression: acute and six month outcomes. *Brain stimulation*, 7(2), 219-225.
- Stahl, S. M. (2013a). Antidepressants *Stahl's essential psychopharmacology: neuroscientific basis and practical applications* (pp. 284-369). Cambridge: Cambridge university press.
- Stahl, S. M. (2013b). Mood disorder *Stahl's essential psychopharmacology: neuroscientific basis and practical applications* (pp. 237-283). Cambridge: Cambridge university press.
- Stanton, R., & Reaburn, P. (2014). Exercise and the treatment of depression: a review of the exercise program variables. *Journal of Science and Medicine in Sport*, 17(2), 177-182.
- Thase, M. E. (2009). Mood Disorders: Neurobiology. In B. J. Sadock, V. A. Sadock, & P. Ruiz (Eds.), *Kaplan & Suddock's Comprehensive Textbook of*

- Psychiatry* (9 ed., Vol. 1, pp. 1664-1674). Philadelphia: Lippincott Williams & Wilkins.
- Vasques, P. E., Moraes, H., Silveira, H., Deslandes, A. C., & Laks, J. (2011). Acute exercise improves cognition in the depressed elderly: the effect of dual-tasks. *Clinics*, 66(9), 1553-1557.
- Wada, T., Ishine, M., Sakagami, T., Kita, T., Okumiya, K., Mizuno, K., . . . Matsubayashi, K. (2005). Depression, activities of daily living, and quality of life of community-dwelling elderly in three Asian countries: Indonesia, Vietnam, and Japan. *Archives of gerontology and Geriatrics*, 41(3), 271-280.
- Wang, H.-Y., Crupi, D., Liu, J., Stucky, A., Cruciat, G., Di Rocco, A., . . . Ghilardi, M. F. (2011). Repetitive transcranial magnetic stimulation enhances BDNF-TrkB signaling in both brain and lymphocyte. *Journal of Neuroscience*, 31(30), 11044-11054.
- Ware, J., Kosinski, M., & Keller, S. (2001). SF-36 physical and mental health summary scales. *a user's manual*, 1994.