

**Malaysian Applied Biology**  
Malaysian Applied Biology  
Malaysian Applied Biology



Published by the Malaysian Society of Applied Biology  
ISSN 0025-7179

**Malaysian Applied Biology**  
Malaysian Applied Biology  
Malaysian Applied Biology



Published by the Malaysian Society of Applied Biology  
ISSN 0025-7179

## MAIN MENU

Home

Editorial Board

Journal Policy

Subscription

Instruction to Authors

Submission

F.A.Q.

Archive

## LATEST MABJ ISSUE

Vol 49(3) October 2020



Published by the Malaysian Soc  
www.mabjournal.com

ISSN 0126-8643

[Table of content](#)

Latest news!

Malaysian Applied Biology is listed in the databases and indexed in Thomson Reuters Master Journal List, Elsevier, Mycite, Biosis, Zoological Records, UDLedge Life Science

## Editorial Board



### Chief Editor:

**Prof. Dr. Wickneswari Ratnam**

Department of Biological Sciences and Biotechnology,  
Faculty of Science and Technology,

Universiti Kebangsaan Malaysia,  
UKM Bangi 43650, Selangor, Malaysia

Tel: 603 89213840

Fax: 603 89253357

Email: [wicki@ukm.edu.my](mailto:wicki@ukm.edu.my)

### Associate Editors:

**Prof. Dr. K Sudesh Kumar A/L C Kanapathi Pillai**

School of Biological Sciences,  
Universiti Sains Malaysia,  
USM Penang 11800, Penang, Malaysia

Email: [ksudesh@usm.my](mailto:ksudesh@usm.my)

**Prof. Dr. Aziz Ahmad**

Department of Biological Sciences,  
Faculty of Science and Technology,  
Universiti Malaysia Terengganu,  
UMT Kuala Terengganu 21030, Terengganu, Malaysia

Email: [aaziz@umt.edu.my](mailto:aaziz@umt.edu.my)

**Assoc. Prof. Dr. Nazlina Ibrahim**

Department of Biological Sciences and Biotechnology,  
Faculty of Science and Technology,  
Universiti Kebangsaan Malaysia,  
Bangi 43600, Selangor

Email: [nazlina@ukm.edu.my](mailto:nazlina@ukm.edu.my)

**Assoc. Prof. Dr. Ramlah Zainudin**

Centre for Pre-University Studies/Department of Zoology,  
Universiti Malaysia Sarawak (UNIMAS),  
94300 Kota Samarahan, Sarawak

Email: [zramlah@unimas.my](mailto:zramlah@unimas.my)

### Managing Editor:

**Dr. Sreelakshmi Sankara Narayanan**

Food Science and Nutrition Programme,  
School of Bioscience,  
Faculty of Health & Medical Sciences,  
Taylor's University Lakeside Campus,  
Petaling Jaya, Selangor, Malaysia

Email: [msabjournal@gmail.com](mailto:msabjournal@gmail.com)

Index, CNKI, J-Gate and  
CABI.

Malaysian Applied  
Biology is indexed in  
Scopus since issue  
41(1) 2012.

According to MyCite  
2014 report, MABJ  
ranked 95 out of 142  
Malaysian journals in  
terms of yearly impact  
factor.

**Journal Administrator:**

**Dr. Mohd Ikmal Asmuni**

Department of Biological Sciences and Biotechnology,  
Faculty of Science and Technology,  
Universiti Kebangsaan Malaysia,  
UKM Bangi 43650, Selangor, Malaysia  
Email: [mohdikmal@siswa.ukm.edu.my](mailto:mohdikmal@siswa.ukm.edu.my)

**\* All inquiries should be directed to the Chief Editor.**



**MAIN MENU**

Home  
 Editorial Board  
 Journal Policy  
 Subscription  
 Instruction to Authors  
 Submission  
 F.A.Q.  
 Archive

**LATEST MABJ ISSUE**

Vol 49(3) October 2020



Published by the Malaysian Soc  
 www.mabjournal.com

ISSN 0128/8643

[Table of content](#)**Latest news!**

Malaysian Applied Biology is listed in the databases and indexed in Thomson Reuters Master Journal List, Elsevier, Mycite, Biosis, Zoological Records, UDLedge Life Science

**Volume 48(3): June 2019****Journal of the Malaysian Society of Applied Biology**

Cover : "[Pollen grains under Scanning Electron Microscope of Byttneria species](#)". Photo courtesy of Noraini Talip.

**Review Article****Food Security and Sustainability: Malaysia Agenda**

*Tapsir, S., Engku Elini, E.A., Roslina, A., Noorlidawati, A.H., Mohd Hafizudin, Z., Hairazi, R. and Rosnani, H.*

| [Abstract](#) | [Full Text PDF](#) ( 69 KB) | pp. 1 |

**Research Articles****Curative potential of *Zingiber zerumbet* on paracetamol-induced hepatotoxicity**

*Nursirhan Athirah Sharkawi, Nur Hilwani Ismail, Nurul Husna Md Jani, Dzulsuhaimi Daud, Nooraain Hashim, Azani Salleh and Harita Hashim*

| [Abstract](#) | [Full Text PDF](#) ( 113 KB) | pp. 11 |

**Pollen morphology and harmomegathic characters of *Byttneria* Löfl. species (Sterculiaceae S. S: subfam. Byttnerioideae)**

*Amirul-Aiman, A.J., Noraini, T., Nurul-Aini, C.A.C., Chung, R.C.K., Phupumirat, W., Ruzi, A.R., Bunawan, H., Idris, S. and Suhaniza, R.*

| [Abstract](#) | [Full Text PDF](#) ( 984 KB) | pp. 19 |

**Chemical composition and potential adulterants in coconut milk sold in Kuala Lumpur**

*Shafrina Azlin-Hasim, Quee Lee Siang, Fadlina Yusof, Mohamad Khairi Zainol and Hayati Mohd Yusof*

| [Abstract](#) | [Full Text PDF](#) ( 154 KB) | pp. 27 |

**Antimicrobial resistance of *Escherichia coli* isolated of ulam from supermarkets and wet markets in Kuala Terengganu, Malaysia**

*Anis Athirah Bahri, Wahida Salleh, Mohd Nizam Lani and Wan Zawiah Wan Abdullah*

| [Abstract](#) | [Full Text PDF](#) ( 150 KB) | pp. 35 |

Index, CNKI, J-Gate and CABI.

Malaysian Applied Biology is indexed in Scopus since issue 41(1) 2012.

According to MyCite 2014 report, MABJ ranked 95 out of 142 Malaysian journals in terms of yearly impact factor.

### **Case-control retrospective study on cardiovascular disease risk and quality of life among Malay adults**

*Mo Min Erl, Asma' Ali, Arnie A. Rahim, Anam Aman, Sakinah Harith and Hayati Mohd Yusof*

| [Abstract](#) | [Full Text PDF](#) ( 117 KB) | pp. 43|

### **Optimization of enzymatic hydrolysis condition of Angelwing Clam (*Pholas orientalis*) meat using Alcalase® to obtain maximum degree of hydrolysis**

*Amiza Mat Amin and Soong Kang Cheng*

| [Abstract](#) | [Full Text PDF](#) ( 72 KB) | pp. 55|

### **Fish characteristics and distribution of potential fishing zones in the Malacca Strait using hydroacoustic assessment during the Southwest Monsoon season**

*Domu Simbolon, Bisman Nababan and La Elson*

| [Abstract](#) | [Full Text PDF](#) ( 875 KB) | pp. 63|

### **Enzyme inhibitory activities of marine sponges against cholinesterase and 5 $\alpha$ -reductase**

*Suciati, Karma Rabgay, Yunda Fachrunniza, Tongchai Saesong, Tri Aryono Hadi, Tutik Sri Wahyuni, Aty Widjawardiyanti and Kornkanok Ingkaninan*

| [Abstract](#) | [Full Text PDF](#) ( 230 KB) | pp. 77|

### **Comparison of genetic diversity of leptin gene between wild goat and domestic goat breeds in Iran**

*Narges Esmaeili Aliabad Vazir, Ali Esmaeilizadeh Koshkooieh, Ahmad Aytolahi Mehrjerdi, Mohammadreza Mohammadabadi, Olena Ivanivna Babenko, Maryna Vitaliivna Bushtruk, Serhii Vasyliovych Tkachenko, Ruslana Volodymyrivna Stavetska and Nataliia Ihorivna Klopenko*

| [Abstract](#) | [Full Text PDF](#) ( 479 KB) | pp. 85|

### **Contamination level and prevalence of foodborne pathogen *Enterobacteriaceae* in broiler and backyard chicken meats sold at traditional markets in Surabaya, Indonesia**

*Ratna Yulistiani, Danar Praseptiangga, Supyani and Sudibya*

| [Abstract](#) | [Full Text PDF](#) ( 409 KB) | pp. 95|

### **Anti-viral activity of *Phyllanthus niruri* against hepatitis C virus**

*Tutik Sri Wahyuni, Dzul Azmi, Adita Ayu Permanasari, Myrna Adianti, Lydia Tumewu, Tri Widiandani, Chie Aoki Utsubo, Aty Widjawardiyanti, Achmad Fuad and Hak Hotta*

| [Abstract](#) | [Full Text PDF](#) ( 198 KB) | pp. 105|

**Limnoecology and carp fish species peak spawning timing in haor basin of Bangladesh**

*Md. Foysal Ahmad, Muhammad Anamul Kabir, Zulhisyam Abdul Kari, Md. Tariqul Alam and Lee Seong Wei*

| [Abstract](#) | [Full Text PDF](#) ( 175 KB) | pp. 113|

**Molecular systematic position of the Sarawak Malay badger, *Mydaus javanensis***

*Badrul Munir Md-Zain, Nor Rahman Aifat, Muhammad Abu Bakar Abdul-Latiff, Nur Syafika Mohd-Yusof and Ahmad Ampeng*

| [Abstract](#) | [Full Text PDF](#) ( 360 KB) | pp. 125|

**Effect of light intensity on the photosynthesis and stomatal density of selected plant species of Gunung Ledang, Johor**

*Aisha Idris, Alona Cuevas Linatoc and Mohd Fadzelly Bin Abu Bakar*

| [Abstract](#) | [Full Text PDF](#) ( 193 KB) | pp. 133|

**Detection threshold and sensory preference for salty taste among Malay, Chinese and Indian students in Health Campus, Universiti Sains Malaysia**

*Marina, A.M., Nurul Husna, A.R. and Ruhaya, H.*

| [Abstract](#) | [Full Text PDF](#) ( 262 KB) | pp. 141|

**Production and toxicity evaluation of Rhamnolipids produced by *Pseudomonas* strains on L6 and Hepg2 cells**

*Arularasu Muthaliar Tamothran, Vigneswari Sevakumaran and Kesaven Bhubalan*

| [Abstract](#) | [Full Text PDF](#) ( 460 KB) | pp. 149|

**Characterization of linamarin-utilizing bacterial strains associated with detoxification of cyanogens in waste effluents**

*Ogunyemi Adewale, K., Samuel Titilola, A., Amund Olukayode, O. and Ilori Mathew O.*

| [Abstract](#) | [Full Text PDF](#) ( 610 KB) | pp. 157|

**Genetic diversity of the cave roosting dusky fruit bat, *Penthetor lucasi* from Sarawak**

*Yuvarajan Manivannan, Mohd Ridwan Abd Rahman, Roberta Chaya Tawie Tingga and Faisal Ali Anwarali Khan*

| [Abstract](#) | [Full Text PDF](#) ( 291 KB) | pp. 167|

**Leaf anatomical responses to drought stress condition in hybrid sugarcane leaf (*Saccharum officinarum* 'KK3')**

Worasitikulya Taratima, Thapakorn Ritmaha, Nuntawoot Jongrunklang,  
Sayam Raso and Pitakpong Maneerattanarungroj

| [Abstract](#) | [Full Text PDF](#) ( 1,242 KB) | pp. 181|

**Antifungal activity of dichloromethane and hexane extracts of four  
Malaysian seaweed species against *Ganoderma boninense***

Syamimi Diyana Abdul Aziz, Nur Fazirah Jafarah, Suriana Sabri, Mohd.  
As'wad Abdul Wahab and Zetty Norhana Balia Yusof

| [Abstract](#) | [Full Text PDF](#) ( 388 KB) | pp. 189|

---

CSS  XHTML



## Indexed Review Journal

Semin Thromb Hemost is covered in main abstracting and indexing services worldwide

[thieme.com](#)

[OPEN](#)

## Malaysian Applied Biology

Country [Malaysia](#) -  SCIMAGO INSTITUTIONS RANKINGS

# 6

Subject Area and Category [Agricultural and Biological Sciences](#)  
[Agricultural and Biological Sciences \(miscellaneous\)](#)

Publisher [Malaysian Society of Applied Biology](#)

H Index

Publication type [Journals](#)

ISSN [01268643](#)

Coverage [1980, 1986-1987, 2012-2019](#)

**Scope** MABJ considers all fields of biology and applied biology. There are basically two types of articles: Full research article and research note. A research note is a short communication that doesn't contain abstract and section subheadings. The reviewers can also suggest that a manuscript to be published as research note if they feel that the research is not adequate for a full paper. We occasionally publish review article by invitation and research articles from MSAB Symposium.

 [Homepage](#)

[How to publish in this journal](#)

[Contact](#)

 [Join the conversation about this journal](#)

## Indexed Review Journal

Semin Thromb Hemost is covered in main abstracting and indexing services worldwide

[thieme.com](#)

[OPEN](#)




# Thieme Publishers

The journal has an impact factor and is covered in main abstracting and indexing services


[thieme.com](http://thieme.com)

OPEN


## Quartiles




## SJR




## Citations per document




## Total Cites Self-Cites



## External Cites per Doc



## % International Collaboration



## Citable documents

## Non-citable documents

## Cited documents

## Uncited documents

**Malaysian Applied Biology**← Show this widget in  
your own websiteJust copy the code below  
and paste within your html  
code:`<a href="https://www.scimagojr.com" data-bbox="259 196 408 209">`

## Submit your manuscript - Read now

This journal is Open Access [thieme.com](http://thieme.com)

Metrics based on Scopus® data as of April 2020

**Dr. Laurensius Lehar** 1 year ago

Can the title below be included in Malaysian Applied Biology.

"POTENTIAL OF CROP IN INCREASING SOIL ORGANIC CARPON LEGUM COVER IN SUSTAINABLE DRYLANDS" please inform me via email. Thank you

reply

**Dwi Rosalina** 2 years ago

how much fee publication if i want submit my article in this journal?

thank you

best regards

reply

**Leave a comment**

Name

Email

(will not be published)

## ANTI-VIRAL ACTIVITY OF *Phyllanthus niruri* AGAINST HEPATITIS C VIRUS

TUTIK SRI WAHYUNI<sup>1,2\*</sup>, DZUL AZMI<sup>1</sup>, ADITA AYU PERMANASARI<sup>2</sup>, MYRNA ADIANTI<sup>2,4</sup>, LYDIA TUMEWU<sup>2</sup>, TRI WIDIANDANI<sup>3</sup>, CHIE AOKI UTSUBO<sup>5</sup>, ATY WIDYAWARUYANTI<sup>1,2</sup>, ACHMAD FUAD<sup>1,2</sup> and HAK HOTTA<sup>6</sup>

<sup>1</sup>Department of Pharmacognocny and Phytochemistry, Faculty of Pharmacy, Airlangga University, Surabaya 60115, Indonesia

<sup>2</sup>Institute of Tropical Disease, Airlangga University, Surabaya, 60115, Indonesia

<sup>3</sup>Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Airlangga University, Surabaya 60115, Indonesia

<sup>4</sup>Tradisional Medicine Study Program, Faculty of Vocational Studies, Airlangga University, Surabaya 60115, Indonesia

<sup>5</sup>Department of International Health, Kobe University Graduate School of Health Sciences, 7-10-2, Tomogaoka, Suma-ku, Kobe 654-0142, Japan

<sup>6</sup>Faculty of Clinical Nutrition and Dietetics, Konan Women's University, 6-2-23 Morikita-machi, Higashinada-ku, Kobe 658-0001, Japan

\*E-mail: tutik-s-w@ff.unair.ac.id

Accepted 21 May 2019, Published online 30 June 2019

### ABSTRACT

Hepatitis C virus (HCV) infection is a global problem that causes liver disease and hepatocellular carcinoma. Although the current standard treatment provided a significant improvement on response rate with sustain virology response more than 90%, however, the high cost was remaining limited access to this therapy, resistance emergence and serious side effects which provide the necessities to find the new anti-HCV agents. The current study, we evaluated the ethanol extract of *Phyllanthus niruri* for its anti-HCV activities. Anti-HCV activity was determined by *in vitro* culture cells of Huh 7it. Anti-HCV activity of *P. niruri* extract revealed strong inhibition against HCV with IC<sub>50</sub> values of 4.14 µg/mL and yield stronger activity in the entry step of the HCV life cycle. Moreover, the *P. niruri* extract enhanced anti-HCV activity of simeprevir (NS3 protease inhibitor) with increase the activity up to 4-fold compared to a single treatment of simeprevir. Docking analysis was performed to predict the interaction phyllanthin and hypophyllantin, known compounds of *P. niruri* against HCV receptor. Both of phyllantin and hypophyllantin were mediated a strong interaction with 4GAG, a protein that involved in entry step of HCV. These results suggested that the ethanol extract of *P. niruri* may be good candidates for the development of anti-HCV drugs.

**Key words:** Hepatitis C virus, antiviral activity, *Phyllanthus niruri*, entry step

### INTRODUCTION

Hepatitis C virus (HCV) infection is a serious health problem and a potential cause of substantial morbidity and mortality. The infections could lead to chronic liver diseases, such as hepatitis, liver disorder and hepatocellular carcinoma. The prevalence of HCV infection is infected around 71 million people worldwide (The Polaris Observatory HCV Collaborators; European Union HCV Collaborators). There is no vaccine of HCV that have been

developed. The current treatment of HCV used oral IFN-free regimen, includes a combination of direct-acting antiviral agents (DAAs), is capable of providing a sustained virological response (SVR) for more than 90% (Gonzalez-Grande *et al.*, 2016). However, the viral resistance issue, side effects, and economical view due to the high cost of current treatment agents remained the necessities to develop complementary and/or alternative drugs for the treatment of HCV.

Medicinal plants are rich with many chemical substances such as flavonoids, terpenoids, lignans, polyphenolics, coumarins, saponins, fureyl com-

\* To whom correspondence should be addressed.

pounds, alkaloids, polylines, proteins and peptides, that have been reported to possess various bioactivities including antiviral. Those compounds and their derivatives have been determined to possess anti-HCV activities that have promise for development into anti-HCV agents (Calland *et al.*, 2012; Jassim & Naji 2003; Khachatoorian *et al.*, 2012; Wahyuni *et al.*, 2016). Alkaloid and coumarin compounds of *R. angustifolia*, pseudane IX and chalepin, respectively, possess a strong inhibition against HCV (Wahyuni *et al.*, 2014). Honokiol, a lignan compound from *Magnolia officinalis*, reported inhibiting HCV with an  $IC_{50}$  value of 4.5  $\mu$ M (Lan *et al.*, 2012), while a lignan compound from *Swietenia macrophylla*, 3-hydroxy caruilignan C, reduced HCV protein and RNA level with  $IC_{50}$  value of 10.5  $\mu$ M (Wu *et al.*, 2012).

*Phyllanthus niruri* belongs to the family of Euphorbiaceae and is widely distributed in tropical and subtropical regions. It has been traditionally used for many kinds of disease including to reduce fever, treat jaundice and liver diseases. The various species of *Phyllanthus* family such as the *P. amarus*, *P. niruri*, *P. urinaria* and *P. orbicularis* have been reported to demonstrate potential inhibitory effect against a broad spectrum of viruses. They showed to possess potential inhibition in human immunodeficiency virus (HIV), herpes simplex virus (HSV) and hepatitis B virus (HBV) (Forero *et al.*, 2008; Tan *et al.*, 2013). *Phyllanthus amarus* has been demonstrated to have potential inhibition on HCV replication (Ravikumar *et al.*, 2011). Nirtetralin B, a lignan isolated compounds of *P. niruri* exhibited to suppress the secretion of the HBV antigens (Liu *et al.*, 2014). Other lignan compounds, phyllanthin and hypophyllanthin were reported to mediate anti-viral activities against hepatitis B virus and human immunodeficiency virus (HIV) (Bagalkotkar *et al.*, 2006; Venkateswaran *et al.*, 1987). This study was evaluated the anti-HCV activity of *P. niruri* by *in vitro* culture cells and further analyzed the mechanism of action. To predict the mechanism of known compounds, docking analysis of phyllanthin and hypophyllanthin was done to evaluate the interaction with protein in Protein Data Base ([www.rcsb.org](http://www.rcsb.org)) that involve in HCV activity.

## MATERIALS AND METHODS

### Extraction and sample preparation

The herb of *P. niruri* was harvested from Bandung, West Java, Indonesia. The plant was verified by a licensed botanist of Botanical garden, Purwodadi, Indonesia. The dried powder of the herb was pulverized and extracted by maceration method with 96% of ethanol. The obtained filtrates were evaporated to yield the ethanolic extracts of *P. niruri*. The stock solution

was prepared by dissolving the extract in dimethyl sulfoxide (DMSO) to obtain a stock concentration of 100 mg/mL. Serial dilutions of extracts were prepared to yield the concentrations of extracts 100, 50, 10, 1, 0.1 and 0.01  $\mu$ g/mL.

### Cells and Hepatitis C virus preparation

Huh7it cells (Aoki *et al.*, 2014) were cultivated in Dulbecco's Modified Eagle Medium (GIBCO Invitrogen, Carlsbad, CS, USA) supplemented with 10% Fetal Bovine Serum (Biowest, Nualle, France), 0.15 mg/mL Kanamycin (Sigma-Aldrich, St. Louis, MO, USA) and non-essential amino acids (GIBCO-Invitrogen) in 5% CO<sub>2</sub> at 37°C. The culture cells were cultivated and maintained by periodically re-feeding with a new medium. The adapted HCV variant was propagated in Huh7it (Apriyanto *et al.*, 2016). Culture supernatant from the infected cells was collected at day 2 and day 5 post-infection and concentrated using Amicon Ultra centrifugal filter unit. Virus titers were determined for antiviral assay (Hafid *et al.*, 2017).

### Anti-Hepatitis C activity assay

Antiviral activity assay was conducted as described previously (Aoki *et al.*, 2014; Hafid *et al.*, 2017; Wahyuni *et al.*, 2013; Wahyuni *et al.*, 2014). Huh7it cells ( $5.4 \times 10^4$ ) were seeded for 24 hours. The HCV at the multiplication of infection (MOI) of 0.1 in the presence of different concentrations of the sample were inoculated to the culture cells. The cells were treated with a serial concentration of *P. niruri* extract. The mixture of extract and virus was incubated for 2 hours for virus absorption and the cells were rinsed with the medium. Infected-cells were further incubated in the medium containing the same concentration of sample for 46 hours. Mode of action analysis was performed by time-of-addition experiments with three series of experiments. First, the culture was treated with the extract both in pre- and post- inoculation. Second, the culture was only treated with the extract at inoculation steps (2 hours). The third extract was added only after inoculation to examine the action of a substance in the post-entry steps of the HCV life cycle. Culture supernatants were collected for virus titration. The percentage inhibitory was determined and the 50% inhibitory effect ( $IC_{50}$ ) was calculated by SPSS probit analysis (Wahyuni *et al.*, 2013; Wahyuni *et al.*, 2014). Combination treatment to evaluate the effect of *P. niruri* extract was done by added the extract to the serial concentration of simeprevir, the NS3 protein inhibitor.

### Virus titration and immunostaining

Huh7it-1 cells ( $2 \times 10^4$  cells/well) were seeded in a 96-well plate and incubated for 24 hours. Virus supernatants were diluted in the medium and

inoculated onto the Huh7it culture cells and incubated for 4 hours. After virus absorption, the cells were cultured with medium containing 0.4% methylcellulose (Sigma-Aldrich) following 41 hours incubation. Infected cells were analyzed with immunostaining using anti-HCV patient anti-serum (250-time dilution on 2% BlockAce/1%BSA/PBS) and HRP-goat antihuman Ig antibody (300× on 2% lockAce/1%BSA/PBS). The HCV antigen positive cells were visualized with Metal Enhanced DAB substrate kits (Thermo Fisher Scientific, Rockford, USA). The infected cells were counted under microscopes and calculated the percentage inhibition.

#### Cytotoxicity analysis

The cytotoxicity analysis was conducted to determine whether the extract mediated any cytotoxicity effects. MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide) assay was done by inoculating 100, 50, 10, 1, 0.1 and 0.01 µg/mL of extract in 96 wells plate culture cells which have seeded for 24 hours. After 48 h incubation, the medium was replaced with MTT reagent containing medium and incubated for 4 hours. Absorbance sample was evaluated under a microplate reader at 450 and 600 nm, which is correlated with the amount of cell viability. The percentage of cell toxicity was calculated by comparing with untreated cells and further determine its 50% cytotoxic concentration (CC<sub>50</sub>) values (Apriyanto *et al.*, 2016).

#### Docking analysis of phyllanthin and hypophyllanthin

The ligand was prepared by making 2D and 3D structures of the phyllanthin and hypophyllanthin using ChemBioOffice program Ultra 11.0 and its energy was minimized using MMF94. The docking analysis continued by Molegro Virtual Docking ver 5.5 program Ver 5.5, resulted in rerank score describing the minimal energy by the ligand in interaction with the receptor.

## RESULTS

#### Anti-hepatitis C virus activity and cytotoxicity assay of *Phyllanthus niruri* extract

Ethanol extract of *P. niruri* was examined for antiviral activities against hepatitis C virus JFH1. The result showed that *P. niruri* possesses potential inhibition against hepatitis C virus without any cytotoxicity. Dose dependent on percentage inhibition and cell viability of *P. niruri* was shown in Table 1. The 50% inhibitory concentrations (IC<sub>50</sub>) was determined by probit analysis and yield the IC<sub>50</sub> value of 4.14 µg/mL without any cytotoxic effect with the 50% cytotoxic concentrations (CC<sub>50</sub>) >100 µg/ml and selectivity indexes (SI: CC<sub>50</sub>/IC<sub>50</sub>) > 24.2.


#### Mode of Action analysis of *Phyllanthus niruri* extract

To evaluate that actions of extract in the HCV life cycle, mode of action analysis of *P. niruri* extract was performed. The three parallel experiments were exerted, first the cells culture were treated with the plant extract in both pre- and post-inoculation, second cells were treated with extract only during inoculation to determine the action of extract in the entry step and third, cells were treated only post-inoculation to evaluate the action of extract in the post entry step. Pre-inoculation treatment obtained higher inhibition than post-inoculation treatment. It indicated that the extract of *P. niruri* mainly inhibits at the entry step. The percentage of HCV inhibition in the entry step was 70% while post-entry step was less than 50% (Figure 1). This result indicated that the mechanism of action of *P. niruri* was predicted in the attachment with some host receptor of entry to the host cell via endocytosis process.

**Table 1.** Dose dependent on percentage inhibition and cell viability of *Phyllanthus niruri*

Concentration (µg/mL)	% HCV inhibition	% Cell viability
0.01	13.3 ± 4.7*	99.8 ± 0.4
0.1	22.5 ± 4.1	99.0 ± 0.9
1	25.5 ± 5.8	98.8 ± 1.2
10	44.3 ± 3.3	98.7 ± 1.2
50	88.0 ± 5.1	98.7 ± 1.3
100	100.0 ± 0.0	98.3 ± 1.5

\* Data represent means ± SEM of data from three independent experiments.



**Fig. 1.** Extract of *Phyllanthus niruri* (30 µg/ml) inhibits HCV mainly in the entry step. Three kinds of experiments were done in parallel. Entry: infected-culture cells were treated with extract only in pre-inoculation, Post entry: infected-culture cells were treated only post-inoculation, both: infected-culture cells were treated with extract in the pre-and post-inoculation. The percentage HCV inhibition, both: 88%, entry: 70%:43%. Data represent means ± SEM of data from three independent experiments.

### Combination treatment of *Phyllanthus niruri* extract and simeprevir (NS3 protease inhibitor)

Simeprevir is an antiviral drug that acts to inhibit NS3 protein. Combination treatment by adding 25  $\mu\text{g/mL}$  of *P. niruri* extract was obtained increasing HCV inhibition (Figure 2). The potency of combination treatment was increased in 4-fold with the  $\text{IC}_{50}$  value  $3.54 \text{ nM} \pm 0.05$ , while the  $\text{IC}_{50}$  value of simeprevir single treatment was  $11.23 \text{ nM} \pm 0.36$ . *P. niruri* extract showed to reduce HCV NS3 protein level under western blot analysis (Figure 3).

### Docking analysis of known compounds from *Phyllanthus niruri*


Phyllanthin and hypophyllanthin are known as isolated compounds from *P. niruri*. To predict the interaction of those compounds docking analysis was performed. Several proteins from Protein Data Base ([www.rcsb.org](http://www.rcsb.org)) were evaluated to examine their interaction with Phyllanthin and hypophyllanthin. The result demonstrated a strong interaction between phyllanthin and hypophyllanthin with 4GAG. The rerank score value of phyllanthin and hypophyllanthin were  $-110.125$  and  $-88.5645$  kcal/mol, respectively, while the rerank score of standard ligands was  $-45.0229$  kcal/mol. This result indicated the strong interaction between phyllanthin and hypophyllanthin to the 4GAG receptor. The receptor of 4GAG was reported to be a protein that involves in the entry step of HCV, neutralizing antibody AP33 in complex with E2 epitope (Colpitts and Baumert 2016; Zhu *et al.*, 2014).

The hydrogen bonding and steric van der Waals interaction were involved in the interaction between compounds to the receptor. Hydrogen bonding of phyllanthin with His 164 and Gln 42, and the steric van der Waals between phyllanthin with, Asp 167, Val 163, His 164, Gln42, Lys39, Pro40 and Gly41 contributed the binding interaction of 4GAG and phyllanthin (Figure 4A). Hypophyllanthin revealed weaker interaction than phyllanthin with hydrogen bonding to His 164 and steric van der Waals-bond to Asp167, Val163, His164 and Thr165 (Figure 4B). While the standard ligand revealed hydrogen bonding to His 164 and Asp 167 and steric van der Waals to Asp 167. The interaction of phyllanthin and 4GAG was clearly described in the 3D profile (Figure 5).


## DISCUSSION

Medicinal plants and many natural sources are rich with chemical substances that responsible in many bioactivities including antiviral. We previously summarized potential plants that have anti-HCV activity (Wahyuni *et al.*, 2016). Our current study reported anti-HCV activities of *P. niruri* against


HCV JFH1. Plant with the genus of *Phyllanthus* was reported to have many bioactivities such as antimicrobial, antioxidant, anticancer, anti-inflammatory, anti-plasmodial, hepatoprotective and antiviral (Calixto *et al.*, 1998; Kaur *et al.*, 2017). This study found that ethanolic extract of *P. niruri* inhibited HCV with  $\text{IC}_{50}$  value of  $4.14 \mu\text{g/mL}$  without showing cytotoxicity. The extract of *P. niruri* may interfere with the inhibition effect in the several steps of the HCV life cycle. Further study to examine the effect of *P. niruri* extract in the entry or post-entry step of the HCV life cycle was evaluated by mode-of-action analysis. The result was described that *P. niruri* possesses a dominant inhibition in the entry steps of the HCV life cycle (Figure 1). The steps in HCV life cycle include entry into the host cells, uncoating and replication of the viral genome, translation of virus proteins, and assembly and release of the virion (Lindenbach *et al.*, 2005; Ploss & Dubuisson 2012). In the attachment/entry step, the HCV lipoviral




**Fig. 2.** Extract of *Phyllanthus niruri* (25  $\mu\text{g/mL}$ ) enhanced anti-hepatitis C activity of simeprevir. In parallel, culture cells were treated with a serial concentration of simeprevir only and the other was treated with a combination of simeprevir and extract. Data represent means  $\pm$  SEM of data from three independent experiments.



**Fig. 3.** Reduction of HCV NS3 protein level by *Phyllanthus niruri* extract. Huh 7.5it cells were infected with HCV and treated with ethanol extract of *Phyllanthus niruri* (50 and 75  $\text{mg/mL}$ ) and the untreated control (CTR) were subjected to western blot analysis.  $\beta$ -actin served as an internal control to verify equal amounts of sample loading.



**Fig. 4.** Hydrogen bond interaction (dashed blue-line) and Steric-Van der Walls bond interaction (dashed red-line) between Standard Ligand and A) phyllanthin, (B) hypophyllanthin, on the active site of HCV protein (4GAG.pdb).



**Fig. 5.** The 3D profile of docking interaction of phyllanthin (green color) with 4GAG protein.

particles attach to the cell surface and interact with glycosaminoglycans (GAG), low-density lipoprotein receptor (LDLR), scavenger receptor class B type 1 (SRB1) and CD81. Then, the viral interaction with CLDN1 results in internalization of the virus via clathrin-mediated endocytosis. Following acidification of the endosome and subsequent fusion of viral and endosomal membranes, the viral genome is released into the cytoplasm. On the other hand, the lateral movement of HCV-CD81 causes the transmission of the virus by cell-to-cell contact (Lindenbach & Rice, 2013; Zeisel *et al.*, 2015).

Combination treatment was done by adding the extract of *P. niruri* to the serial concentrations of simeprevir, an NS3 protein inhibitor which acts in the replication step of HCV. The addition of *P. niruri* extract increased anti-HCV activity of simeprevir in almost 4-fold with the  $IC_{50}$  value 3.54

nM  $\pm$  0.05, while the  $IC_{50}$  value of simeprevir single treatment was 11.23 nM  $\pm$  0.36. These results indicated that *P. niruri* extract enhanced the anti-HCV activity of simeprevir. *P. niruri* was demonstrated to possess a reduction effect on HCV NS3 protease level at the dose of 50  $\mu$ g/mL of extract (Figure 3) which indicated that *P. niruri* may also involve in the replication step of HCV life cycle.

The various metabolites in the plants do contribute to anti-HCV activities such as lignan, terpenoids, flavonoids, alkaloid, tannin and steroid (Geethangili & Ding 2018; Kaur *et al.*, 2017). Niranthin and hinokinin isolated compounds from *P. amarus* have strong anti-hepatitis B activities while corilagin was reported to block HCV NS3 protease and NS5B RNA-dependent-RNA-polymerase (Huang *et al.*, 2003; Reddy *et al.*, 2018). *P. niruri* possessed various compounds, such as quercetin, niruriflavone, rutin, ellagic acid,

phyllanthin, hypophyllanthin, niranthin, lintetralin, phyltetralin, etc. (Bagalkotkar *et al.*, 2006). Phyllanthin and hypophyllanthin are the lignan compounds that were reported as major compounds of *P. niruri* (Calixto *et al.*, 1998). To predict the mechanism-of-action of phyllanthin and hypophyllanthin to the HCV receptor, docking analysis was done by Molegro Virtual Docking ver 5.5 program. Phyllanthin and hypophyllanthin possess smaller re-rank value to compare to the ligand, it indicated the stronger interaction to 4GAG receptor, a protein that involves in the entry step of HCV, neutralizing antibody AP33 in complex with E2 epitope (Colpitts & Baumert 2016). The interaction of phyllanthin and hypophyllanthin to the receptor 4GAG resulting in an entry inhibition of HCV to the hepatocyte cells. The interaction of phyllanthin more stable than hypophyllanthin. Thus, it predicted to have higher biological activity.

## CONCLUSION

These results obtained strong activity of *P. niruri* extract against the hepatitis C virus that showed stronger inhibition in the entry step and enhances the activity of simeprevir. Phyllanthin and hypophyllanthin, known compounds of *P. niruri* predicted to possess strong interaction with 4GAG, a protein receptor that involves in the entry step of HCV. These results suggest that *P. niruri* would be a good candidate to develop anti-hepatitis C agents.

## ACKNOWLEDGEMENT

We gratefully acknowledge Research Institute and Innovation, Airlangga University through a *Riset Mandat* Grant.

## REFERENCES

- Aoki, C., Hartati, S., Santi, M.R., Lydwina, L., Firdaus, R., Hanafi, M., Kardono, L.B.S., Shimizu, Y., Sudarmono, P. & Hotta, H. 2014. Isolation and identification of substances with anti-hepatitis c virus activities from *Kalanchoe pinnata*. *International Journal of Pharmacy and Pharmaceutical Sciences*, **6**: 211-215.
- Apriyanto, D.R., Aoki, C., Hartati, S., Hanafi, M., Kardono, L.B., Arsianti, A., Louisa, M., Sudiro, T.M., Dewi, B.E., Sudarmono, P., Soebandrio, A. & Hotta, H. 2016. Anti-Hepatitis C Virus Activity of a Crude Extract from Longan (*Dimocarpus longan* Lour.) Leaves. *Japanese Journal of Infectious Diseases*, **69**(3): 213-220.
- Bagalkotkar, G., Sagineedu, S.R., Saad, M.S. & Stanslas, J. 2006. Phytochemicals from *Phyllanthus niruri* Linn. and their pharmacological properties: a review. *The Journal of Pharmacy and Pharmacology*, **58**(12): 1559-1570.
- Calixto, J.B., Santos, A.R.S., Filho, V.C. & Yunes, R.A. 1998. A review of the plants of the genus *Phyllanthus*: Their chemistry, pharmacology, and therapeutic potential. *Medicinal Research Reviews*, **18**(4): 225-258.
- Calland, N., Dubuisson, J., Rouille, Y. & Seron, K. 2012. Hepatitis C virus and natural compounds: a new antiviral approach?. *Viruses*, **4**(10): 2197-2217.
- Colpitts, C.C. & Baumert, T.F. 2016. Hepatitis C virus cell entry: a target for novel antiviral strategies to address limitations of direct acting antivirals. *Hepatology International*, **10**: 741-748
- European Union HCV Collaborators. Hepatitis C virus prevalence and level of intervention required to achieve the WHO targets for elimination in the European Union by 2030: a modelling study 2017. *The Lancet Gastroenterology & Hepatology*, **2**(5): 325-336.
- Forero, J.E., Avila, L., Taborda, N., Tabares, P., Lopez, A., Torres, F., Quinones, W., Bucio, M.A., Mora-Perez, Y., Rugeles, M.T., Joseph-Nathan, P. & Echeverri, F. 2008. *In vitro* anti-influenza screening of several Euphorbiaceae species: structure of a bioactive Cyanoglucoside from *Codiaeum variegatum*. *Phytochemistry*, **69**(16): 2815-2819.
- Geethangili, M. & Ding, S.-T. 2018. A Review of the Phytochemistry and Pharmacology of *Phyllanthus urinaria* L. *Frontiers in Pharmacology* **9**: 1109-1109.
- Gonzalez-Grande, R., Jimenez-Perez, M., Gonzalez Arjona, C. & Mostazo, Torres, J. 2016. New approaches in the treatment of hepatitis C. *World Journal of Gastroenterology*, **22**(4): 1421-1432.
- Hafid, A.F., Aoki-Utsubo, C., Permanasari, A.A., Adianti, M., Tumewu, L., Widyawaruyanti, A., Wahyuningsih, S.P.A., Wahyuni, T.S., Lusida, M.I. & Soetjipto, Hotta, H. 2017. Antiviral activity of the dichloromethane extracts from *Artocarpus heterophyllus* leaves against hepatitis C virus. *Asian Pacific Journal of Tropical Biomedicine*, **7**(7): 633-639.
- Huang, R.L., Huang, Y.L., Ou, J.C., Chen, C.C., Hsu, F.L. & Chang, C. 2003. Screening of 25 compounds isolated from *Phyllanthus* species for anti-human hepatitis B virus *in vitro*. *Phytotherapy Research*, **17**(5): 449-453.



- Jassim, S.A. & Naji, M.A. 2003. Novel antiviral agents: a medicinal plant perspective. *Journal of Applied Microbiology*, **95**(3): 412-427.
- Kaur, N., Kaur, B. & Sirhindi, G. 2017. Phytochemistry and Pharmacology of *Phyllanthus niruri* L.: A Review. *Phytotherapy Research*, **31**(7): 980-1004.
- Khachatoorian, R., Arumugaswami, V., Raychaudhuri, S., Yeh, G.K., Maloney, E.M., Wang, J., Dasgupta, A. & French, S.W. 2012. Divergent antiviral effects of bioflavonoids on the hepatitis C virus life cycle. *Virology*, **433**(2): 346-355.
- Lan, K.H., Wang, Y.W., Lee, W.P., Lan, K.L., Tseng, S.H., Hung, L.R., Yen, S.H., Lin, H.C. & Lee, S.D. 2012. Multiple effects of Honokiol on the life cycle of hepatitis C virus. *Liver International*, **32**(6): 989-997.
- Lindenbach, B.D., Evans, M.J., Syder, A.J., Wolk, B., Tellinghuisen, T.L., Liu, C.C., Maruyama, T., Hynes, R.O., Burton, D.R., McKeating, J.A. & Rice, C.M. 2005. Complete replication of hepatitis C virus in cell culture. *Science*, **309**(5734): 623-626.
- Lindenbach, B.D. & Rice, C.M. 2013. The ins and outs of hepatitis C virus entry and assembly. *Nature Reviews Microbiology*, **11**(10): 688-700.
- Liu, S., Wei, W., Li, Y., Lin, X., Shi, K., Cao, X. & Zhou, M. 2014. *In vitro* and *in vivo* anti-hepatitis B virus activities of the lignan nirtetralin B isolated from *Phyllanthus niruri* L. *Journal of Ethnopharmacology*, **157**: 62-68.
- Ploss, A. & Dubuisson, J. 2012. New advances in the molecular biology of hepatitis C virus infection: towards the identification of new treatment targets. *Gut*, **61**(Suppl 1): i25-35.
- Ravikumar, Y.S., Ray, U., Nandhitha, M., Perween, A., Raja Naika, H., Khanna, N. & Das, S. 2011. Inhibition of hepatitis C virus replication by herbal extract: *Phyllanthus amarus* as potent natural source. *Virus Research*, **158**(1-2): 89-97.
- Reddy, B.U., Mullick, R., Kumar, A., Sharma, G., Bag, P., Roy, C.L., Sudha, G., Tandon, H., Dave, P., Shukla, A., Srinivasan, P., Nandhitha, M., Srinivasan, N. & Das, S. 2018. A natural small molecule inhibitor corilagin blocks HCV replication and modulates oxidative stress to reduce liver damage. *Antiviral Research*, **150**: 47-59.
- Tan, W.C., Jaganath, I.B., Manikam, R. & Sekaran, S.D. 2013. Evaluation of antiviral activities of four local Malaysian *Phyllanthus* species against herpes simplex viruses and possible antiviral target. *International Journal of Medical Sciences*, **10**(13): 1817-1829.
- The Polaris Observatory HCV Collaborators. Global prevalence and genotype distribution of hepatitis C virus infection in 2015: a modelling study. 2017. *The Lancet Gastroenterology & Hepatology*, **2**(3): 161-176.
- Venkateswaran, P.S., Millman, I. & Blumberg, B.S. 1987. Effects of an extract from *Phyllanthus niruri* on hepatitis B and woodchuck hepatitis viruses: *in vitro* and *in vivo* studies. *Proceedings of the National Academy of Sciences of the United States of America*, **84**(1): 274-278.
- Wahyuni, T.S., Tumewu, L., Permanasari, A.A., Apriani, E., Adianti, M., Rahman, A., Widyawaruyanti, A., Lusida, M.I., Fuad, A., Soetjipto, D., Nasronudin, D., Fuchino, H., Kawahara, N., Shoji, I., Deng, L., Aoki, C. & Hotta, H. 2013. Antiviral activities of Indonesian medicinal plants in the East Java region against hepatitis C virus. *Virology Journal*, **10**(1): 259.
- Wahyuni, T.S., Utsubo, C.A. & Hotta, H. 2016. Promising Anti-Hepatitis C Virus Compounds from Natural Resources. *Natural Product Communications*, **11**(8): 1193-1200.
- Wahyuni, T.S., Widyawaruyanti, A., Lusida, M.I., Fuad, A., Soetjipto, Fuchino, H., Kawahara, N., Hayashi, Y., Aoki, C. & Hotta, H. 2014. Inhibition of hepatitis C virus replication by chalepin and pseudane IX isolated from *Ruta angustifolia* leaves. *Fitoterapia*, **99**: 276-283.
- Wu, S.F., Lin, C.K., Chuang, Y.S., Chang, F.R., Tseng, C.K., Wu, Y.C. & Lee, J.C. 2012. Anti-hepatitis C virus activity of 3-hydroxy caruilignan C from *Swietenia macrophylla* stems. *Journal of Viral Hepatitis*, **19**(5): 364-370.
- Zeisel, M.B., Crouchet, E., Baumert, T.F. & Schuster, C. 2015. Host-targeting agents to prevent and cure hepatitis C virus infection. *Viruses*, **7**(11): 5659-5685.
- Zhu, Y.-Z., Qian, X.-J., Zhao, P. & Qi, Z.-T. 2014. How hepatitis C virus invades hepatocytes: the mystery of viral entry. *World Journal of Gastroenterology*, **20**(13): 3457-3467.