

REFERENSI

- Adina, M., Revest, P. dan Shortland, P., 2010. *The Nervous System*. 2nd ed. London: Churchill Livingstone, pp.141-158.
- ASHA. (2015). 'Type, degree and configuration of hearing loss'. *Audiology Information Series*, p1–2.
- Ballo, I. R., Kaunang, T. M. D., Munayang, H., dan Elim, C. (2012). Profil Lanjut Usia dengan Depresi yang Tinggal di Balai Penyantunan Lanjut Usia Cerah Manado. *Jurnal Biomedik*, 4(1), p59-67.
- Conradsson, M., Rosendahla, E., Littbranda, H., Gustafson, Y., Olofsson, B., dan Lövheim, H. (2013). Usefulness of the Geriatric Depression Scale 15-item version among very old people with and without cognitive impairment. *Aging and Mental Health*, 17(5), p638–645.
- Depkes. (2010). *Telinga Sehat Pendengaran Baik* [Online]. Tersedia di: <http://www.depkes.go.id/article/print/840/telinga-sehat-pendengaran-baik.html>
- Ernest, N., dan Halton, M.. (2010). *Depression The Treatment And Management of Depression In Adults* (2nd ed). Leicester & London: The British Psychological Society & The Royal College of Psychiatrists.
- Fischer, N., Weber, B., dan Riechelmann, H. (2016). Presbyakusis – Age Related Hearing Loss. *Laryngo-Rhino-Otologie*, 95(07), p497–510.

- Georgakis, M. K., Papadopoulos, F. C., Protogerou, A. D., Pagonari, I., Sarigianni, F., Biniaris-Georgallis, S.-I., *et al.* (2016). Comorbidity of Cognitive Impairment and Late-Life Depression Increase Mortality: Results From a Cohort of Community Dwelling Elderly Individuals in Rural Greece. *Journal of Geriatric Psychiatry and Neurology*, 29(4), p195–204.
- Hatmanti, N. M., dan Muzdalifah L. (2019). Hubungan Tingkat Depresi dengan Kejadian Insomnia pada Lanjut Usia di Griya Werdha Jambangan Surabaya. *Journal Health of Science*, 12(1), p67-77.
- Indawati, R., Kuntoro, Notobroto, H. B., Qomaruddin, M.B., Mahajudin, M. S., dan Asiyah, S.N. (2016). Screening Performance of the Geriatric Depression Scale(GDS-15) for Elderly in the Community, Indonesia. *International Journal of Research in Advent Technology*, 4(12), p11-14.
- Istiqomah, S. N., dan Imanto, M. (2019). Hubungan Gangguan Pendengaran dengan Kualitas Hidup Lansia. *Majority*, 8(2), p234-239.
- Jones, D. A., Victor, C. R., Vetter, N. J. (1984). Hearing difficulty and its psychological implications for the elderly. *Journal of epidemiology and Community Health*, 38, p75-78.
- Lawrence, B. J., Jayakody, D. M. P., Bennet, R. J., Eikelboom, R. H., Gasson, N., Friedland, P. L. (2019). Hearing Loss and Depression in Older Adults:A Systematic Review and Meta-analysis. *The Gerontological Society of America*, 20(20), p1-18.
- Mener, D. J., Betz, J., Genther, D. J., Chen, D., Lin, F. R. (2013). Hearing Loss and Depression in Older Adults. *J Am Geriatr Soc*, 61(9), p1627-1629.

- Mulyaningsih, E. F., Purnami, N., dan Ahadiyah, T. H. (2015). Hubungan Antara Skor Kuesioner Hearing Handicap Inventory for Elderly Screening pada Penderita Presbikusis. *Jurnal THT-KL*, p54-65.
- Nilforoush, M. H., Sepehrnejad, M., dan Habibi, Z. (2019). *Beck Depression Inventory-II in Hearing Impaired Elderly Patients: A Presbycusis Study*. p231–236.
- Njoto, E. N. (2014). Mengenal Depresi pada Usia Lanjut Penggunaan *Geriatric Depression Scale (GDS)* untuk Menunjang Diagnosis. *CDK-217*, 41(6), p.472-474.
- Ogundiran, O., AO, O., BA, E., dan AC, O. (2017). *Presbycusis-Related Depression and Hearing Aid Use – A Review*. 5, p3821–3825.
- Okereke, O. I. (2015). Prevention of late-life depression: Current clinical challenges and priorities. In *Prevention of Late-Life Depression: Current Clinical Challenges and Priorities*. 1st ed. Boston: Humana Press.
- Parasari, G. A. T., dan Lestari, M. D. (2015). Hubungan Dukungan Sosial Keluarga dengan Tingkat Depresi pada Lansia di Kelurahan Sading. *Jurnal Psikologi Udayana*, 2(1), p68-77.
- Pimentel, A. F., Afonso, R. M., Pereira, H. (2012). Depression and Social Support in Old Age. *Psicologia, Saude & Doencas*, 13(2), p311-327.
- Rutherford, B. R., Brewster, K., Golub, J. S., Kim, A. H., dan Roose, S. P. (2018). Sensation and psychiatry: Linking age-related hearing loss to late-life depression and cognitive decline. *American Journal of Psychiatry*, 175(3), p215–224.

- Saunders AZ, Stein AV, dan Shuster NL. (1990). 'Audiometry' in Walker HK, Hall WD, Hurst JW, editors. *Clinical Methods: The History, Physical, and Laboratory Examinations* (3rd ed.). Boston: Butterworths. Chapter 133. Tersedia di: <https://www.ncbi.nlm.nih.gov/books/NBK239/>
- Silviliyana, M., Maylasari, I., Agustina, R., Ramadani, K. D., Sulistyowati, R., Annisa, L., dan Dewi, F. W. R. (2018). *Statistik Penduduk Lanjut Usia 2018*. p14
- Wancanta, J., Alexandrowicz, R., Marquart, B., Weiss, M., Friedrich, F. (2006). The criterion validity of the Geriatric `Depression Scale: a systematic review. *Acta Psychiatr Scand*, 114, p.398-410.
- World Health Organization (2000). Global burden of hearing loss in the year 2000. *World Health Organisation*, (4), p1–30.
- World Health Organization. (2012). WHO Global Estimates on Prevalence Hearing Loss. [online]. Available at: https://www.who.int/pbd/deafness/news/GE_65years.pdf
- World Health Organization. (2017). World Health Organization Report - Depression and Other Common Mental Disorders: Global Health Estimates. *World Health Organization*, p1–24.
- World Health Organization. (2018). WHO Global Estimates on Prevalence of Hearing Loss. [online] Available at: <https://www.who.int/pbd/deafness/estimates/en/>
- World Health Organization. (2019). *Deafness and Hearing Loss* [Online]. Available at: <https://www.who.int/news-room/fact-sheets/detail/deafness-and-hearing-loss>