

# Body Mass Index, Waist-Hip Ratio and Fasting Blood Glucose Levels Amongst University Students

*by Rury Tiara Oktariza*

---

**Submission date:** 23-Jul-2021 10:58AM (UTC+0800)

**Submission ID:** 1622955227

**File name:** and\_Fasting\_Blood\_Glucose\_Levels\_Amongst\_University\_Students.pdf (277.14K)

**Word count:** 3335

**Character count:** 16443

18

## BODY MASS INDEX, WAIST-HIP RATIO AND FASTING BLOOD GLUCOSE LEVELS AMONGST UNIVERSITY STUDENTS

Rury Tiara Oktariza<sup>2</sup>, Viskasari P Kalanjati<sup>1\*</sup>, Ni Wajan Tirthaningsih<sup>1</sup>

<sup>1</sup>Department of Anatomy and Histology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia,

<sup>2</sup>Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

### ABSTRACT

Overweight and obesity have been correlated to the higher risk for developing metabolic diseases in later life, such as DM type 2. This study researched the body mass index (BMI), the waist to hip ratio (WHR) and the fasting blood glucose levels (FBG) in the seemingly healthy university students to comprehend these variables amongst the youth in Indonesia in 2019. The BMI and WHR of 150 male and female students aged 18-22 years old of the IKBW, Kediri were measured by standardised anthropometry. The FBG was measured after 8 hours-minimum of fasting from the capillary blood drop using glucometer. Data was then analysed using SPSS 17 with level of significance of  $P < 0.05$ . According to Asia-Pacific BMI classification, students were 30.7% obese with males significantly higher than females ( $P = 0.016$ ), 18% overweight (significantly higher in females,  $P = 0.04$ ), 36% normal and 15.3% underweight. When compared between genders, the WHR was significantly higher in males ( $P < 0.001$ ); while there were no significant differences in FBG ( $P = 0.6$ ). 4 males and 5 females with FBG  $\geq 100$  mg/dl, whereas others were within normal limits. There were positive significant correlations between BMI and WHR in males and females ( $r = 0.777$ ,  $P < 0.001$ ;  $r = 0.54$ ,  $P < 0.001$ , respectively). There was a significant positive correlation between the BMI and FBG with  $r = 0.217$ ,  $P = 0.008$ ; and between the WHR and FBG with  $r = 0.21$ ,  $P = 0.01$  amongst all students. In this study, male students have significantly higher BMI and WHR than females. Higher FBG was well observed in students with either higher BMI or WHR.

**Keywords:** BMI; diabetes mellitus; obesity

### ABSTRAK

Obesitas dan kelebihan berat badan berkorelasi dengan tingginya resiko mengidap berbagai penyakit metabolik di kemudian hari, termasuk DM tipe 2. Penelitian ini bertujuan menganalisis hubungan indeks massa tubuh (IMT), rasio lingkar pinggang-panggul (WHR) dan kadar glukosa darah puasa (GDP) pada mahasiswa laki-laki dan perempuan usia 18-22 tahun di IKBW, Kediri tahun 2019. IMT dan WHR dari 150 mahasiswa dinilai dengan metode antropometri terstandarisasi. GDP dari darah kapiler diukur menggunakan glukometer minimal 8 jam pasca puasa. Data dianalisis dengan menggunakan SPSS 17 dengan tingkat signifikansi  $P < 0,05$ . Berdasarkan klasifikasi IMT Asia Pasifik, 30,7% subjek mengalami obesitas (IMT laki-laki lebih besar dibandingkan perempuan;  $P = 0,016$ ), 18% gemuk (IMT lebih besar pada perempuan,  $P = 0,04$ ), 36% normal dan 15,3% kurus. WHR lebih besar pada laki-laki ( $P < 0,001$ ) meskipun tidak terdapat perbedaan bermakna antara GDP laki-laki dan perempuan ( $P = 0,6$ ). Didapatkan 4 laki-laki dan 5 perempuan memiliki GDP  $\geq 100$  mg/dl. Terdapat korelasi yang bermakna antara IMT dan WHR pada laki-laki dan perempuan ( $r = 0,777$ ,  $P < 0,001$ ;  $r = 0,54$ ,  $P < 0,001$ ). Terdapat korelasi positif dan bermakna antara IMT dan GDP dengan  $r = 0,217$ ,  $P = 0,008$ ; dan antara WHR dan GDP dengan  $r = 0,21$ ,  $P = 0,01$  dari seluruh mahasiswa. Pada penelitian ini, IMT dan WHR mahasiswa laki-laki lebih besar secara bermakna daripada perempuan. GDP yang tinggi cenderung ditemukan pada mahasiswa dengan IMT atau WHR yang lebih besar.

**Kata kunci:** IMT; diabetes mellitus; obesitas

**Correspondence:** Viskasari P Kalanjati, Department of Anatomy and Histology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia. E-mail: viskasari-p-k@fk.unair.ac.id

pISSN:2355-8393 • eISSN: 2599-056x • doi: 10.20473/fmi.v57i1.14661

• Fol Med Indones. 2020;57:53-57 • Received 31 Jul 2019 • Accepted 16 Jan 2020

• Open access under CC-BY-NC-SA license • Available at <https://e-journal.unair.ac.id/FMI/>

28

## INTRODUCTION

The prevalence of overweight and obesity are increasing thus becoming a global health concern (Mehdad et al 2012, Poobalan & Aucott 2016). These conditions have

also been reported amongst the youth, which might increase the risk to suffer various metabolic diseases in later life (Poobalan & Aucott 2016). Several metabolic diseases correlated to obesity i.e. type-2 diabetes mellitus (T2DM) (Stroud et al 2015, Portero et al 2014).

Early identification of overweight and obesity in the young adults is warranted. World Health Organisation (WHO) has recommended the measurement of body mass index (BMI) as the simplest way for defining obesity (Mishra et al 2014). However, the BMI is an indirect measurement for body fat because the fat mass and the lean body mass components are not differentiated, as such in the measurement of the waist-hip ratio (WHR). WHR that is more accurate in measuring body fat distribution, can predict visceral fat/central obesity (apple-shaped or pear-shaped obesity), after adjusting to the BMI (Doustjalal et al 2016, Gadekar et al 2018, Fu et al 2015). In this study, we analysed the BMI and WHR in the seemingly healthy male and female university students and correlated these variables to the fasting blood glucose levels (FBG) as one indicator of prediabetes.

## MATERIALS AND METHODS

The study was conducted amongst 74 male and 76 female students of the Faculty of Dentistry, Institut Ilmu Kesehatan Bhakti Wiyata (IIKBW), Kediri, East Java, Indonesia aged 18-22 years old. Ethical clearance was obtained from the Research and Ethical Committee of Universitas Airlangga, Surabaya, Indonesia. After informed consent and information for consent were signed, all subjects were fasting overnight from 10.00 pm until the study began at approximately 07.00 am in the next morning. The weight, height, waist circumference and hip circumference were measured using standard anthropometry. Body weight was measured to the nearest 0.1 kg using Omron HN289 (Omron, Japan) digital weighing scale placed on a firm, flat ground with subjects wearing light clothing and no shoes; belts and other accessories were removed and pockets emptied. Height was measured to the nearest 0.1 cm using a wall-mounted tape measure (GEA, Indonesia) with the subjects standing erect, barefoot, heels together and looking straight ahead in the Frankfurt plane.

The BMI was calculated as weight in kilograms divided by height squared in metres. Subjects with BMI of less than 18.5 kg/m<sup>2</sup> were classified as underweight, 18.5-22.9 kg/m<sup>2</sup> were classified as normal weight, 23.0-24.9 kg/m<sup>2</sup> were classified as overweight, 25.0-29.9 kg/m<sup>2</sup> were classified as obese I and BMI greater than or equal to 30 kg/m<sup>2</sup> were defined as obese II (WHO 2000). Waist circumference and hip circumference (both in cm) was measured using the tape measure SECA 201 (SECA, Germany) according to WHO guidelines (WHO 2011). Waist circumference was measured to the nearest 0.1 cm in standing position at the midpoint between the lower margin of the last palpable rib and the top of the iliac crest and at the end of normal expiration. Hip

circumference was measured to the nearest 0.1 cm at the widest diameter around the hip and having the greater trochanter as the landmark. The WHR was calculated as waist circumference divided by the hip circumference. The waist-hip ratio cut-off point for Asian was 0.90 and 0.80 for male and female, respectively. Apple-shaped obesity was defined as high BMI (BMI  $\geq$  23 kg/m<sup>2</sup>) with high WHR (WHR  $>$  0.90 for male and  $>$  0.80 for female), while pear-shaped obesity was defined as high BMI (BMI  $\geq$  23 kg/m<sup>2</sup>) with low WHR (WHR  $<$  0.90 for male and  $<$  0.80 for female) (Fu et al 2015, WHO 2000, WHO 2011). Capillary blood sample was taken to measure the FBG using Accu-Chek Active® glucometer (Germany); after the index or middle finger had been cleaned with alcohol swab.

The finger site was then dried and finger-prick done with single-use lancet or a lancing device to draw blood for each subject. Fasting blood glucose (FBG) was classified according to the recommendation of Perkumpulan Endokrinologi Indonesia (PERKENI) (2015); FBG  $<$ 100 mg/dL is normal fasting glucose; FBG 100-125 mg/dL was prediabetes and FBG  $\geq$  126 mg/dL was provisional diagnosis of diabetes. According to American Diabetes Association (2005), prediabetes is similar to impaired fasting glucose (PERKENI 2015, ADA 2005). The data were analyzed using SPSS 17.0 for Windows. For difference of variables in male and female students, t-test or Mann-Whitney U test for independent samples were used to compare mean scores, and the Chi-square was used to compare frequencies. Pearson correlations of BMI, WHR and FBG of the subjects were assessed. P  $<$ 0.05 was regarded as significant.

## RESULTS

Data on various anthropological parameters and fasting blood glucose were given in Table 1. According to Asia-Pacific BMI classification, students were 30.7% obese with males significantly higher than females (P = 0.016), 18% overweight (significantly more in females, P = 0.04), 36% normal and 15.3% underweight. However, the overall BMI showed no significant differences between males and females (P = 0.41). The apple-shaped obesity was higher in females (18.4%) than in males (14.9%), while pear-shaped obesity was higher in males (36.5%) than in females (27.6%) although not significantly different (P  $>$ 0.05). When compared between genders, the WHR was significantly higher in males (P  $<$ 0.001); while there was no significance in FBG (P = 0.6).

Table 1. Anthropometry parameters and fasting blood glucose of males and females

Variables	All	Males (n = 74)	Females (n = 76)	p value
Weight (kg) ( $\mu$ (SD))	62.3 (14.9)	67.9 (15.9)	56.7(11.5)	<0.001
Height (cm) ( $\mu$ (SD))	162.1 (8.3)	168.2 (5.8)	156.1 (5.6)	<0.001
BMI (kg/m <sup>2</sup> ) ( $\mu$ (SD))	23.6 (4.8)	23.9 (5.1)	23.3 (4.5)	0.411
Underweight (n (%))	23 (15.3)	12 (16.2)	11 (14.5)	0.945
Normal (n (%))	54 (36)	24 (32.4)	30 (39.5)	0.467
Overweight (n (%))	27 (18)	8 (10.8)	19 (25)	0.040
Obese I & II (n (%))	46 (30.7)	30 (40.5)	16 (21)	0.016
BMI $\geq$ 23 kg/m <sup>2</sup> (n (%))	73 (48.7)	38 (51.4)	35 (46.1)	0.627
WC (cm) ( $\mu$ (SD))	78.3 ( 12.4)	82.1 (13.8)	74.6 (9.6)	0.001
HC (cm) ( $\mu$ (SD))	97.1 (9.2)	97.2 (9.8)	96.9 (8.7)	0.865
WHR ( $\mu$ (SD))	0.80 (0.07)	0.84 (0.07)	0.77 (0.05)	<0.001
Apple-shaped <sup>a</sup> (n (%))	25 (16.7%)	11 (14.9%)	14 (18.4%)	0.715
Pear-shaped <sup>b</sup> (n (%))	48 (32%)	27 (36.5%)	21 (27.6%)	0.324
FBG (mg/dl) ( $\mu$ (SD))	87.8 (7.3)	88.1 (7.5)	87.5 (7.3)	0.600
FBG $\geq$ 100 mg/dl (n (%))	9 (6%)	4 (5.4%)	5 (6.6%)	1

10  
BMI: body mass index, WC: waist circumference, HC: hip circumference, WHR: waist-hip ratio, FBG: fasting blood glucose. <sup>a</sup>BMI  $\geq$  23 kg/m<sup>2</sup>, WHR > 0.90 for male and > 0.80 for female. <sup>b</sup>BMI  $\geq$  23 kg/m<sup>2</sup>, WHR < 0.90 for male and < 0.80 for female.

Table 2. Pearson's correlation coefficients (r) of BMI with WHR

Subject	r value	p value
Overall (Males & Females)	0.617	<0.001
Male	0.777	<0.001
Female	0.54	<0.001

BMI: body mass index, WHR: waist-hip ratio

Table 3. Pearson's correlation coefficients (r) of FBG with BMI and WHR

Subject	FBG	
	BMI	WHR
Overall (Males & Females)		
r value	0.217	0.210
p value	0.008	0.010
Male		
r value	0.226	0.232
p value	0.053	0.046
Female		
r value	0.203	0.207
p value	0.079	0.072

BMI: body mass index, WHR: waist-hip ratio, FBG: fasting blood glucose

## DISCUSSION

From this study we found the overweight and obesity have becoming a common issue either in male or in female students. The prevalence is higher when compared to the previous study conducted in Indonesia in 2014-2015 (34.3%) (Pengpid & Peltzer 2017), and to other studies in Pakistan (31.1%), Thailand (20.8%) and Bangladesh (20%) (Khan et al 2016, Pengpid & Peltzer 2015, Howlader et al 2018). In our study, the prevalence of the obesity was slightly higher in males compared to

in females. This might be due to the genetics and/ or environment, such as restrictive diet life style amongst females is arguably more popular to stay slim (Khan et al 2016, Chao et al 2012, Al-Kilani et al 2012). The BMI and WHR differences were found between genders; from both we could determine the fat deposition pattern in the body (Fu et al 2015, WHO 2000, WHO 2011). In male students, the pear-shaped obesity was more commonly found than in females. On the other hand, the apple-shaped obesity was more common amongst females, which indicates the central type obesity that has been reported to a higher risk to develop the cardiovascular disease than those with pear-shaped (Gadekar et al 2018, Fu et al 2015, WHO 2000). Apple-shaped obesity is considered more dangerous than pear-shaped obesity because of the accumulation of fat in the deep abdominal area around the visceral organs that can lead to the development of arterial hypertension and insulin resistance which may lead to the heart diseases, dyslipidemia and T2DM in later life (Gadekar et al 2018, Klop et al 2013, Rolland et al 2007).

We found that students with higher BMI, or WHR, would have significant positive correlation to the higher FBG ( $\geq$ 100 mg/dl). This showed an indication of an impaired fasting glucose, a condition which might lead to the prediabetes if it was not well managed. Higher FBG levels were more common in females than in males. The prevalence of high FBG found in the current study was higher to the previous study in Eastern China due to the difference in the study methodology and sampling (Hao et al 2014, Somers et al 2006, Huang et al 2004). However, in Bangladesh and South Africa, there were no significant correlations between the FBG and the BMI amongst the university students (Howlader et al 2018, Yang et al 2002). Students with higher BMI



tend to have higher WHR; an elevated free fatty acid inflow to the liver tend to occur, and generated the abdominal adiposity. This would contribute to the insulin sensitivity impairment through the secretion of adipokines that might impair the glucose tolerance (Gadekar et al 2018, Klop et al 2013). Furthermore, the adiponectin is lower in the obese patients; a condition that might give some degree of higher risks to develop T2DM. Adiponectin was another type of adipokines which could improve the insulin secretion and sensitivity also the fatty acid oxidation to prevent the atherosclerosis and T2DM by decreasing the inflammatory process (Somers et al 2006, Huang et al 2004, Yang et al 2002).

## CONCLUSION

This study has resulted that male students had significantly higher BMI and WHR than the female students. Higher FBG was significantly observed in students with higher BMI and/ or WHR.

## REFERENCES

- ADA (2005). Diagnosis and classification of diabetes mellitus. *Diabetes Care* 28, S37-42.
- Al-Kilani H, Waly M, Yousef R (2012). Trends of obesity and overweight among college students in Oman: a cross sectional study. *Sultan Qaboos Univ Med J* 12, 69-76.
- Chao CY, Shih CC, Wang CJ, Wu JS, Lu FH, Chang CJ, et al (2012). Low socioeconomic status may increase the risk of central obesity in incoming university students in Taiwan. *Obes Res Clin Pract* 8, e212-219.
- Doustjalal SR, Gujjar KR, Sharma R, Nurfatiha M, Firzanah RN, Syukri MKM, et al (2016). Correlation between body mass index (BMI) and waist to hip ratio (WHR) among undergraduate students. *Pak Journal of Nutr* 15, 618-624.
- Fu J, Hofker M, Wijmenga C (2015). Apple or pear: size and shape matter. *Cell Metab* 21, 507-508.
- Gadekar T, Dudeja P, Basu I, Vashisht S, Mukherji S (2018). Correlation of visceral body fat with waist-hip ratio, waist circumference and body mass index in healthy adults: a cross sectional study. *Med J Armed Forces India*. Available from: <https://doi.org/10.1016/j.mjafi.2017.12.001>. Accessed May 25, 2019.
- Hao C, Zhang C, Chen W, Shi Z (2014). Prevalence and risk factors of diabetes and impaired fasting glucose among university applicants in Eastern China: findings from a population-based study. *Diabet Med* 31, 1194-1198.
- Howlader MH, Islam AM, Rahman MM, Mili US, Nur RS, Kibria MG (2018). Correlation between body mass index and fasting blood glucose level in a selected group of teenagers in Bangladesh. *Medicine Today* 30, 74-77.
- Huang KC, Lue BH, Yen RF, Shen CG, Ho SR, Tai TY, et al (2004). Plasma adiponectin levels and metabolic factors in nondiabetic adolescents. *Obes Res* 12, 119-124.
- Khan ZN, Assir MZ, Shafiq M, Chaudhary AE, Jabeen A (2016). High prevalence of preobesity and obesity among medical students of Lahore and its relation with dietary habits and physical activity. *Indian J Endocrinol Metab* 20, 206-210.
- Klop B, Elte J, Cabezas M (2013). Dyslipidemia in obesity: mechanisms and potential targets. *Nutrients* 5, 1218-1240.
- Mehdad S, Hamrani A, El Kari K, El Hamdouchi A, Barakat A, El Mizibri M, et al (2012). Body mass index, waist circumference, body fat, fasting blood glucose in a sample of Moroccan adolescents aged 11-17 years. *J Nutr Metab* 2012, 510458.
- Mishra S, Behera A, Behera B, Dash S (2014). Effect of body mass index on gender difference in lung functions in Indian population. *Int J Clin Exp Physiol* 1, 229-231.
- Pengpid S, Peltzer K (2015). Prevalence of overweight and underweight and its associated factors among male and female university students in Thailand. *HOMO* 66, 176-186.
- Pengpid S, Peltzer K (2017). The prevalence of underweight, overweight/obesity and their related lifestyle factors in Indonesia, 2014-2015. *AIMS Public Health* 4, 633-649.
- PERKENI (2015). Konsensus pengelolaan dan pencegahan diabetes mellitus tipe 2 di Indonesia 2015. Jakarta, Pengurus Besar Perkumpulan Endokrinologi Indonesia, p 12-13.
- Poobalan A, Aucott L (2016). Obesity among young adults in developing countries: a systematic overview. *Curr Obes Rep* 5, 2-13.
- Portero McLellan KC, Wyne K, Villagomez ET, Hsueh WA (2014). Therapeutic interventions to reduce the risk of progression from prediabetes to type 2 diabetes mellitus. *Ther Clin Risk Manag* 10, 173-188.
- Rolland YM, Haren MT, Patrick P, Banks WA, Malmstrom TK, Miller DK, et al (2007). Adiponectin levels in obese and non-obese middle-aged African-American women. *Obes Res Clin Pract* 1, 27-37.
- Somers A, Rusford E, Hassan M, Erasmus R (2006). Screening for diabetes mellitus in learners residing in the Belhar, Delft and Mfuleni communities of Cape Town, Western Cape, South Africa. *S Afr Fam Pract* 48, 16-16d.

Stroud C, Walker LR, Davis M, Irwin CE (2015). Investing in the Health and Well-Being of Young Adults. *J Adolesc Health* 56, 127–129.

WHO (2000). The Asia-Pacific perspective: redefining obesity and its treatment. Sydney, Health Communication. Available from: <http://www.wpro.who.int/nutrition/documents/docs/Redefiningobesity.pdf>. Accessed May 26, 2019.

WHO (2011). Waist circumference and waist-hip ratio: report of a WHO expert consultation, Geneva 8-11 December 2008. Geneva, WHO Document Production Services, Available from: [http://whqlibdoc.who.int/publications/2011/9789241501491\\_eng.pdf](http://whqlibdoc.who.int/publications/2011/9789241501491_eng.pdf). Accessed May 26, 2019.

Yang WS, Lee WJ, Funahashi T, Tanaka S, Matsuzawa Y, Chao CL, et al (2002). Plasma adiponectin levels in overweight and obese asians. *Obes Res* 10, 1104–1110.

# Body Mass Index, Waist-Hip Ratio and Fasting Blood Glucose Levels Amongst University Students

---

## ORIGINALITY REPORT

---

22%

SIMILARITY INDEX

17%

INTERNET SOURCES

18%

PUBLICATIONS

0%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** Wenzhi Cai, Juan Wang, Li Wang, Jingxin Wang, Li Guo. "Prevalence and risk factors of urinary incontinence for post-stroke inpatients in Southern China", *Neurourology and Urodynamics*, 2015 1%

Publication
- 2** Roya Kelishadi. "Short- and long-term relationships of serum ghrelin with changes in body composition and the metabolic syndrome in prepubescent obese children following two different weight loss programs", *Clinical Endocrinology*, 2/12/2008 1%

Publication
- 3** Hui Zhao, Jun Ma, Qiru Zhou, Wei Chen, Wei Zhu, Zekun Cai, Hongqiang Lei, Yu Deng, Lin Xu, Jian Qiu. "Investigating the differences of body mass index and waist circumference in the follow-up assessment of patients to cardiac rehabilitation with acute coronary syndrome", *Australasian Physical & Engineering Sciences in Medicine*, 2016 1%

4

Youn-Soo Cha, Ji-Ae Yang, Hyang-Im Back, Soo-Ran Kim, Min-Gul Kim, Su-Jin Jung, Won O Song, Soo-Wan Chae. "Visceral fat and body weight are reduced in overweight adults by the supplementation of , a fermented soybean paste ", Nutrition Research and Practice, 2012

Publication

---

1 %

5

K V Haderslev. "Vitamin D status and measurements of markers of bone metabolism in patients with small intestinal resection", Gut, 5/1/2003

Publication

---

1 %

6

Sukru Uguz, Gulsah Seydaoglu, Cem Doğan, Banu Yazgan Inanc, Ebru Yurdagul, Rasim Somer Diler. "Short-term antidepressant treatment of comorbid dental anxiety in patients with panic disorder", Acta Odontologica Scandinavica, 2009

Publication

---

1 %

7

Ardhendu Bhusan Praharaj, Ramesh Kumar Goenka, Sujata Dixit, Manoj Kumar Gupta, Shantanu Kumar Kar, Sapna Negi. "Lacto-Vegetarian Diet and Correlation of Fasting Blood Sugar with Lipids in Population Practicing Sedentary Lifestyle", Ecology of Food and Nutrition, 2017

Publication

1 %



---

8	Ben Brumpton, Arnulf Langhammer, Pål Romundstad, Yue Chen, Xiao-Mei Mai. "General and abdominal obesity and incident asthma in adults: the HUNT study", European Respiratory Journal, 2013 Publication	1 %
9	<a href="http://www.medsci.org">www.medsci.org</a> Internet Source	1 %
10	<a href="http://www.tdx.cat">www.tdx.cat</a> Internet Source	1 %
11	<a href="http://drc.bmj.com">drc.bmj.com</a> Internet Source	1 %
12	<a href="http://r2kn.litbang.kemkes.go.id">r2kn.litbang.kemkes.go.id</a> Internet Source	1 %
13	<a href="http://www.ijcmr.com">www.ijcmr.com</a> Internet Source	1 %
14	Susan Quine, Stephen Morrell, Hal Kendig. "Outlooks for the future in independently living older Australians", Aging & Mental Health, 2008 Publication	<1 %
15	<a href="http://ukzn-dspace.ukzn.ac.za">ukzn-dspace.ukzn.ac.za</a> Internet Source	<1 %
16	<a href="http://www.duo.uio.no">www.duo.uio.no</a> Internet Source	<1 %

---

17 Meity Ardiana, Anwar Santoso, Hanestya O. Hermawan, Ricardo A. Nugraha, Budi S. Pikir, I Gde Rurus Suryawan. "Acute Effects of Cigarette on Endothelial Nitric Oxide Synthase, Vascular Cell Adhesion Molecule 1 and Aortic Intima Media Thickness "Cigarette smoke-induced pro-atherogenic changes"", Cold Spring Harbor Laboratory, 2021  
Publication

---

18 [doaj.org](https://doaj.org)  
Internet Source <1 %

---

19 [jdmdonline.biomedcentral.com](https://jdmdonline.biomedcentral.com)  
Internet Source <1 %

---

20 [journals.lww.com](https://journals.lww.com)  
Internet Source <1 %

---

21 [www.e-sc.org](https://www.e-sc.org)  
Internet Source <1 %

---

22 [www.mdpi.com](https://www.mdpi.com)  
Internet Source <1 %

---

23 [www.researchsquare.com](https://www.researchsquare.com)  
Internet Source <1 %

---

24 [indianjnephrol.org](https://indianjnephrol.org)  
Internet Source <1 %

---

25 [link.springer.com](https://link.springer.com)  
Internet Source <1 %

---

26	<a href="http://www.dovepress.com">www.dovepress.com</a> Internet Source	<1 %
27	N. K. Vikram. "Correlates of Type 2 diabetes mellitus in children, adolescents and young adults in north India: a multisite collaborative case-control study", Diabetic Medicine, 3/2006 Publication	<1 %
28	Reis, J.P.. "The relation of leptin and insulin with obesity-related cardiovascular risk factors in US adults", Atherosclerosis, 200809 Publication	<1 %
29	<a href="http://academic.oup.com">academic.oup.com</a> Internet Source	<1 %
30	<a href="http://digital.library.adelaide.edu.au">digital.library.adelaide.edu.au</a> Internet Source	<1 %
31	<a href="http://he01.tci-thaijo.org">he01.tci-thaijo.org</a> Internet Source	<1 %
32	<a href="http://repository.unib.ac.id">repository.unib.ac.id</a> Internet Source	<1 %
33	<a href="http://www.asean-endocrinejournal.org">www.asean-endocrinejournal.org</a> Internet Source	<1 %
34	<a href="http://www.frontiersin.org">www.frontiersin.org</a> Internet Source	<1 %
35	"Sex and Gender Factors Affecting Metabolic Homeostasis, Diabetes and Obesity", Springer	<1 %

---

36 Dieu-My T Tran, Angela Silvestri-Elmore. <1 %  
"Healthcare-seeking behaviours in college  
students and young adults: a review", Journal  
of Research in Nursing, 2020  
Publication

---

37 Sung Reul Kim, Ju - Hee Nho, Hye Young Kim. <1 %  
"Influence of Type D personality on quality of  
life in university students: The mediating  
effect of health - promoting behavior and  
subjective health status", Psychology in the  
Schools, 2020  
Publication

---

38 addi.ehu.es <1 %  
Internet Source

---

39 banglajol.info <1 %  
Internet Source

---

40 collantropol.hr <1 %  
Internet Source

---

41 mafiadoc.com <1 %  
Internet Source

---

42 repository.nwu.ac.za <1 %  
Internet Source

---

43 www.scialert.net <1 %  
Internet Source

---

44

[www.uob.edu.bh](http://www.uob.edu.bh)

Internet Source

&lt;1 %

45

[www.verywellhealth.com](http://www.verywellhealth.com)

Internet Source

&lt;1 %

46

"Biochemistry of Cardiovascular Dysfunction in Obesity", Springer Science and Business Media LLC, 2020

Publication

&lt;1 %

47

Atika Khalaf, Iman Al Hashmi, Omar Al Omari. "The Relationship between Body Appreciation and Self-Esteem and Associated Factors among Omani University Students: An Online Cross-Sectional Survey", Journal of Obesity, 2021

Publication

&lt;1 %

48

Elaine C. Rush, Ismael Freitas, Lindsay D. Plank. "Body size, body composition and fat distribution: comparative analysis of European, Maori, Pacific Island and Asian Indian adults", British Journal of Nutrition, 2009

Publication

&lt;1 %

49

[repository.unair.ac.id](http://repository.unair.ac.id)

Internet Source

&lt;1 %



Exclude quotes On

Exclude matches Off

Exclude bibliography On

# Body Mass Index, Waist-Hip Ratio and Fasting Blood Glucose Levels Amongst University Students

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---