

Potential Extract Ethanol Citrus Amblycarpa as a Bioinsecticide Against Aedes Aegypti Larvae

by Poedji Hastutiek

Submission date: 05-Aug-2021 03:14PM (UTC+0800)

Submission ID: 1627966641

File name: ential_Extract_Ethanol_Citrus_Amblycarpa_as_a_Bioinsecticide.pdf (994.75K)

Word count: 4653

Character count: 24101

Potential Extract Ethanol Citrus Amblycarpa as a Bioinsecticide Against Aedes Aegypti Larvae

1
Kasman¹, Nuning Irnawulan¹ Ishak*², Poedji Hastutiek³, Endang Suprihati³

¹Departement of Epidemiology and Biostatistics, Faculty of Public Health, Islamic University of Kalimantan, Banjarmasin, Indonesia

²Department of Occupational Health Safety and Environmental Health, Faculty of Public Health, Islamic University of Kalimantan, Banjarmasin, Indonesia

³Department of Veterinary Parasitology, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

ABSTRACT

Introduction: Utilization of compounds active plant into an alternative control of mosquito when it and a front for a mosquito hospitable environment and do not cause interference health. Kuit Limau Plant (Citrus amblycarpa) is local citrus from South Kalimantan which has bioinsecticide potential which can be used in mosquito control.

Objective: This study aims to determine the effectiveness of the ethanol extract of the Peel and Citrus amblycarpa leaves as natural larvicides of Aedes aegypti larvae.

Methods: An experimental study with a completely randomized design of Citrus amblycarpa ethanol extract (500, 1000, 2000, 3000, and 4000 ppm). The positive control used abate and the negative control used distilled water and tween 80. The number of repetitions of each concentration was three times with 25 larvae in each test glass. Larval mortality was analyzed using probit analysis at a 95% confidence level.

Results: Extract Peel requires a time that is shorter (5.3 hours) to kill the larvae of Ae. Aegypti compared with leaf extract (42.8 hours). Is not there a difference significant mortality of larvae in the group leather 4000 ppm with a group of control.

Conclusion: The ethanol extract of Citrus amblycarpa rind has potential as a larvicide for Aedes aegypti.

Keywords: Ethanol extract, Bioinsecticide, Citrus amblycarpa, Aedes aegypti

Correspondence:

Nuning Irnawulan Ishak
Department of Occupational Health Safety and Environmental Health, Faculty of Public Health, Islamic University of Kalimantan, Jl. Adhyaksa No.2 Kayutangi, Banjarmasin, South Kalimantan, Indonesia,
Email address : nuning.fkm@gmail.com

INTRODUCTION

Dengue hemorrhagic fever is a tropical and subtropical disease (1) (2) (3) which is transmitted through the bite of the mosquito Aedes aegypti or Aedes albopictus and is caused by the Dengue virus (4)(5) DHF cases in Indonesia that were reported in 2019 were 138,127 cases. This number increased compared to 2018 of 65,602 cases. Deaths due to dengue fever in 2019 also increased compared to 2018, from 467 to 919 deaths. The incidence rate of dengue fever in 2019 is 51.53 per 100,000 population. This figure shows an increase compared to the previous two years, namely in 2017 and 2018 when the Incidence Rate of DHF was 26.1 and 24.75 per 100,000 population (6). The condition of the occurrence of dengue fever which is evident encourages the need to control the Aedes aegypti mosquito. One method that can be done to break the cycle and kill mosquito larvae is to use insecticides (Susilowati, Darmanto, and Aminah, 2018). The continuous use of synthetic insecticides (temephos (abate), malathion, cypermethrin, lambdasihalothrin, and deltamethrin) against the mosquito vector Aedes aegypti causes resistance to mosquitoes (8) (9), bioactive properties that are difficult to degrade in nature can pollute the environment, the toxic substances in synthetic insecticides will also have an impact on human health problems (10) (Susilowati, Darmanto, and Aminah, 2018). Natural larvae are an alternative in controlling mosquitoes by utilizing plants that are more environmentally friendly so that they can reduce the use of synthetic insecticides and anticipate negative impacts on health (11).

There are 2,400 species of bioinsecticide-producing plants in Indonesia that are toxic to insects (12). One of the local and abundant plants in the South Kalimantan Region, Indonesia with potential as a bioinsecticide against Aedes aegypti is the Citrus amblycarpa plant. This plant has active secondary metabolites such as flavonoids, tannins,

saponins, alkaloids (13) (14). The fresh extract of Citrus amblycarpa fruit Peel has been shown to kill Aedes aegypti instar III larvae in just 7 (seven) hours in all concentrations (15). This study aims to determine the effectiveness of the ethanol extract of Citrus amblycarpa in killing Aedes aegypti larvae.

MATERIALS AND METHODS

This research is experimental research with a completely randomized design. In this study, various concentrations of the ethanol extract of the Peel and Citrus amblycarpa leaves were used, namely 500 ppm, 1000 ppm, 2000 ppm, 3000 ppm, and 4000 ppm. The positive control used abate and negative control used distilled water and Tween 80. The number of repetitions of each concentration was three times. The number of larvae in each test glass was 25 according to WHO standards.

Provision of Citrus amblycarpa samples

A sampling of Citrus amblycarpa, which are local oranges typical of South Kalimantan, were taken in several places in South Kalimantan. Samples are taken and picked directly from the farmers' plantations to maintain its freshness. The leaves and peel are sorted then cleaned of dirt by washing with clean water, drained and spread on a newspaper so that the water is absorbed, weighed as much as 2.5 kg wet weight, aerated for 7 (seven) days by placing it in a place protected from the sun. This activity will be carried out at the Basic Laboratory of the Islamic University of Kalimantan, Banjarmasin, Indonesia.

Ethanol extract preparation

The ethanol extract was made at the Laboratory of Basic Medicine, Faculty of Veterinary Medicine, Airlangga University. The dried leaves and Peel of the Citrus amblycarpa are then mashed to form a powder. One kg of leaf and Peel Simplicia was macerated with ethyl alcohol p.a solvent for three days, filtered every day and the

5 filtrate was collected and evaporated using a rotary evaporator with a decrease in pressure until a dry extract was obtained.

Larvicide Testing


The experimental animal used was the third instar larvae of the *Ae. aegypti* results from rearing in the Entomology Laboratory of the Indonesian Tanah Bumbu Research Institute for 900 instars III larvae. The count of the dead larvae was carried out every hour for 24 hours of immersion, recorded in tabular form. The data obtained were analyzed statistically using the ANOVA test to determine the difference in larva mortality between each group and Probit analysis to determine the killing power of Peel extract and Citrus amblycarpa Leaf against *Aedes*

aegypti instar III larvae expressed in Lethal Time (LT₅₀ and LT₉₀).

RESULTS AND DISCUSSION

Comparison of the Effectiveness of Citrus amblycarpa Peel and Leaf Extract

12 Graph 1 shows the comparison of mortality of *Aedes aegypti* larvae after 24 hours of treatment. The mortality of larvae with ethanol extract of Citrus amblycarpa leaves was higher than that of larvae with ethanol extract of Citrus amblycarpa leaves. Larvae mortality in the control group (+) showed 100% larval mortality while in the control group (-) there was 4% larvae mortality.



Graph 1. Comparison of Larvae Mortality

Probit Analysis

3 Table 1 shows the results of the probit analysis to calculate the Lethal Time of the ethanol extract of Citrus amblycarpa leaves. LT₅₀ shows the time taken to kill 50% of larvae. At

2 a concentration of 4000 ppm it takes 5.3 hours to kill 50% of larvae and it takes 13.2 hours to kill 90% of larvae. In the control + group, it took 3.2 to kill 50% of larvae, and it took 5 hours to kill 90% of larvae.

Table 1. Value of LT₅₀ and LT₉₀ Mortality of Aedes aegypti Larvae Instar III Treatment Group Ethanol Extract Citrus amblycarpa Peel.

Concentration	LT ₅₀ (hour)	95% CI		LT ₉₀ (hour)	95% CI	
		Min	Maks		Min	Maks
500 ppm	402.8	127.4	96x10 ⁴	5,775	598.1	40x10 ⁹
1000 ppm	46.8	36.5	65.3	337.7	189.1	917.5
2000 ppm	18.3	15.8	20.8	78.0	63.1	112.1
3000 ppm	8.3	7.0	9.3	25.8	23.5	28.9
4000 ppm	5.3	4.4	6.1	13.2	11.8	14.7
Kontrol +	3.2	2.1	3.8	5.0	4.3	5.8
Kontrol -	44x10 ³	1x10 ³	58x10 ¹⁵	15x10 ⁶	3x10 ⁴	8x10 ²⁷

3 Table 2 shows the results of the probit analysis to calculate the Lethal Time of the ethanol extract of Citrus amblycarpa leaves. LT₅₀ shows the time taken to kill 50% larvae. At a concentration of 4000 ppm, it takes 42.81 hours to kill

2 50% of larvae and 136.49 hours to kill 90% of larvae. In the control + group, it took 3.19 hours to kill 50% of larvae, and it took 4.98 hours to kill 90% of larvae.

Table 2. Value of LT₅₀ and LT₉₀ Mortality of Aedes aegypti Larvae Instar III Treatment Group of Citrus amblycarpa Leaves Ethanol Extract

Concentration	LT ₅₀ (hour)	95% CI		LT ₉₀ (hour)	95% CI	
		Min	Maks		Min	Maks
500 ppm	1.692	253	2x10 ⁹	35.966	1.529	1x10 ¹⁵
1000 ppm	1.695	229	2x10 ⁸	29.532	1.226	4x10 ¹²

2000 ppm	74,1	46,8	408,8	229,9	96,4	6.216
3000 ppm	43,9	36,2	60,5	119,9	80,7	241,7
4000 ppm	42,8	35,4	56,6	136,5	92,2	261,3
Kontrol +	3,2	2,1	3,8	4,9	4,3	5,8
Kontrol -	44x10 ³	1x10 ³	58x10 ¹⁵	15x10 ⁶	3x10 ⁴	8x10 ²⁷

Prevention of the spread of DHF can be done in various ways, but until now the most effective way is to break the chain of transmission through vector control. Control that has been done a lot is by carrying out chemical control, namely by using synthetic insecticides. Synthetic insecticides work more effectively and quickly than biological and physical controls. However, repeated and long-term use can cause poisoning to humans and livestock, pollute the environment, and cause insect resistance.

21
Another alternative that can be used to control the dengue mosquito vector with a more environmentally friendly method is to use vegetable insecticides. Vegetable insecticides can kill or interfere with pests and diseases through a unique way of working, namely through a combination of various methods or singly. (16). Secondary metabolite compounds from plants are an alternative to insecticides that have a toxic effect on insects but have no side effects and are relatively safe for the environment and human health because they have a small risk. (17) (18). The use of natural insecticides from plants is an alternative to current insecticides, as well as vector control in the

13 re (19).
This study aims to determine the effectiveness of the 25 anol extract of the skin and leaves of Citrus amblycarpa in killing Aedes aegypti larvae. The larvicidal activity test was carried out using 25 larvae with their respective concentrations in the ethanol extract 16 the skin and Citrus amblycarpa leaves starting from a concentration of 500 ppm, 1000 ppm, 2000 ppm, 3000 ppm, and 4000 ppm. The solvent media used were aquadest and tween 80. The addition of Tween 80, in this case, served to increase the solubility of the ethanol extract of the skin and Citrus amblycarpa leaves in water. Extracts and fractions were tested against larvae of the Aedes aegypti mosquito, instar III, the selection of the third instar as the test phase because the larvae are active foraging and their morphological shape is more perfect than instar I and II larvae. (20). The positive control used in this test was temephos 0.01gr / 100ml, while the negative control used in this test was aquadest and tween 80. The results of the larvicidal activity test can be seen in Figure 1.

This study used various concentrations of the ethanol extract of the skin and Citrus amblycarpa leaves that were tested in each group of larvae with 3 times the observed repetition for 24 hours of observation. The mortality of 18 larvae increased with increasing concentration. This proves that the higher the concentration and the lo 14 r the exposure time, the higher the mortality of larvae. This research is in line with the results of research (21) that the increasing the concentration of the ethanol extract of kirinyuh leaves, the toxic substances contained in the larva's living media also increase so that the number of toxic substances that enter the larva's body also 16 reases so that the larvae it's easier to experience death. The high mortality rate of test larvae can be caused by the presence of chemical compounds in this extract that play a role in biological activity in larval growth and 3 velopment. The results of the study (15), also stated that at a concentration of 4.0 ml/100ml and 5.0 ml/100ml had 100% de 3 h after 6 hours of measurement, with an LT₅₀ value of 5.0%

concentration was 2.58 hours and an LT₉₉ value is 5.86 hours. So, from the results of the probit lethal time and lethal concentration tests, it was concluded that the greater the concentration is given and the lo 12 r the treatment time, it could increase the number of Aedes aegypti instar III larvae that died with the condition that the larvae were seen experiencing convulsions and bending the body.

The ability of the ethanol extract of Citrus amblycarpa peel to affect the mortality of Aedes aegypti larvae is due to the presence of certain active ingredients such as terpenoid / free steroid compounds, polyphenols, and saponins (22). Saponins are compounds that are included in terpenoid compounds. The activity of these saponins in the insect's body is to bind free sterols in the digestive tract of food where the sterol itself is a substance that functions as a precursor to the hormone ecdysone so that by decreasing the number of free sterols in the insect's body it will disrupt the process of skin change (molting) in insects. (23). Saponin compounds have the ability to damage cell membranes so that they can cause corrosion on the walls of the larvae digestive tract as stomach poison (24). The results of GC-MS shows the main components 8 the ethanol extract of citron biscuits are compounds 2H-1-Benzopyran-2-one, known as coumarin compound (25).

Coumarin is a secondary metabolic compound from the phenolic compound group which generally comes from higher plants and is rarely found in microorganisms. Coumarin can be found in almost all parts of plants from roots, stems, leaves, flowers, and fruit (26). Coumarin compounds are known for several pharmacological properties including insecticidal activity. The results of research (27), coumarin with a content of 27.7% is a class of compounds contained in the essential oil of lime peel (Citrus aurantifolia). This compound has the potential as a mosquito larvicide (28) (29). One of the chemical components contained in the Citrus sinensis plant is coumarin which has biological activity as an insecticide (30). 10 is highly toxic to insects (31). This compound is very toxic to M. persicae aphids with a value (LC₅₀₍₉₀₎ values of 1.3(1.9) mg L⁻¹) (32). One of the compound components of soursop (A. muricata) leaf and seed extract is that coumarin has the potential to be used as a vegetable pesticide to control cheap and environmentally friendly pest attacks. (33). 1

The ability of the ethanol 9 ract of Citrus amblycarpa leaves to affect the mortality of Aedes aegypti larv 1 is due to the presence of certain active ingredients such as terpenoids / free steroids, flavonoids, polyphenols, and saponins. (14). Flavonoid compounds are one of the toxic compounds for insecticides because of their ability to enter the larva's body through the respiratory system and cause weakness in the nerves. (34). The body position of the larvae that changes from normal can also be caused by flavonoid compounds due to how they enter through the siphon, causing damage so that the larvae have to align their position with the water surface to make it easier to take oxygen (35). Flavonoids have an antifeedant action by inhibiting the taste receptors in the mouth area of the larvae, which will result in the larvae failing to get a taste stimulus, so the larvae are unable to recognize the food

around them. The low feeding activity of larvae causes the energy for larval development to decrease so that the growth process is also inhibited.

The results of the study (36) show that the ethanol extract of cassava leaves (*Manihot utilissima* Pohl) with the chemical compounds of saponins and flavonoids has *Aedes aegypti* larvicidal activity with an LC₉₀ value of 2,613% and larvicidal activity of ethanol extract of cassava leaves at a concentration of 3% and 3.5%. There was no significant difference in *Aedes aegypti* larvae compared to temefos. The GC-MS results showed that the main component of the Citrus amblycarpa leaf ethanol extract was Maragenin I compound (14). Maragenin I is a triterpenoid compound derivative (37) (38). Maragenin I compounds can control insect growth and have potential as an insecticide (39). This compound is a derivative of the triterpenoid/strophanthidin class, so it can be concluded that this compound has the potential as a biopesticide, is toxic to the *Aedes aegypti* mosquito. (40).

The results of this study also showed a comparison of the effectiveness of an ethanol extract from the Citrus amblycarpa plant on its ability as a vegetable larvicide to kill *Aedes aegypti* larvae. The difference is seen from the number of larvae deaths, with the largest concentration of treatment, namely 4000 ppm, the ethanol extract of Citrus amblycarpa skin is more effective in killing larvae with a mortality percentage of 96%, while the ethanol extract of Citrus amblycarpa leaves only kills larvae with a mortality percentage of 17.3%. This is in line with Jubaedah's research (2020), based on a literature study of several types of citrus plants based on chemical compounds that have the potential to be natural larvicides of *Aedes aegypti*, with the highest potential found in the skin of sweet orange (*Citrus aurantium*) fruit compared to other parts. The use of the ethanol extract of the skin and Citrus amblycarpa leaves has the potential to become a natural larvicide and in the future, it can be applied to the community as a substitute for the use of chemical larvicides in killing *Aedes aegypti* larvae, but before application, it is necessary to do more in-depth studies and further research related to other plant extract formulations that can synergize with Citrus amblycarpa plant is like from the leaves which are indeed very small percentage in killing *Aedes aegypti* larvae.

CONCLUSION

Peel Extract can kill *Aedes aegypti* Larvae. Up to 96%. Bark extract requires less time and concentration to kill *Aedes aegypti* larvae compared to Leaf Extract. There was no significant difference in larval mortality between the 4000 ppm skin group and the control group. The ethanol extract of Citrus amblycarpa skin has potential as a larvicide for *Aedes aegypti*.

CONFLICT OF INTEREST STATEMENT

We declare that we have no conflicts of interest.

ACKNOWLEDGEMENTS

Thanks to the Directorate of Research and Community Service, Directorate General of Research and Development Strengthening of the Ministry of Research, Technology and Higher Education for funding this research. The author would like to thank the Chairperson of the LPPM Kalimantan Islamic University, and the Chair of the Department of Parasitology, Faculty of Veterinary Medicine, Airlangga University, the Tanah Bumbu Research and Development Center for Health, as well as the entire research team who participated in this research.

REFERENCES

1. Tosepu R, Tantrakarnapa K, Nakhapakorn K, Worakhunpiset S. Climate variability and dengue hemorrhagic fever in Southeast Sulawesi Province, Indonesia. *Environ Sci Pollut Res*. 2018;25(15):14944-52.
2. Mapalagamage M, Handunnetti S, Premawansa G, Pillainathan S, Fernando T, Kanapathipillai K, et al. Is total serum nitrite and nitrate (NOx) level in dengue patients a potential prognostic marker of dengue hemorrhagic fever? *Dis Markers*. 2018;2(2):1-9.
3. WHO. Better Environmental Management for Control of Dengue. WHO [Internet]. 2017; Available from: <https://www.who.int/heli/risks/vectors/denguecontrol/en/>
4. Retnaningrum OTD, Martini M, Raharjo M. Incidence of Dengue Hemorrhagic Fever (DHF) In Semarang Coastal Area: Epidemiology Descriptive Case and Bionomic Vector. *Indones J Trop Infect Dis*. 2019;7(6):144-9.
5. Maharani A, Sucipto TH, Setyawati H, Untoro YM, Sholihah NF, Churrotin S. In Vitro Antiviral Activity of Morin Compound against Dengue Virus Type 1 in Vero Cells. *Syst Rev Pharm*. 2020;11(9):830-3.
6. Ministry of Health of the Republic of Indonesia. Indonesia Health Profile 2019. Jakarta: Ministry of Health of the Republic of Indonesia; 2020.
7. Susilowati RP, Darmanto W, Aminah NS. "Morizena" Against *Aedes aegypti* DEATH. *Indones J Trop Infect Dis*. 2018;7(2):50-5.
8. Ridha MR, Nisa K. Larva *Aedes aegypti* is Tolerant to Temepos in Banjarbaru. *Vektora*. 2010;III(2):93-111.
9. Chansang A, Champakaew D, Junkum A. Potential of Natural Essential Oils and Cinnamaldehyde as Insecticides Against the Dengue Vector *Aedes aegypti* (Diptera: Culicidae). *Southeast Asian J Trop Med Public Heal*. 2018;49(1):6-22.
10. M. Rasyid Ridha, Wulan Sembiring, Abdullah Fadilly SS. Entomological Indicators and Resistance Status of Dengue Hemorrhagic Fever (*Aedes aegypti* L) Vectors of Insecticides in Banjarbaru City. In: *Prosiding Seminar Nasional Seri 8*. 2018. p. 128-42 (In Indonesian).
11. Poedji Hastutiek AS and RHP. Permot (*Passiflora Foetida* linn.) Leaf Extracts as Bioinsecticide Against *Aedes aegypti* Larvae. *Southeast Asian J Trop Med Public Heal*. 2017;48(6):1169-74
12. Anwar C, Syukur KMY, Dalilah D, Salni S, Novrikasari N. The Efficacy of Red Ginger Fraction (*Zingiber officinale* Roscoe var. rubrum) as Insecticidal *Aedes aegypti*. *Biosci Med*. 2018 Apr;2(2):31-41.
13. Irwan A, Mustikasari K, Ariyani D. Preliminary Chemical Examination of Leaves, Skins and Kuit Limes: Jeruk Lokal Kalimantan Selatan. *Sains dan Terap Kim*. 2017;11(2):71-9 (In Indonesian).
14. Kasman K, Ishak NI, Hastutiek P, Suprihati E, Mallongi I. Identification of Active Compounds of Ethanol Extract of Citrus amblycarpa leaves by Analysis of Thin-layer Chromatography and Gas Chromatography-Mass Spectrometry as Bioinsecticide Candidates for Mosquitoes. *Open Access Maced J Med Sci*. 2020;8(T2):1-6.
15. Ishak NI. Efektivitas Ekstrak Kulit Buah Limau Kuit (*Citrus Amblycarpa*) sebagai Larvasida *Aedes Aegypti* Instar III Effectiveness of Lime Skin Extract (*Citrus Amblycarpa*) as Natural Larvicide *Aedes Aegypti* Instar III. *J MKMI*. 2019;15(3):302-10.

16. Mirnawaty S dan BJ. A Test on the Effectiveness of Lansium Peel Extract (*Lansium Domesticum*) as Mosquito Electric Repellent Against *Aedes aegypti* Mosquitoes. *J Akad Kim.* 2012;1(4):147–52.
17. Perumalsamy H, Jang MJ, Kim J, Kadarkarai M, Ahn Y. Larvicidal activity and possible mode of action of four flavonoids and two fatty acids identified in *Milletia pinnata* seed toward three mosquito species. *Parasites Vectors Biomed Cent.* 2015;8(237):1–14.
18. Azevedo FR de, Maciel GC, Silva GBO e, Mesquita F de O, Alves ACL. Larvicidal Activity of Native Plant Extracts From the Araripe National Forest on *Aedes aegypti*. *J Agric Sci.* 2019;11(7):105.
19. Muangmoon R, Junkum A, Chaithong U. Natural Larvicides of Botanical Origin Against Dengue Vector *Aedes aegypti* (DIPTERA: CULICIDAE). *Southeast Asian J Trop Med Public Health.* 2018;49(2):227–39.
20. Cahyati WH. Biolarvicidal Effects of Papaya Leaves Juice Against *Aedes Aegypti* Linn Larvae. *J Int Dent Med Res.* 2019;12(2):780–6.
21. Husnawati, Prapti Sedijani NL. *The Effectiveness of Kirinyuh Leaf Ethanol Extract (Eupatorium Odoratum L) Against Mortality of Aedes Aegypti Mosquito Larvae as Enrichment of Animal Physiology Practicum.* Mataram; 2018.
22. Kasman, Ishak NI, Hastutiek P, Suprihati E. Identifikasi Senyawa Aktif Ekstrak Etanol Kulit Jeruk Limau Kuit (*Citrus Amblycarpa*) Dan Potensi Sebagai Bioinsektisida Pada Nyamuk *Aedes Aegypti*. In: Sustainable Strategic For Disaster Management In Wetland Area. Banjarbaru: Program Studi Kesehatan Masyarakat Fakultas Kedokteran Universitas Lambung Mangkurat; 2019. p. 350–7.
23. Muh. Saleh, Andi Susilawaty, Syarfaini M. Uji Efektivitas Ekstrak Kulit Buah Jeruk Nipis (*Citrus aurantifolia*) Sebagai Insektisida Hayati Terhadap Nyamuk *Aedes aegypti*. *J Hig.* 2017;3(1):30–6.
24. Abu Naim, Indasah N. Effectiveness of Giving Red Lemongrass (*Cymbopogon Citratus*) as a Mosquito *Aedes Aegypti* Larvaside. In: The 2nd Strada International Conference on Health. 2020. p. 287–95.
25. Kasman, Ishak NI, Hastutiek P, Suprihati E. GC-MS Analysis Citrus amblycarpa Peel Ethanol Extract and Potential as a Bioinsecticides against Mosquitoes. In: Kumpulan Abstrak Seminar Nasional 2019 “Percepatan Hilirisasi Hasil Penelitian di Era Industri 40” [Internet]. Bogor: Lembaga Penelitian dan Pengabdian kepada Masyarakat Institut Pertanian Bogor; 2019. p. 104. Available from: https://lppm.ipb.ac.id/publikasi/prosiding-seminar-ppm/?page_id=208&id=21&abstrak=777
26. Sukmayati Alegantina AI. Identification and Determination of Coumarine Compounds in Methanol Extract of *Artemisia annua* L. *Chromatography Of Thin Layer - Densitometri.* *Bul Penelit Kesehat.* 2010;38(1):17–28.
27. Sandoval-Montemayor NE, García A, Elizondo-Treviño E, Garza-González E, Alvarez L, del Rayo Nacho-Corona M. Chemical composition of hexane extract of *Citrus aurantifolia* and anti-*Mycobacterium tuberculosis* activity of some of its constituents. *Molecules.* 2012 Sep;17(9):11173–84.
28. Narayanaswamy VK, Gleiser RM, Kasumbwe K, Aldhubiab BE, Attimarad M V, Odhav B. Evaluation of Halogenated Coumarins for Antimosquito Properties. *Sci World J.* 2014;1–6.
29. Hamdan DI, Mohamed ME, Abdulla RH, Mohamed SM. Anti-inflammatory, insecticidal and antimicrobial activities and chemical composition of the essential oils of different plant organs from navel orange (*Citrus sinensis* (L.) Osbeck var. Malesy) grown in Egypt. *J Med Plants Res.* 2013;7(18):1204–15.
30. Favela-hernández JM, González-santiago O, Ramírez-cabrera MA, Esquivel-ferriño PC, Camacho-corona MR. Chemistry and Pharmacology of *Citrus sinensis*. *Molecules.* 2016;21(247):1–24.
31. Abdelatif HM, Ahmed S, Ahmed M. Insecticidal and Antibacterial Activity of Citrus Fruits ' Peels and Juices. Sudan: Department of Zoology, Faculty of Science University of Khartoum; 2004. 1–148 p.
32. Pavela R, Maggi F, Benelli G. Coumarin (2H-1-benzopyran-2-one): a novel and eco-friendly aphicide. *Nat Prod Res.* 2019 Sep;1–6.
33. Tando E. Review: Potensi Senyawa Metabolit Sekunder dalam Sirsak (*Annona muricata*) dan Srikaya (*Annona squamosa*) sebagai Pestisida Nabati untuk Pengendalian Hama dan Penyakit pada Tanaman. *J Biotropika.* 2018;6(1):21–7.
34. Supenah P, Taiman T, Sas OA. The effect of orange water of lemon (*Citrus Limon* (L.) Osbeck) as a larvasid of *Aedes Aegypti* in third instar. *J Phys Conf Ser.* 2019;1360(1).
35. Koneri R, Pontororing HH. Extract Test Of Mahoni Seeds (*Swietenia macrophylla*) Against *Aedes aegypti* Larvae Vector of Blood Female Disease. *J MKML.* 2016;12(4):216–23 (In Indonesian).
36. Ervina N. Ethanol Extract Activity Test of Singkong Leaves (*Manihot utilisima* Pohl) as Larvacides *Aedes aegypti*. In: Naskah Publikasi. 2014. p. 1–17.
37. P.J Hylands A. S. Maragenin I, II, and III, New Pentacyclic Triterpenes From *Marah Macrocarpus*. *Tetrahedron.* 1978;35:417–20.
38. Itokawa H, Nakajima H, Ikuta A, Iitaka Y. Two triterpenes from the flowers of *Camellia japonica*. *Phytochemistry.* 1981;20(11):2539–42.
39. Srivastava US, Jaiswal AK. An insect growth regulatory factor in *Cuscuta reflexa* Roxby. *Natl Acad Sci Lett.* 1990;13(9):361–3.
40. I gusti Agung Ayu Kusuma Wardani, Fitria Megawati PS. (Effectiveness of Electric Liquid From Gumitir Extract (*Tagetes erecta* L.) AS *Aedes aegypti* Repellent). *J Ilm Medicam.* 2019;5(1):1–5.

Potential Extract Ethanol Citrus Amblycarpa as a Bioinsecticide Against Aedes Aegypti Larvae

ORIGINALITY REPORT

19%

SIMILARITY INDEX

16%

INTERNET SOURCES

15%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	www.id-press.eu Internet Source	5%
2	Grewal, P.S.. "Rapid Changes in Thermal Sensitivity of Entomopathogenic Nematodes in Response to Selection at Temperature Extremes", <i>Journal of Invertebrate Pathology</i> , 199607 Publication	2%
3	journal.unhas.ac.id Internet Source	1%
4	repository.uin-malang.ac.id Internet Source	1%
5	kesmas.ulm.ac.id Internet Source	1%
6	jurnal.poltekeskupang.ac.id Internet Source	1%
7	Proceeding of LPPM UPN "Veteran" Yogyakarta Conference Series 2020 – Engineering and Science Series, 2020	1%

8	lppm.ipb.ac.id Internet Source	1 %
9	Nurlailah Nurlailah, Anny Thuraidah. "Biolarvasides of Cananga odorata Flower and Zyzygium polyanthum Leaves Ethanol Extract on Aedes aegypti Larva", Medical Laboratory Technology Journal, 2020 Publication	1 %
10	www.mysciencework.com Internet Source	<1 %
11	www.mdpi.com Internet Source	<1 %
12	Tri Wahyuni Sukesi, - Sulistyawati, Eva Hendrawati, Surahma Asti Mulasari. "Effectivity of Bacterial Suspension Bacillus thuringiensis Var Israelensis in Killing Aedes aegypti L. Mosquito Larvae", Bangladesh Journal of Medical Science, 2019 Publication	<1 %
13	ecampus.poltekkes-medan.ac.id Internet Source	<1 %
14	zombiedoc.com Internet Source	<1 %
15	Fouad El-Akhal, Abdelhakim El Ouali Lalami, Raja Guemmouh. "Larvicidal activity of	<1 %

essential oils of Citrus sinensis and Citrus aurantium (Rutaceae) cultivated in Morocco against the malaria vector Anopheles labranchiae (Diptera: Culicidae)", Asian Pacific Journal of Tropical Disease, 2015

Publication

16

repository.kmou.ac.kr

Internet Source

<1 %

17

jurnal.unismuhpalu.ac.id

Internet Source

<1 %

18

www.bioscmed.com

Internet Source

<1 %

19

Valentine C. Mbatchou, David P. Tchouassi, Rita A. Dickson, Kofi Annan et al. "Mosquito larvicidal activity of Cassia tora seed extract and its key anthraquinones aurantio-obtusin and obtusin", Parasites & Vectors, 2017

Publication

<1 %

20

eber.uek.krakow.pl

Internet Source

<1 %

21

jurnal.fkmumi.ac.id

Internet Source

<1 %

22

librepo.stikesnas.ac.id

Internet Source

<1 %

23

res.mdpi.com

Internet Source

<1 %

24

www.tandfonline.com

Internet Source

<1 %

25

Kuntum Ekawati Nurdin, Loisa Ratna Yuvita Olla, Stefany Febiand Feoh, Anita Dwi Putri Galla et al. "Effectivity Test of 96% from Soe (Citrus sinensis L.) Sweet Orange Rind Ethanol Extract as Biolarvaside", JURNAL INFO KESEHATAN, 2019

Publication

<1 %

26

Edible Medicinal and Non Medicinal Plants, 2014.

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On