

ASSOCIATION OF SERUM MAGNESIUM LEVELS WITH GLYCEMIC CONTROL IN PATIENTS WITH TYPE 2 DIABETES MELLITUS

by Sony Wibisono

Submission date: 13-Apr-2021 03:25PM (UTC+0800)

Submission ID: 1557966309

File name: Artikel.pdf (272.05K)

Word count: 2044

Character count: 10985



ISSN: 0975-833X

Available online at <http://www.journalcra.com>

International Journal of Current Research
Vol. 12, Issue, 01, pp.9531-9533, January, 2020

DOI: <https://doi.org/10.24941/ijer.35638.01.2020>

INTERNATIONAL JOURNAL
OF CURRENT RESEARCH

RESEARCH ARTICLE

ASSOCIATION OF SERUM MAGNESIUM LEVELS WITH GLYCEMIC CONTROL IN PATIENTS WITH TYPE 2 DIABETES MELLITUS

1,*Sony Wibisono Mudjanarko and 2,3Huwainan Nisa Nasution

¹Endocrine and Metabolic Division of the Department of Internal Medicine Airlangga University, Indonesia

²Department of Internal Medicine Airlangga University, Indonesia

³Department of Internal Medicine Muhammadiyah University of North Sumatera, Indonesia

ARTICLE INFO

Article History:

Received 12th October, 2019
Received in revised form
28th November, 2019
Accepted 09th December, 2019
Published online 30th January, 2020

Key Words:

Serum magnesium levels,
Glycemic control,
HbA1c,
Type 2 diabetes mellitus.

*Corresponding author:

Sony Wibisono Mudjanarko

Copyright © 2020, Sony Wibisono Mudjanarko and Huwainan Nisa Nasution. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Sony Wibisono Mudjanarko and Huwainan Nisa Nasution, 2020. "Association of serum magnesium levels with glycemic control in patients with type 2 diabetes mellitus", International Journal of Current Research, 12, (01), 9531-9533.

INTRODUCTION

The prevalence of type 2 diabetes mellitus (T2DM) has been increasing exponentially with high rates of diabetes-related morbidity and mortality due to complications (Bergman *et al.*, 2012). Hypomagnesemia is one of the conditions that can accelerate complication in patients with T2DM (Chhabra *et al.*, 2017). The prevalence of hypomagnesemia in patients with type 2 diabetes varies between 25-39% (Gunther, 2010). Several studies have shown that hypomagnesemia is significantly associated with poor glycemic control and increased incidence of diabetes-related complications within a certain period of time (Sales *et al.*, 2011; Dasgupta *et al.*, 2012; Gupta *et al.*, 2014; Sadiq and Nawaz, 2014; and Shariff, 2015). Magnesium plays an important role in carbohydrate metabolism as a critical cofactor for the mechanism of glucose transport in membrane cells and the activities of various enzymatic reactions involved in carbohydrate oxidation (Hans *et al.*, 2002). Magnesium deficiency has been associated with worsening of plasma glycemic control due to an impairment of both, insulin secretion and insulin action (Chhabra *et al.*, 2017; Gommers *et al.*, 2016).

ABSTRACT

Background: Hypomagnesemia is one of the conditions that can accelerate diabetes complications in patients with type 2 diabetes mellitus (T2DM) by altered insulin function and secretion. Hypomagnesemia has been linked to poor glycemic control due to increased insulin resistance and impaired of insulin secretion. **Objective:** The aim of this study is to analyze the association between serum magnesium levels and glycemic control in patients with T2DM who treated with metformin or pioglitazone. **Methods:** Serum magnesium levels and HbA1c were examined from a total of 41 patients with T2DM treated with metformin ≥ 750 mg/day for at least 3 weeks or pioglitazone ≥ 15 mg/day for at least 4 weeks with a body mass index (BMI) of < 30 kg/m². An examination of HbA1c was analyzed using Variant and D10 with National Glycohaemoglobin Standardization Program (NGSP) certification. The serum magnesium level was analyzed using Roche/Hitachi Cobas C 311/501 System. **Results:** The mean value of serum magnesium levels was 2.04 ± 0.19 mg/dl and the mean of HbA1c was $7.44 \pm 1.57\%$. There was significant negative correlation between serum magnesium levels with HbA1c ($r = -0.449$, p -value 0.003). Low serum magnesium levels was correlated with poor glycemic control. **Conclusion:** Serum magnesium levels in patients with T2DM treated with metformin or pioglitazone was inversely correlated to glycemic control. Serum magnesium levels monitoring is important to reduce the risk of diabetes-related complications.

MATERIALS AND METHODS

It is an observational analytic study with cross-sectional design conducted at Private Practice of Internists of Metabolic and Diabetes Endocrinology Consultants in Surabaya, Indonesia. The study was approved by the Committee of the Health Research Ethics of the Faculty of Medicine, Airlangga University, Surabaya. The inclusion criteria were patients with T2DM, more than 18 years, use of metformin ≥ 750 mg/day for at least 3 weeks or pioglitazone ≥ 15 mg/day for at least 4 weeks and the body mass index (BMI) < 30 Kg/m². The exclusion criteria included impaired renal function with serum creatinine > 1.2 mg/dl for women and > 1.5 mg/dl for men, pregnant or lactating women, acute infections or inflammation, gastrointestinal and chronic liver diseases, history of drinking alcohol or smoking, taking proton pump inhibitors, diuretics, aminoglycoside drugs, amphotericin B, vitamin and mineral supplementation, steroids and history of getting therapy of cetuximab, erlotinib, cisplatin, carboplatin, cyclosporine or tacrolimus. The examinations of HbA1c and serum magnesium levels were done. HbA1c was analyzed using Variant and D10 with National Glycohaemoglobin Standardization Program (NGSP) certification. The serum magnesium level was

analyzed using Roche/Hitachi Cobas C 311/501 System. All data were analyzed using SPSS version 20.0 software. Bivariate analysis of correlations among the variables was performed by Rank Spearman test. The results were presented in a correlation coefficient (r-value), and the significant p-value was < 0.05 and the confidence interval was 95%.

RESULTS

The study involved 41 subjects of patients with T2DM who met the inclusion and exclusion criteria. The characteristics of the subjects are presented in Table 1. The mean of serum magnesium levels was 2.04 ± 0.19 mg/dl. The mean of HbA1c was $7.44 \pm 1.57\%$. Bivariate analysis by using with Rank Spearman test related to the association of serum magnesium levels with HbA1c. Serum magnesium levels had a significant association with HbA1c with p-value 0.003 and correlation coefficient -0.449. These data are shown in Table 2. Figure 1 shows a graph of the correlation between serum magnesium level and HbA1c.

Table 1. Characteristics of study participants

Characteristics	n = 41
Female, n (%)	30 (73.2%)
Age (years), mean (SD)	60.3 (11.5)
Duration of type 2 DM (years), mean (SD)	11.8 (10.11)
Dyslipidemia, n (%)	25 (65.9)
Use of statin, n (%)	27 (65.9)
Use of antihypertensive, n (%)	5 (12.2)
Use of insulin, n (%)	13 (31.7)
BT (Kg/m ²), mean (SD)	25.4 (2.8)
Fasting plasma glucose (mg/dl), mean (SD)	131.8 (40.95)
Postprandial plasma glucose (mg/dl), mean (SD)	185 (78.43)
HbA1c (%), mean (SD)	7.44 (1.57)
Total cholesterol (mg/dl), mean (SD)	167.41 (37.90)
Triglyceride (mg/dl), mean (SD)	127.9 (52.9)
LDL (mg/dl), mean (SD)	95.94 (34.08)
HDL (mg/dl), mean (SD)	47.64 (15.41)
Serum creatinine (mg/dl), mean (SD)	0.79 (0.18)
eGFR (ml/minute), mean (SD)	86.8 (17.62)
Serum magnesium (mg/dl), mean (SD)	2.04 (0.19)

Table 2. Association of serum magnesium levels with fasting plasma glucose

Independent variable	Dependent variable	r-value	p-value
Serum magnesium	HbA1c	-0.449	0.003




Figure 1. Graph of correlation of serum magnesium levels with HbA1c

DISCUSSION

The characteristics of the subjects were dominated by women. It is different from Pokharel *et al.* (2017) in Nepal and Gupta *et al.* (2014) in India which the subjects were mostly men. This result is in line with previous data in Indonesia taken from Riskesdas 2013 indicated that the prevalence of diabetes was higher in women (Gupta *et al.*, 2014; Pokharel *et al.*, 2017). The mean of age was 60 years and the dominant age group was 46-60 years (46.3%). A different result was shown by Gupta *et al.* (2014) in which dominated by the age group of 34-45 years. The study is in line with the previous study of Riskesdas 2013 which dominated by the age group of 45-64 years. The mean of serum magnesium levels in this study was 2.04 ± 0.19 mg/dl. A similar result was shown by Tarigan *et al.* (2015) in Jakarta which the serum magnesium levels < 1.7 mg/dl was 13.2% and ≥ 1.7 mg/dl was 86.8%, mostly in normomagnesemia group (Tarigan *et al.*, 2015). This study also in line with Pokharel *et al.* (2017) with the mean serum magnesium levels was 2.10 ± 0.16 mg/dl (Pokharel *et al.*, 2017). The different results were shown in the previous studies in India by Gupta *et al.* (2014) with the mean of serum magnesium levels were 1.2 mg/dl (Gupta *et al.*, 2014). The mean HbA1c shown in this study was $7.44 \pm 1.57\%$. Different results were shown by Pokharel *et al.* (2017) and Gupta *et al.* (2014) with an average of $8.4 \pm 1.2\%$ and $8.2 \pm 0.45\%$. The different result probably caused by this study was conducted in a private clinic and the majority of subjects has been well-controlled (82.9%). The second reason was the subjects has been treated with metformin or pioglitazone resulted in lower HbA1c. In this study, the serum magnesium levels had a significant negative correlation to HbA1c with r-0.449 and p-value 0.003. It is similar to Gupta *et al.* (2014) with r-0.29; p < 0.0001 and Pokharel *et al.* (2017) with r-0.299; p < 0.01. Low serum magnesium levels in patients with T2DM is associated with poor glycemic control. The variation in the study may be due to several factors including different research design, the use of metformin or pioglitazone, and other confounding factors such as diet, physical activities, and dyslipidemia. The limitations and weaknesses of this study were uncontrolled confounding variables such as diet, physical activity, and dyslipidemia and unprovided overall antidiabetic drug equation.

Conclusion

Serum magnesium levels in patients with T2DM treated with metformin or pioglitazone was inversely correlated to glycemic control. Serum magnesium levels monitoring is important to reduce the risk of diabetes-related complications.

REFERENCES

- Bergman M. Pathophysiology of prediabetes and treatment implications for prevention of type 2 diabetes mellitus. *Endocrine*. 2012; 43(3): 504-513. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23132321>
- Chhabra S, Chhabra S, Ramessur K, Chhabra N. Hypomagnesemia and its implications in type 2 diabetes mellitus-A review article. *Webmed Central Diabetes*. 2012. Available from: https://www.webmedcentral.com/wmcpdf/Article_WMC004005.pdf [Accessed 14 June 2017].
- Gunther T. The biochemical function of Mg²⁺ in insulin secretion, insulin signal transduction and insulin resistance. *Magnesium Research*, 2010; 23(1): 5-18.

- Sales CH, Pedrosa LFC, Lima JG, Lemos TMAM, Colli C. Influence of magnesium status and magnesium intake on the blood glucose control in patients with type 2 diabetes. *Clinical Nutrition*, 2011; 30: 359-364.
- Dasgupta A, Sarma D, Saikia UK. Hypomagnesemia in type 2 diabetes mellitus. *Indian Journal of Endocrinology and Metabolism*. 2012; 16(6).
- Gupta NK, Sharma S, Kaushik GG, Gupta B. Association between indexes of insulin sensitivity/resistance and serum magnesium levels in overweight diabetic subjects. *Indian Journal of Clinical Practice*, 2014; 24 (10): 932-937.
- Sadiq F, Nawaz MR. Hypomagnesemia in patients with type 2 diabetes. *Pakistan Journal of Medial Science*, 2014; 8 (4).
- Rao P, Shariff MG. Serum magnesium levels in type 2 diabetic patients with microalbuminuria and normoalbuminuria. *International Journal of Scientific Study*, 2015; 3(4).
- Hans CP, Sialy R, Bansal DD. Magnesium deficiency and diabetes mellitus. *Current Science*, 2002; 83 (12).
- Gommers LMM, Hoenderop JGJ, Bindels RJM, De Baaji JHF. Hypomagnesemia in type 2 diabetes: A vicious circle?. *Diabetes*, 2016; 65: 3-13. WMC003878.
- Pokharel DR, Khadka D, Sigdel M, Yadav N, Kafle R, Sapkota RM, et al. The association of serum magnesium level with poor glycemic control and renal functions in Nepalese patients with type 2 diabetes mellitus. *Diabetes and metabolic syndrome: clinical research and reviews*, 2017; Available at: <http://dx.doi.org/10.1016/j.dsx.2017.03.028>.
- Infodatin (Pusat Data dan Informasi Kementerian Kesehatan RI). Situasi dan analisis diabetes. 2014. Available from: <http://www.depkes.go.id/resources/download/pusdatin/infodatin/infodatin-diabetes.pdf>.
- Tarigan TJE, Marbun MBH, Harimurti K. Korelasi kadar magnesium serum dengan albuminuria pada pasien diabetes mellitus tipe 2. *Journal Kedokteran Indonesia*. 2015; 3 (2): 115-119.

ASSOCIATION OF SERUM MAGNESIUM LEVELS WITH GLYCEMIC CONTROL IN PATIENTS WITH TYPE 2 DIABETES MELLITUS

ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

15%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Nènè Oumou Kesso Barry, Michel Assane Ndour, Moustapha Djite, Demba Diedhiou et al. "Association of Magnesium Status with Poor Glycemic Control and Microangiopathic Complications (Neuropathy and Nephropathy) in Diabetic Patients in Dakar", *Advances in Biochemistry*, 2019
Publication 5%
- 2 www.mdpi.com
Internet Source 2%
- 3 eprints.covenantuniversity.edu.ng
Internet Source 2%
- 4 "Poster Session Clinical", *European Journal of Heart Failure*, 2013.
Publication 1%
- 5 Hasan Korkmaz, Ilgin Karaca, Mustafa Koç, Orhan Önalın, Mücahit Yılmaz, Mehmet Nail Bilen. "Early Effects of Treatment with Nebivolol and Quinapril on Endothelial

Function in Patients with Hypertension", Endothelium, 2009

Publication

6

NH Rekha, MS Bharath, SP Channakeshava.
"Study of Prevalence of Hypomagnesemia in
Patients with Type II Diabetes Mellitus", The
Journal of Medical Sciences, 2017

Publication

1 %

7

Kalinga B.E., Shruthi C. Bhojashettar, Salman
Beig, Niranjan M.. "Study of Serum
Magnesium and Its Correlation with HbA1c in
Patients with Type 2 Diabetes Mellitus",
Journal of Evidence Based Medicine and
Healthcare, 2020

Publication

1 %

8

Haruka Tamura, Yoshinobu Kondo, Kohei Ito,
Masanori Hasebe, Shinobu Satoh, Yasuo
Terauchi. "Comparison of the effects of
empagliflozin and glimepiride on endothelial
function in patients with type 2 diabetes: A
randomized controlled study", Research
Square, 2019

Publication

1 %

9

Rosnani Pangaribuan, Jürgen G Erhardt,
Veronika Scherbaum, Hans K Biesalski.
"Vitamin A capsule distribution to control
vitamin A deficiency in Indonesia: effect of
supplementation in pre-school children and

1 %

compliance with the programme", Public Health Nutrition, 2007

Publication

10

Vahideh Behrouz, Ali Dastkhosh, Golbon Sohrab. "Overview of dietary supplements on patients with type 2 diabetes", Diabetes & Metabolic Syndrome: Clinical Research & Reviews, 2020

Publication

1 %

11

K EGUCHI, Y MATSUI, S SHIBASAKI, J ISHIKAWA, S HOSHIDE, S ISHIKAWA, T KABUTOYA, J SCHWARTZ, T PICKERING, K SHIMADA. "Changes in Self-Monitored Pulse Pressure Correlate With Improvements in B-Type Natriuretic Peptide and Urinary Albumin in Treated Hypertensive Patients", American Journal of Hypertension, 2007

Publication

1 %

12

S Sushma, MS Kusuma Devi, S Naveen. "The Study of Relationship between Serum Magnesium and Glycated Hemoglobin in Clinically Uncomplicated Type 2 Diabetes Mellitus Patients", Journal of Health Sciences & Research, 2015

Publication

1 %

13

bmcpublikealth.biomedcentral.com

Internet Source

1 %

14

Internet Source

1 %

15

repositorio.ufsm.br

Internet Source

1 %

16

www.researchsquare.com

Internet Source

1 %

17

Tri Wahyuliati, Agung Pranoto, Samekto Wibowo. "Serum Level Changes of Neurotrophin-3 after Performing Diabetic Foot Exercise in Diabetic Neuropathy", International Journal of Public Health Science (IJPHS), 2015

Publication

1 %

18

serval.unil.ch

Internet Source

1 %

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On

ASSOCIATION OF SERUM MAGNESIUM LEVELS WITH GLYCEMIC CONTROL IN PATIENTS WITH TYPE 2 DIABETES MELLITUS

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
