

A Preliminary Study Troponin T and Reg3 β in Children with Left-to-Right Shunt Congenital Heart Disease with Heart Failure

by I Ketut Alit Utamayasa

Submission date: 20-Mar-2021 10:32AM (UTC+0800)

Submission ID: 1537516421

File name: t-to-Right_Shunt_Congenital_Heart_Disease_with_Heart_Failure.pdf (402.97K)

Word count: 2507

Character count: 13410

A Preliminary Study: Troponin T and Reg3 β in Children with Left-to-Right Shunt Congenital Heart Disease with Heart Failure

I. Ketut Alit Utamayasa¹, Mahrus A. Rahman¹, Taufiq Hidayat¹, Rendi Aji Prihaningtyas²

¹Lecturer, ²Pediatric Resident, Department of Child Health, Faculty of Medicine, Universitas Airlangga, Dr. Soetomo General Hospital, Surabaya

Abstract

Background: Congenital heart disease (CHD) cause heart failure and myocardial injury. Troponin in the heart is a biomarker of myocardial injury in adults and children. Studies that examine troponin T and Reg3 β in children with left-to-right shunt CHD with heart failure are still limited.

Objective: This study aims to analyse the troponin T and Reg3 β in children with left-to-right shunt CHD with heart failure compared to children without heart failure.

Method: This study was a case control study of children with left-to-right shunt CHD with heart failure and children with left-to-right shunt CHD without heart failure performed with non-random sampling consecutive techniques at the Dr. Soetomo General Hospital, Surabaya in April–June 2019. The diagnosis of left-to-right shunt CHD was determined based on echocardiographic examination. All subjects with left-to-right shunt CHD were evaluated using the Paediatric Heart Failure Score. Troponin T examination was carried out using a one-dimensional electrophoresis technique, which is 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). Reg3 β examination was carried out by the ELISA method. Data analysis was performed with an independent sample t test using the SPSS.

Results: This study involved 11 children, consisting of 7 children with left-to-right shunt CHD with heart failure and 4 children with left-to-right shunt CHD without heart failure. Most children (72.7%) were female, 3 children (27.3%) were \leq 5 years old, 5 children (45.4%) were 5–10 years old and 3 children (27.3%) were $>$ 10 years old. There was a significant increase in the Troponin T and Reg3 β in children with left-to-right shunt CHD with heart failure as compared to children without heart failure.

Conclusion: Troponin T and Reg3 β can be used as biomarkers in children with left-to-right shunt CHD with heart failure.

Keywords: Congenital Heart Disease, Troponin T, Reg3 β , Heart Failure.

22

Introduction

Congenital heart disease (CHD) is a congenital disorder that often occurs in children¹. Excessive pressure and volume in children with CHD creates the risk of injury to the myocardium². Troponin in the heart is a biomarker of myocardial injury in adults³. Injury to the myocardium is the cause of elevated troponin levels in 60% of cases⁴. Studies in children mention the role of troponin as a diagnostic marker of myocarditis in children⁵.

Corresponding Author:

I Ketut Alit Utamayasa,

Department of Child Health, Faculty of Medicine,
Universitas Airlangga, Dr. Soetomo General Hospital,
Jl. Mayjen Prof. Dr. Moestopo No. 47, Airlangga,
Surabaya, East Java, Indonesia
Contact No.: +62 31 5501681
e-mail: alit_tusari@yahoo.com

Injury to the myocardium raises the body's response by triggering proliferation, phagocytosis and M2 macrophage polarization⁶. This role is mediated by Reg3 β , which is increased due to cardiac inflammation⁷ and provides a protective mechanism when cardiac injury and stress occur⁶. Previous studies have mentioned an increase in the level of injured Reg3 β ⁸.

Several studies of biomarkers of myocardial injury have been carried out to predict the prognosis and improve outcomes in patients. Troponin T is associated with the degree of myocardial damage and can predict morbidity and mortality due to heart disease⁹. Troponin T is associated with prognosis and mortality in heart failure¹⁰. Another study stated that Reg3 β can be used as a prognostic factor in mortality in patients with acute coronary syndrome⁷. Studies that examine troponin T and Reg3 β in children with left-to-right shunt CHD with heart failure are still limited.

Material and Method

This research was a case control study conducted at the Paediatric Cardiology Outpatient Clinic, Emergency Room and Paediatric Ward, Dr. Soetomo General Hospital, Surabaya in April–June 2019. Subjects were children with left-to-right shunt CHD with heart failure with a comparison group, children without heart failure. The diagnosis of left-to-right shunt CHD was determined by echocardiographic examination. The types of cardiac abnormalities categorized as left-to-right shunt CHD included ventricular septal defect (VSD), atrial septal defect (ASD) and patent ductus arteriosus (PDA). Inclusion criteria in this study were age between 5 and 10 years and meeting the clinical criteria for heart failure according to the Paediatric Heart Failure Score. Exclusion criteria in this study were children who scheduled for surgery within the next month, impaired renal function, hyperkalaemia with serum potassium levels > 5.5 mEq/L and unstable clinical conditions, such as receiving intravenous inotropes, ventilator pneumonia and sepsis. Sampling was done by consecutive non-random sampling techniques. All subjects with left-to-right shunt CHD were evaluated using the Paediatric Heart Failure Score.

Troponin T measurement was carried out using a one-dimensional electrophoresis technique, which is 12% SDS-PAGE. Reg3 β measurement was performed using the ELISA method.

The mean difference between the two groups was evaluated by the independent sample t test if the data were normally distributed and the Mann-Whitney U test if the data were not normally distributed. The normality of data distribution was tested with the Shapiro-Wilk test. Differences in troponin T and Reg3 β in children with left-to-right shunt CHD with heart failure and children without heart failure were analysed using an independent sample t test. Data were analysed using the Statistical Package for Social Sciences (SPSS).

Findings: A total of 11 children were involved in this study, consisting of 8 (72.7 %) females, 3 children (27.3 %) \leq 5 years old, 5 children (45.4 %) 5–10 years old and 3 children (27.3 %) > 10 years old (Table 1).

Table 1. Subject Characteristics

Variable	Case n (%)	Control n (%)
Sex		
Female	5 (71.4 %)	3 (75 %)
Male	2 (28.6 %)	1 (25 %)
Age		
\leq 5 years old	2 (28.6 %)	1 (25 %)
5–10 years old	3 (42.8 %)	2 (50 %)
> 10 years old	2 (28.6 %)	1 (25 %)




Figure 1. Troponin T

Figure 1 showed the measurement of Troponin T using 12% SDS-PAGE. This study showed that there was a significant increase in 37 kDa protein distribution (Troponin T). The protein band profile obtained from the SDS-PAGE electrophoresis showed differences in

the synthesized protein bands. The group with left-to-right shunt CHD with heart failure had higher Troponin T levels as compared to the group without heart failure (Table 2).

Table 2. Level of Reg3 β and Troponin T

	Case (Mean \pm Standard Deviation)	Control (Mean \pm Standard Deviation)
Reg3 β (ng/mL)	19.063 \pm 0.619	9.978 \pm 0.678
Troponin T	42.600 \pm 15.545	6.831 \pm 5.61

Reg3 β ELISA results showed that there was a significant increase in Reg3 β levels in the left-to-right shunt CHD group with heart failure as compared to controls (Table 2).

Discussion

CHD is a congenital disorder that often occurs in children and accounts for up to 1/3 of cases of congenital abnormalities in children¹. CHD occurs in 6 to 10 per 1000 live births, with an average of 8 out of 1000 live births¹¹. The highest prevalence of CHD occurs in Asian countries, which is 9.3 out of 1000 live births¹. CHD is divided into several types, including right-to-left shunt CHD, obstructive heart disease, cyanotic heart disease and miscellaneous heart disease. The pathogenesis of CHD is multifactorial, including genetic factors or chromosomal abnormalities, environmental factors, maternal infections, smoking and alcohol during pregnancy, pregnancy with diabetes mellitus and obesity during pregnancy or interactions of all these factors¹².

Heart failure can be a complication of CHD. Left-to-right shunts of CHD that can generally cause heart failure include VSD, ASD, atrioventricular septal defect (AVSD) and patent ductus arteriosus (PDA) with moderate to large diameter defects. Children with CHD have a risk of myocardial injury due to excess pressure and volume².

Troponin is a single homogeneous protein consisting of four main protein fractions using SDS-PAGE, namely fraction 2, fraction 3 and fraction 4¹³. There are three forms of troponin, namely troponin T (TnT), troponin I (TnI) and troponin C (TnC)¹⁴. Troponins in the heart are distinguished by regions with different amino acid sequences. Fraction 2 (~24 kDa) is called TnI ('I' for inhibitory), which inhibits the activity of Mg²⁺-dependent actomyosin ATPase in the absence of Ca²⁺. Fraction 3 (~37 kDa) is bound to tropomyosin, so it is called TnT

('T' for tropomyosin), which connects tropomyosin and troponin complexes. Fraction 4 (~20 kDa) is bound to Ca²⁺ and is referred to as TnC ('C' for calcium), which regulates the activity of thin-filaments¹³.

Troponin and tropomyosin work to regulate muscle contraction¹⁴. Troponin levels increase significantly in children with VSD and ASD as compared to healthy children. This condition indicates a significant increase in volume and pressure because left-to-right shunts in CHD can cause damage to the myocardium¹⁵. Troponin correlates with oxygen saturation and ejection fraction in children with CHD².

This study found that children with left-to-right shunt CHD with heart failure had higher levels of troponin T as compared to children without heart failure. Troponin T is associated with myocardial injury in children. However, studies in new-borns have shown that troponin T levels were neither related to the type of heart failure nor the type of heart abnormality¹⁶.

Reg (Regenerating gene) is a protein that was first isolated from rat cDNA and consists of three subtypes, namely types I, II and III. Reg3 consists of Reg3 α , Reg3 β and Reg3 γ ¹⁷. Reg3 β was first found in mice models with pancreatitis¹⁸. Previous studies have suggested that Reg3 β was associated with the intensity of inflammation in the heart and increased in cases of acute coronary syndrome⁷.

In this study, there was an increase in Reg3 β levels in children with left-to-right shunt CHD with heart failure. Previous studies have suggested that Reg3 β was involved in the mechanism of protection against cardiac injury and stress⁶. Reg3 β levels increase in cardiac ischemia. Healing myocardium after injury requires macrophages⁸. Reg3 β plays a role in the repair of myocardial injury by triggering proliferation, phagocytosis and polarization of M2 macrophages⁶.

Conclusion

Troponin T and Reg3 β can be used as biomarkers in children with left-to-right shunt CHD with heart failure.

Conflict of Interest: The authors declare that there is no conflict of interest regarding this research.

Source of Funding: The authors received no specific grants from any funding agency in the public, commercial, or not-for-profit sectors.

Ethical Clearance: This study was approved by the Ethical Committee of Dr. Soetomo General Hospital, Surabaya No. 1198/KEPK/V/2019.

Reference

1. van der Linde D, Konings EEM, Slager MA, Witsenburg M, Helbing WA, Takkenberg JJM, et al. Birth Prevalence of Congenital Heart Disease Worldwide. *Journal of the American College of Cardiology*. 2011 Nov;58(21):2241–7.
2. Hafez MO, Morsy SM, Mahfoz RA, Ali AR. Myocardial Injury in Children with Unoperated Congenital Heart Diseases. *Cardiology Research and Practice*. 2015;2015:1–5.
3. Myhre PL, O'Meara E, Claggett BL, de Deus S, Jarolim P, Anand IS, et al. Cardiac Troponin I and Risk of Cardiac Events in Patients With Heart Failure and Preserved Ejection Fraction. *Circulation: Heart Failure* [Internet]. 2018 Nov [cited 2019 Dec 15];11(11). Available from: <https://www.ahajournals.org/doi/10.1161/CIRCHEARTFAILURE.118.005312>
4. McCarthy CP, Raber I, Chapman AR, Sandoval Y, Apple FS, Mills NL, et al. Myocardial Injury in the Era of High-Sensitivity Cardiac Troponin Assays: A Practical Approach for Clinicians. *JAMA Cardiology*. 2019 Oct 1;4(10):1034.
5. Eisenberg MA, Green-Hopkins I, Alexander ME, Chiang VW. Cardiac Troponin T as a Screening Test for Myocarditis in Children: Pediatric Emergency Care. 2012 Nov;28(11):1173–8.
6. Zhou S, Jiang H, Wang H, Lu H, Chen R, Xu H, et al. Reg3 β from cardiomyocytes regulated macrophage migration, proliferation and functional skewing in experimental autoimmune myocarditis. *Am J Clin Exp Immunol*. 2018;7(2):8–15.
7. Lörchner H, Widera C, Hou Y, Elsässer A, Warnecke H, Giannitsis E, et al. Reg3 β is associated with cardiac inflammation and provides prognostic information in patients with acute coronary syndrome. *International Journal of Cardiology*. 2018 May;258:7–13.
8. Lörchner H, Pöling J, Gajawada P, Hou Y, Polyakova V, Kostin S, et al. Myocardial healing requires Reg3 β -dependent accumulation of macrophages in the ischemic heart. *Nature Medicine*. 2015 Apr;21(4):353–62.
9. Lipshultz SE, Rifai N, Sallan SE, Lipsitz SR, Dalton V, Sacks DB, et al. Predictive Value of Cardiac Troponin T in Pediatric Patients at Risk for Myocardial Injury. *Circulation*. 1997 Oct 21;96(8):2641–8.
10. Kociol RD, Pang PS, Gheorghiade M, Fonarow GC, O'Connor CM, Felker GM. Troponin Elevation in Heart Failure. *Journal of the American College of Cardiology*. 2010 Sep;56(14):1071–8.
11. Wren C. The Epidemiology of Cardiovascular Malformations. In: Moller JH, Hoffman JIE, editors. *Pediatric Cardiovascular Medicine* [Internet]. Oxford, UK: Wiley-Blackwell; 2012 [cited 2019 Dec 21]. p. 268–75. Available from: <http://doi.wiley.com/10.1002/9781444398786.ch19>
12. Nakanishi T, Markwald RR, Baldwin HS, Keller BB, Srivastava D, Yamagishi H, editors. *Etiology and Morphogenesis of Congenital Heart Disease: From Gene Function and Cellular Interaction to Morphology* [Internet]. Tokyo: Springer; 2016 [cited 2019 Dec 21]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK500252/>
13. Park KC, Gaze DC, Collinson PO, Marber MS. Cardiac troponins: from myocardial infarction to chronic disease. *Cardiovascular Research*. 2017 Dec 1;113(14):1708–18.
14. Lippi G, Targher G, Franchini M, Plebani M. Genetic and biochemical heterogeneity of cardiac troponins: clinical and laboratory implications. *Clinical Chemistry and Laboratory Medicine* [Internet]. 2009 Jan 1 [cited 2019 Dec 10];47(10). Available from: <https://www.degruyter.com/view/j/cclm.2009.47.issue-10/cclm.2009.322/cclm.2009.322.xml>
15. Sugimoto M, Ota K, Kajihama A, Nakau K, Manabe H, Kajino H. Volume Overload and Pressure Overload due to Left-to-Right Shunt-Induced Myocardial Injury. *Circulation Journal*. 2011;75(9):2213–9.
16. Tarkowska A, Furmaga-Jabłońska W. The Evaluation of Diagnostic Role of Cardiac Troponin T (cTnT) in Newborns with Heart Defects. *The Scientific World Journal*. 2012;2012:1–6.

17. Narushima Y, Unno M, Nakagawara K, Mori M, Miyashita H, Suzuki Y, et al. Structure, chromosomal localization and expression of mouse genes encoding type III Reg, RegIII α , RegIII β , RegIII γ . *Gene*. 1997 Feb;185(2):159–68.
18. Iovanna J, Orelle B, Keim V, Dagorn JC. Messenger RNA sequence and expression of rat pancreatitis-associated protein, a lectin-related protein overexpressed during acute experimental pancreatitis. *J Biol Chem*. 1991 Dec 25;266(36):24664–9.

A Preliminary Study Troponin T and Reg3 β in Children with Left-to-Right Shunt Congenital Heart Disease with Heart Failure

ORIGINALITY REPORT

17%

SIMILARITY INDEX

8%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Ana L. Neves, Tiago Henriques-Coelho, Adelino Leite-Moreira, José C. Areias. "Cardiac injury biomarkers in paediatric age: Are we there yet?", Heart Failure Reviews, 2016
Publication 3%
- 2 link.springer.com
Internet Source 1%
- 3 www.coursehero.com
Internet Source 1%
- 4 www.termedia.pl
Internet Source 1%
- 5 Kazumori, H.. "Localization of Reg receptor in rat fundic mucosa", The Journal of Laboratory and Clinical Medicine, 200202
Publication 1%
- 6 Laura Trementino, Marina Cardinaletti, Carolina Concettoni, Giorgia Marcelli et al. "Salivary cortisol is a useful tool to assess the

early response to pasireotide in patients with Cushing's disease", Pituitary, 2014

Publication

7	edoc.ub.uni-muenchen.de Internet Source	1 %
8	openaccess.hacettepe.edu.tr:8080 Internet Source	1 %
9	Terry Brissac, Eriel Martinez, Katherine L Kruckow, Ashleigh N Riegler et al. "Capsule protects against intracellular killing and enables vascular endothelial cell translocation during invasive pneumococcal disease", Cold Spring Harbor Laboratory, 2021 Publication	1 %
10	journals.plos.org Internet Source	1 %
11	rgd.mcw.edu Internet Source	1 %
12	www.firstaidteam.com Internet Source	1 %
13	www.pubmedcentral.nih.gov Internet Source	1 %
14	Alyaa A. Kotby, Manal M. Abd Al Aziz, Adel H. Husseiny, Marwa M. Al-Fahham. "Detection of Early Myocardial Injury in Children with Ventricular Septal Defect Using Cardiac	<1 %

Troponin I and Two-Dimensional Speckle Tracking Echocardiography", Pediatric Cardiology, 2020

Publication

15

biblio.ugent.be

Internet Source

<1 %

16

jaha.ahajournals.org

Internet Source

<1 %

17

Marc Thiriet. "Vasculopathies", Springer Science and Business Media LLC, 2018

Publication

<1 %

18

Michael E. Zenilman. "Comparison of reg I and reg III Levels During Acute Pancreatitis in the Rat", Annals of Surgery, 11/2000

Publication

<1 %

19

Michiaki Abe, Koji Nata, Takako Akiyama, Nausheen J. Shervani et al. "Identification of a novel Reg family gene, Reg III δ , and mapping of all three types of Reg family gene in a 75 kilobase mouse genomic region", Gene, 2000

Publication

<1 %

20

Navin Suthahar, Laura M.G. Meems, Jennifer E. Ho, Rudolf A. de Boer. "Sex-related differences in contemporary biomarkers for heart failure: a review", European Journal of Heart Failure, 2020

Publication

<1 %

21 Steven E Lipshultz. "Cardiovascular risk factors, monitoring, and therapy for HIV-infected patients :", AIDS, 04/2003
Publication <1 %

22 docplayer.net
Internet Source <1 %

23 www.nejm.org
Internet Source <1 %

24 Kyung Chan Park, David C Gaze, Paul O Collinson, Michael S Marber. "Cardiac troponins: from myocardial infarction to chronic disease", Cardiovascular Research, 2017
Publication <1 %

25 M. Gironella. "Experimental acute pancreatitis in PAP/HIP knock-out mice", Gut, 8/1/2007
Publication <1 %

26 Mohamed O. Hafez, Saed M. Morsy, Ragab A. Mahfoz, Ahmed R. Ali. "Myocardial Injury in Children with Unoperated Congenital Heart Diseases", Cardiology Research and Practice, 2015
Publication <1 %

27 Pathophysiology and Pharmacotherapy of Cardiovascular Disease, 2015.
Publication <1 %

28

Internet Source

<1 %

29

"Etiology and Morphogenesis of Congenital Heart Disease", Springer Science and Business Media LLC, 2016

Publication

<1 %

30

Purnami Nyilo, Manyakori Serafika Permoni Putri. "The Association of Reactive Oxygen Species Levels on Noise Induced Hearing Loss of High Risk Workers in Dr. Soetomo General Hospital Surabaya, Indonesia", Indian Journal of Otolaryngology and Head & Neck Surgery, 2018

Publication

<1 %

31

www.dovepress.com

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

A Preliminary Study Troponin T and Reg3 β in Children with Left-to-Right Shunt Congenital Heart Disease with Heart Failure

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
