



RISK FACTOR OF BACTEREMIA IN CHILDREN WITH PNEUMONIA



Retno Asih, Zuhrotul Aini, Landia Setiawati

Division of Respiriology Department of Child Health, Medical Faculty
Airlangga University/Dr. Soetomo Hospital – Surabaya

BACKGROUND

Pneumonia is known as a frequent cause of morbidity and mortality among children in developing countries. In children, it caused predominantly by bacteria. Bacteremia has been associated with severity and mortality of pneumonia. Identifying factors caused by bacteremia is important to prevent severity and mortality of pneumonia.

OBJECTIVE

The objective of this study was to identify risk factors of bacteremia in children with pneumonia.

METHODS

- Method: Retrospective study
- Sample: Children with pneumonia
- Setting: Dr. Soetomo Surabaya Hospital
- Time: January 2007 – December 2008
- Tool: Chi-Square test, Fisher's exact test

The diagnosis of pneumonia in this study was made based on clinical symptoms and abnormality on chest radiography presenting of lower respiratory tract infection. Sex age, clinical characteristics, laboratory and radiology findings were collected by medical records review.

RESULTS

Table 1. Risk Factors Bacteremia of children with pneumonia at dr. Soetomo Hospital January 2007 – December 2008

Interval from onset of symptoms to hospital admission, n (%)				
< 3 days	36 (8.2)	199 (43.4)	1.35 (0.69-2.67)	0.386
> 3 = 5	31 (7.1)	168 (38.3)	0.97 (0.48-1.97)	0.933
= 5	8 (1.8)	5 (1.1)	22.69 (6.96-73.9)	0.000*
Nutritional state				
Well nourished	26 (5.9)	255 (58.2)	1.52 (0.71-3.23)	0.279
Moderate malnourished	8 (1.8)	142 (32.4)	0.52 (0.23-1.18)	0.113
Severely malnourished	3 (0.7)	4 (0.9)	9.05 (1.94-42.13)	0.001*
Temperature				
< 36.5°C or > 38.5°C	8 (1.8)	116 (26.5)	0.70 (0.31-1.59)	0.397
Laboratory findings				
Hb < 10g/dl	21 (4.8)	230 (52.5)	2.44 (1.04-5.71)	0.034*
Leukocyte < 5.000/cmm or > 20.000/cmm	16 (3.7)	100 (22.8)	2.41 (1.21-4.84)	0.011*
CRP positive	25 (5.7)	304 (69.4)	0.67 (0.26-1.72)	0.404
Blood Gas Analysis				
pH < 7.35	6 (1.4)	69 (15.8)	0.85 (0.29-2.44)	0.766
pO ₂ < 80mmHg	3 (0.7)	100 (22.8)	4.25 (1.18-15.4)	0.018*
pCO ₂ > 50mmHg	5 (1.1)	27 (6.2)	0.48 (0.15-1.60)	0.236
SpO ₂ < 95%	7 (1.6)	89 (20.3)	1.19 (0.27-5.31)	0.858
Thorax X-ray				
Lobar consolidation	7 (1.6)	14 (3.2)	1.19 (0.27-5.31)	0.833

CONCLUSION

Risk factors bacteremia in children with pneumonia included age under 1 year, symptoms more than 5 days, severe malnourished, anemia, leucocyte count less than 5.000/mm³ and more than 20.000/mm³ and pO₂ less than 80mmHg. Bacteremia has been associated with severity and mortality of pneumonia. Identifying factors caused bacteremia is important to prevent severity and mortality of pneumonia.

Keywords: risk factors, bacteremia, pneumonia

Risk Factor of Bacteremia in Children with Pneumonia

Retno Asih, Zuhrotul aini, Landia Setiawati
Division of Respirology Department of Child Health ,Medical Faculty
Airlangga University/Dr. Soetomo Hospital - Surabaya

ABSTRACT

Background :

pneumonia is known as a frequent cause of morbidity and mortality among children in developing countries. In children, it caused predominantly by bacteria. Bacteremia has been associated with severity and mortalitas of pneumonia. Identify factors caused bacteremia important to prevent severity and mortalitas of pneumonia.

Objective : The objective of this study was to identify risk factors of bacteremia in children with pneumonia.

Methods : A retrospective study was conducted in children with pneumonia in Dr. Soetomo Surabaya Hospital from January 2007 to December 2008. Blood cultures be performed on all of this patients. Factors associated with bacteremia were identified following review of medical records include clinical features, laboratory , radiology and blood culture results.

Results : Frequency of bacteremia was 8,2 % (36 patients) of 438 children with pneumonia. Interval from onset of symptoms to hospital admission more than 5 days (22.69 CI 95 %), severe malnourished (OR 9.05 CI 95 %), anemia (OR 2.44 CI 95 %), leucocyt counts less than 5000/mm³ and more than 20.000/mm³ (OR 2.41 CI 95 %) and pO₂ less than 80mmHg (OR 4.25 CI 95 %) were at increased risk of bacteremia in children with pneumonia.

Conclusion : Risk factors bacteremia in children with pneumonia included age under 1 year, symptoms more than 5 days ,severe malnourished ,anemia, leucocyt counts less than 5000/mm³ and more than 20.000/mm³ and pO₂ less than 80mmHg

Keyword : Risk factors, bacteremia, pneumonia

INTRODUCTION

Pneumonia is an acute inflammatory disease of the lung parenchyma involving the distal terminal bronchioles, respiratory bronchioles, alveoli ducts, alveoli sacs, and alveoli .. This disease is one of the most common infections in the pediatric age group. In developing countries, researchers estimate that more than 150 million new cases occur annually in children < 5 years. It is well known as a frequent cause of morbidity and a leading cause of mortality among children.¹ One in five of childhood deaths in developing countries have been ascribed to acute respiratory tract infections (ARI) and 90% of these deaths are due to pneumonia.²

Pneumonia in developing countries is caused predominantly by bacteria.¹ The frequency of bacteremia in pneumonia patients varies from 4% to 18%.³ Bacteremic pneumonia are potentially life-threatening in children.⁴

Predictor of bacteremia in pneumonia included recent antibiotic treatment, comorbidity disease, increase respiratory rate, temperature, pulse and laboratory abnormalities.³

The objective of this study was to identify risk factors of bacteremia in children with pneumonia. A retrospective study was conducted in children with pneumonia in Dr. Soetomo Surabaya Hospital from January to December 2009. Blood cultures be performed on all of this patients. Factors associated with bacteremia were identified following review of medical records include clinical features, laboratory , radiology and blood culture results. Identify factors caused bacteremia important to prevent severity and mortalitas of pneumonia

SUBJECTS AND METHODS

. A retrospective study was conducted in children age 1 months until 5 years with diagnosis community acquired pneumonia Dr. Soetomo Surabaya Hospital from January 2007 to December 2008. The diagnosis of pneumonia in this study was made based on clinical symptoms and abnormality on chest radiography presenting of lower respiratory tract infection. Sex, age, clinical characteristics, laboratory and radiology findings were collected by medical records review.

Patients were defined as bacteremic if a blood culture drawn of presentation to the hospital before antibiotic treatment grew an organism, and not defined as a contaminant.

From the time of admission, we noted duration of symptoms, clinical manifestation, nutritional state, laboratory and radiological findings. We classified temperature into 2 categories, less than 36,5°C, or more than 38,5°C and between 36,5 C and 38,5°C. Nutritional state were classified into 3 categories, well nourished if ideal body weight more than 90 %, moderate malnutrition if ideal body weight between 70 %-90 % and severe malnutrition if ideal body weight less than 70 %.

We collected laboratory finding includes hemoglobin level, leucocyt level, C Reactive Protein and blood gas analyse. Anemia if hemoglobin level less than 10 gr/dl. Leucocyt counts classified into less than 5000/cmm, between 5000/cmm and 20.000/cmm and more than 20.000/cmm.

X² tests, Fisher's exact test and odds ratio and relative risks with 95% confidence intervals were used to determine whether an association was significant (p<0.05).

RESULTS

We identified 440 patients with pneumonia who met the inclusion criteria for the study. Blood culture was performed on 438 patients, bacteremia was detected in 36 patients (8.2%) and 73 patients (16.7%) had a contaminated bacteria (Table 1)

Table 1. Characteristic of children with Pneumonia at Dr. Soetomo Hospital, January 2007 -December 2008.

Age (months) mean ± SD (range)	12.2 ± 11.2
1-12 month, n (%)	293 (66.9)
13-36 month	126 (28.8)
> 36 month	19 (4.3)
Sex ratio (male : female)	1.6 : 1
Interval from onset of symptoms to hospital admission, n (%)	
< 3 days	226 (51.6)
>3-<5	199 (45.4)
≥5	13(3.0)
Nutritional state, n (%)	
Well nourished	281(64.2)

Moderate malnourished	150 (34.2)
Severely malnourished	7 (1.6)
Temperature, n(%)	
< 36.5 C or > 38.5 C	124 (28.3)
36.5 C - 38.5 C	314 (71.7)
Laboratory findings	
Hb < 10 g/dl	251 (57.3)
Leucocyte < 5,000/cmm or > 20,000/cmm	116 (26.5)
CRP positive	329 (75.1)
Blood Gas Analysis	
pH < 7.35	75 (17.1)
pO ₂ < 80 mmHg	103 (23.5)
pCO ₂ > 50mmHg	32 (7.3)
SpO ₂ < 95 %	96 (21.9)
Thorax X-ray, n(%)	
Normal	60 (13.7)
Lobar consolidation	21 (4.8)
Patchy infiltrate	354 (80.8)
Pleural effusion	3 (0.7)
Blood culture, n (%)	
Sterile	329 (75.1)
Bacteremia	36 (8.2)
Contaminant	73 (16.7)

Of the total cases, ratio male and women was 1.6 :1. The mean age was 12.2 months with the majority at the age of 1-12 months as many as 293 (66.9%) children. The majority interval from initial symptoms until it is brought to the hospital was less than 3 days (51.6%) children. Most nutritional status is well nourished as much as 281 (64.2%) children.

Anemia was found in 251 (57.3%) children, levels of leukocytes less than 5000/mm³ or more than 20.000/mm³ many as 116 (26.5%) children, positive CRP was found in 329 (75.1%) of children and acidosis obtained in 75 (17.1%) children. Preview photos thoracic infiltrates was found in 354 (80.8%) children.

Table 2. Based line characteristic of children with Pneumonia at dr. Soetomo Hospital January 2007 – December 2008

Variable	Bacteremia n (%)	No Bacteremia n (%)	p value
Sex			
Male	25 (5.7)	248 (56.6)	0.169
Female	11 (2.5)	154 (35.2)	
Age group (years)			
1-12 month	27 (6.2)	266 (60.7)	0.307
13-36 month	9 (2.1)	117 (26.7)	0.431
> 36 month	2 (0.5)	17 (38.8)	0.548

There were no significant difference of based line characteristic children with bacteremia positive and bacteremia negative include sex and age.(Table 2)

Table 3. Risk Factors Bacteremia of children with Pneumonia at dr. Soetomo Hospital January 2007 – December 2008

Variable	Bacteremia positive n (%)	Bacteremia negative n (%)	OR (95% CI)	p value
Interval from onset of symptoms to hospital admission, n (%)				
< 3 days	36 (8.2)	199 (43.4)	1.35(0.69-2.67)	0.386
>3-<5	31 (7.1)	168 (38.3)	0.97(0.48-1.97)	0.933
≥5	8 (1.8)	5 (1.1)	22.69(6.96-73.9)	0.000*
Nutritional state, n (%)				
Well nourished	26 (5.9)	255 (58.2)	1.52(0.71-3.23)	0.279
Moderate malnourished	8 (1.8)	142 (32.4)	0.52(0.23-1.18)	0.113
Severely malnourished	3 (0.7)	4 (0.9)	9.05(1.94-42.13)	0.001*
Temperature < 36.5 C or > 38.5 C	8 (1.8)	116 (26.5)	0.70(0.31-1.59)	0.397
Laboratory findings Hb < 10 g/dl	21(4.8)	230(52.5)	2.44(1.04-5.71)	0.034*

Leucocyte < 5,000/cmm or > 20,000/cmm	16(3.7)	100(22.8)	2.41(1.21-4.84)	0.011*
CRP positive	25(5.7)	304(69.4)	0.67(0.26-1.72)	0.404
Blood Gas Analysis				
pH < 7.35	6(1.4)	69(15.8)	0.85(0.29-2.44)	0.766
pO ₂ < 80mm Hg	3(0.7)	100(22.8)	4.25(1.18-15.4)	0.018*
pCO ₂ > 50mmHg	5(1.1)	27(6.2)	0.48(0.15-1.60)	0.236
SpO ₂ < 95 %	7(1.6)	89(20.3)	1.19(0.27-5.31)	0.858
Thorax X-ray				
Lobar consolidation	7(1.6)	14(3.2)	1.19(0.27-5.310)	0.833

Table 3 shows risk factors several variables on the occurrence of bacteremia in patients with pneumonia of children. From the table, the risk factors of pneumonia were the interval from onset of symptoms to hospital admission more than 5 days, severely malnutrition, anemia, leukocyte less than 5000/mm³ or more of 20.000/mm³ and pO₂ less than 80 %.

DISCUSSION

The aetiology of pneumonia in developed countries is predominantly viral, associated with a low case fatality rate, whereas in developing countries bacteraemia is common and associated with a high case fatality rate. Poor sanitation, overcrowding, inadequate nutrition, insufficient vaccination coverage, low levels of education, and accumulation of other diseases have been suggested as reasons for the differences in aetiology and mortality.⁵

To reduce mortality from pneumonia in developing countries the problem has to be addressed from a number of aspects, social and environmental as well as medical.⁵

In this comparative study of bacteremic and non bacteremic patients, we found that bacteremia was detected in 8.2 % children. The previous study reported frequency of bacteremia in patients pneumonia varies from as low as 4% to as high as 14 to 18%.

We found that bacteremic and non-bacteremic pneumonia patients not differed in baseline characteristics. The study by Spooner et al from Papua New Guinea(PNG) identified firstborn children and female children to have increased bacteremia and mortality risk in children with pneumonia.⁵ Meanwhile Jover et al, in research on adult patients with pneumonia, bacteremia get no difference between men and women.⁶

We have identified six independent factors of bacteremia in children with pneumonia. Five of these significantly associated with bacteremia. Include interval from onset of symptoms to hospital admission has significantly associated with bacteremia, the interval from onset of symptoms to hospital admission more than 5 days, severely malnutrition, anemia, leukocyte less than 5000/mm³ or more of 20.000/mm³ and pO₂ less than 80 %. Spooner, et al identified that poor feeding, cyanosis, bronchial breathing, and a temperature > 38C were all associated with bacteraemia.⁵

Interval from onset of symptoms to hospital admission more than 5 days has significantly associated with bacteremia. Previous study reported that history of fever for more than 7 days significantly increased the chance of dying.⁵ The onset to hospital admission associated with prior use of antibiotic. Metersky et al, reported that the risk of bacteremia could be predicted by assessing the prior use of antibiotics.³

Malnourished children are particularly at risk as demonstrated in this study. A study in PNG reported that malnourished patients had a significantly higher risk of dying.⁵

In our study, bacteremic patients were more likely to be anemic than non-bacteremic patients and to have abnormal leucocyt counts and pO₂ less than 80 %. Brandeburg found that bacteremic patients were more likely to have anemia, lower albumin, and elevated blood urea and serum creatinine levels.⁷ While Metersky reported that blood urea nitrogen more than 30 mg/dl (11 mmol/L), sodium less than 130 mmol/L and WBC less than 5,000/mm³ or more than 20,000/mm³ were independent predictors of bacteremia in community-acquired patients with pneumonia.³

Results of radiological findings in association with bacteraemia have been analysed. They showed that a peripheral homogeneous opacity was the best predictor of bacteraemia.⁶ Jover reported that although not statistically significant, pleural effusion

was more frequent in bacteremic patients.⁶ In this study, the most radiological finding was infiltrate.

There are some limitations in this study. The main weakness of our study is its retrospective design. Collection of some data was therefore incomplete. Another limitation could be the lower number of non-bacteremic cases compared to bacteremic cases.

CONCLUSION

Risk factors bacteremia in children with pneumonia included age under 1 year, symptoms more than 5 days, severe malnourished, anemia, leucocyt counts less than 5000/mm³ and more than 20.000/mm³ and PO₂ less than 80 mmHg.

Bacteremia has been associated with severity and mortalitas of pneumonia. Identify factors caused bacteremia important to prevent severity and mortalitas of pneumonia.

REFERENCES

1. Masria S. Pattern of Bacteria Causing Pneumonia in Children and Its Sensitivity to Some Antibiotics. Proc ASEAN Congr Trop Med Parasitol 2008; 3:121-4
2. Narayanan M, Falade AG. Clinical Risk Factors for Death in Children with Pneumonia. International Child Health Review Collaboration. 13th October 2006
3. Metersky ML, Ma A, Bratzler DW, Houck PM. Predicting Bacteremia in Patients with Community-Acquired Pneumonia. Am J Respir Crit Care Med 2004; 169: 342-7
4. Toikka P, Virkki R, Mertsola J, Ashorn P, Eskola J, Ruuskanen O. Bacteremic Pneumococcal pneumonia in children. Clin Infect Dis 1999; 29:568-72
5. Spooner V, Barker J, Tulloch S, Lehmann D, Marshall TF, Kajo M, Alpers MP. Clinical signs and risk factors associated with pneumonia in children admitted to Goroka Hospital, Papua New Guinea. J Trop Pediatr. 1989;35:295-300

6. Jover F, Cuadrado JM, Andreu L, Martínez s, Canizares R, Tabla Vo et al. A comparative study of bacteremic and non-bacteremic pneumococcal pneumonia. *Eur J of Intern Med* 2008; 19:15-21
7. Brandenburg JA, Marrie TJ, Coley CM, Singer DE, Obrosky S, Kapoor WN, et al. Clinical presentation, processes and outcomes of care for patients with pneumococcal pneumonia. *J Gen Intern Med* 2000;15:638-47.