


OVOZOA

Departemen Reproduksi Veteriner
Fakultas Kedokteran Hewan Unair





























VOL 8, NO 2 (2019)

OVOZOA : JOURNAL OF ANIMAL REPRODUCTION

TABLE OF CONTENTS

ORIGINAL ARTICLE

- | | | |
|---|--|---|
|  | HUBUNGAN ANTARA BODY CONDITION SCORE (BCS) DENGAN PRODUKSI SUSU SAPI PERAH FRIESIAN HOLSTEIN (FH) |  |
| | | 89-93 |
|  | 10.20473/ovz.v8i2.2019.89-93 | |
|  | Melsa Netika, Roesno Darsono, Budi Utomo, Imam Mustofa, Ismudiono Ismudiono, Tri Wahyu Suprayogi | |
|  | HUBUNGAN ANTARA UMUR DENGAN CALVING INTERVAL, DAYS OPEN, DAN SERVICE PER CONCEPTION SAPI FRIESIAN HOLSTEIN (FH) |  |
| | | 94-99 |
|  | 10.20473/ovz.v8i2.2019.94-99 | |
|  | Hanifah Muslimah Ananda, Wurlina Wurlina, Nove Hidajati, Mas'ud Hariadi, Abdul Samik, Tjuk Imam Restiadi | |
|  | PENGARUH EKSTRAK DELIMA MERAH (Punicagranatum L.) TERHADAP PENINGKATAN PERSENTASE KAPASITASI DAN REAKSI AKROSOM SPERMATOZOA TIKUS PUTIH (Rattus norvegicus) |  |
| | | 100-104 |
|  | 10.20473/ovz.v8i2.2019.100-104 | |
|  | Baktiar Budi Satrio, Mas'ud Hariadi, Retno Sri Wahjuni, Budi Utomo, Sri Agus Sudjarwo, Tita Damayanti L | |
|  | KORELASI ANTARA LINGKAR DADA, PANJANG BADAN DAN TINGGI GUMBA TERHADAP VOLUME SEMEN DAN KONSENTRASI SPERMATOZOA SAPI MADURA |  |
| | | 105-109 |
|  | 10.20473/ovz.v8i2.2019.105-109 | |
|  | Dilla Aranda Mayola, Rimayanti Rimayanti, Sunaryo Hadi Warsito, Suzanita Utama, Hana Eliyani, Erma Safitri | |
|  | PERANAN VITAMIN B12 DALAM PENGECER SUSU SKIM KUNING TELUR TERHADAP MOTILITAS DAN VIABILITAS SPERMATOZOA DOMBA SAPUDIPOST THAWING |  |
| | | 110-115 |
|  | 10.20473/ovz.v8i2.2019.110-115 | |
|  | Mirza Asafaizh Arini, Tri Wahyu Suprayogi, Sunaryo Hadi Warsito, Trilas Sardjito, Suherni Susilowati, Indah Norma Triana | |
|  | PENGARUH PENAMBAHAN EKSTRAK TEH HIJAU (Camellia sinensis) DALAM BAHAN PENGECER SUSU SKIM TERHADAP KUALITAS SPERMATOZOA DOMBA SAPUDI YANG DISIMPAN PADA SUHU DINGIN |  |
| | | 116-121 |
|  | 10.20473/ovz.v8i2.2019.116-121 | |
|  | Maulinda Agchirilia Vladika, Sunaryo Hadi Warsito, Suherni Susilowati, Tatik Hernawati, Rochmah Kurnijasanti, Tjuk Imam Restiadi | |
|  | PENGARUH PENAMBAHAN EKSTRAK TEH HIJAU (Camellia sinensis) DALAM BAHAN PENGECER SUSU SKIM KUNING TELUR TERHADAP KUALITAS SPERMATOZOA DOMBA SAPUDI YANG DISIMPAN PADA SUHU DINGIN |  |
| | | 122-126 |

People

Contact

Editorial Team

Reviewers

Instruction for Author

Guide for authors

Online Submission

Download Documents

Journal Policy

Focus and Scope

Section Policies

Publication Frequency

Publication Ethics

Indexing

Article Processing Charge

Peer Reviewers Process

Open Access Statement

Archiving

Plagiarism Screening

ORCID Id Policy

History

Copyright

old website

Tools




grammarly



OPEN JOURNAL SYSTEMS

doi 10.20473/ovz.v8i2.2019.122-126


 Wahyu Retno Swari, Emy Koestanti Sabdoningrum, Wurlina Wurlina, Suherni Susilowati, Rochmah Kurnijasanti, Erma Safitri


 **PENGARUH PENAMBAHAN EKSTRAK TEH HIJAU (*Camellia sinensis*) DALAM BAHAN PENGECER KUNING TELUR SITRAT TERHADAP KUALITAS SPERMATOZOA DOMBA SAPUDI YANG DISIMPAN PADA SUHU DINGIN**



127-131

doi 10.20473/ovz.v8i2.2019.127-131

 Itrah Khoirunnisa, Suherni Susilowati, Lilik Maslachah, Tri Wahyu Suprayogi, Rochmah Kurnijasanti, Hermin Ratnani

 **PENGARUH PENGECER KUNING TELUR AYAM DENGAN AIR KELAPA MUDA TERHADAP INTEGRITAS MEMBRAN PLASMA DAN ABNORMALITAS SPERMATOZOA DOMBA SAPUDI**
PENGARUH PENGECER KUNING TELUR AYAM DENGAN AIR KELAPA MUDA TERHADAP INTEGRITAS MEMBRAN PLASMA DAN ABNORMALITAS SPERMATOZOA DOMBA SAPUDI



139-143

doi 10.20473/ovz.v8i2.2019.139-143

 Mayrena Ayu Cesaria, A.T.Soelih Estoepongastie, Suherni Susilowati, Tatik Hernawati, Sri Pantja Madyawati, Indah Norma Triana


CASE REPORT

 **DAMPAK PEMBERIAN ROTI AFKIR TERHADAP EFISIENSI REPRODUKSI SAPI PERAH Friesian Holstein DI WILAYAH KERJA KPSP SETIA KAWAN NONGKOJAJAR, PASURUAN**



144-148

doi 10.20473/ovz.v8i2.2019.144-148


 Ricko Nurdianto S., Sri Hidanah, Sri Pantja Madyawati, Abdul Samik, wurlina wurlina, Tri Wahyu Suprayogi

 **PERBANDINGAN BODY CONDITION SCORE, SERVICES PER CONCEPTION, CALVING INTERVAL, DAN DAYS OPEN SAPI FRIESIAN HOLSTEIN (FH) DI KEMITRAAN GREENFIELD DENGAN KUD TANI WILIS SENDANG**



149-153

doi 10.20473/ovz.v8i2.2019.149-153

 Aghid Cahya P., Sri Mulyati, Suwarno Suwarno, Imam Mustofa, Pudji Srianto, Budi Utomo

 **PENGARUH pH LENDIR MUKOSA VAGINA SAAT BIRAH TERHADAP PERSENTASE KEBUNTINGAN (Conception Rate) PADA SAPI PERAH DI KUD TANI WILIS KABUPATEN TULUNGAGUNG DAN KSU TUNAS SETIA BARU KABUPATEN PASURUAN**



154-158

doi 10.20473/ovz.v8i2.2019.154-158


 Andaru Rizki, Pudji Srianto, Endang Suprihati, Trilas Sardjito, Ismudiono Ismudiono, Mohammad Anam Al Arif

 **CONCEPTION RATE DAN SERVICE PER CONCEPTION PADA SAPI BALI HASIL INSEMINASI BUATAN DI KABUPATEN SUMBAWA BARAT TAHUN 2017**



159-163

doi 10.20473/ovz.v8i2.2019.159-163


 Asih Deskayanti, Trilas Sardjito, Agus Sunarso, Pudji Srianto, Tri Wahyu Suprayogi, Herry Agoes Hermadi

 **PERSENTASE KEJADIAN KAWIN BERULANG SAPI PERAH YANG TELAH DI IB (INSEMINASI BUATAN) TINGKAT PETERNAK DESA NYAWANG KIDUL DI KUD TANI WILIS KECAMATAN SENDANG KABUPATEN TULUNGAGUNG**



164-168

doi 10.20473/ovz.v8i2.2019.164-168

 Arda Wahyu Setyadi, Pudji Srianto, Oky Setyo Widodo, Wurlina Wurlina, Abdul Samik, Tjuk Imam Restiadi

 **HUBUNGAN MORFOMETRI AMBING TERHADAP PRODUKSI SUSU KUDA DI DAERAH BIMA NUSA TENGGARA BARAT**



169-174

doi 10.20473/ovz.v8i2.2019.169-174

 Mujahid Mujahid, Imam Mustofa, Benjamin Chr Tehupuring, Tjuk Imam Restiadi, Hana Eliyani, Hermin Ratnani

Journal Help

USER

Username

Password

Remember me

Login

NOTIFICATIONS

[View](#)

[Subscribe](#)

[View My Stats](#)



Ovozoa: Journal of Animal Reproduction by Unair is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#).



This journal is supported by



Pusat Pengembangan Jurnal dan Publikasi Ilmiah
Universitas Airlangga
copyright@2017 Template PPJPI

PENGARUH EKSTRAK DELIMA MERAH(*Punicagranatum L.*) TERHADAP PENINGKATAN PERSENTASE KAPASITASI DAN REAKSI AKROSOM SPERMATOZOA TIKUS PUTIH (*Rattusnorvegicus*)

INFLUENCE OF POMEGRANATE EXTRACT (*Punicagranatum L.*) ON IMPROVEMENT OF PERCENTAGE OF SPERMATOZOA CAPACITATION AND ACROSOME REACTION OF WHITE RAT (*Rattusnorvegicus*)

Baktiar Budi Satrio¹⁾, *Mas'ud Hariadi²⁾, Retno Sri Wahjuni³⁾, Budi Utomo⁴⁾ Sri Agus Sudjarwo⁵⁾, Tita Damayanti L.⁶⁾

¹⁾Student, ^{2,4,6)}Department of Veterinary Reproduction, ^{3,5)}Basic Veterinary Medicine Faculty of Veterinary Medicine, Universitas Airlangga

*Corresponding author: email:masudhariadi@yahoo.co.id; baktiarsatrio@gmail.com

ABSTRACT

The aim of this research is to know the effect of pomegranate (*Punicagranatum L.*) on the improvement of spermatozoa quality that is increase of percentage of capacitation and acrosome reaction in white rat (*Rattusnorvegicus*). This research used 20 healthy white rat (*Rattusnorvegicus*) with average age 2-3 months old, 200-300 g. Research design using completely randomized design (CRD) with four treatments and 5 replications. Rats were randomly divided into 4 treatment groups. Rat in the negative control group not treated just kept alone, solvent control group were given CMC Na 0,5% and is exposed to heat, the treatment groups 1, 2, and 3 induced pomegranate extract standardized to 40% ellagic acid at a dose of 75mg / kg, 150mg / kg and 300mg /kg perorally with doses 1 ml/rat/day gastric for along 14 days and exposed to heat. Data were analyzed using One Way Analysis of Variant (ANOVA) followed by Duncan ($p < 0,05$). The result showed that there was significant effect from therapy on the quality of rats spermatozoa especially to capacitation and acrosome reaction. The treatment from treatment 2 group showed the best result in maintaining the quality of rats spermatozoa with 27.6 ± 3.04 values for capacitation and $6,80 \pm 1,30$ for acrosome reactions. Extract of pomegranate (*Punicagranatum L.*) standardized to 40% ellagic acid has a significant change in improve the quality of spermatozoa, namely an increase in the percentage of capacitation and acrosome reactions in white rats (*Rattusnorvegicus*) exposed to heat.

Keywords : Pomegranate extract, Capacitation, Acrosome Reaction

Research Background

The quality of reproduction is in-fluenced by the ability of spermatozoa to fertilize. Fertilization plays a role in the process of capacitation, where spermatozoa cells can penetrate the ovum has an im-portant role in improving reproductive qua-lity (Lestari, 2014). Fertilization involves a process of capacitation and acrosome reac-tions. The process of capacitation is a series of changes that occur to prepare the sperm to meet and interact with the ovum at the ti-me of fertilization. The process of capaci-tation includes the ability to spermatozoa for motile or movement, the ability to fer-tilize and the removal of cytoplasmic dro-plet. Spermatozoa cannot afford to fertilize without a capacitation process (Bearden and Fuquay, 2000).

Acrosome reaction is the reaction of re-lease of enzymes from the acrosome to pe-netrate the oocyte layers by induction by zone proteins. Acoromic reactions occur only in spermatozoa that have intact mem-branes. In addition, the acrosome reaction takes place before the spermatozoa pene-trate into the zona pelucida / ZP (Grudzinskas and Yovich, 1995 in Anwar 2014).

Spermatozoa require reactive oxygen species (ROS) at low concentrations to in-duce the process of capacitation and acro-some reactions (Sikka, 2004) and bind to the zona pellucida (Sanocka and Kupisz, 2004) so that fertilization proceeds well. Physiologically free radicals are present in sperm (Zavos *et al.* 1998) and the onset of free radicals in the body is offset by en-

ogenous defense mechanisms, by producing substances that have an anti-free radical effect called antioxidants. However, as the ROS level rises above the body's antioxidant defense system, oxidative stress occurs (Saleh *et al.*, 2003). To reduce ROS needs to be provided additional antioxidants (Makker *et al.*, 2009).

Pomegranate (*Punica granatum L*) are one source of antioxidants from plants with high polyphenol content (Aviram *et al.*, 2014). Anthocyanin is one of the pomegranate compound that have a powerful antioxidants that can prevent various damages caused by oxidative stress so as to protect cells from free radicals (Yanjun *et al.*, 2009).

Research on the effect of antioxidants in pomegranate to the process of sperma-tozoa capacitation has never been done. Therefore this research was conducted find out the effect of giving pomegranate to spermatozoa by using experimental animal White rat (*Rattus norvegicus*)

Materials and Methods

This research was carried out for 4 weeks in May to June 2017. The research was conducted at three locations. Laboratorium Farmakologi Kedokteran Hewan, Departemen Biokimia Fakultas Kedokteran, and Institute Tropical Disease (ITD). This research used 20 healthy white rat (*Rattus norvegicus*) with average age 2-3 months old, 200-300 g body weight (BW) and bought from Pusat Veterinaria Farma (PUSVETMA) Surabaya.

Research material used in this study is white rat (*Rattus norvegicus*), white mouse semen, pomegranate (*Punica granatum L*) extract powder that contains 40% ellagic acid produced by Xi'an Biof Bio-Technology Co., Ltd. (Room 1-1111, High-tech Venture Park, No. 69 Jinye Road, Gaoxin Distric of Xian, People Republic of China), *FlourescentIsotiocianant* (FITC) Staining, PBS Aquabides, NaCl physiological, For-maldehyde, Ethanol 70 % and Glycerol.

Research tools used in this study is a scaled-semen container tube, epifluorecent microscope, object glass, glass cover, test tube, tube rack, petri dish, bath, 1 ml and 3 ml syringe pipette, thermometer, aluminum foil, tissue paper and camera digital.

White Rat aged 2-3 months weighing 200-300 grams were adapted for 7 days. Experimental animals consisted of 1 control group, and 3 treatment groups. Each group consisted of 5 white rats and randomly selected. Before giving pomegranate extract, experimental animals exposed to the sun at 09.45 WIB for 15 minutes to get free radical directly. The control group (K) was given only CMC Na 5% oral solvent, T1 group, T2, T3 white rat were treated with excretion of pomegranate orally with multilevel dose, i.e. dose 75 mg / kgBW, 150 mg / kgBW, and 300mg / kgBW once daily For 14 days.

At the end of the treatment all of the rat were terminated with ether, then the abdominal surgery was performed and a collection of testicular organ was performed (Aulanni'am *et al.*, 2007). Spermatozoa taken from the epididymis, exactly 1 cm below chorda spermatica. In the place is clamped and cut. The cut part is removed by sperm by squeezing on the epididymis then given physiological NaCl and stirred to become homogeneous.

Testing of capacitation status was performed by *FlourescentIsotiocianant* (FITC) Staining. Briefly the FITC preparation process is as follows: (1) 100 µl of treated semen is fed into aluminum foil enclosed eppendorf tube and added with 100 µl dye FITC, vortex for 1 min, and add 8 µl FITC fixative vortex solution for 1 min, (2) Taken 10 µl (mixture 1 and 2) and placed on top of the glass object, added with 10 µl of DABCO solution then mixed with a micropipette tip, then covered with a glass cover and carefully pressed with thickly coated tissue palms. The side of the glass cover is adhesive with cutex.

(3) The observations were performed with an epifluorescence microscope using an ultra violet light source. FITC staining on spermatozoa exhibits 2 different forms of fluorescent flux, ie (1) The same fluorescent distribution in the spermatozoa (non-capacitated) head, and (2) Fluorescent is concentrated in the acrosomal region indicating the spermatozoa is capacitated. Observation of capacitation status is done after the next 24 hours (Utomo, 2011).

Examination of acrosome reaction were done by fixated semen with 4% formaldehyde, then washed by adding 3 ml PBS and

centrifuged 1500 rpm for 10 min, supernatant was removed and added with 0.3 ml of FITC con A (Sigma) with 10 µg / ml concentration in PBS dulbeccos. Staining is done for 25 minutes at room temperature, then washed 2 times with 1500 rpm centrifugation for 10 minutes. The supernatant was removed and the precipitate was scratched on the slide labs of the specimen, dripping with 90% glycerol. Further specimens were observed in the epifluorescent microscope with excitation B (excitation 490 rpm with 525 nm emission) to determine fluorescence in FITC spermatozoa. Observations show: (a) spermatozoa with intact acrosome, and (b) spermatozoa without acrosom (Utomo, 2011). The data obtained were arranged in one table, then the white mouse capacitated results were analyzed using One Way Analysis of Variant (ANOVA) then proceed to Duncan to know the real difference between the treatments (Kusriningrum, 2008).

Result and Discussion

Experimental result of capacitation and acrosome reaction of rat (*Rattus norvegicus*) which was given pomegranate extract (*Punicagranatum L.*) after stained with FITC (Fluorescent Isotiocianat) and was observed with epifluorescent microscope 400x magnification as shown in Tabel 1.

Table 1. Experimental result of capacitation and acrosome reaction of rat (*Rattus norvegicus*) spermatozoa

Treatment	Spermatozoa Capatitation (X ± SD)	Spermatozoa Reaction Acrosome (X ± SD)
K	15,40 ^a ± 1,81	2,20 ^a ± 0,83
T1	17,60 ^a ± 1,81	3,00 ^a ± 0,70
T2	27,60 ^b ± 3,04	6,80 ^c ± 1,30
T3	25,00 ^b ± 1,58	5,20 ^b ± 0,83

Different superscripts in the same column showed significant differences ($p < 0.05$). K= Control treatment, Rats were given CMC solvent Na 0.5%, T1 = Treatment 1, White rats given pomegranate extract dose 75 mg / kg BW, T2 = Treatment 2, White rats given pomegranate extract dose 150 mg / kg BW, T3 = Treatment 3, White rats given pomegranate extract dose 300 mg / kg BW




Figure 1 Results of observation of capacitation and white rat spermatozoa acrosome reaction of white rat with FITC staining using epifluorescent microscope (A: Non-Capacitation, B: Capacitation, C: Acrosome Reaction).

Based on the results of the Analysis of Variance (ANOVA) test followed by Duncan test on the average of spermatozoa capacitation and acrosome reaction of white rats examination showed that there was a noticeable difference ($P < 0.05$). The results of examination of percentage of spermatozoa capacitation on K(15,40) and T1 (17,60) have no significant differences, T3 (25,00) and T2 (27,60) also have no significant difference, but K and T1 differ significantly with T2 and T3. In the Acrosome Reaction results, K (2,20) and T1 (3,00) had no significant differences, T2 (6,80) differed significantly with T3 (5,20), K and T1 had significant differences with T2 and T3.

Normally, free radical levels are maintained in physiological levels by antioxidants. Endogenous antioxidants that play a role in maintaining free radical levels are glutathione (GSH), superoxide dismutase (SOD), and catalase. GSH plays an important role in coordinating antioxidants in the body to counteract free radicals. However, if the production of excessive free radicals, there remains a part of the remaining ROS. To absorb the remaining ROS needs to be provided additional antioxidants (Makker, 2009).

Pomegranate is a potent source of exogenous antioxidants. All parts of pomegranate plants contain polyphenols and have antioxidant activity. The main group of phytochemicals in pomegranates is

the polyphenols found in the fruit. Polyphenols comprise flavonoids, hydrolysable tannins and condensed tannins (Yuniarti, 2012). Hydrolyzable tannins (HTs) as a major part of polyphenols in pomegranate and punicalaginis the most dominant part of HTs responsible for nearly half of the pomegranate juice antioxidant activity. Punicalagin is one of ellagitannins compounds found in many fruit and pomegranate stalks. Punicalagin contained in pomegranate has antioxidant activity up to 89% (Gil *et al.*, 2000).

In this study it was proved that the results of treatment using pomegranate extracts had a significant effect on increasing the amount of capacitation and spermatozoa acrosome reaction. With the results of the average research conducted can show the maximum level of use of the pomegranate extension. The average result of capacitation in this study showed that the percentage of spermatozoa control treatment capacitation showed a yield of 15.40 ± 1.81 . In the dosing group 75mg / kg BW showed a result of 17.60 ± 1.81 it can be concluded that the giving of pomegranate extract with a dose of 75 mg / kg BW can maintain spermatozoa capacitation of white rats. In the treatment with the dose of 150 mg / kg BW showed 27.60 ± 3.04 can be seen that with the provision of pomegranate extract 150 mg / kg BW proved abler to maintain spermatozoa capacitation to reach the ovum, because the success of fertility is determined by the spermatozoa capacitation status, membrane quality and penetration power against zona pellucida (Ba'a, 2009). In the examination of the percentage of capacitation with a dose of 300 mg / kg BW resulted in a $25,00 \pm 1.58$ percentage, the percentage result at the treatment of 300 mg / kg BW dose was decreased from a better dose of 150 mg / kg BW. This is thought to be due to the content in flavonoids and tannins contained in the pomegranate given has exceeded the optimal limit in dosing. One of the antioxidant content of tannin that can cause sperm clot, alkaloid suppress the secretion of reproductive hormone that is testosterone so that spermatogenesis process is disturbed (Winarno, 1997).

The result of percentage of acrosome reaction in this study showed that treatment with dosage of 150 mg / kg BW with a va-

lue of 6.80 ± 1.30 shows the highest rate of acrosome reaction results with different doses. At dosing 75 mg / kg BW increased compared with control treatment that was not given pomegranate extract. But the ANOVA statistic test followed by the Duncan test showed insignificant results. This suggests that administration of pomegranate extract at doses of 75 mg / kg BW may increase the acrosome reaction but the increase is not statistically significant. This increase shows that pomegranate extract acts as an exogenous antioxidant capable of neutralizing free radicals (Reynertson, 2007). This means that the polyphenols contained in the pomegranate are influential in maintaining the integrity of the spermatozoa membrane and by donating hydrogen atoms to free radicals and converting them into non-reactive forms (Bravo, 1998).

From the results it is clear that by giving the dose of pomegranate extract 150 mg / kg BW per oral give the biggest result that is $6,80 \pm 1,30$ which is different with treatment without giving extract of pomegranate. In the treatment of doses of 300 mg / kg BW showed a decrease in yield compared with a dose of 150 mg / kg BW. The decrease can be caused by the head of the spermatozoa cannot be protected by cytoplasmic membranes that have been damaged by imbalance doses and free radicals so it will be easy to suck the color so it can be concluded that the percentage of sperm that included death (Nugraheni, 2003).

Conclusion

The conclusion that can be drawn from this research is the effect of giving pomegranate extract (*Punicagranatum L.*) standardized 40% ellagic acid with doses of 75 mg / kg BW, 150 mg / kg BW and 300 mg / kg BW can improve the quality of spermatozoa, namely an increase in the percentage of capacitation and acrosome reactions in white rats (*Rattusnorvegicus*). The optimum dosage generated from this study was by administering a dose of 150 mg / kg BW.

References

Anwar, G.R.F. 2014. Pengaruh Pemberian infusa Kulit Manggis (*Garcinia mangostana L.*) Terhadap Reaksi

- Akrosom Spermatozoa Tikus Putih (*Rattus norvegicus*) [Skripsi]. Fakultas Kedokteran Hewan. Universitas Airlangga.
- Aulanni'am, M.A., dan Rosmaidar. 2007. Efek Antifertilitas Fraksi Air Biji Pinang (*Areca catechu*) Sebagai Agen Apoptosis Pada Sel-sel Jaringan Testis *Rattus norvegicus*. Jurnal Media Kedokteran Hewan. 23(3): 179-183.
- Aviram, M. 2014. Methods of Using Pomegranate Extract for Treating Diabetes Related Atherosclerotic Complications in Human. Google Patents.
- Ba'a, L.O. 2009. Peran D-fruktosa dan Kuning Telur dalam Proses Penghambatan Kapasitasi dan Kerusakan Membran Spermatozoa Kambing. Program Pasca Sarjana Unibraw. Malang.
- Bearden, H. J. and Fuquay, J. W. 2000. Applied Animal Reproduction. 5th Ed. Prentice Hall. New Jersey.
- Bravo, L. 1998. Polyphenols: Chemistry, Dietary Sources Metabolism, and Nutritional Significance. Nutrition Reviews. 56:317-333.
- Gil, M.I.Tomas-Barberan, F.A., Hess-Pierce, B., Haircoft, D.M., Kader, A.A. 2000. Antioxidant Activity of pomegranate juice and its Relationship with Phenolic composition and Processing. J Agric Food Chem. 48(10): 10-15.
- Kusriningrum. 2008. Perancangan Percobaan. Airlangga University Press. Surabaya.
- Lestari, V. P. 2014. Pengaruh Pemberian Infusa Kulit Manggis (*Garcinia mangostana* L) Terhadap Peningkatan Presentase Kapasitas Spermatozoa Tikus Putih (*Rattus norvegicus*) [Skripsi]. Fakultas Kedokteran Hewan. Universitas Airlangga.
- Makker, K., Agarwal, A., Sharma, R. 2009. Oxidative Stress and Male Infertility. Indian J Mes Res. 129: 67-357.
- Nugraheni T., Okid parama A., T widiyani. 2003. Pengaruh Vitamin C Terhadap Perbaikan Spermatogenesis dan Kualitas Spermatozoa Mencit (*Mus musculus* L.) Setelah pemberian ekstrak tembakau (*Nicotiana tabacum* L.). Biofarmasi 1(1): 13-7
- Reynertson, K. A. 2007. Phytochemical Analysis of Bioactive Constituents From Edible Myrtaceae Fruit. [Dissertation]. The City University of New York, New York.
- Saleh, R. A., Agarwal, A., Nada, E. A., El-Tonsy, M. H., Sharma, R. K., Meyer, A. 2003. Negative Effect of Increased Sperm DNA Damage in Relation to Seminal Oxidative Stress in Men With Idiopathic and Male Factor Infertility. Fertil Steril. 79(3): 160-1597.
- Sanocka, D. and Kurpisz, M. 2004. Reactive Oxygen Species and Sperm Cell. Reprod. Biol. Endocrinol. 2:12.
- Sikka, S. C. 2004. Role of Oxidative Stress and Antioxidant in Andrology and Assisted Reproductive Technology. J. Androl. 25: 5-18.
- Utomo, B. 2011. Suplementasi Akrosin Pada Semen Kambing Peranakan Etawa Pasca Thawing Terhadap Peningkatan Kualitas dan Potensi Spermatozoa [Disertasi]. Universitas Airlangga. Surabaya.
- Winarno, F.G. 1997. Kimia Pangan dan Gizi. Gramedia Pustaka. Jakarta
- Yuniarti, W.M. 2012. Efek Antifibrotik Ekstrak Buah Delima (*Punica granatum* L.) pada Fibrosis Hati [Disertasi]. Universitas Airlangga. Surabaya
- Yanjun, Z., Dana, K., Robert, D., Rypo, L., dan David, W. 2009. International Multidimensional Authenticity Specification (IMAS) Algorithm for Detection of Comercial Pomegranate Juice Adulteration. J. Agric Food Chem. 57(6): 2550-2557.
- Zavos, P. M. and Zarmakoupis-Zavoz, P. N. 1999. Impact of Cigarette Smoking on Human Reproduction: Its Effect on Male and Female Fecundity. Technology. 6: 9-16.