
galley proof Correlation between Mother's Behavior with Periodontal Status

Acta Medica Philippina UP Manila <actamedicaphilippina.upm@up.edu.ph>
To: mega-m-p@fkg.unair.ac.id, symposium@fkg.unair.ac.id

Tue, Nov 26, 2019 at 10:00 AM

Nov. 26, 2019

Mega Moeharyono Puteri
Department of Pediatric Dentistry
Faculty of Dental Medicine
Universitas Airlangga

Dear Dr. Puteri,

Attached is the galley proof of your manuscript "Correlation between Mother's Behavior with Periodontal Status and Periodontal Treatment Needs in Autistic Children". If you have corrections, kindly email us as soon as possible (our new email address actamedicaphilippina.upm@up.edu.ph).

May we also respectfully request you to include Reference #26 in the main text (you have 26 references in the list)

Thank you very much!

Sincerely,

Dr. Abigail C. Pascual-Domingo



Correlation between Mother's Behavior with Periodontal Status.pdf
191K

Correlation between Mother's Behavior with Periodontal Status and Periodontal Treatment Needs in Autistic Children

Mega Moeharyono Puteri, Tania Saskianti and Ridha Rizki Ananda

Department of Pediatric Dentistry, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia

ABSTRACT

Background. Autism is a form of a neurological disorder. Autistic children have problems related to physical, psychological, and mental barriers that can hinder their ability to achieve optimal dental health status. Therefore, in maintaining the dental health of autistic children, it is influenced by parents teaching skills and habits in autistic children's life. From previous study, there were about 17.4% autistic children in Saudi Arabia suffering from bleeding of the gingiva. Periodontal disease is often found in autistic children.

Objective. This study analyzes the relationship between mother's behavior with periodontal status and periodontal treatment needs of autistic children.

Methods. Analytical observational study with cross-sectional approach at AGCA Centre Surabaya with a total sample of 34 pairs of autistic children and their mothers. This study used the HU-DBI questionnaire which consisted of knowledge, attitude, and mother's action and oral examination of autistic children with the CPITN index.

Results. Of the autistic children, 55.8% had healthy periodontal status. The knowledge, attitudes, and actions of mothers were high. Statistical results with Spearman correlation test obtained a value of $p > 0.05$ on aspects of knowledge, attitudes, and actions towards the CPITN index and periodontal treatment needs.

Conclusion. There was no significant correlation between the mother's behavior and the periodontal status and periodontal treatment needs of autistic children in managing their oral health.

Key Words: periodontal tissue, CPITN index, mother's behavior, autism

INTRODUCTION

Autism can be defined as a neurological development disorder that causes children to appear unable to relate to others, as if they live in their own world.¹ The main characteristics of autism are related to problems of social interaction with others, communication disorders, repetitive behavior, and cognitive instability.² The Centers for Disease Control and Prevention (CDC) in the United States in March 2013 reported that the prevalence of autism increased to 1:50 in the past year. When the number of births in Indonesia is 6 million per year, the number of autistic children in Indonesia increases by 0.15% or 6,900 children per year.³

In Saudi Arabia, 17.4% of autistic children have been reported to have gingival bleeding.⁴ Meanwhile, Da Silva et al. (2017) state that oral health problems that are often found in autistic children are periodontal disease.⁵ Meanwhile, the dominant periodontal disease that is found

Paper presented at the Joint Scientific Meeting in Special Care Dentistry, July 5, 2019, Amerta Room, 4th Floor, main campus of Universitas Airlangga, Surabaya, Indonesia.

Corresponding author: Mega Moeharyono Puteri
Department of Pediatric Dentistry
Faculty of Dental Medicine
Universitas Airlangga
Jl. Mayjen. Prof. Dr. Moestopo No. 47 Surabaya 60132 - Indonesia
Email: mega-m-p@fkg.unair.ac.id

in children and adolescents is gingivitis.⁶ Gingivitis is an inflammation that only involves gingival tissue with no loss of attachment or bone.⁷

Due to the problems of social interaction with others, communication disorders, and cognitive instability, autistic children cannot maintain their oral hygiene individually.⁸ Kalyoncu and Tanboga (2017) state that there was no difference between the prevalence of periodontal disease in autistic children so that the incidence of periodontal disease in autistic children is not much different from normal children.⁹ Another previous study also mentions that autistic children can maintain their good oral hygiene despite the limitations they have but their oral hygiene is based on family members healthy lifestyle habits.¹⁰

Autistic children's behavior in maintaining dental health such as tooth brushing habit and not being afraid to go to the dentist can be influenced by families/parents that teach skills and habits in autistic children's daily life.¹⁰ For this reason, oral health of autistic children is very dependent on their parents.¹¹ The mother is the closest person to a child who always educates, teaches, and trains children's skills, especially in terms of maintaining oral hygiene. A child tends to take a behavior similar to his/her mother's behavior due to a process of imitation of a model that he/she considers important. Mother's behavior regarding oral health can be used to predict the status or condition of the child's oral health.¹²

A few other studies describe the correlation between mother's behavior with periodontal status and periodontal treatment needs in autistic children. For this reason, the researchers wanted to examine the correlation between mother's behavior towards the periodontal status in children with autism and their periodontal treatment needs.

MATERIALS AND METHODS

This was an observational analytic study using a cross-sectional study design. The study was conducted at Special Education AGCA Centre Surabaya in September - October 2018. The population in this study were all children diagnosed with autism in the Special Education AGCA Centre Surabaya, i.e., 34 students along with parents of autistic children according to the selection criteria for the research; each study participant had signed an informed consent form and the research protocol had been approved by an ethics committee. Criteria for selection of research subjects include: children diagnosed with autism, aged 3-14 years, in a healthy physical condition, coming accompanied by parent or caregiver, and being cooperative. The sampling technique used was total sampling.

Data collection was obtained from the results of the HU-DBI modification questionnaire (Hiroshima University - Dental Behavioral Inventory) regarding the behavior of maintaining oral health by mothers as parents of autistic children and examining the status of periodontal tissue of autistic children using the CPITN Index.¹³ Informed

consent was given by mothers as respondents to be asked for research approval; respondents were interviewed directly to answer the questionnaire, and the status of periodontal of autistic children was examined using the CPITN Index. The results obtained were processed and presented in table form. Data analysis was performed using the Spearman correlation test to analyze the relationship between the mother's behavior and the status of periodontal health and periodontal treatment needs in autistic children.

RESULTS

Data collected from questionnaire were in the form of the general characteristics of respondents and the relationship between mother's behavior and health of periodontal tissue and periodontal care needs of autistic children. In this study group, most of autistic children were at the age of 7-9 years in the mixed dentition stage at 50%, followed by the age group >10 years at 26.47%, and the age group 3-6 years at 23.5%.

The educational background of the mothers with college education was 79.41%, followed by the group of mothers with primary education (Elementary/Junior High/Senior High School) which was 20.58%. In this study, it was found that 47% of mothers did not have jobs, which was the majority of the total number of samples, followed by mothers who worked as private employees at 32.35%, mothers who worked as civil employee at 8.8%, and mothers with other jobs. Most mothers who had independent income above 3 million rupiahs consisted of 50% of the total sample; mothers who did not have independent income were 29.41%; mothers with income range of 1 million - 3 million rupiahs were 17.64%; and mothers who had independent income under 1 million rupiahs were 2.94%.

Table 1 shows that most autistic children had a score of 0 or healthy periodontal tissue, i.e., 19 children (55.8%). Then, there were 8 children with autism who experienced bleeding on probing (23.5%), and children whose teeth had sub/supra gingival calculus on probing consisted of 7 (20.5%). Table 2 shows that most children did not need periodontal treatment, i.e., 19 children (55.8%). In addition, 8 children (23.5%) needed an improvement in oral hygiene, while 7 children (20.5%) needed dental health education and professional scaling. In this study, a HU-DBI questionnaire was used to measure the behavior of mothers in caring for and maintaining the health of their children's dental and oral health. Table 3 shows the distribution of mother's behavior in maintaining the children's dental and oral health. It is known that behavior is divided into three aspects, namely aspects of knowledge, attitudes, and actions. In the aspect of knowledge, it has the highest average value as compared to the other two aspects, namely all respondents had high scores or had average of 100%. Furthermore, the attitude aspect had a low value of 8.8% and a high value of 91.1%. In addition, the aspect of action had low value of 47% and high value of 53%.

Table 1. Periodontal status in autistic children with the CPITN index

Score of CPITN	Frequency	Percentage (%)
0	19	55.8
1	8	23.5
2	7	20.5
3	0	0
Total	34	100

Table 2. Periodontal treatment needs in autistic children with the CPITN index

Treatment needs	Frequency	Percentage (%)
0	19	55.8
1	8	23.5
2	7	20.5
3	0	0
Total	34	100

Table 3. Distribution of mother's behavior in maintaining children's dental and oral health

Aspect	Frequency	Percentage (%)
Knowledge		
High	34	100
Low	0	0
Attitude		
High	31	91.1
Low	3	8.8
Action		
High	18	53
Low	16	47

Table 4. Spearman correlation between mother's behavior and CPITN score

Score	Knowledge		Attitude		Action	
	Sig (p)	r	Sig (p)	r	Sig (p)	r
Treatment	0.5	0.10	0.4	-0.32	0.7	-0.54
Needs	0.5	0.10	0.5	-0.10	0.6	-0.85

Once the results of the study were analyzed, it was found that the significance value (p) was > 0.05, so there was no correlation seen between the mother's behavior and the periodontal status (CPITN score), both from the aspects of knowledge, attitudes, and actions of the mothers and correlations with children's treatment needs (Table 4).

DISCUSSION

In this study, the number of autistic children examined was mostly male, i.e., 61.7%, while females consisted of only 38.3%. This is consistent with Nugraheni's research in 2012 stating that the prevalence of autism ranged from 1-2 per 1000 inhabitants and mostly found in men than women (4:1).¹⁴ This was related to differences in hormone

production. From a study in autistic person ranging from 6-20 years old, it was found that men produce more testosterone hormone while women produce more estrogen hormones. Both hormones have the opposite function of a brain regulating gene called alpha retinoic acid related orphan receptor or RORA. The testosterone hormone can inhibit the performance of RORA while estrogen increases the work of RORA. If RORA's work is hampered, there will be brain coordination problems, for example, a gene should have the duty to protect nerve cells from stress and inflammation; meanwhile, if the performance is hampered, the cell cannot work properly.¹⁵

Periodontal status in autistic children examined using the CPITN index showed that 44% of autistic children had problems in their periodontal tissue. A total of 19 children from 34 total research subjects of autistic children had healthy periodontal tissue conditions, so that the periodontal treatment needs of autistic children in this study also amounted to a small number, i.e., 44%, which required an increase in oral hygiene and professional scaling.

The results of this study showed that the majority of independent income of mothers was in the range of 1-3 million rupiahs, i.e., 24 people. This is in accordance with the research conducted by Duncan and Bonner in 2014, stating that a person's ability to pay for dental care for dental insurance coverage is directly related to the individual's income. High income individuals have a greater chance of receiving better dental care.¹⁶ Income received by mothers can be an addition to families to improve socio-economic status. Families with high socio-economic status will find it easier to obtain health services. Most of the samples in this study came from parents who had high socio-economic status, so they had a greater chance of getting health services in achieving better dental health.¹⁷

Furthermore, the profession of the mother is also influential. Of the total, 47% of the mothers of autistic children in this study did not have a job or were housewives, so they had more time to pay attention to maintaining the dental health of their children considering each autistic child has limitations in maintaining his/her own dental health and is very dependent to parents especially the mother. This is because parents are role models for their children and are decision makers. Parental decisions are influenced by their attitudes which reflect children's oral health. If the mother's behavior regarding oral and dental health is good, her child's behavior will also be good.¹⁷

All respondents had a high level of knowledge, i.e., 100%. This was consistent with the research conducted by Susi, Bachtiar and Azmi in 2012, stating that there was a relationship between the level of knowledge of parents about dental and oral health with the behavior of maintaining dental health in children. Majority of respondents, where 79.41% of them had college education, while the rest of the respondents who had primary education (SD/SMP/SMA) only consisted of 20.58%.¹⁸ Education can affect individual

health status. The higher the level of education of a person, the easier it is to receive information.¹⁹ Someone who has a higher education will have better knowledge in maintaining and caring for his teeth.¹⁶

A total of 91.1% respondents had a good attitude in maintaining the dental health of their children. This was consistent with the research conducted by Oktarina in 2016, stating that there was a correlation between the level attitudes of parents about dental and oral health with the behavior of maintaining dental health of their children.¹⁹ Education will enhance information and influence the development of individual attitude towards newly introduced values.¹⁹ According to Notoatmodjo (2007), attitudes are the readiness of individuals to act according to their feelings and thoughts based on the values believed. So that attitude can be learned, instead of taken from birth; it is not settled, it can change. Attitude is the beginning of behavior that affects individual actions. This can be influenced by parents' initiative in preventing and promoting dental health problems.²⁰ The higher the initiative of parents in terms of maintaining health, the greater the action that will be formed and the healthier the periodontal tissue the child will be.

The results of this study in the aspect of action showed that 91.1% of respondents had good actions in maintaining the dental and oral health. This was consistent with the research conducted by Nurjannah in 2016 stating that there was a correlation between actions with the level of dental and oral hygiene.²¹ This is due to the level of full awareness of respondents regarding dental and oral health which has an impact on the hygiene status of their teeth and mouth. The higher level of the mother awareness, the more actions produced by the mother will be so that the child's optimal dental and oral health can be achieved.

After analyzing the data, this study showed that there was no significant relationship between the mother's behavior and the periodontal status and periodontal treatment needs in autistic children ($p > 0.05$) in aspects of knowledge, attitudes, and actions. This is consistent with the study of Oktarina, Tumaji and Roosihermatie in 2016 stating that there was no significant relationship between mother's behavior (knowledge, attitudes, and actions) with children's oral and dental health.¹⁹ This can be caused by knowledge, attitudes, and actions which are not being the only factors that influence a person's behavior. In Oktarina et al's research, (2016), they noted Blum's theory which states that the health status of a person's mouth and society is influenced by four important factors, namely heredity, environment (physical and cultural), behavior, and health services. Mother's parenting also affects the behavior of mothers in caring for and maintaining the dental health of children. In this study, it was shown that the health of the periodontal tissues of autistic children tended to be healthy so that they did not require large maintenance needs.¹⁹

In this study, 44.1% of parents of autistic children used Indonesian's national health insurance and social security

(BPJS). Besides mother's income and profession, health insurance also has a significant impact on dental care. The scope of dental insurance services is an important factor in deciding to seek dental care. The use of dental insurance can reduce or even eliminate the cost of dental and oral care which causes high demand for dental and oral health care.²² The scope of dental and oral services guaranteed by BPJS Health includes: extraction of primary teeth (topical, infiltration), permanent tooth extraction without complications, teeth scaling (1x a year), etc.²³ Ease of access to dental health service facilities provided can encourage mothers to take their children to the dentist in terms of care for the child's teeth.

Special education of AGCA Centre located in Surabaya, East Java is a school for children who have special needs, such as autistic children, cerebral palsy, etc. AGCA Centre has an accreditation which is included in the superior schools and is located in elite high socioeconomic locations. The facilities and services provided are good for the costs required are not small so that not all autistic children can attend the AGCA Centre; only parents who have high socioeconomic status can send their autistic children to AGCA. This can lead to uniformity or homogeneous data so that there is no big difference in the periodontal status of autistic children, so the majority of autistic children have a healthy periodontal tissue condition with little maintenance needs. In addition, environmental factors for autistic children can also affect the health of the periodontal tissue rather than only from the mother, most of the time the child is used to going to the AGCA Centre daily. Information about maintaining the dental health of children can also come from schools, teachers, and the internet from the gadgets being played by children.

Oral and dental health of autistic children can be affected due to limited understanding of the importance of maintaining oral health, difficulties in communicating oral health needs, consumption of anticonvulsant drugs that affect gingival health, and fear of oral health procedures. In addition, autistic children tend to be hyperactive and have uncontrolled pulling of muscles. Autistic children also have low levels of salivary secretions which can also lead to low oral hygiene.²⁴

In addition, the difficulty in processing sensory stimuli is something that is found in many people with autism. Research by Cermak shows that the children's being more receptive to sensory responses is an important factor that influences their response to receive dental and oral care.²⁴ Sensory stimuli that can be found in dental practice, such as bright neon lights from the dental chair, touches in and around the oral cavity in dental care actions, the atmosphere and smell of oral care products in the practice of dentists or hospitals, have the potential negative impact in these children, making it difficult for dentists to provide care. This can cause an increase in stress and anxiety of autistic children, and cause uncooperative behavior in receiving dental and oral care.²⁵

Autistic children also tend to have self-injurious behaviors (SIB), which are located in the head-neck area, which can include head banging, face tapping, and gingival picking. In addition, they sometimes also have unusual parafunctional habits including bruxism, stretching their tongue forward, chewing things carelessly like gravel, cigarette butts, or pens. Children with autism also tend to have a preference for sweet and cariogenic foods. This can cause serious problems in the oral and dental health of children with autism.²⁵ This is in contrast to the results obtained in this study that most autistic children have healthy periodontal tissue. This can be caused by the level of awareness from mothers of autistic children in maintaining their children's dental and oral health. School environment also supports children so that they can always maintain and care for the oral health of children. This study examined the mother knowledge, attitudes, and actions to maintain their children's dental and oral health. Further study is necessary to examine the correlation between mothers' educational background and socio-economic status to obtain deeper information with Periodontal Status and Periodontal Treatment Needs in Autistic Children.

CONCLUSION

The results of research showed that there was no significant correlation between knowledge, attitudes, and actions of mothers in maintaining dental and oral health and the periodontal status and periodontal treatment needs in autistic children.

Statement of Authorship

All authors participated in data collection and analysis, and approved the final version submitted.

Author Disclosure

All authors declared no conflict of interest.

Funding Source

None.

REFERENCES

- Jaber M. Dental caries experience, oral health status and treatment needs of dental patients with autism. *J Appl Oral Sci.* 2011; 19(3):212-7.
- Morales-Chávez MC. Oral health assessment of a group of children with autism disorder. *J Clin Pediatr Dent.* 2017; 41(2):147-9.
- R. Arfiriana, and F. Dieny. Relationship to casein-free gluten-free diet score with autism behaviour score. *Journal of Nutrition College.* 2014;3(1): 34-42.
- Murshid EZ. Diet, Oral Hygiene Practices and Dental Health in Autistic Children in Riyadh, Saudi Arabia. *OHDM.* 2014; 13(1):91-6.
- da Silva SN, Gimenez T, Souza RC, Mello-Moura ACV, Raggio DP, Morimoto S, et al. Oral Health Status of Children and Young Adults with Autism Spectrum Disorder: Systematic Review and Meta - Analysis. *Int J Paediatr Dent.* 2017; 27(5):388-98.
- Varas F, Zillmann G, Munoz A, Hassi J, Yevenes I, Echeverria S, et al. Periodontal status and treatment needs of children from 6 to 8 years old in the Santiago Metropolitan Region of Chile. *Rev Odonto Cienc.* 2011; 26(1):10-5.
- William V. Stenberg Jr., 6th ed. *Periodontal Problems in Children and Adolescents.* Pediatric Dentistry. 2019: 371-378.
- Kumar S, Sharma J, Duraiswamy P, Kulkarni S. Determinants for oral hygiene and periodontal status among mentally disabled children and adolescents. *J Indian Soc Pedod Prev Dent.* 2009. 27(3):151-7.
- Kalyoncu IÖ, Tanboga I. Oral health status of children with autistic spectrum disorder compared with non-authentic peers. *Iran J Public Health.* 2017; 46 (11):1591-3.
- Nonong YH, Setiawan A, Dewi FD Navaneetha C. Oral health knowledge among parents of autistic child in Bandung-Indonesia. *Dental Journal.* 2014. 47(3):146-52.
- Sengkey MM, Pangemanan DHC, Mintjelungan CN. Dental and oral hygiene status autistic children in Manado. *Jurnal e-Gigi.* 2015; 3(2):235-40.
- Pintauli S. Analysis of behavior relationship of dental and oral health care on dental and oral health status of elementary and junior high school students in Medan. *Jurnal Pendidikan dan Kebudayaan.* 2010; 16(4):376-90.
- Dagli RJ, Tadakamadla S, Dhanni C, Duraiswamy P, Kulkarni S. Self reported dental health attitude and behavior of dental students in India. *J Oral Sci.* 2008; 50(3):267-72.
- Nugraheni SA. Revealing the Wilderness of Autism. *Psychology Bulletin.* 2012; 20(1): 9-17.
- Deshinta R, Hardiani RS, Dewi EI. The Effect of Glenn Doman Method towards Autistic Social Interaction Ability in SLB-B and Autistic TPA. *E-jurnal Pustaka Kesehatan.* 2015; 3(1):66-73.
- Duncan L, Bonner A. Effects of Income and Dental Insurance Coverage on Need for Dental Care in Canada. *J Can Dent Assoc.* 2014; 80:e6.
- Bahaguna R, Jain A, Khan SA. Knowledge and attitudes of parents regarding child dental care in an Indian population. *Asian J Oral Health Allied Sci.* 2011; 1:9-12.
- Susi BH, Azmi U. Relationship of parents' socio-economic status with caries in the child's primary teeth. *Majalah Kedokteran Andalas.* 2012; 36(1):96-105.
- Oktarina, Tumaji, Roosiermiatic B. Correlation of mother factors and their kindergarten's oral health status in Kelurahan Kemayoran , Krembangan Sub district, Surabaya City. *Bulletin Research in Health System.* 2016; 19(4):226-35.
- Kumar G, Dileep C L, Ahuja R. Oral health of preschool children in Dhanbad District, Jharkhand: A gander into the maternal behavior and practices. *J Family Med Prim Care.* 2016; 5(4):829-33.
- Nurjannah. Relationship of Knowledge, Attitude, and Practice of Dental and Oral Health to Oral Hygiene Status in Students Junior High School (SMP) /MTs Pondok Pesantren Putri Ummul Mukminin. *Essay. Faculty of Dentistry. Hasanuddin University;* 2016. pp 1-64
- Danial Z, Kalantar Motamedi MH, Mirhashemi S, Khoshmohabat H, Danial N. Universal Health Coverage and Oral Health. *J Interdiscipl Med Dent Sci.* 2016; 4:e106.
- Social Insurance Administration Organization (BPJS). *Practical Guide to Dental Services & Dental Prosthesis for the Indonesia National Health Insurance Participants.* BPJS Kesehatan. 2014; 9-11.
- Blomqvist M, Bejerot S, Dahllöf G. A cross-sectional study on oral health and dental care in intellectually able adult with autism spectrum disorder. *BMC Oral Health.* 2015; 15:81.
- Cermak SA, Duker LIS, Williams ME, Dawson ME, Lane CJ, Polido JC. Sensory Adapted Dental Environments to Enhance Oral Care for Children with Autism Spectrum Disorders: A Randomized Controlled Pilot Study. *J Autism Dev Disord.* 2015; 45(9):2876-88.
- Naidoo M, Singh S. The oral health status of children with autism spectrum disorder in KwaZulu-Nata, South Africa. *BMC Oral Health.* 2018; 18:165.

REVISI ARTIKEL PUBLIKASI ACTA MEDICA PHILIPINA

mega moeharyono puteri <mega-m-p@fkg.unair.ac.id>
To: Symposium FKG Universitas Airlangga <symposium@fkg.unair.ac.id>

Sun, Oct 20, 2019 at 4:53 PM

Kepada
Yth Panitia Artikel Ilmiah
Symposium FKG Universitas Airlangga

Bersama dengan ini saya kirimkan cover letter, author declaration, ethic dan revisi naskah.
atas bantuan dan kerjasama yang diberikan terimakasih.

wassalam wr wb.

Mega Moeharyono Puteri, DDS., Sp.KGA., Ph.D
Lecturer Staff Department of Pediatric Dentistry
Board Members for Indonesia Pediatric Association (IDGAI)
Faculty of Dental Medicine
Universitas Airlangga - Campus A
Jl. Mayjen Prof Dr Moestopo no 47
Surabaya, Indonesia

Representative staff for WUACD
(World Universities Association for Community Development)
Management office Universitas Airlangga - Campus C
Jl. Dr. Ir. H. Soekarno No.200,
Mulyorejo, Kec. Mulyorejo, Surabaya,
Jawa Timur 60115
Indonesia



Virus-free. www.avg.com

[Quoted text hidden]

5 attachments



etik RD.jpg
1149K

 **Revisi Rev 1 MG RD.docx**
63K

 **Reply Template Acta Medica Rev 1 MG RD.doc**
24K

 **Acta Medica Philipina Author Declaration MG TS and Ridha TTD.pdf**

266K



Cover letter Acta Medica MG RD (1).PDF

4345K



**UNIVERSITAS AIRLANGGA FACULTY OF DENTAL MEDICINE
HEALTH RESEARCH ETHICAL CLEARANCE COMMISSION**

ETHICAL CLEARANCE CERTIFICATE

Number : 192/HRECC.FODM/VII/2018

Universitas Airlangga Faculty Of Dental Medicine Health Research Ethical Clearance Commission has studied the proposed research design carefully, and therefore, shall herewith certify that the research entitled :

“HUBUNGAN PERILAKU IBU TERHADAP KESEHATAN JARINGAN PERIODONTAL DAN KEBUTUHAN PERAWATAN PERIODONTAL ANAK AUTIS”

Principal Researcher : RIDHA RIZKI ANANDA

Unit/Institution/Place of Research : - SLB AGCA Center, Jl. Ngagel Jaya Tengah
3 No. 21 Surabaya
- SD Muhammadiyah 16, Jl. Barata Jaya I No.
11 Surabaya

CERTIFIED TO BE ETHICALLY CLEARED



Surabaya, July 31th, 2018
Chairman,

Prof. Dr. M. Rubianto, drg.,MS.,Sp.Perio(K)
Official No.195009081978021001

To,

The Editor of Acta Medica Philippina

Sub: Submission of Manuscript for publication

Dear Sir,

We intend to publish an article entitled “***Correlation Between Mother's Behavior With Periodontal Status and Periodontal Treatment Needs In Autistic Children***” in your esteemed journal as an Original Article.

I/We have been sufficiently involved in this work to take public responsibility for its validity and final presentation as an original publication. The article had written/informed consent for publication from the involved subject/s, had conformed with ethical standards, and/or had been reviewed by the appropriate ethics committee.

This paper has not been published and is not under simultaneous consideration for publication elsewhere. This paper has been subjected to ethics review Universitas Airlangga Faculty of Dental Medicine Health Research Ethical Clearance Commission (Number: 192/HRECC.FODM/VII/2018), July 31st 2019 and has been endorsed favorably by the said board or committee. I/We hereby confer all copyright ownership/s to the Acta Medica Philippina in the event that this work is published in this journal.

We hereby transfer, assign, or otherwise convey all copyright ownership, including any and all rights incidental thereto, exclusively to the journal, in the event that such work is published by the journal.

Thanking you,

Yours' sincerely,

Signature

Corresponding contributor:

Mega Moeharyono Puteri, Department of Pediatric Dentistry, Faculty of Dental Medicine, Universitas Airlangga, Jl. Mayjend. Prof. Dr. Moestopo No. 47 Surabaya 60132, Indonesia. E-mail: mega-m-p@fkg.unair.ac.id

(A manuscript for an original article should not exceed 25 typewritten pages, including tables, figures, illustrations and references).

Title of the article:

Correlation Between Mother's Behaviour with Periodontal Status and Periodontal Treatment Needs In Autistic Children

Running title

Author (s)

1 Mega Moeharyono Puteri, drg., Ph.D., Sp.KGA(K)

Tania Saskianti, drg., Ph.D., Sp.KGA(K)

2 Ridha Rizki Ananda

Department(s) and institution(s)

1 Staff Department of Pediatric Dentistry, Faculty of Dental Medicine, Universitas Airlangga Surabaya – Indonesia

2 Students Faculty of Dental Medicine, Universitas Airlangga Surabaya-Indonesia

Corresponding Author:

Name Mega Moeharyono Puteri

Address Jl. Mayjend. Prof. Dr. Moestopo No. 47 Surabaya 60132, Indonesia

Phone numbers +62 812 98771488

Facsimile numbers -

E-mail address mega-m-p@fkg.unair.ac.id

Total number of pages:

Total number of photographs:

Word counts

for abstract: 200

for the text:

Presentation at a meeting: Joint Scientific Meeting in Special Care Dentistry

Organisation

Revisi Rev 1 MG RD

Place Surabaya Indonesia

Date July 5 th, 2019

Conflicting Interest (If present, give more details): -

Title of the article:

Correlation Between Mother's Behaviour with Periodontal Status and Periodontal Treatment Needs in Autistic Children

Abstract (Max 200 words):

Background:

Autism is a form of a neurological disorder which causes brain function didn't work normally. Periodontal disease is often found in autistic children. Autistic children have problems related to physical, psychological and mental barriers that can hinder their ability to achieve optimal dental health status. Because of their limited ability, they cannot maintain their oral hygiene. Oral health of autistic children is depending on their parents. This study analyses the relationship between mother's behaviour with periodontal status and periodontal treatment needs of autistic children.

Methods:

Analytical observational study with cross-sectional approach at SLB AGCA Centre Surabaya with a total sample of 34 autistic children and their mothers. This study uses the method of answering the HU-DBI questionnaire by mothers and oral examination of autistic children with the CPITN index.

Results:

55.8% of autistic children had a healthy periodontal status. The knowledge, attitudes and actions of mothers were high, statistical results with Spearmen correlation test obtained a value of $p > 0.05$ on aspects of knowledge, attitudes and actions towards the CPITN index and periodontal treatment needs.

Revisi Rev 1 MG RD

Conclusion:

There was no significant relationship between mother's behaviour with periodontal status and periodontal treatment needs of autistic children in managing their oral health.

Keywords:

Periodontal tissue, CPITN index, Mother's Behaviour, Autism

Introduction:

Autism can be defined as a neurological development disorder that causes children to appear unable to relate to others, as if they live in their own world.¹ The main characteristics of autism are related to problems of social interaction with others, communication's disorders, repetitive behaviour, repetition of behaviour, and Cognitive instability.² The Prevention of Centre Disease Control (CDC) in the United States in March 2013 reported that the prevalence of autism increased to 1:50 in the past year. If the birth rate in Indonesia is 6 million per year, the number of autistic children in Indonesia increases by 0.15% or 6,900 children per year.

In Saudi Arabia, 17.4% of autistic children having a gingival bleeding.⁴ Whereas according to Da Silva et al in 2017, oral health problems that are often found in autistic children are periodontal disease.⁵ Whereas, the dominant periodontal disease that happened in children and adolescents are gingivitis.⁶ Gingivitis is an inflammation that only involves gingival tissue around the teeth.⁷

Due to the limited ability of autistic children, they cannot maintain their oral hygiene.⁸ However, according to Kalyoncu and Tanboga's research in 2017, there was no difference between the prevalence of periodontal disease in autistic children so that the incidence of periodontal disease in autistic children is not much different from normal children.⁹ Most other autistic children can maintain their good oral hygiene behind the limitations they have.¹⁰

Behaviour to maintaining the dental health of autistic children can be influenced by families/parents of autistic children where parents teach skills and habits in of autistic children's life everyday.¹⁰ For this reason, oral health of autistic children is very dependent on their parents.¹¹ Mother as the closest person to a child who always educates, teaches, and trains children's skills, especially in terms of maintaining oral

hygiene. A child tends to take an attitude similar to his mother's attitude due to a process of imitation or imitation of a model that he considers important. Mother's behaviour regarding to oral health can be used as an example by children so that it can be used to predict the status or condition of their child's oral health.¹²

Until now there have been no studies that describe the relationship between mother's behaviour with periodontal status and periodontal treatment needs in autistic children in Surabaya, for this reason the researchers wanted to examine the relationship between mother's behaviour towards the periodontal status in children with autism and their periodontal treatment needs.

Materials and Methods:

This study was an observational analytic study using a cross-sectional study design. The study was conducted at SLB AGCA Centre Surabaya in September - October 2018. The population in this study were all children that diagnosed with autism in the SLB, amounting to 34 people along with parents of autistic children according to the criteria of the research subject. Criteria for the research subject were used children diagnosed with autism aged 3-14 years, in a healthy physical condition, coming accompanied by parents or caregiver, and being cooperative. The sampling technique used is total sampling.

Data collection was obtained from the results of answering the HU-DBI questionnaire (Hiroshima University - Dental Behavioural Inventory) regarding the behaviour of maintaining oral health by mothers as parents of autistic children and examining the status of periodontal tissue using the CPITN Index. Informed consent was read for mothers as respondents to be asked for research approval, respondents

were interviewed directly to answer questions in the questionnaire, and the status of periodontal tissue was examined using the CPITN Index. The results obtained are processed and presented in table form. Data analysis was performed using the spearman correlation test to analyse the relationship between mother's behaviour towards the status of periodontal and the periodontal treatment needs in autistic children.

Results:

This research was conducted on SLB AGCA Centre students in Surabaya with a total sample of 34 autistic children and their biological parents as respondents in September - October 2018. After the results of the study were analysed it was found that the significance value (p) was > 0.05 so there was no correlation between mother's behaviour and the periodontal status (CPITN score) both from the aspects of knowledge, attitudes, and actions and correlations with treatment needs.

Discussion:

In this study, the percentage of autistic children examined was more amount in male, that was 61.7% and females 38.3%. This is consistent with Nugraheni's research in 2012, which stated that the prevalence of autism ranged from 1-2 per 1000 inhabitants, with the distribution of men more than women (4:1).¹³ This was related to differences in hormone production. In men, more produce the hormone testosterone while women produce more of the hormone estrogen. Both of these hormones have the opposite function of a brain regulating gene called alpha retinoic acid related orphan receptor or RORA. The hormone testosterone can inhibit the performance of RORA while estrogen increases the work of RORA. If RORA's work is hampered, there will be brain coordination problems, for example, a gene should have the duty to protect nerve cells from stress and inflammation, but if the performance is hampered, the cell cannot work properly.¹⁴

Periodontal status in autistic children examined using the CPITN index showed that as many as 44% of autistic children had problems in their periodontal tissue, 23.5% of each had bleeding on probing and 20.5% had sub gingiva and supra gingiva calculus. A total of 19 children from 34 total research subject of autistic children had healthy periodontal tissue conditions, so that the periodontal treatment needs of

autistic children in this study also amounted to a small number, requiring treatment by 44%, which requires an increase in Oral Hygiene and professional scaling.

In the results of the study found that the majority of independent income of mothers in the range of 1-3 million as many as 24 people while those who do not have independent income as many as 10 people. This is in accordance with the research conducted by Duncan and Bonner in 2014, that a person's ability to pay for dental care for dental insurance coverage is directly related to the individual's income. High income individuals have a greater chance of receiving better dental care.¹⁵ Income received by mothers can be an addition to families to improve socio-economic status. Families with high socio-economic status will find it easier to obtain health services. Most of the samples in this study come from parents who have high socio-economic status so they have a greater chance of getting health services in achieving higher dental health.¹⁶

In addition to income, the profession of the mother can also influence. 47% of mothers of autistic children in this study did not work or as housewives, so they have more time to pay attention to their autistic children in maintaining the dental health of their children considering this autistic child has limitations in maintaining his own dental health so it is very dependent to parents especially their mothers. This is because parents are role models for their children and decision makers. Parental decisions are influenced by their attitudes which reflect children's oral health. If the mother's behaviour regarding oral and dental health is good, then her child's behaviour will also be good.¹⁶

All respondents have a high level of knowledge as much as 100%. This is consistent with the research conducted by Susi, Bachtiar and Azmi in 2012, that there was a relationship between the level of knowledge of parents about dental and oral

health with the behaviour of maintaining dental health in children. This can be caused by the majority of respondents, number of 79.41% having the last education, in college, while respondents who were educated most recently in primary education (SD / SMP / SMA) were only 20.58%.¹⁷ Education can affect individual health status. The higher the level of education of a person, the easier it is to receive information.¹⁸ Someone who has a higher education will have better knowledge in maintaining and caring for his teeth.¹⁷

A total of 91.1% respondents had a good attitude in maintaining the dental health of their children, while as many as 8.8% had a bad attitude. This is consistent with the research conducted by Oktarina in 2016, that there was a relationship between the level attitudes of parents about dental and oral health with the behaviour of maintaining dental health in children.¹⁸ Education will enhance information and influence the development of individual attitude towards newly introduced values.¹⁸ According to Notoatmodjo (2007), attitudes are the readiness of individuals to act according to their feelings and thoughts based on the values believed. So that attitude can be learned, not taken from birth, not settled it can change. Attitude is the beginning of behaviour that affects individual actions. This can be influenced by parents' initiative in preventing and promoting dental health problems.¹⁹ The higher of the initiative parents in terms of maintaining health, the greater the action that will be formed, the healthier the periodontal tissue the child has.

The results of the study on the aspects of action showed that as many as 91.1% of respondents had good actions in maintaining the health of their teeth and mouth, while as many as 8.8% had bad actions. This is consistent with the research conducted by Nurjannah in 2016 which stated that there was a relationship between actions with the level of dental and oral hygiene.²⁰ This is due to the level of full

awareness of respondents regarding dental and oral health which has an impact on the hygiene status of their teeth and mouth. The higher level of the mother awareness, the more actions produced by the mother so that the child's dental and oral health can be achieved.

After analysing the data, this study showed that there was no significant relationship between mother's behaviour with the periodontal status and periodontal treatment needs in autistic children ($p > 0.05$) both in aspects of knowledge, attitudes, and actions. This is consistent with the study of Oktarina, Tumaji and Roosihermiatie in 2016, that there was no significant relationship between mother's behaviour (knowledge, attitudes, and actions) with children's oral and dental health.¹⁸ This can be caused by knowledge, attitudes, and actions not being the only factors that influence a person's behaviour. In Oktarina's research, et al. (2016), as Blum's theory states that the health status of a person's mouth and society is influenced by four important factors, namely heredity, environment (physical and cultural), behaviour, and health services. Mother's parenting also affects the behavior of mothers in caring for and maintaining the dental health of children. In this study, it was shown that the health of the periodontal tissues of autistic children tended to be healthy so that they did not require large maintenance needs.¹⁸

In this study 44.1% of parents of autistic children used BPJS. Besides mother's income and profession, health insurance also has a significant impact on dental care. The scope of dental insurance services is an important factor in deciding to seek dental care. The use of dental insurance can reduce or even eliminate the cost of dental and oral care which causes high demand for dental and oral health care.²¹ The scope of dental and oral services guarantee by BPJS Health includes: extraction of primary teeth (topical, infiltration), permanent tooth extraction without complications, teeth

scaling (1x a year), etc.²² Ease of dental health service facilities provided, can encourage mothers to take their children to the dentist in terms of care for the child's teeth.

SLB AGCA Centre is a school for children who have special needs, such as autistic children, cerebral palsy, and others. SLB AGCA Centre has an accreditation which is included in the superior schools and is located in elite high socioeconomic locations. The facilities and services provided are good for that the costs required are not small so that not all autistic children can attend the AGCA Centre, only parents who have high socioeconomic status can send their autistic children to AGCA. This can lead to uniformity or homogeneous data so that there is no big difference in the periodontal status that is owned by autistic children, so the majority of autistic children have a healthy periodontal tissue condition with little maintenance needs. In addition, environmental factors for autistic children can also affect the health of the periodontal tissue rather than only from the mother, most of the time the child is used to go to the AGCA Centre daily. Information about maintaining the dental health of children can also come from schools, teachers and the internet from the gadgets played by children.

Oral and dental health of autistic children can be affected due to limited understanding of the importance of maintaining oral health, difficulties in communicating oral health needs, and consumption of anticonvulsant drugs that affect gingival health and fear of oral health procedures, autistic children tend to be hyperactive and have movement uncontrolled pulling muscles. Autistic children also have low salivary secretions which can also cause low oral hygiene in autistic children.²³

In addition, the difficulty in processing sensory stimuli is something that is found in many people with autism. Research by Cermak, Sharon et al. (2015) show that children who were more receptive to sensory responses are important factors that influence their response to receiving dental and oral care. Sensory stimuli can be found in dental practice, such as bright neon lights from the dental chair, touches in and around the oral cavity in dental care actions, the atmosphere and smell of oral care products in the practice of dentists or hospitals, have the potential to have a negative impact in these children, increasing their negative response making it difficult for dentists to provide care. This can cause an increase in stress and anxiety psychology of autistic children, behavioural disorders that cause uncooperative behaviour in receiving dental and oral care.²⁴

Autistic children also tend to have self-injurious behaviours (SIB), which are located in the head-neck area, which can include head banging, face tapping, and gingival picking. In addition, they sometimes also have unusual parafunctional habits including bruxism, stretching their tongue forward, chewing things carelessly like gravel, cigarette butts, or pens that are repeated. Children with autism also tend to have a preference for food, namely sweets and cariogenic foods. This can cause serious problems in the oral and dental health of children with autism.²⁵ This is in contrast to the results obtained in this study that most autistic children have healthy periodontal tissue. This can be caused by the level of awareness of mothers of high autistic children in maintaining the health of their child's teeth and mouth and school environment factors also support children so that they can always maintain and care for the oral health of children.

Conclusion:

Based on the results of research that has been done shows that there was no significant relationship between knowledge, attitudes, and actions of mothers in maintaining dental and oral health with the periodontal status and periodontal treatment needs in autistic children. In this study suggest for further research can be conducted in order to obtain deeper information or can be added with other variables.

Acknowledgement: -

References:

- Please use reference manager, such as Mendeley.
- References in the text should be identified by Arabic Numerals in superscript on the same line as the preceding sentence.
- References should be typed double-spaced on a separate sheet. They should be numbered consecutively in the order by which they are mentioned in the text. They should not be alphabetized.
- All references should provide inclusive page numbers.
- Journal abbreviations should conform with those used in PubMed.
- A maximum of six authors per article can be cited; beyond that, name the first three and add “ et al.”
- The style/punctuation approved by Acta Medica conforms to that recommended by the ICMJE, which is the ANSI standard style used by the NLM, and should follow the format of the examples shown below:

Journal

Putala J, Metso AJ, Metso TM, et al. Analysis of 1008 consecutive patients aged 15 to 49 with first-ever ischemic stroke. *Stroke*. 2009;40(4):1195-203.

Website/Internet

Centers for Disease Control and Prevention, Falls among older adults: summary of research findings [Online]. 2005 [cited 2006 Jan]. Available from <http://www.cdc.gov/ncipc/pubres/toolkit/SummaryOfFalls.htm>.

Book

Dumitru D, Amato AA, Zwarts MJ. Nerve conduction studies. In: Dumitru D, Amato AA, Zwarts MJ, eds. *Electrodiagnostic medicine*, 2nd ed. Philadelphia: Hanley and Belfus, Inc; 2002. pp. 159–217.

Tables and Figures:

Tables:

1. Cite all tables consecutively in the text and number them accordingly. Create tables preferably using a spreadsheet program such as MS Excel with one table per worksheet. Tables should not be saved as image files. The content of tables should include a table number (Arabic) and title above the table, and explanatory notes and legends as well as definitions of abbreviations used below. Recommended font is Arial Narrow size 8.
2. Each table must be self-explanatory, being a supplement rather than a duplicate of information in the text. The use of too many tables is discouraged.

Tabel:

1. Kutip semua tabel secara berurutan dalam teks dan beri nomor sesuai. Buat tabel lebih disukai menggunakan program spreadsheet seperti MS Excel dengan satu tabel per lembar kerja. Tabel tidak boleh disimpan sebagai file gambar. Isi tabel harus mencakup nomor tabel (Arab) dan judul di atas tabel, dan catatan penjelasan dan legenda serta definisi singkatan yang digunakan di bawah ini. Font yang disarankan adalah Arial Narrow size 8.
2. Setiap tabel harus jelas, menjadi pelengkap daripada duplikat informasi dalam teks. Penggunaan terlalu banyak tabel tidak disarankan.

Figures:

FIGURES AND GRAPHS

1. Figures or graphs should be identified by Arabic Numeral/s with titles and explanations underneath. The numbers should correspond to the order in which the figures/graphs occur in the text. It is recommended that figures/graphs also be submitted as image files (preferably as .jpeg or .gif files) of high resolution.
2. All identifying data of the subject/s or patient/s under study such as name, case numbers, etc., particularly in case reports, should be removed.

ILLUSTRATIONS AND PHOTOGRAPHS

1. Should at least be 800 x 600 dpi.
2. Computer-generated illustrations which are not suited for reproduction should be professionally redrawn or printed on good quality laser printers. Photocopies are not acceptable.
3. All lettering for illustration should be done professionally and should be of adequate size to retain even after size reduction.
4. For photomicrographs, the stain used (ex. H & E) and magnification (ex. X400) should be included in the description

Reply template Acta medica Rev 1 MG RD

**JOINT SCIENTIFIC MEETINGS in SPECIAL CARE DENTISTRY 2019
FACULTY OF DENTAL MEDICINE, UNIVERSITAS AIRLANGGA
SCIENTIFIC ARTICLE COMMITTEE**

Reply to the reviewers' comments

Title: Correlation Between Mother's Behaviour with Periodontal Status and Periodontal Treatment Needs in Autistic Children

Author:

Reviewer Number (1/2)	Original comments of the reviewer	Reply by the author(s)	Changes done on page number and line number
2	Please describe more about the connection between mother's behavior and autism child actions/making a choice.	Therefore in maintaining the dental health of autistic children influenced by parents teaching skills and habits in autistic children's life. From previous study it has been found that there were about 17,4 % autistic children in Saudi Arabian suffering from bleeding from bleeding at gingiva.	Page number : 1 Line number : 9
2	Please add the problem statement. Is periodontal disease the main problem in autistic children?	Periodontal disease is often found in autistic children. This study analyses How is the relationship between mother's behaviour with periodontal status and periodontal treatment needs of autistic children	Page number : 1 Line number : 11
2	Does this mean "34 pairs of autistic children and mothers"?	Yes, with a total sample 34 pairs of autistic children and their mothers.	Page number : 1 Line number : 16
2	Mention the statistical analysis conducted in this study	Analytical observational study with cross-sectional approach	Page number : 1 Line number : 15
2	Briefly explain about this kind of questionnaire.	answering the HU-DBI questionnaire which is consists of knowledge, attitude, and mother's action	Page number : 1 Line number : 17

Reply template Acta medica Rev 1 MG RD

2	Reference?	R. Arfiriana, and F. Dieny. Relationship to casein-free gluten-free diet score with autism behaviour score. Journal of Nutrition College. 2014;3(1): 34–42.	Page number : 2 Line number : 40
1	What do you mean by “if”	When the birth rate in Indonesia is 6 million per year, the number of autistic children in Indonesia increases by 0.15% or 6,900 children per year	Page number : 2 Line number : 38
1	You mean cervical area?	No, Gingivitis is an inflammation that only involves gingival tissue with no loss of attachment or bone.	Page number : 2 Line number : 44
1	Meaning?	Due to the main characteristics of autism which had problems of social interaction with others, communication’s disorders, and Cognitive instability, they cannot maintain their oral hygiene individually	Page number : 2 Line number : 46
1	Meaning?	Certain autistic children’s can maintain their good oral hygiene behind the limitations they have.	Page number : 2 Line number : 51
1	Whose behaviour?	Autistic Children’s behaviour to maintaining the dental health such as tooth brushing habit and not afraid to go to dentist can be influenced by families/parents where parents teach skills and habits in of autistic children’s life everyday	Page number : 2 Line number : 53
1	Unclear sentence	Mother’s behaviour regarding to oral health can be used to predict the status or condition of their child’s oral health.	Page number : 2 Line number : 59
1	Even so, the reason of study should be something scientific based, and not because of “no data”	There were still a few studies that describe the relationship between mother’s behaviour with periodontal status and periodontal treatment needs in autistic	Page number : 3 Line number : 63

Commented [mp1]: Ternyata ini punya Rifmie Arfiriana Pratiwi, Fillah Fithra Dieny

Tetapi dalam mensitasinya mereka sudah menulis... how to cite this jurnal

Namanya jadi seperti tersebut .

Sdh saya ganti di reference

Commented [mp2]: Ridha

Boleh tolong diberitahu certain itu yang seperti apa bentk autisnya. Saya coba browsing di jurnal reff nya. Jika tidak salah punya dokter yetty nonong , saya tidak bisa menemukan certain yang bagaimana. Bisa tolong digarisi di jurnalnya. Supaya bisa saya bantu utk memahami maksud jurnalnya.

Reply template Acta medica Rev 1 MG RD

		children, for this reason the researchers wanted to examine the relationship between mother's behaviour towards the periodontal status in children with autism and their periodontal treatment needs	
1	= Sekolah Luar Biasa? Please translate into english	The study was conducted at Special Education AGCA Centre Surabaya	Page number : 3 Line number : 68
2	It looks like your study was experimenting on human, do this study approved by ethical committee? Did you have an informed consent signed by the respondents or their guardian?	The population in this study were all children that diagnosed with autism in the Special Education AGCA Centre, amounting to 34 people along with parents of autistic children according to the criteria of the research subject which is among them previously had signed an informed consent form and had been approved by ethical committee	Page number : 3 Line number : 71
2	Give the reference. Is there any previous study used this kind of questionnaire?	the HU-DBI modification questionnaire (Hiroshima University - Dental Behavioural Inventory) regarding the behaviour of maintaining oral health by mothers as parents of autistic children and examining the status of periodontal tissue using the CPITN Index.	Page number : 3 Line number : 77
1	Of who?	the status periodontal tissue of autistic children using the CPITN Index	Page number : 3 Line number : 80
1	Periodontal tissue	the status periodontal of autistic children was examined using the CPITN Index.	Page number : 3 Line number : 82
1	???	Data collected from questionnaire in the form of general characteristics of respondents and the relationship between mother's behaviour with health of periodontal tissue and periodontal care needs of autistic children	Page number : 4 Line number : 90

Reply template Acta medica Rev 1 MG RD

1	Meaning	Then followed by 8 children (23.5%) needed an increase in oral hygiene while 7 children (20.5%) needed dental health education and professional scaling.	Page number : 4 Line number : 109
1	Rewrite this part	Furthermore, the attitude aspect has a low value of 8.8% and a high value is 91.1%. The last aspect is the aspect of action that is low value of 47%, high value is 53%.	Page number : 4 Line number : 117

**ACTA MEDICA PHILIPPINA
AUTHOR DECLARATION**

I/We have been sufficiently involved in this work to take public responsibility for its validity and final presentation as original publication.

I/We declare that this manuscript has not been published and is not under simultaneous consideration for publication elsewhere.

I/We declare that this manuscript has been subjected to ethics review by Universitas Airlangga Faculty of Dental Medicine Health Research Ethical Clearance Commission (Number: 192/HRECC.FODM/VII/2018), July 31st 2019 and has been endorsed favourably by the said board or committee.


I/We declare that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

I/We declare that the manuscript has been read and approved by all authors and that the order of authors listed in the manuscript has been approved by all of us.


I/We confer all copyright ownership to Acta Medica Philippina in the event that this manuscript is published in this journal.

I/We understand that the Corresponding Author is responsible for communicating with the other authors about the status of submitted manuscripts, submission of revisions and final approval of proofs.


Signed by all authors as follows:



Mega Moeharyono Puteri



Tania Saskianti



Ridha Rizki Ananda

To,

The Editor of Acta Medica Philippina

Sub: Submission of Manuscript for publication

Dear Sir,

We intend to publish an article entitled "Correlation Between Mother's Behaviour with Periodontal Status and Periodontal Treatment Needs in Autistic Children" in your esteemed journal as an Original Article.

I/We have been sufficiently involved in this work to take public responsibility for its validity and final presentation as an original publication. The article had written/informed consent for publication from the involved subject/s, had conformed with ethical standards, and/or had been reviewed by the appropriate ethics committee.

This paper has not been published and is not under simultaneous consideration for publication elsewhere. This paper has been subjected to ethics review by (Universitas Airlangga Faculty of Dental Medicine Health Research Ethical Clearance Commission, July 31st 2018) and has been endorsed favorably by the said board or committee. I/We hereby confer all copyright ownership/s to the Acta Medica Philippina in the event that this work is published in this journal.

We hereby transfer, assign, or otherwise convey all copyright ownership, including any and all rights incidental thereto, exclusively to the journal, in the event that such work is published by the journal.

Thanking you,

Yours' sincerely,



(Mega Moeharyono Puteri)

Corresponding contributor:

Name : Mega Moeharyono Puteri
Affiliation : Faculty of Dental Medicine Universitas Airlangga
Address : Jl. Mayjen Prof Dr Moestopo No 47 Surabaya, Indonesia
Phone Number : +62 31 5030255
Fax : +62 31 5026288
E-mail : mega-m-p@fkg.unair.ac.id