

---

## Special Care in Dentistry - Manuscript ID 3088.R2

1 message

---

**Special Care in Dentistry Editorial Office** <onbehalf@manuscriptcentral.com>

Sun, Nov 3, 2019 at 5:12  
PM

Reply-To: scdeditorial@wiley.com

To: mega-m-p@fkg.unair.ac.id

Cc: mega-m-p@fkg.unair.ac.id, fadila.kemala.dwi-2014@fkg.unair.ac.id, teguh-b-w@fkg.unair.ac.id

03-Nov-2019

Dear Miss Puteri:

Your revised manuscript entitled "Oral Health Behavior and its Association with the Caries Index in Visually-Impaired Children" by Puteri, Mega; Ruslan, Fadila; Wibowo, Teguh, has been successfully submitted online and is presently being given full consideration for publication in Special Care in Dentistry.

Co-authors: Please contact the Editorial Office as soon as possible if you disagree with being listed as a co-author for this manuscript.

Your manuscript ID is 3088.R2.

For your reference: the manuscript number of the PREVIOUS manuscript version is: 3088.R1.

Please mention the above manuscript ID in all future correspondence or when calling the office for questions. If there are any changes in your street address or e-mail address, please log in to ScholarOne Manuscripts at <https://mc.manuscriptcentral.com/scid> and edit your user information as appropriate.

You can also view the status of your manuscript at any time by checking your Author Center after logging in to <https://mc.manuscriptcentral.com/scid>.

Thank you for submitting your manuscript to Special Care in Dentistry.

Sincerely,  
Special Care in Dentistry Editorial Office



[Home](#)

[Author](#)

[Review](#)

## Author Dashboard

**3 Manuscripts with Decisions**

[Legacy Instructions](#)

[5 Most Recent E-mails](#)

[Before You Submit](#)

## Manuscripts with Decisions

ACTION	STATUS	ID	TITLE	SUBMITTED	DECISIONED
	EIC: Diz Dios, Pedro ADM: Varghese, Sindhu <ul style="list-style-type: none"> <li>Accept (05-Nov-2019)</li> </ul> <i>Archiving completed on 21-Jan-2020</i> <a href="#">view decision letter</a> <a href="#">Contact Journal</a>	3088.R2	Oral Health Behavior and its Association with the Caries Index in Visually-Impaired Children <i>Files Archived</i>	03-Nov-2019	05-Nov-2019
a revision has been submitted (3088.R2)	EIC: Diz Dios, Pedro ADM: Varghese, Sindhu <ul style="list-style-type: none"> <li>Minor Revision (29-Sep-2019)</li> <li>a revision has been submitted</li> </ul>	3088.R1	Oral Health Behavior and its Association with the Caries Index in Visually Impaired Children <i>Files Archived</i>	17-Sep-2019	29-Sep-2019






ACTION	STATUS	ID	TITLE	SUBMITTED	DECISIONED
	<p><i>Archiving completed on 21-Jan-2020</i></p> <p><a href="#">view decision letter</a></p> <p><a href="#">✉ Contact Journal</a></p>				
<p>a revision has been submitted (3088.R1)</p>	<p>EIC: Diz Dios, Pedro ADM: Varghese, Sindhu</p> <ul style="list-style-type: none"> <li>▪ Major Revision (21-Jul-2019)</li> <li>▪ a revision has been submitted</li> </ul>	<p>3088</p>	<p>Oral Health Behavior and its Association with the Caries Index in Visually Impaired Children</p> <p><i>Files Archived</i> </p>	<p>20-Jun-2019</p>	<p>21-Jul-2019</p>
	<p><i>Archiving completed on 21-Jan-2020</i></p> <p><a href="#">view decision letter</a></p> <p><a href="#">✉ Contact Journal</a></p>				

© Clarivate | © ScholarOne, Inc., 2023. All Rights Reserved.

ScholarOne Manuscripts and ScholarOne are registered trademarks of ScholarOne, Inc.

ScholarOne Manuscripts Patents #7,257,767 and #7,263,655.

 [@Clarivate for Academia & Government](#) |  [System Requirements](#) |  [Privacy Statement](#) |  [Terms of Use](#)

# Oral Health Behavior and its Association with the Caries Index in Visually-Impaired Children

## ABSTRACT

**AIM:** To assess the correlation between oral health behavior and the Caries Index (CI) among visually-impaired children.

**METHODS AND RESULTS:** An analytical cross-sectional study was conducted from September to October 2017 in four schools in Surabaya, Indonesia. A total of 34 visually-impaired and 34 non-visually-impaired children were selected and included in the study using a total sampling method. The oral health knowledge of, attitude towards and practice of subjects were assessed by means of questionnaire. The author subsequently performed an intraoral examination of all subjects. The presence and extent of dental caries were recorded using dmft and DMFT scoring and categorized according to WHO classification. Visually-impaired children were shown to have a low CI of 1.5. A Spearman's statistical test showed that, while knowledge significantly affected the CI of visually-impaired children ( $p < 0.05$ ), attitude ( $p = 0.98$ ) or practice ( $p = 0.42$ ) did not.

**CONCLUSION:** A correlation clearly exists between oral health knowledge relating to CI in visually-impaired children. However, there appeared to be no relationship between attitudes towards

and actions in maintaining dental health through a CI in visually-impaired children.

**Keywords:** Caries; oral health behavior; visually-impaired children.

## **INTRODUCTION**

Vision loss, also known as visual impairment, is a functional limitation which renders individuals unable to obtain information from their surroundings by means of their sight.<sup>1</sup> According to the WHO (2010), approximately 285 million people suffer from this medical condition: 39 million people experienced total vision loss, while the remaining 246 million suffered impaired vision. Of these cases, 19 million cases affected children.<sup>2</sup>

Children with visual impairment demonstrate limited ability to perform everyday activities, one of them being an appropriate personal oral hygiene regime.<sup>3</sup> This is due to the fact that it is more challenging for people with defective sight to gain access to dental care.<sup>4</sup> Consequently, children with this disability run greater risk of developing dental caries due to the difficulties they experience in cleaning plaque and recognizing the early symptoms of tooth decay, such as discoloration (white spots).<sup>5</sup>

Dental caries, a condition in which the localized tooth surface consisting of the enamel, dentin and cementum is destroyed, is known to be caused by multiple risk factors, including the host, agent (microorganisms), substrate and time. Moreover, it is also widely accepted that the behavior, environment and access to

health care on the part of individuals all exacerbate the risk of developing caries.<sup>6,7</sup>

The dental and oral health of an individual can be indicative of their knowledge of, attitude towards and practice in maintaining oral hygiene which can be classified into three types of behavior: health maintenance, seeking access to healthcare and maintaining a salubrious environment.<sup>8</sup>

Despite the improving efforts of dental health professionals to enhance public oral health, according to Gokhale *et al.*, the incidence of dental caries continues to increase.<sup>9</sup> In a study conducted by John *et al.*, the caries index (CI) of children with visual impairment in Chennai, India was high at 4.5.<sup>10</sup> On the other hand, Reddy *et al.*, reported a low CI of 1.1 in visually impaired children. In Medan, Indonesia, children with visual impairment was reported by Girsang to have an even higher CI of 5.5.<sup>11</sup> Based on these studies, it remains debatable whether a correlation between quality of sight and CI exists. Therefore, further information regarding the association between these two variables is required. This research investigates the correlation between oral health behavior and CI in children with compromised vision.

## **METHODS**

This research employed an observational cross-sectional study method approved by the Ethical Research Committee of the Faculty of Dental Medicine, Universitas XXX (192/HRECC.FODM/IX/2017).

This research was conducted between September and October 2017 in four Surabaya-based schools: *Sekolah Dasar Luar Biasa A* (SDLB-A), *Sekolah Menengah Pertama Luar Biasa A* (SMPLB-A), *Sekolah Dasar Negeri Pacarkeling IV* and *Sekolah Menengah Pertama Negeri 29*. A total sample of 68 subjects (34 visually-impaired, 34 non-visually-impaired) constituted the research population. The inclusion criteria for members of the visually-impaired group were as follows: children aged 6-16 years old presenting total vision loss who were willing and cooperative participants in the research. Meanwhile, children aged between 6 and 16 years with no vision loss who had agreed to participate in the research and proved cooperative were eligible for inclusion in the non-visually-impaired group. Informed consent was obtained from children's legal guardians for being included in the study.

Two primary forms of data, questionnaires and intraoral examination, were employed during the research. The questionnaire comprised three sections: knowledge of, attitude towards, and practice in maintaining oral hygiene (see Appendix)



and had previously been tested for validity (Pearson sig. values for all items <0.05) and reliability (Cronbach's Alpha = 0.705).<sup>12</sup> The questionnaire was completed by interviewing the subject to elicit his/her knowledge, attitudes and practice regarding oral hygiene. A score of 1 was assigned to every correct answer, while 0 was assigned to ones which were incorrect. Those who scored higher than 4.6 on the 'knowledge' and 'attitude' categories were classified as high/good, while a score of <4.6 was deemed to be low. Meanwhile, in the 'oral hygiene practice' section, a score of >5.4 was classified as good and <5.4 as poor.

Intraoral examinations, performed on the subjects using a dental mirror and sterile probe, were undertaken to assess the severity of dental caries, the results of which were recorded in a Caries Index (CI) form. The computed results were classified in terms of severity according to WHO categories: very low (CI <1.2), low (CI 1.2-2.6), moderate (2.7-4.4), high (4.5-6.5) and very high (>6.5).

A 21<sup>st</sup> version SPSS was used to analyze the data collected by means of both questionnaires and intraoral examinations. A P-value of <0.05 was considered statistically significant.

## RESULTS

The mean CI of the visually-impaired group members can be seen to be low at 1.5 and observed to be lower among its older members, as shown in Table 1. A low incidence of dental caries (CI = 2.0) was reported among subjects in the 6-12 years age group (n=17), while the remaining 17 subjects in the 13-16 years age group possessed a CI of 1.1 indicating very low severity. A Mann-Whitney test found the difference in CI between the two age groups to be statistically insignificant ( $p$ -value = 0.12).

The non-visually-impaired group was observed to have low CI at 1.7. A Mann-Whitney test found that the difference in CI between visually-impaired and non-visually-impaired groups was not statistically significant ( $p$ -value = 0.71). Therefore, there was no difference in CI between the two groups.

A summary of the knowledge of, attitude towards and practice in maintaining the oral health of visually-impaired subjects is reported in Table 2. Of the 34 visually-impaired subjects, 82.4% (n=28) possessed considerable knowledge regarding dental and oral health, whilst that of the remaining 17.6% was considered poor. 23 subjects (67.6%) possessed a positive attitude towards dental and oral health. Finally, the practice in maintaining oral health was good in 58.8% of the subjects (n=20).

The relationship between the three domains, such as knowledge of, attitude towards and practice in maintaining oral health and CI in **visually-impaired** subjects was analyzed using a Spearman's Rank Test. The correlation between knowledge of oral and dental health and CI was statistically significant ( $p$ -value  $< 0.05$ ). The association between attitude towards and practice in maintaining oral health and CI were shown to be statistically insignificant ( $p$ -value=0.98 and 0.42, respectively).

## DISCUSSION

In this study, the dental CI among **visually-impaired** children were reported to be of low severity, a finding matching that of a study conducted by Reddy *et al.* (2011) in which a low category CI was recorded in **visually-impaired** children.<sup>13</sup> This study also found a decrease in CI within older age groups, although the relationship was not statistically significant mirroring the findings of Kiswaluyo (2013) that a decline in CI was observable in older individuals due to the fact that they demonstrate greater awareness of dental and oral health care.<sup>14</sup> Despite this, a study published by Singh *et al.* (2017) concluded that no significant difference in CI existed in individuals between the ages of 9 - 15 because of their capacity to memorize, synthesize and apply the information retrieved being equal.<sup>15</sup>

This comparison of CI in **visually-impaired** and **non-visually-impaired** children showed no statistically significant difference, a result confirming that of a study by Herwis *et al.* (2013). Since the two groups demonstrated similar habits with regard to maintaining oral health, such as the level of sugar consumption and frequency of brushing, no significant difference in CI between groups was observed.<sup>16</sup>

A total of 82.4% of **visually-impaired** subjects possessed good knowledge regarding dental and oral health maintenance. This was possibly due to their subjects having been formally educated in special needs schools under the management of *Yayasan Pendidikan Anak Buta* (YPAB) Surabaya which received health advice from the community health clinics (*Puskesmas*). Moreover, nowadays, information about dental and oral health is easily accessible (e.g. via television or advice from dental health professionals).<sup>17</sup>

67.6% of the participating **visually-impaired** subjects demonstrated a positive attitude towards maintaining dental and oral health. The paper written by Notoatmodjo (2010) argued that the cognitive ability of an individual can influence his/her attitude. Those aware of the importance and means of maintaining dental and oral hygiene are more likely to take appropriate action.<sup>8</sup>

Of the **visually-impaired** group members, 58.8% followed good practice in maintaining oral health. According to Notoatmodjo (2010), this is easier for an individual when access to a dentist, dental nurse, and healthcare facilities is available.<sup>8</sup>

The correlation between the knowledge that a subject possesses regarding the maintenance of oral health and CI is statistically significant. This finding matches that of the study by Marimbun

*et al.* (2016) which posited the existence of a significant relationship between knowledge of oral health and CI in **visually-impaired** subjects. A satisfactory understanding of dental and oral health would positively influence the attitude and actions of an individual with regard to oral health maintenance. Appropriate knowledge of dental and oral cleansing procedures represents the basis for maintaining oral hygiene.<sup>18</sup>

The attitude towards and practice of maintaining dental health were shown not to have a significant relationship with the CI. A study in 2009 also reported to have found no significant correlation between attitude and the dental CI.<sup>19</sup> In keeping with the results contained in this paper, Ahmad *et al.* found that there was no causal link between oral hygiene maintenance and the severity of dental caries.<sup>20</sup> The absence of these relationships may be attributable to the presence of enabling factors.

Enabling factors, for instance parenting practice and access to healthcare facilities, could influence the attitude of an individual. The majority of **visually-impaired** children studying at YPAB were observed to be very dependent on their parents or caregivers in performing daily activities. In addition to the three domains focused on in this study, deficient motor skills can also contribute to difficulties in maintaining dental

health.<sup>21</sup> Therefore, children who are **visually-impaired** require the support of others. Parents or caregivers play a vital role in promoting children's dental and oral health, such as motivating and assisting them to follow effective oral and dental hygiene practices and ensuring that they attend regular dental health check-ups. A good CI and effective dental care practice could be the result of accessible healthcare facilities, such as a Puskesmas. According to Kencana (2014), health education or counseling could positively **influence** the behavior of an individual, including that relating to oral health.<sup>22</sup>

Based on the result of the study, a firm conclusion regarding the relationship between knowledge of oral health and the dental CI of **visually-impaired** children can be drawn. However, there appeared to be no relationship between attitudes towards and actions in maintaining dental health through a CI in **visually-impaired children**. This was **possibly** due to the presence of enabling factors such as access to healthcare and parenting methods. It is expected that visually impaired children have a better oral health behavior by promotion of oral health in community-based or group activities among visually impaired children so that the caries index will be decreased. Further studies are recommended to include large group of visually

impaired children and should focus on the enabling factors and  
reinforcing factor, as this study focused on the predisposing  
factors only.



## REFERENCES

1. Permana H, Indahyani D, Yustisia Y. Kelarutan kalsium email pada saliva penderita tunanetra. *Dentofasial*. 2014;13(3):150-4.
2. World Oral Health Organization. Visual impairment and blindness 2014 [Cited 2017 Jan 1]. Available from: <http://www.who.int/blindness/en/>
3. Prashanth S, Bhatnagar S, Das UM, Gopu H . Oral health knowledge, practice, oral hygiene status, and dental caries prevalence among visually-impaired children in Bangalore. *J Indian Soc Pedod Prev Dent*. 2011;29(2):102-5.
4. Ozdemir-Ozenen D, Sungurtekin E, Cildir S, Sandali N. A Comparison of the oral health status of children who are blind and children who are sighted in Istanbul. *JVIB*. 2012;106(6):362-7.
5. Samnieng P, Seehaumpai P, Wichachai S, Yosookh P. Oral health status and treatment need of visual impairment in Phitsanuloke, Thailand. *J Dent Indones*. 2014;21(2):63-7.
6. Mariati N. Pencegahan dan perawatan karies rampan. *Jurnal Biomedik*. 2015;7(1):23-8.
7. Miftakhun N, Salinkun, Sunarjo L, Mardiati E. Faktor eksternal penyebab terjadinya karies gigi pada anak pra

- sekolah di PAUD Strawberry RW 03 kelurahan Bangetayu Wetan kota Semarang tahun 2016. Jurnal Kesehatan Gigi. 2016;3(2):27-34.
8. Notoatmodjo S. Ilmu perilaku kesehatan. Edisi Revisi. Jakarta: PT. Rineka Cipta; 2010. pp 20-32
  9. Gokhale N, Sivakumar N, Svsg N, Abinash M. Dental caries and body mass index in children of Nellore. J Orofac Sci. 2010;2(2):4-6
  10. John J, Daniel B, Paneerselvam D, Rajendran G. Prevalence dental caries, oral hygiene knowledge, status, and practices among visually-impaired individuals in Chennai, Tamil Nadu. Int J Dent. 2017;2017(9419648):1-6.
  11. Girsang EN. Perbandingan oral hygiene dan karies gigi pada anak tunanetra dan tidak tunanetra usia 12 dan 15 tahun di Medan. Medan: Universitas Sumatera Utara. 2003. pp. 27
  12. Suharsimi A. Manajemen penelitian. Edisi revisi 6. Jakarta: PT. Rineka. 2009. pp14.
  13. Reddy K, Sharma A. Prevalence of oral health status in visually-impaired children. J Indian Soc Pedod Prev Dent. 2011;29(1):15-7
  14. Kiswaluyo. Hubungan usia dan jenis kelamin dengan kejadian karies gigi siswa sekolah dasar Summersari dan Puger

- kabupaten Jember. *Prodentia Journal of Dentistry*.  
2013;1(2):47-54.
15. Singh A, Dhawan P, Gaurav V, Rastogi P, Singh S. Assessment of oral health-related quality of life in 9-15 year old children with visual impairment in Uttarakhand, India. *Dent Res J*. 2017;14(1):43-9.
  16. Herwis K, Ali A, Elturki, Khamis. Oral health status of visually-impaired and sighted groups of children In Benghazi: A comparative study. *Libyan Dental Journal*. 2013;3:1-5.
  17. Sharda AJ, Shetty S. Relationship of peridontal status and dental caries with oral health knowledge, attitude, and behaviour among professional students in India. *Int J Oral Sci*. 2009;1(4):196-206.
  18. Marimbun BE, Mintjelungan CN, Pangemanan DHC. Hubungan tingkat pengetahuan tentang kesehatan gigi dan mulut dengan status karies gigi pada penyandang tunanetra. *Jurnal e-GiGi*. 2016;4(2):177-82.
  19. Warni Linda. Hubungan perilaku murid SD kelas V dan VI pada kesehatan gigi dan mulut terhadap status karies gigi di wilayah kecamatan Delitua kabupaten Deli Serdang. Thesis Universitas Sumatera Utara; 2009. p 20-25

20. Ahmad MS, Jindal MK, Khain S, Hshmi H. Oral health knowledge, practice , oral hygiene status and dental caries prevalence among visually-impaired students in residential institute of Aligarh. *J Dent Oral Hyg.* 2009;1(2):22-6.
21. Suresan V, Das D, Jnaneswar A, Jha K, Kumar G, Subramaniam GB. Assessment of dental caries, oral hygiene status, traumatic dental injuries and provision of basic oral health care among visually-impaired children of Eastern Odisha. *J Indian Soc Pedod Prev Dent.* 2017;35(4):284-90.
22. Kencana IGS. Peranan perawat gigi dalam pemeliharaan kesehatan gigi dan mulut pada anak berkebutuhan khusus (disabled children). *Jurnal Kesehatan Gigi.* 2014;2(2):260-5.

**TABLES**

**Table 1.** CI of visually-impaired group and non-visually-impaired group

	<b>Age (Year)</b>	<b>N</b>	<b>CI score (Category)</b>	<b>Mean CI score (Category)</b>	<b>Sig. (P-value)</b>
<b>Visually- impaired</b>	6-12	17	2 (Low)		
	13-16	17	1.1 (Very low)	1.5 (Low)	0.12
					0.71
<b>Non- visually- impaired</b>	6-16	34	-	1.7 (Low)	-

**Table 2.** Correlation between knowledge of, attitude towards and practice in maintaining dental and oral health as indicated by the CI of visually-impaired group members

Category	Domain					
	Knowledge		Attitude		Practice	
	High/Good	Low/Poor	High/Good	Low/Poor	High/Good	Low/Poor
	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)
	28	6	23	11	20	14
	(82.4%)	(17.6%)	(67.6%)	(32.4%)	(58.8%)	(41.2%)
Spearman's Rank Test (P-value)						
CI of						
visually-						
impaired		0.03*		0.98		0.42
group						

\*statistically significant