

# Coumarins and Carbazole Alkaloid from *Clausena excavata* Roots and Investigation of their $\alpha$ - glucosidase Inhibitory Activity

*by Nanik Siti Aminah*

---

**Submission date:** 27-Aug-2021 05:01PM (UTC+0800)

**Submission ID:** 1636750937

**File name:** C07.\_Artikel\_COUMARINE\_\_CARBAZOLE\_TJNPR\_2020.pdf (980.06K)

**Word count:** 3182

**Character count:** 15841

**Coumarins and Carbazole Alkaloid from *Clausena excavata* Roots and Investigation of their  $\alpha$ -glucosidase Inhibitory Activity**

13

Tin M. Thant<sup>1,2</sup>, Nanik S. Aminah<sup>1\*</sup>, Alfinda N. Kristanti<sup>1</sup>, Rico Ramadhan<sup>1</sup>, Iffa H. Hasna<sup>3</sup>, Hnin T. Aung<sup>4</sup>, Yoshiaki Takaya<sup>5</sup><sup>1</sup>Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Komplek Kampus C UNAIR, Jl. Mulyorejo, Surabaya, Indonesia<sup>2</sup>Department of Chemistry, Mandalay Degree College, Mandalay, Myanmar<sup>3</sup>Faculty of Medicine, Kampus A Universitas Airlangga Surabaya, St. Mayjen. Prof. Dr. Moestopo 47, Surabaya 60132, Indonesia<sup>4</sup>Department of Chemistry, Mandalay University, Mandalay, Myanmar<sup>5</sup>Faculty of Pharmacy, Meijo University, 150 Yagotoyama, Tempaku, Nagoya, 468-8503 Japan

20

## ARTICLE INFO

## ABSTRACT

## Article history:

Received 13 June 2020

Revised 06 August 2020

Accepted 19 August 2020

Published online 28 August 2020

**Copyright:** © 2020 Thant *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Clausena excavata* Burm.f. has long been associated with medicinal benefits in folk medicine. This study isolated three known compounds which are two coumarins, umbelliferone (**1**), clausenidin (**2**) and one carbazole alkaloid, (2-hydroxy-3-formyl-7-methoxycarbazole (**3**) from the root of the plant. The compounds were characterized using their <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, COSY, HSQC and HMBC spectra and confirmed by literature data. The  $\alpha$ -glucosidase inhibitory activity of the extract, fractions, compounds **2** and **3** were evaluated. The methanol extract exhibited higher inhibition against maltase  $\alpha$ -glucosidase with IC<sub>50</sub> value 0.020 mg/mL than the n-hexane fraction which IC<sub>50</sub> value was 0.041 mg/mL.

**Keywords:** *Clausena excavata*, Umbelliferone, Clausenidin, 2-hydroxy-3-formyl-7-methoxycarbazole,  $\alpha$ -glucosidase, Antidiabetic.

36

## Introduction

Diabetes mellitus (DM) is a chronic metabolic disease that has now been recognized as one of the most challenging health problems<sup>1,2</sup>. Main symptom of DM is hyperglycemia (high blood glucose levels) which often time leads to severe diabetic complications over time, including coronary artery disease, stroke, peripheral artery disease, retinopathy, nephropathy, and neuropathy.<sup>3,4</sup> To treat patients with DM is essential to control the level of postprandial blood glucose. This is found to be helpful to prevent any further complications caused by DM.  $\alpha$ -Glucosidase inhibitors are a class of drugs with the ability to treat DM patients by preventing the carbohydrates digestion and consequently deferring glucose absorption as well as suppressing the postprandial hyperglycemia. Therefore, the inhibitors of  $\alpha$ -glucosidase are usually suggested as vital agents to treat patients with DM. They can be used alone or in combination with other antidiabetic agents for the treatment of type 2 diabetes mellitus (T2DM) and can also be used for patients with type 1 diabetes mellitus (T1DM).<sup>5</sup>

Alpha-Glucosidase ( $\alpha$ -Glucosidase) inhibitors, which are classified as third generation oral hypoglycemic agents have shown vital inhibition of postprandial blood glucose in T2DM patients, and have been recommended as a first line therapy by the International Diabetes Federation and the American Association of Clinical Endocrinologists. Hence,  $\alpha$ -glucosidase is considered a safe natural inhibitor with little or no side effects which has gradually attracts broad interests for use as a main ingredient in the management of T2DM around the world.<sup>5</sup>

\*Corresponding author. E mail: [nanik-s-a@fst.unair.ac.id](mailto:nanik-s-a@fst.unair.ac.id)  
Tel: +62-31-5936501

**Citation:** Thant TM, Aminah NS, Kristanti AN, Ramadhan R, Hasna IH, Aung HT, Takaya Y. Coumarins and Carbazole Alkaloid from *Clausena excavata* Roots and Investigation of their  $\alpha$ -glucosidase Inhibitory Activity. Trop J Nat Prod Res. 2020; 4(8):334-337. [doi.org/10.26538/tjnpr/v4i8.2](https://doi.org/10.26538/tjnpr/v4i8.2)

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

Alpha-Glucosidase serves as enzymes that catalyze the exo-hydrolysis of 1,4- $\alpha$ -glycosidic linkages which lead to the release of  $\alpha$ -glucose. Inhibition of this enzyme has the ability to reduce carbohydrates absorption from the digestive tract. As a result, it reduces postprandial glucose level.<sup>6</sup>

*Clausena excavata* is a plant of the family Rutaceae. It typically grows in India, China and Southeast Asia.<sup>7,8</sup> The plant is categorized as a wild shrub and several of its components are used in traditional medicine.<sup>7</sup> Previous studies indicate that *C. excavata* possesses some pharmacological properties.<sup>9</sup> Previously reported constituents are the carbazole alkaloids, coumarins, flavonoids and limonoids.<sup>9-13</sup> In the course of continuing the search for  $\alpha$ -glucosidase inhibitors from natural sources we hereby report on the evaluation of the  $\alpha$ -glucosidase inhibitory activity of the extract, fractions and isolated compounds from *C. excavata*. This is the first report of  $\alpha$ -glucosidase inhibitory activity of isolated compounds and fractions from *C. excavata* roots.

## Materials and Methods

## General experiment procedure

1D and 2D NMR experiments were performed on a Bruker 600 MHz (<sup>1</sup>H) and 151 MHz (<sup>13</sup>C) spectrometer in solvent CDCl<sub>3</sub>. The chemical shifts were reported in ppm with referenced to tetramethyl silane as internal standard. Vacuum liquid chromatography and analytical preparative thin layer chromatography were performed on Kieselgel 60 (F<sub>254</sub>, Merck). Merck Kieselgel 60 (40-63  $\mu$ m) was employed for column chromatography. Spectrophotometric measurements for the  $\alpha$ -glucosidase inhibition were recorded on a TECAN Infinite 50 microplate reader spectrophotometer

## Plant material

The roots of *C. excavata* were collected in Naypyitaw region, Myanmar in July 2016. The plant was identified by the Botany Department, Mandalay University, Myanmar with the specimen voucher number UM-22032018.

**Extraction and isolation**

The roots were air dried and then were cut into small pieces. The powder material (3.6 kg) was extracted with 95% ethanol (12.0 L) to yield a crude extract. After removing the solvent, 100 g of the extract was dissolved in methanol (300 ml) then it was extracted thrice using n-hexane in ratio 1:1(v/v). The methanol fraction (80.4 g) was subjected to Vacuum Liquid Chromatography (VLC) over silica gel eluting step-wise with mixture of n-hexane and ethyl acetate (EtOAc) to yield seven combined fractions (CF-1 to CF-7). Fraction CF-6 (25.6 g) was further fractionated by VLC with a gradient of n-hexane and ethyl acetate (n-hexane, 90:10-10:90, ethyl acetate) to obtain sub-fractions AF-1 to AF-7. AF-2 (80.60 mg) was further purified by silica gel column chromatography using three solvent systems of n-hexane-chloroform-ethyl acetate with the ratio of 100:70: 5-10: 5-20. A total of 67 fractions were collected and fractions 41-58 yielded a white amorphous solid and they were combined and purified in ethyl acetate to produce 4.5 mg of compound 1. AF-1 was subjected to column chromatography using n-hexane-ethyl acetate (n-hexane, 90:1-10:100, EtOAc) to obtain compounds 2 (9.8 mg) and 3 (28.6 mg) as pale yellow solids.

 **$\alpha$ -Glucosidase inhibitory activity**

The  $\alpha$ -glucosidase inhibitory activity was investigated using a protocol previously described by <sup>10</sup>. The inhibition against rat intestinal maltase and sucrase by the isolated compounds and fractions were determined using the aforementioned protocol. A 10  $\mu$ L of phosphate buffer (pH 6.9, 30  $\mu$ L), 20  $\mu$ L of the substrate solution (maltose: 10 mM; sucrose: 100 mM) in 0.1 M phosphate buffer, glucose kit (80  $\mu$ L) and the acetone extract of rat intestinal crude enzyme solution (20  $\mu$ L) were mixed. The reaction mixture was then incubated at 37°C for either 10 min (for maltose) or 40 min (for sucrose). The concentration of glucose released from the reaction mixture was detected by the glucose oxidase method using a glu-kit (Human, Germany). Enzymatic activity was quantified by measuring absorbance at 503 nm. The percent inhibition of reaction was calculated using the formulae;

$$\left[ \frac{(A_0 - A_1)}{A_0} \right] \times 100$$

where  $A_0$  is the absorbance without the sample, and  $A_1$  is the absorbance with the sample. The  $IC_{50}$  value was determined from a plot of percentage inhibition versus sample concentration. Acarbose® was used as standard control and the experiment was performed in triplicate.

In addition, the inhibition against yeast by all isolated compounds was determined by the method described by <sup>11</sup>. Sample (10  $\mu$ L) with the concentrations (5.0, 1.0, 0.20 and 0.04 mg/mL) was mixed with yeast (0.4 U/mL) in 1 mM phosphate buffer (pH 6.9), followed by shaking with microplate shaker about 2 min and pre-incubation at 37°C for 10 min. To the reaction mixture was added 50  $\mu$ L of p-nitrophenyl- $\alpha$ -D-glucopyranoside (PNPG) and then the mixture was incubated at 37°C for 20 min. After the incubation period, the reaction was quenched by adding  $Na_2CO_3$  (1M, 100  $\mu$ L). The release of p-nitrophenoxide from PNPG was detected by a microplate reader at 415 nm (iMark microplate reader).

**Statistical analysis**

All data are given as duplicate measurements and analyzed by non-


linear regression analysis. Quantitative data obtained were analyzed descriptively. All measurement was done in duplicate.

**Results and Discussion**

Compound 1 was obtained as a white amorphous powder. The proton NMR spectrum showed the presence of two cis-olefinic signals at  $\delta_H$  6.17 (1 H,  $J = 9.3$  Hz, H-3) and 7.84 (1 H,  $J = 9.3$  Hz, H-4). The presence of three aromatic protons was indicated by an ABX system at  $\delta_H$  7.44 (d,  $J = 8.5$  Hz, H-5), 6.78 (1H, dd,  $J = 2.2, 8.5$  Hz, -6H) and  $\delta_H$  6.70 (d,  $J = 2.2$  Hz, H-8) (Table 1). The  $^{13}C$ -NMR spectrum of compound 1 showed a cyclic lactone carbonyl at  $\delta_C$  162.4, three quaternary carbons 111.6, 162.2, 155.9, three aromatic CH carbons at 110.7, 129.2, 113.3, and two olefinic CH carbons at 144.7, 102.1 (Table 1). The DQF-COSY spectrum showed neighboring proton connections between  $\delta_H$  6.16 (H-3) and  $\delta_H$  7.84 (H-4), and also  $\delta_H$  6.78 (H-6) with 7.44 (H-5), and 6.70 (H-8). The long range HMBC experiments of compound 1 also confirmed the positions of protons and carbons in the compounds. From the above data, the structure of compound 1 was identified as umbelliferone (Figure 1) and the chemical shift data were in agreement with literature reports.<sup>12</sup> Compound 2 was obtained as a yellowish solid. The  $^1H$ -NMR data of compound 2 revealed protons characteristic of a coumarin moiety by two doublet signals at  $\delta_H$  6.16 and 8.04 (each 1H,  $J = 9.6$  Hz). Moreover, the presence of prenyl moiety was indicated by signals at  $\delta_H$  6.23 (dd,  $J = 17.4, 10.5$  Hz, 1H), 5.08 (dd,  $J = 17.4, 0.9$  Hz, 1H), 5.06 (dd,  $J = 10.5, 0.9$  Hz, 1H), and two germinal methyl group at 1.64 (s, 6H). The existence of a benzopyran ring was shown by a singlet signal representing two methylene protons at  $\delta_H$  2.75 (s, 2H) and another set of germinal methyl groups at  $\delta_H$  1.49 (s, 6H). The  $^{13}C$ -NMR and DEPT-90 and 135 spectra also indicated the presence of a cyclic lactone and ketone carbonyls at  $\delta_C$  160.6 and 198.2, respectively (Table 1). All the NMR data support compound 2 being clausenidin (Figure 1) and its NMR data also matched with literatures reports.<sup>13-14</sup>

Compound 3 was also obtained as a pale yellow solid. The  $^1H$ -NMR spectrum of the compound suggested a carbazole alkaloid type moiety based on the aldehyde proton signal at  $\delta_H$  9.93, N-H at 8.21, a hydrogen bonded OH at 11.42. Two singlets signals at  $\delta_H$  6.84 (s, 1H), and 8.05 (s, 1H) indicated a disubstitution on ring B. An ABX system of signals at 7.85 (d,  $J = 8.3$  Hz, 1H), 6.89 (m, 1H) and 6.88 (d,  $J = 2.2$  Hz, 1H) revealed that position C-7 of ring A was substituted. The  $^1H$ -NMR also showed the presence of one methoxy group at  $\delta_H$  3.93 (s, 3H). The existence of aldehydic carbonyl at  $\delta_C$  195.1 was confirmed by  $^{13}C$ -NMR, together with DEPT-90 and 135. After careful analysis of the 2D NMR and comparison with previous report the compound 3 was identified as 2-hydroxy-3-formyl-7-methoxycarbazole (lasine) (Figure 1).<sup>15</sup>

The  $\alpha$ -glucosidase inhibitory effect of the methanol fraction, n-hexane fraction, and the two of the isolated compounds (2 and 3) were tested using rat intestinal (maltase, sucrase) baker yeast enzymes. The methanol fraction with  $IC_{50}$  of 0.020 mg/mL showed a higher inhibitory activity than the n-hexane fraction with  $IC_{50}$  of 0.041  $\mu$ g/mL against maltase enzymes. But the two fractions (methanol and n-hexane) showed moderated activity against sucrase with  $IC_{50}$  values of 0.063 mg/mL each. Clausenidin (2) and lasine (3) showed no inhibition against yeast enzymes (Table 3).



**Figure 1:** Chemical structures of isolated compounds from *C. excavate*

**Table 1:** The  $^1\text{H}$  (600 MHz),  $^{13}\text{C}$  (151 MHz) NMR spectral data of compounds 1 and 2 in  $\text{CDCl}_3$ 

Position	$\delta_{\text{H}}$ (multi, $J$ values in Hz)		$\delta_{\text{C}}$ (ppm)	
	1	2	1	2
2	-	-	162.4	160.6
3	6.17 (d, $J = 9.3$ Hz, 1H)	6.16 (d, $J = 9.6$ Hz, 1H)	110.7	110.8
4	7.84 (d, $J = 9.3$ Hz, 1H)	8.04 (d, $J = 9.6$ Hz, 1H)	144.7	138.6
4a	-	-	111.6	103.2
5	7.44 (d, $J = 8.5$ Hz, 1H)	-	129.2	160.0
5a	-	-	-	104.0
6	6.78 (dd, $J = 2.3, 9.6$ Hz, 1H)	-	113.3	198.2
7	-	2.75 (s, 2H)	162.2	47.7
8	6.70 (d, $J = 2.2$ Hz, 1H)	-	102.1	80.0
8a	-	1.49 (s, 3H)	155.9	27.2
8b	-	1.49 (s, 3H)	-	27.2
9a	-	-	-	159.9
10	-	-	-	114.5
10a	-	-	-	159.0
1'	-	-	-	41.0
1a'	-	1.64 (s, 3H)	-	29.5
1b'	-	1.64 (s, 3H)	-	29.5
2'	-	6.23 (dd, $J = 17.4, 10.5$ Hz, 1H)	-	149.6
3'	-	-	-	108.4
3a	-	5.08 (dd, $J = 17.4, 0.9$ Hz, 1H)	-	-
3b	-	5.06 (dd, $J = 0.9, 10.5$ Hz, 1H)	-	-

**Table 2:** The  $^1\text{H}$  (600 MHz),  $^{13}\text{C}$  (151 MHz) NMR spectral data of compound 3 in  $\text{CDCl}_3$ 

Position	$\delta_{\text{H}}$ (mult, $J$ Hz)	$\delta_{\text{C}}$ (ppm)
1a	-	145.7
1	6.84 (s, 1H)	96.9
2	-	160.6
3	-	126.8
4	8.05 (s, 1H)	125.9
4a	-	115.4
5a	-	117.9
5	7.85 (d, $J = 8.3$ Hz, 1H)	120.5
6	6.89 (m, 1H)	95.6
7	-	159.2
8	6.88 (d, $J = 2.2$ Hz, 1H)	108.9
8a	-	141.5
7-OCH <sub>3</sub>	3.93 (s, 3H)	55.7
-CHO	9.93	195.1
-NH	8.21	-
-OH	11.42	-

40  
**Table 3:**  $\alpha$ -Glucosidase inhibitory activities of fractions and isolated compounds

Fraction/Compound	$\alpha$ -Glucosidase inhibition		
	Baker's yeast IC <sub>50</sub> (mM)	Maltase IC <sub>50</sub> (mg/mL)*	Sucrase IC <sub>50</sub> (mg/mL)*
n-hexane fraction	-	0.041	0.063
methanol fraction	-	0.020	0.063
2	NI	-	-
3	NI	-	-
Acarbose	0.1030	2.35 ( $\mu$ M)	15.48 ( $\mu$ M)

\* Nonlinear regression analyses were evaluated by Sigma-Plot 12.5  
NI = no inhibition at concentration  $\leq$  5 mg/mL

### Conclusion

Three compounds were obtained from *C. excavata*. The antidiabetic activity of the compounds and fractions were evaluated. The methanol fraction showed higher inhibitory activity than the n-hexane fraction against maltase enzymes. Among the three isolated compounds, 2 and 3 were tested against the yeast enzymes but both did not show any inhibition.

### Conflict of interest

The authors declare no conflict of interest.

### Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

### Acknowledgements

This work was supported by: (1) Airlangga Development Scholarship (ADS) (2) The Botany Department, Mandalay University, Myanmar for plant identified, and (3) The Faculty of Pharmacy, Meijo University, Nagoya, Japan for NMR data measurement.

### References

- Bathula C, Mamidala R, Thulluri C, Agarwal R, Jha KK, Munshi P, Adepally U, Singh A, Thirumalachary M, Sen S. Substituted furopyridinediones as novel inhibitors of  $\alpha$ -glucosidase. RSC Adv. accepted m:1-27.
- Zhao X, Tao J, Zhang T, Jiang S, Wei W, Han H, Shao Y, Zhou G, Yue H. Resveratrols Alleviates Postprandial Hyperglycemia in Diabetic Mice by Competitively Inhibiting  $\alpha$ -Glucosidase. J Agric Food Chem. 2019;67(10):2886-2893.
- Patel DK, Kumar R, Laloo D, Hemalatha S. Natural medicines from plant source used for therapy of diabetes mellitus: An overview of its pharmacological aspects. Asian Pacific J Trop Dis. 2012;2(3):239-250.

29

- Gothai S, Ganesan P, Park SY, Fakurazi S, Choi DK, Arulselvan P. Natural phyto-bioactive compounds for the treatment of type 2 diabetes: Inflammation as a target. Nutrients. 2016;8(8):1-28.
- Chen G, Guo M. Rapid screening for  $\alpha$ -glucosidase inhibitors from *Gymnema sylvestre* by affinity ultrafiltration-hplc-ms. Front Pharmacol. 2017;8(APR):1-8.
- Choi CW, Choi YH, Cha MR, Yoo DS, Kim YS, Yon GH, Hong KS, Kim YH, Ryu SH. Yeast  $\alpha$ -glucosidase inhibition by isoflavones from plants of leguminosae as an in vitro alternative to acarbose. J Agric Food Chem. 2010;58(18):9988-9993.
- Kumar R, Saha A, Saha D. A new antifungal coumarin from *Clausena excavata*. Fitoterapia. 2012;83(1):230-233.
- Peh TH, Lim GK, Taufiq-yap YH, Cheng G, Ee L. A New Cytotoxic Carbazole Alkaloid Isolated from the Stem Bark of Malaysian *Clausena excavata*. Can Chem Trans. 2013;1(3):165-172.
- Seo C, Ahn EK, Kang JS, Lee JH, Oh JS, Hong SS. Excavosides A and B, two new flavonoid glycosides from *Clausena excavata* Burm. f. (Rutaceae). Phytochem Lett. 2017;20(April):93-97.
- Ramadhan R, Phuwapraisirisan P. New arylalkanones from *Horsfieldia macrobotrys*, effective antidiabetic agents concomitantly inhibiting  $\alpha$ -glucosidase and free radicals. Bioorganic Med Chem Lett. 2015;25(20):4529-4533.
- Damsud T, Chanwun T, Kaewpiboon C. Antidiabetic Agents with  $\alpha$ -glucosidase Inhibition and Antioxidant Capacity from the Shoots of *Clausena cambodiana* Guill. Int J Agric Technol. 2017;13(4):449-456.
- Arbab IA, Abdul AB, Aspollah M, et al. *Clausena excavata* Burm. f. (Rutaceae): A review of its traditional uses, pharmacological and phytochemical properties. J Med Plants Res. 2011;5(33):7177-7184.
- Sripisut T, Cheenpracha S, Ritthiwigrom T, Prawat U, Laphookhieo S. Chemical Constituents from the Roots of *Clausena excavata* and Their Cytotoxicity. Rec Nat Prod. 2012;6(4):386-389.
- Su CR, Yeh SF, Liu CM, Damu AG, Kuo TS, Chiang PC, Bastow KF, Lee KH, Wu TS. Anti-HBV and cytotoxic activities of pyranocoumarin derivatives. Bioorganic Med Chem. 2009;17(16):6137-6143.

# Coumarins and Carbazole Alkaloid from *Clausena excavata* Roots and Investigation of their $\alpha$ -glucosidase Inhibitory Activity

## ORIGINALITY REPORT

**25%**  
SIMILARITY INDEX

**19%**  
INTERNET SOURCES

**21%**  
PUBLICATIONS

**0%**  
STUDENT PAPERS

## PRIMARY SOURCES

**1** [etasr.com](http://etasr.com) 1%  
Internet Source

**2** [www.cepceb.ucr.edu](http://www.cepceb.ucr.edu) 1%  
Internet Source

**3** P Mahesh Kumar, Shaheen Begum, Gella Sunil, P Rajyalakshmi Devi, Naga Santhosh Kesana. "Screening of larvicidal activity of nanoparticles synthesized from flower extracts of *Hibiscus vitifolius*", *Journal of Applied Biotechnology & Bioengineering*, 2018 1%  
Publication

**4** Xu Wan, Shao-Yong Zhang, Hui Zhang, Jun Zhai, Jun Huang, An-Liang Chen, Ji-Dong Wang. " Two new tenvermectins from a genetically engineered strain MHJ1011 ", *Journal of Asian Natural Products Research*, 2016 1%  
Publication

5	Internet Source	1 %
6	K N W Tun, N S Aminah, A N Kristanti, R Ramadhan, Y Takaya. " Two Flavonoids From Stem Bark of and Their Antidiabetic and Antioxidant Activities ", IOP Conference Series: Earth and Environmental Science, 2019 Publication	1 %
7	dx.doi.org Internet Source	1 %
8	koreascience.or.kr Internet Source	1 %
9	Khun Nay Win Tun, Nanik Siti Aminah, Alfinda Novi Kristanti, Hnin Thanda Aung, Yoshiaki Takaya. "Natural products isolated from Casimiroa", Open Chemistry, 2020 Publication	1 %
10	mts.intechopen.com Internet Source	1 %
11	www.faqs.org Internet Source	1 %
12	brevets-patents.ic.gc.ca Internet Source	1 %
13	journals.sagepub.com Internet Source	1 %

14	coek.info Internet Source	1 %
15	ediss.uni-goettingen.de Internet Source	1 %
16	www.dovepress.com Internet Source	1 %
17	dokumen.pub Internet Source	1 %
18	protocolexchange.researchsquare.com Internet Source	1 %
19	repository.unair.ac.id Internet Source	1 %
20	real.mtak.hu Internet Source	1 %
21	Guangzhong Yang, Huijin Feng, Yuanchao Li. "Di-and Triterpenoids from Tripterygium wilfordii", Natural Product Letters, 2001 Publication	1 %
22	docplayer.net Internet Source	1 %
23	Wisuttaya Worawalai, Preecha Phuwapraisirisan. "Samin-derived flavonolignans, a new series of antidiabetic agents having dual inhibition against $\alpha$ -	< 1 %



# glucosidase and free radicals", Natural Product Research, 2019

Publication

24

[patents.justia.com](https://patents.justia.com)

Internet Source

<1 %

25

Ahmed I. Khodair, Najim A. Al-Masoudi, Jean-Pierre Gesson. "A New Approach to the Synthesis of Benzothiazole, Benzoxazole, and Pyridine Nucleosides as Potential Antitumor Agents", Nucleosides, Nucleotides and Nucleic Acids, 2003

Publication

<1 %

26

Yun-Song Wang, Bi-Tao Li, Shi-Xi Liu, Zheng-Qi Wen, Jing-Hua Yang, Hong-Bin Zhang, Xiao-Jiang Hao. " Anisucoumaramide, a Bioactive Coumarin from ", Journal of Natural Products, 2017

Publication

<1 %

27

[www.ijert.org](http://www.ijert.org)

Internet Source

<1 %

28

[macau.uni-kiel.de](http://macau.uni-kiel.de)

Internet Source

<1 %

29

[rabm.scholasticahq.com](http://rabm.scholasticahq.com)

Internet Source

<1 %

30

[www.research.lancls.ac.uk](http://www.research.lancls.ac.uk)

Internet Source

<1 %

31 Ming-An Ouyang, Zhen-Dan He, Cui-Ling Wu. " Anti-oxidative activity of glycosides from ", Natural Product Research, 2003  
Publication <1 %

---

32 Abou-Gazar, H.. "Antioxidant lignans from Larrea tridentata", Phytochemistry, 200409  
Publication <1 %

---

33 M. Tourirte, T. Oulih, H. B. Lazrek, J. L. Barascut, J. L. Imbach, N. A. Almasoudi. "Synthesis of 3'-Deoxy-3'-[4-(pyrimidin-1-yl)methyl-1,2,3-triazol-1-yl]thymidine via 1,3-Dipolar Cycloaddition", Nucleosides, Nucleotides and Nucleic Acids, 2003  
Publication <1 %

---

34 Suda Chakthong, Nitima Bindulem, Sirorat Raknai, Sariya Yodwaree, Soonthorn Kaewsanee, Akkharawit Kanjana-Opas. " Carbazole-pyranocoumarin conjugate and two carbazole alkaloids from the stems of ", Natural Product Research, 2016  
Publication <1 %

---

35 moam.info  
Internet Source <1 %

---

36 paginas.estsp.ipp.pt  
Internet Source <1 %

---

37 repository-tnmgrmu.ac.in  
Internet Source <1 %

---

38

[www.arkat-usa.org](http://www.arkat-usa.org)

Internet Source

&lt;1 %

39

Krista A. Gill, Fabrice Berru , Jennifer C. Arens, Russell G. Kerr. " Isolation and Structure Elucidation of Cystargamide, a Lipopeptide from ", Journal of Natural Products, 2014

Publication

&lt;1 %

40

Teerayut Sriyatep, Ittipon Siridechakorn, Wisanu Maneerat, Acharavadee Pansanit et al. " Bioactive Prenylated Xanthonenes from the Young Fruits and Flowers of ", Journal of Natural Products, 2015

Publication

&lt;1 %

41

[link.springer.com](http://link.springer.com)

Internet Source

&lt;1 %

42

[lup.lub.lu.se](http://lup.lub.lu.se)

Internet Source

&lt;1 %

43

[orca.cf.ac.uk](http://orca.cf.ac.uk)

Internet Source

&lt;1 %

44

[www.thieme-connect.com](http://www.thieme-connect.com)

Internet Source

&lt;1 %

45

[www.thieme-connect.de](http://www.thieme-connect.de)

Internet Source

&lt;1 %

46

Mar a Losada-Echeberr a, Mar a Herranz-L pez, Vicente Micol, Enrique Barrajon-Catal n. "Polyphenols as Promising Drugs

&lt;1 %

against Main Breast Cancer Signatures",  
Antioxidants, 2017

Publication

---

47

O. Osadebe Patience, U. Odoh Estella, F. Uzor Philip. "The search for new hypoglycemic agents from plants", African Journal of Pharmacy and Pharmacology, 2014

Publication

---

<1 %

48

Amy M. Hamlin, David Lapointe, Kyle Owens, Richmond Sarpong. " Studies on C - Diterpenoid Alkaloids: Synthesis of the Hetidine Framework and Its Application to the Synthesis of Dihydrovirine and the Atisine Skeleton ", The Journal of Organic Chemistry, 2014

Publication

---

<1 %

49

Chaiyavat Chaiyasut, Bhagavathi Sivamaruthi, Noppawat Pengkumsri, Waranya Keapai et al. "Germinated Thai Black Rice Extract Protects Experimental Diabetic Rats from Oxidative Stress and Other Diabetes-Related Consequences", Pharmaceuticals, 2016

Publication

---

<1 %

50

Clynton W. Halstead, Paul I. Forster, Peter G. Waterman. " Alkaloids from the stem bark of an Australian population of ", Natural Product Research, 2006

Publication

---

<1 %

51

Ernesto Rivera-Becerril, Pedro Joseph-Nathan, Víctor M. Pérez-Álvarez, Martha S. Morales-Ríos. "Synthesis and Biological Evaluation of (-)- and (+)-Debromoflustramine B and Its Analogues as Selective Butyrylcholinesterase Inhibitors", Journal of Medicinal Chemistry, 2008

Publication

&lt;1 %

52

Li HUANG, Zhe-Ling FENG, Yi-Tao WANG, Li-Gen LIN. "Anticancer carbazole alkaloids and coumarins from Clausena plants: A review", Chinese Journal of Natural Medicines, 2017

Publication

&lt;1 %

53

Nanik S. Aminah, Tin M. Thant, Alfinda N. Kristanti, Rico Ramadhan, Hnin T. Aung, Yoshiaki Takaya. " Carbazomarin: A New Potential of  $\alpha$ -Glucosidase Inhibitor From Roots ", Natural Product Communications, 2020

Publication

&lt;1 %

54

Yian Hoon Lee, Candy Choo, Mindani I Watawana, Nilakshi Jayawardena, Viduranga Y Waisundara. "An appraisal of eighteen commonly consumed edible plants as functional food based on their antioxidant and starch hydrolase inhibitory activities", Journal of the Science of Food and Agriculture, 2015

&lt;1 %

55

Zhongzhi Zhu, Jinghe Cen, Xiaodong Tang,  
Jianxiao Li, Wanqing Wu, Huanfeng Jiang.  
"Access to Amidines and Arylbenzimidazoles:  
Zinc-Promoted Rearrangement of Oxime  
Acetates", *Advanced Synthesis & Catalysis*,  
2018

<1 %

Publication

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# Coumarins and Carbazole Alkaloid from *Clausena excavata* Roots and Investigation of their $\alpha$ -glucosidase Inhibitory Activity

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---