

OVOZOA

Journal of Animal Reproduction

Kejadian mastitis subklinis pada kambing perah Peranakan Etawa di desa Bangelan kecamatan Wonosari kabupaten Malang

*Nur Zaini Nanda Mardian, Soeharsono Soeharsono, Nenny Harijani,
Herry Agoes Hermadi, Budiarto Budiarto, Wurlina Wurlina* 60 - 63

Efektivitas pemberian GnRH pada sapi perah yang mengalami hipofungsi ovarium terhadap waktu timbulnya birahi dan angka kebuntingan

*Winadya Reika Ummaisyah, Sri Pantja Madyawati, Retno Sri Wahjuni,
Rimayanti Rimayanti, Wurlina Wurlina, Tjuk Imam Restiadi* 64 - 68

Penambahan alfa-tokoferol dalam pengencer susu skim - kuning telur terhadap kualitas spermatozoa domba Sapudi yang disimpan pada suhu 5°C

*Satya Alysa Cahya Puspita, Suherni Susilowati, Soeharsono Soeharsono,
Trilas Sardjito, Abdul Samik, Indah Norma Triana* 69 - 76

Pengaruh injeksi nikotin terhadap motilitas, viabilitas, dan integritas membran spermatozoa mencit (*Mus Musculus*)

*Kautsar Pandu Pramudita, Widya Paramita Lokapirnasari, Suheni Susilowati,
Epy Muhammad Luqman, Hermin Ratnani, Erma Safitri* 77 - 81

Relationship between the value of the estrous detector measurement result and serum progesterone level in Etawa crossbred goats after estrous synchronization

*Annisa Lita Triagil, Ismudiono Ismudiono, Boedi Setiawan, Tita Damayanti Lestari,
Abdul Samik, Budi Utomo* 82 - 85



Ovozoa: Journal of Animal Reproduction by Unair is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.



**Publisher: Department of Veterinary Reproduction
Faculty of Veterinary Medicine Universitas Airlangga
in collaboration with the Indonesian Association of
Department of Veterinary Reproduction**



VOL 9, NO 3 (2020)

OVOZOA : JOURNAL OF ANIMAL REPRODUCTION

TABLE OF CONTENTS

FRONT MATTER

Front Matter Vol.9 No.3 (2020)



BACK MATTER

Back Matter Vol.9 No.3 (2020)



ORIGINAL ARTICLE



Kejadian mastitis subklinis pada kambing perah Peranakan Etawa di desa Bangelan kecamatan Wonosari kabupaten Malang



60-63



10.20473/ovz.v9i3.2020.60-63



Nur Zaini Nanda Mardian, Soeharsono Soeharsono, Nenny Harijani, Budiarto Budiarto, Herry Agoes Hermadi, Wurlina Wurlina



Efektivitas pemberian GnRH pada sapi perah yang mengalami hipofungsi ovarium terhadap waktu timbulnya birahi dan angka kebuntingan



64-68



10.20473/ovz.v9i3.2020.64-68



Winadya Reika Ummaisyah, Sri Pantja Madyawati, Rimayanti Rimayanti, Wurlina Wurlina, Tjuk Imam Restiadi, Retno Sri Wahjuni



Penambahan alfa-tokoferol dalam pengencer susu skim - kuning telur terhadap kualitas spermatozoa domba Sapudi yang disimpan pada suhu 5°C



69-76



10.20473/ovz.v9i3.2020.69-76



Satya Alysa Cahya Puspita, Suherni Susilowati, Trilas Sardjito, Abdul Samik, Indah Norma Triana, Soeharsono Soeharsono



Pengaruh injeksi nikotin terhadap motilitas, viabilitas, dan integritas membran spermatozoa mencit (Mus Musculus)



77-81



10.20473/ovz.v9i3.2020.77-81



Kautsar Pandu Pramudita, Widya Paramita Lokapirnasari, Suheni Susilowati, Hermin Ratnani, Erma Safitri, Epy Muhammad Luqman



Relationship between the value of the estrous detector measurement result and serum progesterone level in Etawa crossbred goats after estrous synchronization



82-85



10.20473/ovz.v9i3.2020.82-85



Annisa Lita Triagil, Ismudiono Ismudiono, Tita Damayanti Lestari, Abdul Samik, Budi Utomo, Boedi Setiawan

People

Contact

Editorial Team

Reviewers

Instruction for Author

Guide for authors

Online Submission

Download Documents

Journal Policy

Focus and Scope

Section Policies

Publication Frequency

Publication Ethics

Indexing

Article Processing Charge

Peer Reviewers Process

Open Access Statement

Archiving

Plagiarism Screening

History

Copyright

old website

Tools



grammarly



OPEN JOURNAL SYSTEMS

Journal Help

00032465

[View My Stats](#)



Ovozoa: Journal of Animal Reproduction by Unair is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#).



This journal is supported by



USER

Username

Password

Remember me

NOTIFICATIONS

[View](#)
[Subscribe](#)



Pusat Pengembangan Jurnal dan Publikasi Ilmiah
Universitas Airlangga
copyright©2017 Template PPJPI

Relationship between the value of the estrous detector measurement result and serum progesterone level in Etawa crossbred goats after estrous synchronization

Hubungan antara nilai hasil pengukuran detektor estrus dengan progesteron serum pada kambing Peranakan Etawa setelah sinkronisasi estrus

Annisa Lita Triagil¹, Ismudiono Ismudiono^{2*}, Boedi Setiawan³, Tita Damayanti Lestari², Abdul Samik², Budi Utomo²

¹ Student, ² Department of Veterinary Reproduction, ³ Department of Veterinary Clinic, Faculty of Veterinary Medicine, Universitas Airlangga

*Corresponding author, e-mail: ismudiono@fkh.unair.ac.id

Open access under CC BY – SA license, Doi : 10.20473/ovz.v9i3.2020.82-85

Received September 24 2020, Revised November 18 2020, Accepted December 6 2020

Published online December 6 2020

ABSTRACT

This study was conducted to determine the relationship between the estrous detector's reading value (Draminski®, Dramiński S.A, Poland) and serum progesterone levels on Etawa crossbred does. Eighteen healthy non-pregnant Etawa crossbreds does weighing 20-25 kg, in their first parity were used in this study. The does were divided equally into two groups. In the two groups, the does were synchronized their estrus by means of intramuscular and intravulvar injection of 7.5 mg prostaglandin F_{2α} (PGF_{2α}) twice in 11-day intervals. After the second injection, the estrous detector value measurement, visual observation of the estrus sign, and blood sampling were conducted for five days. The blood sample was processed into the serum to examine for progesterone levels by ELISA. The data analysis was using the simple correlation method on SPSS type 23 for windows. The result showed that a positive relationship ($p < 0.05$) with coefficient correlation (r) was 0.609 between the values of the estrous detector measurement with serum progesterone levels. It could be concluded that the estrous detector's low value indicates that the Etawa crossbred was in the estrus phase and followed by the decreasing of progesterone level but not reaching the basal level.

Keywords: estrous detector, Etawa crossbred, progesterone level, synchronization

INTRODUCTION

Etawa crossbred goat is crossed from the Jamnapari from Etawa district of India's Uttarbal region with Kacang goat, a local goat of Indonesia. Etawa crossbred goats have higher milk and meat yields than local goats (Prajoga, 2007). Breeders in improving the quality of livestock need to have artificial insemination. Determination of the right estrous time is essential for artificial insemination (Omontese, 2018).

Draminski's estrous detector is a device that can help detect estrous. This tool shows figures

that describe the quality of vagina mucus through electrical resistance (Izabela et al., 2007; Meydilasari *et al.*, 2020). According to Draminski (2009), a goat is declared estrus if an estrous detector is inserted to examine vaginal mucus showing a number close to 200. Synchronization of estrus was carried out on livestock to obtain estrus simultaneously. The synchronization that is commonly used is prostaglandin (PF_{2α}) (Ismudiono *et al.*, 2009). The prostaglandin mechanism (F_{2α}) as a hormone produced by the endometrium is to lyse the corpus luteum. Lysis of CL resulted in the normal cycle's return to being effective at

inducing estrus (Saoeni, 2008). Progesterone is one of the essential hormones in reproduction, secreted by the luteal cells (Atabany *et al.*, 2001). Progesterone levels decreased during estrus on day 0 to day 21 (Siregar, 2009). The decreased progesterone levels in the estrous cycle are due to the corpus luteum regressed (Sunendar *et al.*, 2008).

MATERIALS AND METHODS

Eighteen, healthy, non-pregnant Etawa Crossbred does, in their first parity, weighing 20-25 kg were used for the study. Does were synchronized their estrus using 7.5 mg PGF2 α (Enzaprost) twice in 11 days intervals. The does were divided equally into two groups. In the first group, the does were injected intramuscularly (IM). Meanwhile, the second group was injected intravulva (IV). After the second injection, the estrous detector value measurement, visual observation of estrus signs, and blood sampling were conducted for five consecutive days.

Vaginal mucus resistance was examined using Draminski estrous detector the following day after the second PGF2 α injection. Vaginal mucus resistance was measured using an estrous detector, which is initiated by applying lubricant to the vaginal probe. A vaginal probe, which was equipped with electrodes, was inserted around three-quarters of the vagina depth, then the value was recorded. Simultaneously, with the estrous detector value

measurement, the estrus sign was observed by bringing the female goat closer to the male goat cage. The three main signs observed were the red color of the vaginal mucous membrane, the amount of cervical mucus, and tail position (up or down). Blood samples were taken from the jugular vein using venoject and vacuum plain tubes. Then the vacuum tube was tilted for 1 hour until a clear coloured serum appeared. Tubes were then centrifuged at 3000 rpm for 15 minutes. Afterward, serum was placed in Eppendorf tubes and stored in a -20⁰C freezer until the level of progesterone was analyzed with ELISA (Goat Progesterone, PROG ELISA KIT, Cat. MBS704614).

Ethics

This study was approved ethically by the Animal Care and Use Committee (ACUC) Faculty of Veterinary Medicine, Universitas Airlangga, with ethical clearance No.2.KE.143.07.2019.

RESULT

The average value of mucus resistance after the second PGF2 α injection can be seen in Table 1. Two out of the nine does in each group (intramuscular and intravulvar injection) exhibited obvious signs of estrus including reddish vaginal mucous membrane, plenty of cervical mucus, and tail up (Table 2). Serum progesterone level is presented in Tabel 3.

Table 1 The value of the estrous detector measurement result of Etawa crossbred goat after the second PGF2 α injection.

	day-1	day-2	day-3	day-4	day-5
IM	347.78 \pm 136.91	353.33 \pm 135.46	296.67 \pm 134.07	325.56 \pm 83.98	353.33 \pm 97.72
IV	348.89 \pm 116.24	317.78 \pm 77.59	267.78 \pm 82.28	283.33 \pm 97.08	307.78 \pm 63.20

IM= intramuscular PGF2 α injection; IV= intravulvar PGF2 α injection; replicates= 9

Table 2 Estrus sign based on visual observation of Etawa crossbred does after the second PGF2 α intramuscular and intravulvar injection

	day-1	day-2	day-3	day-4	day-5
intramuscular					
105					
Vmm	pale	reddish	reddish	reddish	pale
Cm	slight	moderate	a lot	a lot	slight
tail	-	up	up	down	down
125					
Vmm	pale	pale	reddish	reddish	pale
Cm	slight	moderate	a lot	a lot	moderate
tail	-	-	up	down	down
intravulvar					
285					
Vmm	reddish	pale	reddish	pale	pale
Cm	moderate	slight	a lot	a lot	slight
tail	-	-	up	down	down
324					
Vmm	reddish	reddish	reddish	pale	pale
Cm	moderate	a lot	a lot	slight	moderate
tail	-	up	up	down	down

105, 125, 285, 324= does number; Vmm= vaginal mucous membrane; Cm= cervical mucus

Table 3 Serum Progesterone level (ng/mL) of Etawa crossbred does after the second PGF2 α intramuscular and intravulvar injection

does #	day-1	day-2	day-3	day-4	day-5
IM (n=2)					
105	11.106	1.391	1.809	1.589	1.944
125	5.269	0.865	1.162	1.222	1.373
X \pm SD	8.19 \pm 4.13	1.13 \pm 0.37	1.49 \pm 0.46	1.41 \pm 0.26	1.66 \pm 0.40
IV (n=2)					
285	1.006	1.609	3.701	1.787	1.664
324	1.085	1.139	1.599	1.685	1.421
X \pm SD	1.05 \pm 0.06	1.13 \pm 0.37	2.65 \pm 1.49	1.74 \pm 0.07	1.54 \pm 0.17

IM= intramuscular PGF2 α injection; IV= intravulvar PGF2 α injection

DISCUSSION

The Draminski Estrous detector shows a lower number (around 200) in estrous onset on day 0-3 and had an increase after estrous (Setiawan, 2012). On the 3rd day, the female begins to reddish vagina mucous membrane, a lot of cervical mucus, tail up (Table 2) and mounted the other goat, followed by a thick and clear discharge from the vulva (Davilia *et al.*, 2018; Ridlo *et al.*, 2018 and Tanjung *et al.*, 2015).

The device's electrical current resistance causes the estrous detector measurement results at the estrous phase. This is due to changes in the composition in the vagina; that is, the amount of mucus that increases (Balumbi *et al.*, 2019, Meydilasari *et al.*, 2020). The composition contained in vaginal mucus after the second PGF2 α injection consists of sodium, potassium, and ions, which are electrolyte and have conductivity (Tsilinggani *et al.*, 2001), so that the higher the volume or number of ions in the vaginal mucus, the lower electrical resistance will occur (Verma *et al.*, 2014). The

estrous detector measurement result has decreased in the estrous phase and can be caused by the hormone estrogen. The vaginal vasodilation causes an increase in the mucus of electrolyte ions and has a relatively high conductivity so that the resistance to electrical current is low and the value of the estrous detector decreases (Kitwood *et al.*, 1993).

In this study, the serum progesterone level of Etawa crossbred goat at the estrus phase was 1.49 ± 0.46 to 2.65 ± 1.49 ng/mL. It was higher compared to 0.083 ± 0.15 ng/mL of day 0 and 0.084 ± 0.11 of Etawa-Saanen crossbred goats progesterone level (Sitarasemi *et al.*, 2017). The progesterone levels in the estrous phase are influenced by several factors, including nutrition (Tanaka *et al.*, 2003), unreported corpus luteum, BCS, postpartum state, lactation, and the presence of males (McDonald *et al.*, 2014; Suharto *et al.*, 2008; Khanum *et al.*, 2006; Totti, 2009).

The high levels of progesterone after the synchronization of estrous in this research might be caused by nutrition. The goats fed with low nutrition have increased levels of progesterone in the estrous to the luteal phase. Nutrition plays a vital role in reproduction. Low nutrition can cause the hormones gonadotropin and LH to be secreted in small amounts, where they play an essential role in stimulating the ovaries for ovulation (McDonald *et al.*, 2014).

High progesterone levels during the estrous phase can occur as a result of postpartum. Goats in 27.9 ± 14 days postpartum show 2.1 ng / ml on day 0 to third (Khanum *et al.*, 2008). Progesterone level in the luteal phase of postpartum does ranged from 2.6 to 5.4 ng /mL (Gaafar *et al.*, 2005). In the IM PGF2Fa injection Group, does numbers 105 and 125 on the first day showed high levels of progesterone and decreased on the second day. The other two goats (number 285 and 324) show a progesterone fluctuation after IV PGF2Fa injection. It means they were in deference stages of the cycle and might be the does were not affected by the hormone injection. This is due to the first day of the second PGF2Fa injection, the corpus luteum has not been regressed. The second day after the second injection, the corpus luteum starts to lysis and

decreases progesterone concentration (Satiti *et al.*, 2014, Lestari dan Ismudiono, 2014).

Adequate nutrition for goats should be given a diet containing 60-75% TDN with 12-17% protein content. Malnutrition of goats affects the inadequate response of estrous synchronization (Paula *et al.*, 2005). The interval of removal of the sponge in the synchronization and the ovulation rate tend to depend on the amount of body mass lost during malnutrition (Lozano *et al.*, 1997). The less than optimal response to the synchronization causes the corpus luteum to undergo an imperfect regression, where PGF2a should lyse the corpus luteum, so that blood flow in CL decreases dramatically, followed by a decrease in progesterone (Towle, 2002). The imperfect regression corpus luteum causes high levels of progesterone

There was a positive relationship ($r=0.609$, $p<0.05$) between the value of the estrous detector measurement result with progesterone levels. It means a high progesterone level follows a high estrous detector value. The estrous detector's low value indicates that the goat has estrus followed by a decrease in progesterone levels even though it is not in a basal state. The goat, which shows usual symptoms of estrous with progesterone levels above the normal level of estrous phase, is possibly caused by a good body condition score. The animal with good health shows a normal cycle of reproduction (Lestari and Ismudiono 2014). Nevertheless, some goats might behave good estrous but not accompanied by LH surges. The absence of LH surge can be caused by an imperfect corpus luteum and finally showed lower progesterone level (Suharto *et al.*, 2008).

CONCLUSION

In conclusion, there is a positive relationship between the value of the estrous detector and progesterone levels. This relationship has a correlation value of 0.609 and a significant value (2-tailed) of 0.004 <0.05 (positive). That means the estrous detector's low value indicates the Etawa crossbred goats have estrous and progesterone levels and decrease but do not reach the basal level.

ACKNOWLEDGEMENT

The authors grateful to UPT HMT Singosari Malang's head and staff, who have given permission and assistance during the research.

REFERENCES

- Atabany A, Abdulgani IK, Sudono A, Mudikdjo K. 2001. Performa Produksi, Reproduksi dan Nilai Ekonomis Kambing Peranakan Etawa di Peternakan Barokah. *Media Peternakan* 24: 1-7.
- Balumbi M, Setiadi MA, Supriatna I. 2019. Response and Characteristics of Estrous after Estrous Synchronization with Cloprostenol in Friesian Holstein Cow. *Acta Vet Indonesiana* 7: 29-36.
- Davilia FS, Del Bosque Gonzales AS, Barragá HB. 2018. Reproduction in Goat. *intechopen*.70003. <http://dx.doi.org/10.5772/>
- Gaafar KM, Gabr MK, Teleb DF. 2005. The hormonal profile during the estrous cycle and gestation in Damascus goats. *J Small Rumin Res.* 57: 85-97.
- Ismudiono I, Anwar H, Srianto P, Madyawati SP, Samik A, Safitri E. 2009. *Buku Ajar Fisiologi Reproduksi pada Ternak*. Airlangga University Press. Surabaya.
- Izabela M, Angela S, Paula P, Laura U. 2007. Researchs on the Electrical Resistence of Cervical Mucus in Cows . *lucrări științifice zootehnie și biotehnologii*. Timisoara 40:568-57.
- Khanum SA, Hussain M, Kausar R. 2006. Assessment of Reproductive Parameters in Female Dwarf goat (*Capra hircus*) on The Basis of Progesterone Profile. *J Anim Rep Sci.* 102: 267-75.
- Khanum SA, Hussain M, Kausar R. 2008. Progesterone and Estradiol Profile During Estrous Cycle and Gestation in Dwarf Goat (*Capra Hircus*). *Pakistan Vet J.* 28: 1-4.
- Kitwood SE, Philips CJC, Weise M. 1993. Use of a Vaginal Mucus Impedance Meter to Detect Estrus. *J. Theriogenology* 40: 559-69.
- Lestari TD dan Ismudiono I. 2014. Ilmu Reproduksi Ternak. Airlangga University Press. Surabaya.
- Lozano JM, Abecia JA, Forcada F, Zazaraga L, Alfaro B. 1997. Effect of Undernutrition on The Distribution of Progesterone in The Uterus of Ewe During The Luteal Phase of The Estrous Cycle. *Theriogenology* 49: 539-46.
- McDonald P, Edwards RA, Greenhalgh JFD, Morgan CA, Sinclair LA, Wilkinson RG. 2010. *Animal Nutrition*. Seventh Edition. Pearson. London. 388.
- Meydilasari NI, Hernawati T, Hidanah S, Lestari TD, Safitri E, Sardjito T, Ismudiono I. 2020. Hubungan antara skor hasil bacaan estrous detector dengan gambaran sitologi vagina sebagai parameter penentuan estrus pada kambing. *Ovozoa J Anim Reprod.* 9: 7-11.
- Omontese BO. 2018. Estrus Synchronization and Artificial Insemination in Goats. Open access peer-reviewed chapter. <https://doi.org/10.5772/intechopen.74236>
- Paula NRO, Galeati G, Teixeira DIA, Lopes Junior ES, Freitas VJF, Rondina D. 2005. Responsiveness to Progestagen-eCG-Cloprostenol Treatment in Goat Food Restricted for Long Period and Refed. *Reprod Domest Anim.* 40: 108-10.
- Prajoga SBK. 2007. Effect of In-Breeding on Estimation of Respon to Selection of Weaning Weight of Etawa Cross Bred (PE), in The Small Population. *J Anim Sci Padjadjaran Univ.* 7: 170-8. <https://doi.org/10.24198/jit.v7i2.2254>.
- Ridlo MR, Ummami R, Dalimunthe NWY, Ramadani D, Prihanani NI, Andityas M, Widi TSM. 2018. Profil vulva dan suhu tubuh kambing Peranakan Etawa pada sinkronisasi estrus menggunakan Medroxy Progesterone Acetate dan suplementasi Zinc (Zn). *J Nasional Teknologi Terapan.* 2:198-211.
- Saoeni R. 2007. Efek pemberian Prostaglandin F2 α secara intra vaginal spons (IVS) dan intra muskular (IM) terhadap peningkatan kinerja reproduksi domba. *J Animal Production* 9: 129-34.

- Satiti D, Triana IN, Rahardjo AP. 2014. Pengaruh Penggunaan Kombinasi Progesteron (Medroxy Progesterone Acetate) dan Prostaglandin (PGF2 α) Injeksi Terhadap Persentase Estrus dan Kebuntingan pada Domba Ekor Gemuk. *Veterinaria Medika*. 7:126-33.
- Setiawan KD. 2012. Karakteristik Lendir Vagina Kambing Peranakan Etawa (PE) Setelah Sinkronisasi berahi dengan Prostaglandin [Skripsi]. Fakultas Kedokteran Hewan. Institut Pertanian Bogor. 10. <http://repository.ipb.ac.id/handle/123456789/60909>
- Siregar TN. 2009. Estrogen and Progesterone Profile in Aceh Local Goat Estrous Cycle. *J Ked Hewan* 3: 240-47.
- Sitarasemi PI, Widyobroto BP, Bintara S, Widayati DT. 2017. Progesterone and Biochemical Profile of Ettawa-Saanen Crossbreed Goat in Turri Area Yogyakarta-Indonesia. *Int J Dairy Sci*. 12: 289-94.
- Suharto K, Junaidi A, Kusumawati A, Widayati DT. 2008. Comparison of Fertility of Etawa Crossbred Goats in Low versus Ideal Body Condition Score after Estrus Synchronization and Artificial Insemination. *Media Kedokteran Hewan* 24: 49-54.
- Sunendar S, Widayati DT, Junaidi A. 2008. Profil Hormon Progesteron dan Estrogen pada Kambing Peranakan Etawah yang Disinkronisasi Estrus dengan Implan Controlled Internal Drug Release. *Buletin Peternakan* 32: 26-37.
- Tanaka TT, Yamaguchi H, Kamomae H, Kaneda Y. 2003. Nutritionally Induced Body Weight Loss and Ovarian Quiescence in Shiba Goats. *J Reprod Dev*. 49: 113-9.
- Tanjung AD, Setiatin ET, Samsudewa D. 2015. Level of estrogen hormone and estrus performance of different postpartum estrus of Jawa Randu goat. *J Indonesian Trop Anim Agric*. 40: 87-92.
- Totti T. 2009. Studi hubungan hormon Progesteron dengan jumlah korpus luteum pada kambing. *jurnal ilmiah aplikasi isotop dan radiasi* 5: 43-51.
- Towle TA, Tsang PCW, Milvae RA, Newbury MK, McCracken JA. 2002. Dynamic In Vivo Changes in Tissue Inhibitors of Metalloproteinases 1 and 2, and Matrix Metalloproteinases 2 and 9, During Prostaglandin F2 α -Induced Luteolysis in Sheep. *Biol Reprod*. 66:1515-1521.
- Tsilingianni TH, Karagianidis A, Brikas P, Saratis PH. 2001. Physical Properties of Cervical Mucus During Normal and Induced (Progesterone and/or PGF2 α). *Theriogenology* 55: 629-40.
- Verma KK, Prasad S, Kumaresan A, Mohanty TK, Layek SS, Patbandha TK, Chand S. 2014. Characterization of Physico-Chemical Properties of Mucus in Relation to Parity and Conception Rate in Murrah Buffaloes. *Vet world* 7: 467-71.