

# The association of fatty acid desaturase gene polymorphisms on long-chain polyunsaturated fatty acid composition in Indonesian infants

*by Conny Tanjung*

---

**Submission date:** 17-Apr-2021 04:06PM (UTC+0800)

**Submission ID:** 1561709799

**File name:** polyunsaturated\_fatty\_acid\_composition\_in\_Indoesian\_infants.pdf (180.97K)

**Word count:** 8747

**Character count:** 40723

# The association of fatty acid desaturase gene polymorphisms on long-chain polyunsaturated fatty acid composition in Indonesian infants

Conny Tanjung,<sup>1,2</sup> Peter Rzehak,<sup>3</sup> Herawati Sudoyo,<sup>4,5</sup> Muchtaruddin Mansyur,<sup>6</sup> Zakiudin Munasir,<sup>7</sup> Suzanna Immanuel,<sup>8</sup> Roedi Irawan,<sup>9</sup> Eva Reischl,<sup>10</sup> Hans Demmelmaier,<sup>3</sup> Sri Rezeki Hadinegoro,<sup>7</sup> Damayanti Rusli Sjarif,<sup>2,7</sup> and Berthold Koletzko<sup>3</sup>

<sup>1</sup>Pantai Indah Kapuk Hospital, Jakarta, Indonesia; <sup>2</sup>Human Genetic, Indonesian Medical Education and Research Institute, Jakarta, Indonesia; <sup>3</sup>LMU - Ludwig-Maximilians-Universität München, Dr von Hauner Children's Hospital, Division of Metabolic and Nutritional Medicine, University of Munich Medical Centre, Munich, Germany; <sup>4</sup>Faculty of Medicine, Sydney University, Sydney, Australia; <sup>5</sup>Department of Medical Biology, Faculty of Medicine; and <sup>6</sup>Faculty of Community Medicine, Universitas Indonesia, Jakarta, Indonesia; <sup>7</sup>Departments of Pediatrics and <sup>8</sup>Clinical Pathology, Faculty of Medicine, Universitas Indonesia, Dr Ciptomangunkusumo National Referral Hospital, Jakarta, Indonesia; <sup>9</sup>Department of Pediatrics, Faculty of Medicine, Universitas Airlangga, Dr Sutomo General Hospital, Surabaya, Indonesia; and <sup>10</sup>Research Unit of Molecular Epidemiology, Institute of Epidemiology II, Helmholtz Zentrum Muenchen, Munich, Germany

## ABSTRACT

**Background:** Adequate availability of long-chain polyunsaturated fatty acids (LC-PUFAs) is important for human health from pregnancy to adulthood. Previous studies on fatty acid desaturase (*FADS*) gene single-nucleotide polymorphisms (SNPs) have been performed predominantly in Western populations and showed that *FADS* SNPs had a marked impact on LC-PUFA composition in blood and tissues.

**Objectives:** We aimed to investigate the influence of fetal *FADS* genotypes on LC-PUFA composition in umbilical artery plasma lipids in Indonesian infants.

**Design:** We performed a cross-sectional study to assess for these associations.

**Results:** A total of 12 cord plasma n-6 ( $\omega$ -6) and n-3 ( $\omega$ -3) fatty acids were analyzed for associations with 18 *FADS* gene cluster SNPs from 390 women with single parturition from the Indonesian Prospective Study of Atopic Dermatitis in Infants (ISADI). Fetal *FADS* genotypes influenced cord plasma LC-PUFA composition, but, in contrast to previous studies from Western populations, the quantitatively predominant SNPs were associated with lower LC-PUFA content.

**Conclusion:** To our knowledge, this study was the first in South East Asia on *FADS* genotypes and arterial cord blood fatty acids to show an association between fetal LC-PUFA composition and fetal *FADS* SNPs. The *FADS* genotype distribution differs markedly between different geographical populations. This trial was registered at [clinicaltrials.gov](http://clinicaltrials.gov) as NCT02401178. *Am J Clin Nutr* 2018;108:1135–1144.

**Keywords:** *FADS1*, *FADS2*, *FADS3*, polymorphisms, fetus, LC-PUFA composition, Indonesia

## INTRODUCTION

Adequate availability of essential PUFAs and the long-chain PUFA (LC-PUFA) metabolites of both the n-6 and n-3 (families is important in every stage of human life from pregnancy to adulthood, because these fatty acids modulate motor and cognitive function, mental health, cardiovascular function, and inflammatory responses, such as allergies and eczema (1, 2). However, studies on the effects and the magnitude of these effects remain inconclusive (3–5). For example, the authors of a meta-analysis that explored the effects of LC-PUFA supply in infant formula on vision and neurodevelopment concluded that although some studies showed significant benefits, overall, no significant effect was detectable (4). With regards to allergies, limited evidence supports a benefit of maternal n-3 LC-PUFA intake during pregnancy or lactation for reducing allergic disease in children (5). Caution is needed when interpreting current evidence, as many of the included trials had methodologic

Supported by the Mead Johnson Pediatric Nutrition Institute (grant 8670 to CT) and by the Commission of the European Communities with the European Research Council Advanced Grant META-GROWTH (ERC-2012-AdG 322605). The funders had no influence on the design, implementation, analysis and interpretation of the data.

Address correspondence to DRS (e-mail: [djarif1959@gmail.com](mailto:djarif1959@gmail.com)).

Abbreviations used: A, adrenic acid; ALA,  $\alpha$ -linolenic acid; ALSPAC study, Avon Longitudinal Study of Parents and Children; ARA, arachidonic acid; DGLA, dihomo- $\gamma$ -linolenic acid; *FADS*, fatty acid desaturase; FAME, fatty acid methyl ester; HWE, Hardy-Weinberg equilibrium; ISADI, Indonesian Prospective Study of Atopic Dermatitis of Infants; LA, linoleic acid; LC-PUFA, long-chain PUFA; MAF, minor allele frequency; SNP, single-nucleotide polymorphism.

Received March 20, 2018. Accepted for publication June 28, 2018.

First published online November 23, 2018; doi: <https://doi.org/10.1093/ajcn/nqy166>.

limitations such as small sample sizes, high attribution rates, and a lack of intention-to-treat analyses (3).

Studies published in the last decade have demonstrated a marked impact of single-nucleotide polymorphisms (SNPs) in the fatty acid desaturase (*FADS*) gene clusters on the activity of *FADS* enzymes, as well as precursor PUFA and LC-PUFA levels in blood and tissue (6–15). Desaturase enzymes convert precursor PUFAs, namely linoleic acid (LA) and  $\alpha$ -linolenic acid (ALA), through a desaturation and elongation process, to LC-PUFA metabolites such as arachidonic acid (ARA), EPA, and DHA. In European populations, minor allele carriers of *FADS* SNPs had increased substrate amounts (LA and ALA) and decreased product amounts (ARA, EPA, and DHA), which are reflective of reduced desaturase enzyme activity, as shown in numerous studies involving a wide range of subjects (69–4457), and genome-wide associations (9–14, 16, 17).

As most previous studies have been conducted in European populations (9–14), we studied variants of the *FADS1*, *FADS2*, and *FADS3* genes in Indonesian newborns and their relation to LC-PUFA composition in umbilical artery plasma. Our primary endpoint was to characterize the impact of genetic variation in the *FADS1–3* gene clusters on plasma LC-PUFA composition in cord blood. The secondary endpoints were the characterization of the frequency of *FADS1*, *FADS2*, and *FADS3* gene SNPs in the Indonesian population, their relation to the fatty acid composition of umbilical artery plasma lipids, and differences in minor alleles among various populations.

## METHODS

### Study area and population

We performed a cross-sectional analysis among 390 mother-infant pairs from the Indonesian Prospective Study of Atopic Dermatitis in Infants (ISADI). The primary endpoint of the ISADI research was to assess the role of filaggrin (*FLG*) gene mutation, *FADS1/FADS2* gene polymorphisms, and LC-PUFAs in atopic dermatitis patients (18). Our investigation of *FADS1/FADS2* gene polymorphisms and LC-PUFAs in all infants is complete, but genetic exploration of the *FLG* gene is not yet complete. Based on the available data, we explored the impact of *FADS* gene cluster SNPs on the LC-PUFA composition of cord blood lipids.

We used a consecutive sampling procedure to collect participants. The number of participants was calculated using the formula of categorical descriptive sample size. The study was conducted at Kemayoran Primary Health Care, central Jakarta, from April 2014 to December 2015. Participants were apparently healthy newborn infants whose parents agreed to participate in the study, and provided written informed consent.

Inclusion criteria were apparently healthy newborns born at term (37–42 wk of gestation) with a birth weight  $\geq 2500$  g, to mothers with normal gestational history and who had omnivorous diets.

Exclusion criteria were serious congenital anomalies or diseases in the infant, maternal gestational hypertension, gestational diabetes mellitus, or other severe illness, disorders, or the intake of n-3 and n-6 supplements by the mother during pregnancy and breastfeeding, or by the infant.

## Ethics

The present study was conducted according to the International Ethical Guidelines for Biomedical Research Involving Human Subjects. All procedures were approved by the Permanent Medical Research Ethics Committee in Medicine and Health/Faculty of Medicine Universitas Indonesia/Dr Cipto Mangunkusumo Hospital (47/H2.F1/ETIK/2014, extended by letter 148/UN2.F1/ETIK/2015). Written informed consent was obtained from all parents of the study subjects. The study was registered at [clinicaltrials.gov](http://clinicaltrials.gov) as NCT02401178.

## Procedures

### Sample collection and handling

Umbilical artery blood specimens were collected into EDTA tubes directly after birth, but before detachment of the placenta. Specimens were immediately centrifuged. The plasma and buffy coat were frozen at  $-80^{\circ}\text{C}$ , transported on dry ice to LMU, Munich, Germany, and stored at  $-80^{\circ}\text{C}$  until further analysis.

### LC-PUFA measurements

Plasma glycerophospholipid fatty acids were quantified as described previously (18). Briefly, selective preparation of methyl ester derivatives of glycerophospholipid fatty acids was achieved by coprecipitation of triacylglycerols and cholesterol esters with proteins and base-catalyzed transesterification excluding methyl ester synthesis from nonesterified fatty acids. Aliquots (100  $\mu\text{L}$ ) of EDTA plasma were combined with methanol containing dipentadecanoyl phosphatidylcholine as an internal standard. Tubes were centrifuged ( $3030 \times g$ ; 10 min;  $4^{\circ}\text{C}$ ) and the supernatant transferred to another vial. Sodium methoxide solution was added for fatty acid methyl ester (FAME) synthesis from glycerophospholipid fatty acids at room temperature. After 4 min, the reaction was stopped with methanolic HCl, and the FAME was extracted into 1 mL hexane for gas chromatographic analysis. The FAME was quantified by gas chromatography with flame ionization detection (Agilent 7890 GC; Agilent, Waldbronn, Germany), using a 50-m, 0.22-mm inner diameter BPX70 column (SGE, Weiterstadt, Germany). Injection was performed with a programmable temperature vaporizer (Gerstel, M $\ddot{u}$ hlheim, Germany) to avoid preconcentration of hexane extracts before gas chromatography. FAME peaks were identified and calibrated relative to pentadecanoic acid methyl ester (internal standard) by comparison with a FAME standard mixture (GLC-569B; Nu-Check Prep, Inc., Elysian, MN). Agilent Chemstation (revision B.04.03) was used for peak integration. Interassay coefficients of variation were determined by analysis of multiple aliquots of a sample distributed within the analysis of study subjects. Fatty acid results are reported as a percentage of total fatty acids analyzed (% wt/wt) (19). In this analysis, 12 PUFA measurements were included, namely LA (C18:2n-6),  $\gamma$ -linolenic acid (C18:3n-6), eicosadienoic acid (C20:2n-6), dihomo- $\gamma$ -linolenic acid (DGLA/C20:3n-6), ARA (C20:4n-6), adrenic acid (C22:4n-6), docosapentaenoic acid (DPA, C22:5n-6 and C22:5n-3), ALA (C18:3n-3), eicosatrienoic acid (C20:3n-3), eicosapentaenoic acid (EPA, C20:5n-3), and docosahexaenoic acid (DHA, C22:6n-3).



**TABLE 1**  
Fatty acid content (wt%) of cord plasma glycerophospholipids<sup>1</sup>

| Fatty acid  | Girls<br>(n = 190) | Boys<br>(n = 200) | P      | Total<br>(n = 390) |
|-------------|--------------------|-------------------|--------|--------------------|
| n-6         |                    |                   |        |                    |
| 18:2 (LA)   | 8.62 ± 1.15        | 8.74 ± 1.17       | 0.297  | 8.68 ± 1.16        |
| 18:3 (GLA)  | 0.16 ± 0.03        | 0.16 ± 0.04       | 0.678  | 0.16 ± 0.04        |
| 20:2 (EDA)  | 0.49 ± 0.11        | 0.50 ± 0.13       | 0.262  | 0.49 ± 0.12        |
| 20:3 (DGLA) | 5.63 ± 0.91        | 5.60 ± 0.80       | 0.727  | 5.62 ± 0.85        |
| 20:4 (ARA)  | 15.71 ± 1.56       | 15.44 ± 1.71      | 0.101  | 15.57 ± 1.65       |
| 22:4 (DTA)  | 0.65 ± 0.12        | 0.64 ± 0.11       | 0.576  | 0.64 ± 0.12        |
| 22:5 (DPA)  | 1.27 ± 0.34        | 1.29 ± 0.33       | 0.448  | 1.28 ± 0.33        |
| 20:3/18:2   | 0.66 ± 0.12        | 0.65 ± 0.13       | 0.445  | 0.66 ± 0.12        |
| 20:4/20:3   | 2.89 ± 0.71        | 2.84 ± 0.68       | 0.458  | 2.87 ± 0.69        |
| 20:4/18:2   | 1.86 ± 0.35        | 1.81 ± 0.38       | 0.144  | 1.83 ± 0.37        |
| n-3         |                    |                   |        |                    |
| 18:3 (ALA)  | 0.05 ± 0.03        | 0.06 ± 0.02       | 0.023* | 0.05 ± 0.03        |
| 20:3 (ERA)  | 0.13 ± 0.02        | 0.14 ± 0.02       | 0.156  | 0.14 ± 0.02        |
| 20:5 (EPA)  | 0.12 ± 0.07        | 0.11 ± 0.05       | 0.162  | 0.11 ± 0.06        |
| 22:5 (DPA)  | 0.24 ± 0.10        | 0.22 ± 0.08       | 0.180  | 0.23 ± 0.09        |
| 22:6 (DHA)  | 5.17 ± 1.24        | 5.06 ± 1.02       | 0.342  | 5.12 ± 1.14        |
| 20:5/18:3   | 2.49 ± 1.51        | 2.05 ± 1.08       | 0.001* | 2.26 ± 1.32        |

<sup>1</sup>Values of the wt% are means ± SDs. P value is from t test for the difference in measurements between girls and boys. \*Significant differences, P < 0.05. Enzyme activity of D5D was determined by the ratio of ARA/DGLA. Enzyme activity of D6D was determined by the ratio of DGLA/LA. ALA, α-linolenic acid; ARA, arachidonic acid; DGLA, dihomo-γ-linolenic acid; DPA, docosapentaenoic acid; DTA, docosatetraenoic acid; EDA, eicosadienoic acid; ERA, eicosatrienoic acid; GLA, γ-linolenic acid; LA, linoleic acid.

### Genotyping

*FADS1*, *FADS2*, and *FADS3* SNP analysis was performed at the Research Unit of Molecular Epidemiology at Helmholtz Zentrum Munich, Germany, as previously described (20). DNA was extracted from the buffy coat of umbilical artery blood by the Puregene DNA isolation kit (Gentra Systems). Genotyping was performed using iPLEX Gold Chemistry (Sequenom) and matrix-assisted laser desorption ionization-time of flight mass spectrometry, with methods to detect allelic differences. In brief, locations containing certain SNPs were amplified by polymerase chain reaction, using specific primers. After deactivation by alkaline phosphatase, single-base elongation was performed in accordance to the print order. After salt ion removal by ion switch and elongation reaction, the specimen was transferred to a silicone chip and covered with 3-hydroxypicolinic acid. The differences from specific alleles were measured by matrix-assisted laser desorption ionization-time of flight mass spectrometry. Allele recognition from SNPs was performed by Mass ARRAY Typer version 4.0.5 (Sequenom). SNPs for *FADS* genes were selected based on 3 criteria: 1) the SNP has been studied in previous publications; 2) the SNP candidates being considered are SNPs that have already been shown to be associated with LC-PUFA status; and 3) minor allele frequency (MAF) >10%.

### Statistical analysis

All data processing was performed with SAS version 9.4 (SAS Institute Inc, NC), or R 3.2.4 (<http://www.r-project.org>). Allele frequency, Hardy-Weinberg equilibrium (HWE), and linkage disequilibrium were assessed with R-package Genetics (<https://cran.r-project.org/web/packages/genetics/index.html>). The linkage disequilibrium test for each pair of SNP loci was tested with Lewontin's D'. Normal distribution of fatty acid

was assessed by the Kolmogorov-Smirnov test. Regression analysis was performed to explore the associations between the genetic variation of *FADS1* to *FADS3* genes and LC-PUFA composition (21). Reported P values in tables are uncorrected for multiple testing, with \*\*\* indicating significance after Bonferroni correction to account for multiple testing (critical corrected P value is  $P < 1.84 \times 10^{-4}$ , i.e., usual  $\alpha$  of 0.05 divided by the number of tests = 17 SNPs × 16 fatty acid outcomes).

### RESULTS

A total of 390 healthy infants, 190 girls and 200 boys, who met the inclusion criteria were enrolled in this study. The mean ± SD birth weights were 3022 ± 337.2 g for girls and 3132.1 ± 399 g for boys. The mean Apgar scores were 9 at 1 min, and 10 at 5 min. The average maternal age of delivery was 27.5 y. Most parents came from low socioeconomic backgrounds with senior high school education.

No statistical differences in the fatty acids were found between boys and girls for any of the n-6 and n-3 fatty acids, except for ALA and the ratio of EPA to ALA (Table 1). The ratios of ARA to LA, as well as EPA and DHA to ALA, were calculated as markers of the activity of fatty acid desaturation mediated by the enzymes D5D and D6D, respectively. The ratio of DGLA to LA is used to indicate the D6D activity, whereas D5D activity is reflected by the ratio of ARA to DGLA.

The distribution of all 18 *FADS1*, *FADS2*, and *FADS3* SNPs genotyped in this study population is presented in Table 2. Genotyping was done along a 34-kb portion from position 618,393,876 (rs174548) to position 61,888,645 (rs174455). The positions in base pairs were defined according to genome build 108, assembly GRCH38.p2 (<https://www.ncbi.nlm.nih.gov/snp>). This lists major and minor *FADS* alleles, MAF, and P values of the

**TABLE 2**  
Characteristics of SNPs analyzed in the *FADS1–FADS3* gene clusters in Indonesian infants<sup>1</sup>

| Gene         | dbSNP build 108 | Position, bp | Major/minor alleles | MAF, % | HWE <i>P</i> value |       |
|--------------|-----------------|--------------|---------------------|--------|--------------------|-------|
| <i>FADS1</i> | rs174548        | 61,803,876   | C/G                 | 27     | 0.696              |       |
|              | rs174556        | 61,813,163   | A/G                 | 27     | 0.696              |       |
|              | rs174561        | 61,815,236   | G/A                 | 27     | 0.696              |       |
| Intergenic   | rs3834458       | 61,827,449   | del/T               | 22     | 0.456              |       |
|              | rs968567        | 61,828,092   | C/T                 | NA     | NA                 |       |
| <i>FADS2</i> | rs174570        | 61,829,740   | T/C                 | 23     | 0.477              |       |
|              | rs174574        | 61,832,870   | A/C                 | 22     | 0.548              |       |
|              | rs2727271       | 61,835,886   | A/T                 | 25     | 0.685              |       |
|              | rs174576        | 61,836,038   | A/C                 | 22     | 0.368              |       |
|              | rs174578        | 61,838,027   | A/T                 | 22     | 0.368              |       |
|              | rs174579        | 61,838,141   | C/T                 | 28     | 1.000              |       |
|              | rs174602        | 61,856,942   | C/T                 | 41     | 0.465              |       |
|              | rs498793        | 61,857,233   | C/T                 | 15     | 0.421              |       |
|              | rs526126        | 61,857,413   | C/G                 | 22     | 0.056              |       |
|              | rs174575        | 61,834,531   | C/G                 | 30     | 1.000              |       |
|              | Intergenic      | rs174448     | 61,872,101          | T/C    | 48                 | 0.760 |
|              |                 | rs174449     | 61,872,907          | C/T    | 47                 | 0.186 |
|              | <i>FADS3</i>    | rs174455     | 61,888,645          | C/T    | 45                 | 0.474 |

<sup>1</sup>dbSNP, database single-nucleotide polymorphism; del., deleted; *FADS*, fatty acid desaturase; HWE, Hardy-Weinberg equilibrium; MAF, minor allele frequency; NA, not available.

HWE tests. One of the SNPs (rs968567) was monomorphic in the study and, therefore, could not be analyzed further. The other 17 SNPs of the ISADI study were consistent with HWE. The SNPs rs174548, rs174556, and rs174561 had an almost identical MAF of 27% and were highly collinear (the linkage disequilibrium calculation gave a *D'* value of 0.99999). Consequently, 14 SNPs were included in further analyses to avoid informational repetition.

**Table 3** compares the MAF of the *FADS* genes among different populations in Indonesia, the Mexican POSGRAD (Prenatal Omega-3 Supplementation on Child Growth and Development) Study (22), and Europe; data from the latter come from the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study, which included 9 countries (23), the British ALSPAC (Avon Longitudinal Study of Parents and Children) Study (20), the German LISA (Influences of Lifestyle-related Factors on the Immune System and the Development of Allergies in Childhood) Study, and the KOALA (“Kind, Ouders en gezondheid: Aandacht voor Leefstijl en Aanleg”) Birth Cohort Study in the Netherlands (8).

**Tables 4** and **5** list the regression coefficients ( $\beta$  coefficients), related *P* values, and explained variance ( $R^2$ ) of the regression of each fatty acid level, from n-6 and n-3 (% wt/wt), respectively, for each of the 14 analyzed SNPs of the *FADS* gene clusters. The composition of the n-6 fatty acids and ratios of DGLA to LA, ARA to DGLA, and ARA to LA were significantly associated with almost all SNPs of the *FADS1* and most of *FADS2* gene clusters, even after Bonferroni correction of *P* values for multiple testing. However, genetic variants of rs498793, rs526126, and rs174575 of the *FADS2* gene; rs174448 and rs174449 of the intergenic *FADS2* and *FADS3* gene; and rs174455 of the *FADS3* gene, showed no (or only rarely) statistically significant relations with any of the n-6 fatty acids, after correction for multiple testing.

The variance ( $R^2$ ) of fatty acid levels explained by SNPs was highest for DGLA, with 37.9% for genetic variants in the *FADS1* gene, 37.2% for rs3834458 (intergenic *FADS1/2*), and 35.9% for

rs174574 in the *FADS2* gene. The next highest  $R^2$  was 21.8% for regression of ARA in rs3834458 (intergenic *FADS1/2*), followed by ~20% for *FADS1* and some *FADS2* genetic variants.

The explained variance ( $R^2$ ) for the ARA:DGLA ratio was mainly determined by genetic variants of *FADS1* or intergenic *FADS1/2* and some *FADS2* genetic variants, accounting for 41.0% (rs3834458) of variation of the ratio. This finding supports the conclusion that the *FADS1* gene product is involved in D5-desaturation from DGLA to ARA. In addition, the *FADS2* gene product D6D is involved.

The high explained variance ( $R^2$ ) for the ARA:LA ratio for some *FADS2*-related SNPs of 25.5% (rs174574), and the *FADS1/2* intergenic SNP rs3834458 with 24.4%, provides further evidence that the *FADS2* gene product is involved in D6-desaturation from LA to ARA. In contrast, only small amounts of variance ranging from <1% to 2.5% of n-3 fatty acids were explained by *FADS* SNPs in this Indonesian infant population. The highest  $R^2$  of 2.5% was found for the *FADS1/2* intergenic SNP rs3834458 with regards to DHA level, followed by  $R^2 >2.0\%$  for some SNPs of the *FADS2* gene (rs174570 rs174574, rs174576, and rs174578). This finding provides further support that the *FADS2* gene product is involved in D6-desaturation. Interestingly, the explained variance for the EPA:ALA ratio was <1% for all SNPs, except for rs526126 (*FADS2*), with an  $R^2$  of 1.8%.

## DISCUSSION

Our study demonstrated a significant association of cord blood composition of precursor PUFAs and LC-PUFAs with common variants in the *FADS* gene clusters. In this population-based study, we found a higher content of ARA ( $15.57\% \pm 1.65\%$ ) and DHA ( $5.12\% \pm 1.14\%$ ) than other PUFAs in umbilical artery plasma glycerophospholipids. This finding is in agreement with those of the LISA Study (24), and reflects the active and preferential maternal-fetal transfer of LC-PUFAs across the human placenta (25, 26).

**TABLE 3**  
Comparison of common single-nucleotide polymorphisms among different populations<sup>1</sup>

| Gene         | dbSNP build 108 | ISADI    | MAF, % POSGRAD | MAF, % | ALSPAC | MAF, % | KOALA | MAF, % | LISA | MAF, % | HELENA | MAF, % |   |
|--------------|-----------------|----------|----------------|--------|--------|--------|-------|--------|------|--------|--------|--------|---|
| <i>FADS1</i> | rs174545        | —        | —              | —      | —      | —      | G     | 32.9   | G    | 31.1   | —      | —      |   |
|              | rs174546        | —        | —              | —      | —      | —      | T     | 32.9   | T    | 31     | T      | 31     |   |
|              | rs174548        | G        | 27             | G      | 22     | C      | 30.1  | —      | —    | —      | —      | —      |   |
|              | rs174556        | G        | 27             | G      | 24     | A      | 29.6  | T      | 30.7 | T      | 28.5   | —      |   |
|              | rs174561        | A        | 27             | A      | 24     | G      | 29.6  | C      | 30.7 | C      | 28.5   | —      |   |
| Intergenic   | rs3834458       | T        | 22             | NA     | NA     | del    | 33.0  | del    | 33.1 | del    | 31     | —      |   |
|              | rs968567        | T        | NA             | NA     | NA     | T      | 16.9  | —      | —    | —      | T      | 16     |   |
| <i>FADS2</i> | rs174570        | C        | 23             | C      | 27     | T      | 12.6  | —      | —    | —      | T      | 11     |   |
|              | rs174574        | C        | 22             | C      | 20     | A      | 34.0  | —      | —    | —      | —      | —      |   |
|              | rs2727271       | T        | 25             | T      | 40     | T      | 11.6  | —      | —    | —      | —      | —      |   |
|              | rs174576        | C        | 22             | C      | 21     | A      | 34.0  | —      | —    | —      | —      | —      |   |
|              | rs174578        | T        | 22             | T      | 21     | A      | 34.0  | —      | —    | —      | —      | —      |   |
|              | rs174579        | T        | 28             | T      | 38     | T      | 21.4  | —      | —    | —      | —      | —      |   |
|              | rs174602        | T        | 41             | T      | 37     | G      | 20.4  | —      | —    | —      | C      | 22     |   |
|              | rs498793        | T        | 15             | T      | 35     | T      | 39.9  | —      | —    | —      | C      | 40     |   |
|              | rs526126        | G        | 22             | NA     | NA     | G      | 18.1  | —      | —    | —      | G      | 20     |   |
|              | rs174575        | G        | 30             | G      | 40     | NA     | NA    | —      | —    | —      | —      | —      |   |
|              | Intergenic      | rs174448 | C              | 48     | T      | 29     | C     | 35.9   | —    | —      | —      | —      | — |
|              |                 | rs174449 | T              | 47     | T      | 28     | C     | 35.1   | —    | —      | —      | —      | — |
|              | <i>FADS3</i>    | rs174455 | T              | 45     | T      | 22     | C     | 34.8   | —    | —      | —      | —      | — |

<sup>1</sup>ALSPAC, Avon Longitudinal Study of Parents and Children; dbSNP, database single-nucleotide polymorphism; del., deleted; *FADS*, fatty acid desaturase; HELENA, Healthy Lifestyle in Europe by Nutrition in Adolescence; ISADI, Indonesian Study of Atopic Dermatitis of Indonesian Infants; KOALA, "Kind, Ouders en gezondheid: Aandacht voor Leefstijl en Aanleg" Birth Cohort Study; LISA, Influences of Lifestyle-related Factors on the Immune System and the Development of Allergies in Childhood Study; MAF, minor allele frequency; POSGRAD, Prenatal Omega-3 Supplementation on Child Growth and Development.

The extent to which the fetus can produce LC-PUFAs from precursors, i.e., from n-6 LA and n-3 ALA, is unknown. LC-PUFA formation is facilitated by both elongases encoded by *ELOVL* genes and by desaturases encoded by *FADS* genes (27). In our study, we found a close association between *FADS* SNPs and indicators of the activity of *FADS1*- and *FADS2*-encoded enzymes, i.e., blood levels of n-6 LC-PUFAs. As much as 20.4% of the ARA level variation was determined by the rs174548 SNP. Effects of similar size were also found for other SNPs. In contrast, Lattka et al. (20) showed that variations of *FADS* genes had a relatively low impact on LC-PUFA composition in cord venous plasma fatty acid, with a coefficient of variation of 2.99% determined by *FADS* SNPs. Umbilical artery samples may better reflect the effects of *FADS* SNP variation because the umbilical artery transports blood from the fetus back to the placenta.

We assessed 18 SNPs from the *FADS1*-3 gene clusters, but only 14 SNPs were included in further analyses, in order to avoid informational repetition. HWE in this study population had a value of >5%, which indicated that the population is in a constant state and not influenced by mutations, natural selection, nonrandom mating, genetic drift, or gene flow.

Indeed, we noted different *FADS1* and *FADS2* SNPs distributions in our Indonesian population compared with European populations. We then explored minor alleles in different populations. The major alleles of European populations (7, 8, 20, 23) were the minor alleles of the ISADI population. Comparing the minor SNP differences between the British ALSPAC Study (20) and the ISADI Study, we found a 70% difference for all SNPs, except for rs968567, rs2727271, rs174579, rs498793, and

rs526126. On the other hand, 14 out of 15 minor alleles examined were the same among participants of the Indonesian ISADI study and the Mexican POSGRAD Study (22) (namely, rs174548, rs174556, rs174561, rs174570, rs174574, rs2727271, rs174576, rs174578, rs174579, rs174602, rs498793, rs174575, rs174449, and rs174455).

In addition, we found that minor alleles in different populations were the same among participants of the Indonesian ISADI study and the Mexican POSGRAD Study (22) (namely, rs174548, rs174556, rs174561, rs174570, rs174574, rs2727271, rs174576, rs174578, rs174579, rs174602, rs498793, rs174575, rs174449, and rs174455). In addition, we found that minor alleles in different populations were associated with lower levels of substrates (LA and ALA) and higher levels of products (DGLA and ARA) and ratios of DGLA/LA, ARA/DGLA, and ARA/LA, indicating higher desaturase activity. In contrast, the minor alleles of European populations were associated with higher substrate and lower LC-PUFA levels, indicating lower desaturase activity (9, 13–17, 20, 23). The difference in the activity of *FADS* genes of different populations might be the source of heterogeneity in the meta-analysis, as it was not accounted for in almost all of the past trials (3).

Our findings begged the question as to the reason for great variations in the *FADS* gene clusters between populations. Our observations support the theories of natural selection (28–30) and human migration effects (31–33). Indonesia has the world's second-longest coastline. The coastline of the country is ~54,720 km long, hence, fish and other marine mammals are common food sources. This result was similar to the study of Fumagalli et al. (29) in an Inuit population. They found higher amounts of substrate n-3 fatty acids and a decrease in the amounts of EPA and DPA, with no significant effect on DHA. The similar changes could, therefore, be an effect of selection driven by environmental conditions, i.e., a habitual diet rich in LC-PUFAs



**TABLE 4**  
Regression of n-6 levels (wt% of cord plasma glycerophospholipids) on each of 14 SNPs of the *FADS1-3* regions<sup>†</sup>

|              |                    | Fatty acid   |                           |                        |                           |                           |                           |                          |                        |                           |                           |                           |       |                         |  |                             |  |                             |  |  |
|--------------|--------------------|--------------|---------------------------|------------------------|---------------------------|---------------------------|---------------------------|--------------------------|------------------------|---------------------------|---------------------------|---------------------------|-------|-------------------------|--|-----------------------------|--|-----------------------------|--|--|
|              |                    | ln(C18:3n-6) |                           | C20:2n-6               |                           | C20:3n-6                  |                           | C20:4n-6                 |                        | ln(C22:4n-6)              |                           | C22:5n-6                  |       | C20:3n-6<br>6(C18:2n-6) |  | ln(C20:4n-6<br>6(C18:2n-6)) |  | ln(C20:4n-6<br>6(C18:2n-6)) |  |  |
| <i>FADS1</i> | rs174548           | C18:2n-6     | -0.562                    | 0.062                  | -0.046                    | -0.835                    | 1.186                     |                          |                        | 0.055                     | 0.080                     | -0.053                    | 0.233 | 0.144                   |  |                             |  |                             |  |  |
|              |                    | coefficient  |                           |                        |                           |                           |                           |                          |                        |                           |                           |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | P value      | $6.97 \times 10^{-10}***$ | $5.28 \times 10^{-4}$  | $1.89 \times 10^{-6}***$  | $4.51 \times 10^{-42}***$ | $3.71 \times 10^{-21}***$ | $8.72 \times 10^{05}***$ | $2.42 \times 10^{-03}$ | $6.44 \times 10^{-08}***$ | $1.58 \times 10^{-15}***$ | $3.97 \times 10^{-22}***$ |       |                         |  |                             |  |                             |  |  |
| Intergenic   | rs3834458          | C18:2n-6     | 9.2                       | 2.8                    | 5.5                       | 37.9                      | 20.4                      | 3.7                      | 2.1                    | 7.1                       | 40.4                      | 21.3                      |       |                         |  |                             |  |                             |  |  |
|              |                    | coefficient  |                           |                        |                           |                           |                           |                          |                        |                           |                           |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | P value      | $7.39 \times 10^{-12}***$ | $1.66 \times 10^{03}$  | $2.1 \times 10^{-07}***$  | $4.72 \times 10^{-11}***$ | $1.33 \times 10^{-22}***$ | $5.38 \times 10^{05}***$ | $3.21 \times 10^{-04}$ | $1.71 \times 10^{-06}***$ | $2.49 \times 10^{-16}***$ | $1.97 \times 10^{-25}***$ |       |                         |  |                             |  |                             |  |  |
| <i>FADS2</i> | rs174570           | C18:2n-6     | 11.2                      | 2.3                    | 6.5                       | 37.2                      | 21.8                      | 3.9                      | 3.1                    | 5.5                       | 41.0                      | 24.4                      |       |                         |  |                             |  |                             |  |  |
|              |                    | coefficient  |                           |                        |                           |                           |                           |                          |                        |                           |                           |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | P value      | $5.8 \times 10^{-12}***$  | $1.35 \times 10^{03}$  | $1.06 \times 10^{-07}***$ | $6.23 \times 10^{-34}***$ | $1.2 \times 10^{-21}***$  | $5.05 \times 10^{-04}$   | $2.37 \times 10^{-03}$ | $6.06 \times 10^{-05}***$ | $5.29 \times 10^{-10}***$ | $5.7 \times 10^{-25}***$  |       |                         |  |                             |  |                             |  |  |
| rs174574     | C18:2n-6           | 11.3         | 2.4                       | 6.8                    | 31.6                      | 20.9                      | 2.8                       | 2.1                      | 3.8                    | 36.4                      | 23.9                      |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | coefficient  |                           |                        |                           |                           |                           |                          |                        |                           |                           |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | P value      | $3.44 \times 10^{-12}***$ | $2.27 \times 10^{03}$  | $3.9 \times 10^{-07}***$  | $2.06 \times 10^{-39}***$ | $3.45 \times 10^{-22}***$ | $1.74 \times 10^{04}***$ | $4.79 \times 10^{-04}$ | $5.39 \times 10^{-06}***$ | $1.26 \times 10^{-44}$    | $1.33 \times 10^{-25}***$ |       |                         |  |                             |  |                             |  |  |
| rs2727271    | C18:2n-6           | 0.355        | -0.022                    | 0.021                  | 0.403                     | -0.760                    | 1.297                     | 0.056                    | 0.099                  | 0.084                     | 0.230                     | 0.158                     |       |                         |  |                             |  |                             |  |  |
|              |                    | coefficient  |                           |                        |                           |                           |                           |                          |                        |                           |                           |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | P value      | $1.34 \times 10^{-05}***$ | $1.6 \times 10^{01}$   | $1.38 \times 10^{-02}$    | $9.54 \times 10^{-12}***$ | $2.79 \times 10^{-11}***$ | $1.96 \times 10^{-01}$   | $5.37 \times 10^{-02}$ | $4.52 \times 10^{-02}$    | $3.79 \times 10^{-15}***$ | $5.9 \times 10^{-12}***$  |       |                         |  |                             |  |                             |  |  |
| rs174576     | C18:2n-6           | 4.6          | 0.3                       | 1.3                    | 11.1                      | 10.6                      | 0.2                       | 0.7                      | 0.8                    | 14.6                      | 11.3                      |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | coefficient  |                           |                        |                           |                           |                           |                          |                        |                           |                           |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | P value      | $3.48 \times 10^{-12}***$ | $2.86 \times 10^{03}$  | $6.32 \times 10^{-07}***$ | $4.3 \times 10^{-38}***$  | $7.09 \times 10^{-21}***$ | $1.83 \times 10^{04}***$ | $3.15 \times 10^{-04}$ | $1.11 \times 10^{-05}***$ | $1.47 \times 10^{-12}***$ | $7.55 \times 10^{-25}***$ |       |                         |  |                             |  |                             |  |  |
| rs174579     | C18:2n-6           | 0.130        | -0.020                    | 0.018                  | 0.233                     | -0.140                    | 1.251                     | 0.056                    | 0.101                  | 0.046                     | 0.241                     | 0.161                     |       |                         |  |                             |  |                             |  |  |
|              |                    | coefficient  |                           |                        |                           |                           |                           |                          |                        |                           |                           |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | P value      | $1.63 \times 10^{-01}$    | $2.66 \times 10^{-01}$ | $5.54 \times 10^{-02}$    | $6.2 \times 10^{04}$      | $2.92 \times 10^{-01}$    | $6.02 \times 10^{-02}$   | $2.97 \times 10^{-01}$ | $6.54 \times 10^{-02}$    | $3.82 \times 10^{-03}$    | $1.42 \times 10^{-01}$    |       |                         |  |                             |  |                             |  |  |
| rs174602     | C18:2n-6           | 0.2%         | 0.1%                      | 0.7%                   | 2.7%                      | 0.0%                      | 0.0%                      | 0.7%                     | 0.0%                   | 0.6%                      | 1.9%                      | 0.3%                      |       |                         |  |                             |  |                             |  |  |
|              |                    | coefficient  |                           |                        |                           |                           |                           |                          |                        |                           |                           |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | P value      | $7.99 \times 10^{-03}$    | $1.85 \times 10^{-01}$ | $6.04 \times 10^{-02}$    | $3.71 \times 10^{-10}***$ | $6.56 \times 10^{-06}***$ | $2.96 \times 10^{-02}$   | $4.73 \times 10^{-02}$ | $1.64 \times 10^{-03}$    | $8.12 \times 10^{-11}***$ | $2.26 \times 10^{-05}***$ |       |                         |  |                             |  |                             |  |  |
| rs498793     | C18:2n-6           | 0.111        | -0.004                    | 0.028                  | 0.174                     | -0.311                    | 0.013                     | 0.013                    | 0.013                  | 0.013                     | 0.013                     | 0.013                     |       |                         |  |                             |  |                             |  |  |
|              |                    | coefficient  |                           |                        |                           |                           |                           |                          |                        |                           |                           |                           |       |                         |  |                             |  |                             |  |  |
|              |                    | P value      | $3.6 \times 10^{-01}$     | $8.67 \times 10^{-01}$ | $2.35 \times 10^{-02}$    | $4.99 \times 10^{-02}$    | $6.96 \times 10^{-02}$    | $5.14 \times 10^{-01}$   | $3.97 \times 10^{-01}$ | $3.07 \times 10^{-01}$    | $3.15 \times 10^{-02}$    | $1.27 \times 10^{-01}$    |       |                         |  |                             |  |                             |  |  |
|              | R <sup>2</sup> , % | 0.0          | -0.3                      | 1.1                    | 0.7                       | 0.6                       | -0.1                      | -0.1                     | 0.0                    | 0.9                       | 0.3                       |                           |       |                         |  |                             |  |                             |  |  |

(Continued)

TABLE 4  
(Continued)

|          | Fatty acid               |                          |                          |                          |                          |                          |                          |                          |                          |                   |                       |
|----------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|-------------------|-----------------------|
|          | C18:2n-6                 | ln(C18:3n-6)             | C20:2n-6                 | C20:3n-6                 | C20:4n-6                 | ln(C22:4n-6)             | C22:5n-6                 | C20:3n-6/C18:2n-6        | ln(C20:4n-6/C18:2n-6)    | C20:3n-6/C18:2n-6 | ln(C20:4n-6/C18:2n-6) |
| rs526126 | 0.056                    | -0.001                   | 0.008                    | 0.135                    | -0.424                   | 0.009                    | 0.022                    | 0.012                    | -0.053                   |                   |                       |
|          | 5.55 × 10 <sup>-01</sup> | 9.67 × 10 <sup>-01</sup> | 4. × 10 <sup>-01</sup>   | 5.53 × 10 <sup>-02</sup> | 1.68 × 10 <sup>-03</sup> | 5.23 × 10 <sup>-01</sup> | 4.21 × 10 <sup>-01</sup> | 2.45 × 10 <sup>-01</sup> | 5.02 × 10 <sup>-03</sup> |                   |                       |
|          | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              |                   |                       |
|          | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  |                   |                       |
|          | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       |                   |                       |
| rs174575 | -0.2                     | -0.3                     | -0.1                     | 0.7                      | 2.3                      | -0.2                     | -0.1                     | 0.1                      | 1.8                      |                   |                       |
|          | 0.138                    | -0.019                   | 0.015                    | 0.264                    | -0.136                   | -0.031                   | -0.040                   | 0.021                    | -0.059                   |                   |                       |
|          | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              |                   |                       |
|          | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  |                   |                       |
|          | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       |                   |                       |
|          | 0.3                      | 0.1                      | 0.4                      | 3.7                      | 0.0                      | 1.1                      | 0.3                      | 1.0                      | 2.4                      |                   |                       |
|          | 0.129                    | -0.009                   | 0.009                    | 0.133                    | -0.103                   | -0.023                   | -0.065                   | 0.004                    | -0.032                   |                   |                       |
|          | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              |                   |                       |
|          | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  |                   |                       |
|          | 1.18 × 10 <sup>-01</sup> | 5.79 × 10 <sup>-01</sup> | 2.95 × 10 <sup>-01</sup> | 2.94 × 10 <sup>-02</sup> | 3.8 × 10 <sup>-01</sup>  | 6.93 × 10 <sup>-02</sup> | 5.73 × 10 <sup>-03</sup> | 6.44 × 10 <sup>-01</sup> | 5.02 × 10 <sup>-02</sup> |                   |                       |
|          | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              |                   |                       |
|          | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  |                   |                       |
|          | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       |                   |                       |
|          | 0.4                      | -0.2                     | 0.0                      | 1.0                      | -0.1                     | 0.6                      | 1.7                      | -0.2                     | 0.7                      |                   |                       |
|          | -0.158                   | 0.015                    | -0.015                   | -0.206                   | 0.183                    | 0.021                    | 0.065                    | -0.009                   | 0.051                    |                   |                       |
|          | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              |                   |                       |
|          | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  |                   |                       |
|          | 4.99 × 10 <sup>-02</sup> | 3.44 × 10 <sup>-01</sup> | 7.25 × 10 <sup>-02</sup> | 4.85 × 10 <sup>-04</sup> | 1.11 × 10 <sup>-01</sup> | 8.95 × 10 <sup>-02</sup> | 4.72 × 10 <sup>-03</sup> | 2.85 × 10 <sup>-01</sup> | 1.5 × 10 <sup>-03</sup>  |                   |                       |
|          | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              |                   |                       |
|          | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  |                   |                       |
|          | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       |                   |                       |
|          | 0.7                      | 0.0                      | 0.6                      | 2.9                      | 0.4                      | 0.5                      | 1.8                      | 0.0                      | 2.3                      |                   |                       |
|          | -0.173                   | 0.007                    | -0.007                   | -0.162                   | 0.130                    | 0.018                    | 0.066                    | -0.004                   | 0.039                    |                   |                       |
|          | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              |                   |                       |
|          | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  |                   |                       |
|          | 3.5 × 10 <sup>-02</sup>  | 6.45 × 10 <sup>-01</sup> | 4.29 × 10 <sup>-01</sup> | 7.36 × 10 <sup>-03</sup> | 2.68 × 10 <sup>-01</sup> | 1.51 × 10 <sup>-01</sup> | 4.5 × 10 <sup>-03</sup>  | 6.66 × 10 <sup>-01</sup> | 1.75 × 10 <sup>-02</sup> |                   |                       |
|          | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              |                   |                       |
|          | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  |                   |                       |
|          | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       | R <sup>2</sup> , %       |                   |                       |
|          | 0.9                      | -0.2                     | -0.1                     | 1.6                      | 0.1                      | 0.3                      | 1.8                      | -0.2                     | 1.2                      |                   |                       |
|          | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              | coefficient              |                   |                       |
|          | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  | P value                  |                   |                       |
|          | 2.97 × 10 <sup>-02</sup> | 2.97 × 10 <sup>-02</sup> | 2.97 × 10 <sup>-02</sup> | 2.97 × 10 <sup>-02</sup> | 2.97 × 10 <sup>-02</sup> | 2.97 × 10 <sup>-02</sup> | 2.97 × 10 <sup>-02</sup> | 2.97 × 10 <sup>-02</sup> | 2.97 × 10 <sup>-02</sup> |                   |                       |

<sup>1</sup> Outcome is respective fatty acid wt% of cord plasma glycerophospholipids; ln indicates whether the outcome was log transformed to account for severe right skewness of the fatty acid distribution, or entered into the regression equation on the original scale. The β coefficient is the estimated regression coefficient from unadjusted linear regression of each fatty acid on each genetic variant of the FADS gene regions. The P values are uncorrected for multiple testing, with \*\*\*\* indicating significance after Bonferroni correction for multiple testing (critical P value is 1.84 × 10<sup>-4</sup>).



TABLE 5

Regression of n-3 levels (wt% of cord plasma glycerophospholipids) on each of 14 SNPs of *FADS1-3* regions<sup>1</sup>

|              |           |                     | Fatty acid             |                        |                        |                        |                        |                        |
|--------------|-----------|---------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
|              |           |                     | ln(C18:3n-3)           | ln(C20:3n-3)           | ln(C20:5n-3)           | ln(C22:5n-3)           | C22:6n-3               | ln(C20:5n-3/C18:3n-3)  |
| <i>FADS1</i> | rs174548  | $\beta$ coefficient | -0.076                 | 0.015                  | 0.012                  | 0.013                  | 0.0260                 | 0.088                  |
|              |           | <i>P</i> value      | $1.22 \times 10^{-2}$  | $2.21 \times 10^{-01}$ | $7.22 \times 10^{-01}$ | $6.75 \times 10^{-01}$ | $4.34 \times 10^{-03}$ | $3.31 \times 10^{-02}$ |
|              |           | $R^2$ , %           | 1.4                    | 0.1                    | -0.2                   | -0.2                   | 1.8                    | 0.9                    |
| Intergenic   | rs3834458 | $\beta$ coefficient | -0.096                 | 0.014                  | -0.012                 | -0.003                 | 0.318                  | 0.083                  |
|              |           | <i>P</i> value      | $2.83 \times 10^{-03}$ | $3.11 \times 10^{-01}$ | $7.31 \times 10^{-01}$ | $9.36 \times 10^{-01}$ | $9.65 \times 10^{-04}$ | $5.79 \times 10^{-02}$ |
|              |           | $R^2$ , %           | 2.0                    | 0.0                    | -0.2                   | -0.3                   | 2.5                    | 0.7                    |
| <i>FADS2</i> | rs174570  | $\beta$ coefficient | -0.091                 | 0.012                  | -0.001                 | -0.004                 | 0.302                  | 0.090                  |
|              |           | <i>P</i> value      | $3.95 \times 10^{-03}$ | $3.43 \times 10^{-01}$ | $9.8 \times 10^{-01}$  | $8.93 \times 10^{-01}$ | $1.45 \times 10^{-03}$ | $3.69 \times 10^{-02}$ |
|              |           | $R^2$ , %           | 1.9                    | 0.0                    | -0.3                   | -0.3                   | 2.3                    | 0.9                    |
|              | rs174574  | $\beta$ coefficient | -0.100                 | 0.012                  | -0.007                 | -0.010                 | 0.299                  | 0.093                  |
|              |           | <i>P</i> value      | $1.89 \times 10^{-03}$ | $3.89 \times 10^{-01}$ | $8.38 \times 10^{-01}$ | $7.63 \times 10^{-01}$ | $2.15 \times 10^{-03}$ | $3.55 \times 10^{-02}$ |
|              |           | $R^2$ , %           | 2.2                    | -0.1%                  | -0.2                   | -0.2                   | 2.2                    | 0.9                    |
|              | rs2727271 | $\beta$ coefficient | 0.070                  | -0.016                 | 0.000                  | 0.005                  | -0.178                 | -0.071                 |
|              |           | <i>P</i> value      | $8.34 \times 10^{-03}$ | $1.55 \times 10^{-01}$ | $9.97 \times 10^{-01}$ | $8.43 \times 10^{-01}$ | $2.73 \times 10^{-02}$ | $5.32 \times 10^{-02}$ |
|              |           | $R^2$ , %           | 1.5                    | 0.3                    | -0.3                   | -0.2                   | 1.0                    | 0.7                    |
|              | rs174576  | $\beta$ coefficient | -0.101                 | 0.010                  | -0.009                 | -0.008                 | 0.296                  | 0.093                  |
|              |           | <i>P</i> value      | $1.53 \times 10^{-03}$ | $4.43 \times 10^{-01}$ | $8.14 \times 10^{-01}$ | $7.98 \times 10^{-01}$ | $2.23 \times 10^{-03}$ | $3.4 \times 10^{-02}$  |
|              |           | $R^2$ , %           | 2.3                    | -0.1                   | -0.2                   | -0.2                   | 2.1                    | 0.9                    |
|              | rs174579  | $\beta$ coefficient | -0.001                 | 0.11                   | 0.004                  | 0.001                  | -0.033                 | 0.005                  |
|              |           | <i>P</i> value      | $9.76 \times 10^{-01}$ | $3.66 \times 10^{-01}$ | $9.09 \times 10^{-01}$ | $9.86 \times 10^{-01}$ | $7.2 \times 10^{-01}$  | $9.08 \times 10^{-01}$ |
|              |           | $R^2$ , %           | -0.3                   | 0.0                    | -0.3                   | -0.3                   | -0.2                   | -0.3                   |
|              | rs174602  | $\beta$ coefficient | -0.046                 | -0.012                 | -0.012                 | -0.028                 | 0.076                  | 0.034                  |
|              |           | <i>P</i> value      | $8.7 \times 10^{-02}$  | $2.76 \times 10^{-01}$ | $6.85 \times 10^{-01}$ | $2.9 \times 10^{-01}$  | $3.53 \times 10^{-01}$ | $3.58 \times 10^{-01}$ |
|              |           | $R^2$ , %           | 0.5                    | 0.0                    | -0.2                   | 0.0                    | 0.0                    | 0.0                    |
|              | rs498793  | $\beta$ coefficient | 0.034                  | 0.009                  | 0.036                  | 0.018                  | 0.043                  | 0.002                  |
|              |           | <i>P</i> value      | $3.92 \times 10^{-01}$ | $5.65 \times 10^{-01}$ | $4.19 \times 10^{-01}$ | $6.45 \times 10^{-01}$ | $7.17 \times 10^{-01}$ | $9.69 \times 10^{-01}$ |
|              |           | $R^2$ , %           | -0.1                   | -0.2                   | -0.1                   | -0.2                   | -0.2                   | -0.3                   |
|              | rs526126  | $\beta$ coefficient | 0.071                  | 0.020                  | -0.049                 | -0.005                 | -0.137                 | -0.120                 |
|              |           | <i>P</i> value      | $2.22 \times 10^{-02}$ | $1.15 \times 10^{-01}$ | $1.59 \times 10^{-01}$ | $8.84 \times 10^{-01}$ | $1.43 \times 10^{-01}$ | $4.41 \times 10^{-03}$ |
|              |           | $R^2$ , %           | 1.1                    | 0.4                    | 0.3                    | -0.3                   | 0.3                    | 1.8                    |
|              | rs174575  | $\beta$ coefficient | -0.001                 | 0.010                  | 0.015                  | 0.008                  | -0.016                 | 0.015                  |
|              |           | <i>P</i> value      | $9.85 \times 10^{-01}$ | $4.25 \times 10^{-01}$ | $6.55 \times 10^{-01}$ | $7.87 \times 10^{-01}$ | $8.61 \times 10^{-01}$ | $7.02 \times 10^{-01}$ |
|              |           | $R^2$ , %           | -0.3                   | -0.1                   | -0.2                   | -0.2                   | -0.3                   | -0.2                   |
| Intergenic   | rs174448  | $\beta$ coefficient | 0.129                  | -0.009                 | 0.009                  | 0.133                  | -0.103                 | -0.023                 |
|              |           | <i>P</i> value      | $1.18 \times 10^{-01}$ | $5.79 \times 10^{-01}$ | $2.95 \times 10^{-01}$ | $2.94 \times 10^{-02}$ | $3.8 \times 10^{-01}$  | $6.93 \times 10^{-02}$ |
|              |           | $R^2$ , %           | 0.4                    | -0.2                   | 0.0                    | 1.0                    | -0.1                   | 0.6                    |
| Intergenic   | rs174449  | $\beta$ coefficient | -0.020                 | 0.007                  | -0.044                 | -0.028                 | 0.006                  | -0.023                 |
|              |           | <i>P</i> value      | $4.38 \times 10^{-01}$ | $5.02 \times 10^{-01}$ | $1.37 \times 10^{-01}$ | $2.91 \times 10^{-01}$ | $9.38 \times 10^{-01}$ | $5.12 \times 10^{-01}$ |
|              |           | $R^2$ , %           | -0.1                   | -0.1                   | 0.3                    | 0.0                    | -0.3                   | -0.1                   |
| <i>FADS3</i> | rs174455  | $\beta$ coefficient | -0.024                 | -0.002                 | -0.053                 | -0.036                 | -0.007                 | -0.029                 |
|              |           | <i>P</i> value      | $3.77 \times 10^{-01}$ | $8.34 \times 10^{-01}$ | $7.75 \times 10^{-02}$ | $1.71 \times 10^{-01}$ | $9.34 \times 10^{-01}$ | $4.2 \times 10^{-01}$  |
|              |           | $R^2$ , %           | -0.1                   | -0.2                   | 0.5                    | 0.2                    | -0.3                   | -0.1                   |

<sup>1</sup> Outcome is respective fatty acid (wt% of cord plasma glycerophospholipids); ln indicates whether the outcome was log transformed to account for severe right skewness of the fatty acid distribution, or entered into the regression equation on the original scale.  $\beta$  Coefficient is the estimated regression coefficient from unadjusted linear regression of each fatty acid on each genetic variant of the *FADS* gene regions. The *P* values are uncorrected for multiple testing (critical *P* value is  $1.84 \times 10^{-4}$ ). *FADS*, fatty acid desaturase; SNP, single-nucleotide polymorphism.

that did not provide an advantage to individuals with a genetically rapid LC-PUFA synthesis (29). Agricultural diets, on the other hand, would have led to a higher consumption of grains and other plant-derived foods. Alleles that increase the rate of conversion of short-chain PUFAs to LC-PUFAs would, therefore, have been favored (30).

The 93% resemblance between Indonesian and Mexican *FADS* SNPs, and the 70% difference in the ALSPAC Study population, might support the theory of human migration known as "out of Africa" that divides ancestor migration from Africa, either

heading west to Europe or heading east to Asia and onwards to Latin America (31). Karafet et al. (32) found that the Y chromosome of the western Indonesian population was related to haplogroups that may have entered Indonesia during the Paleolithic era from mainland Asia, whereas a study by Moreno-Estrada et al. (33) stated that some Mexican ancestors originated from Eastern Asia. (41)

In addition to the association of *FADS1* and *FADS2* SNPs with LC-PUFA composition, we also assessed for a possible association between rs174455 from the *FADS3* gene and LC-

PUFA composition. No significant association was revealed, but there was a trend in the same direction as *FADS1* and *FADS2*. This finding was similar to a study by Koletzko et al. (14), who found a weak association between rs174455 and changes in substrates and products. Two special SNPs, rs2727271 and rs498793, showed effects in the opposite direction compared to other SNPs; they decreased products, but increased substrates. Findings for these 2 SNPs were also reported by Lattka et al. (20): with only 1 minor allele from rs2727271, there was a 0.76% decrease in ARA level. In addition, the 1 minor allele from rs498793 led to a 0.31% decrease in ARA level.

We found no significant associations between *FADS 1–3* SNPs and n–3 LC-PUFA composition. It appears that maternal dietary EPA and DHA have a greater effect on the cord blood levels of LC-PUFAs than endogenous synthesis.

Sufficient availability of LC-PUFAs is considered essential for early development. In this study, polymorphisms in the *FADS* gene cluster were shown to modify the fatty acid composition of glycerophospholipid plasma of umbilical arterial blood. We can conclude that polymorphisms in fetal *FADS* genes are associated with fetal n–6 LC-PUFA synthesis, in addition to effects of placental transfer of preformed LC-PUFAs. When comparing SNPs of different populations, the similarity of the *FADS1–3* SNP distribution patterns in this Indonesian and a previously studied Mexican population (22) are noteworthy, as both differed from European populations. Both the Indonesian and the Mexican populations showed the minor alleles to be associated with high levels of LC-PUFA synthesis, whereas the major genotypes were associated with low levels of LC-PUFA synthesis. Therefore, the supply of preformed LC-PUFA may be particularly important in these populations to achieve similar plasma and tissue concentrations, as well as related biological effects.

As the major alleles of these populations differ and each SNP differs in outcome, direct extrapolation from observational and intervention studies in European populations to Indonesian populations appears questionable, due to their genetically determined differences in PUFA metabolism. Hence, intervention studies in infant populations in Indonesia or other populations with similar genotype distributions seem necessary.

The authors' contributions were as follows—CT, BK, and DRS: conceived and designed the experiments; CT, PR, MM, HS, SRH, BK, and DRS: analyzed the data; ER: performed the genotyping; HD: examined the PUFAs; CT, PR, BK, DRS, SRH, MM, ZM, HS, SI, and RI: wrote the article; and all authors: read and approved the final manuscript. All the authors declare no conflict of interest regarding this manuscript.

## REFERENCES

- Lattka E, Illig T, Heinrich J, Koletzko B. Do FADS genotypes enhance our knowledge about fatty acid related phenotypes? *Clin Nutr* 2010;29:277–87.
- Lattka E, Illig T, Heinrich J, Koletzko B. FADS gene cluster polymorphisms: important modulators of fatty acid levels and their impact on atopic diseases. *J Nutrigenet Nutrigenomics* 2009;2:119–28.
- Koletzko B. Human milk lipids. *Ann Nutr Metab* 2016;69:28–40.
- Jasani B, Simmer K, Patole SK, Rao SC. Long chain polyunsaturated fatty acid supplementation in infants born at term. *Cochrane Database Syst Rev* 2017;3:CD000376.
- Gunaratne AW, Makrides M, Collins CT. Maternal prenatal and/or postnatal n-3 long chain polyunsaturated fatty acids (LCPUFA) supplementation for preventing allergies in early childhood. *Cochrane Database Syst Rev* 2015;7:CD010085.
- Cho HP, Nakamura MT, Clarke SD. Cloning, expression and nutritional regulation of the mammalian delta-6 desaturase. *J Biol Chem* 1999;274:471–7.
- Rzehak P, Heinrich J, Klopp N, Schaeffer L, Hoff S, Wolfram G, Illig T, Linselsen J. Evidence for an association between genetic variants of the fatty acid desaturase 1 fatty acid desaturase 2 (*FADS1 FADS2*) gene cluster and the fatty acid composition of erythrocyte membranes. *Br J Nutr* 2009;101:20–6.
- Rzehak P, Thijs C, Standl M, Mommers M, Glaser C, Jansen E, Klopp N, Kopperman GH, Singmann P, Postma DS, et al. Variants of the *FADS1 FADS2* gene cluster, blood levels of polyunsaturated fatty acids and eczema in children within the first 2 years of life. *PLoS One* 2010;5:e13261.
- Koletzko B, Lien E, Agostoni C, Bohles H, Campoy C, Cetin I, Decsi T, Dudenhausen JW, Dupont C, Forsyth S, et al. The roles of long-chain polyunsaturated fatty acids in pregnancy, lactation and infancy: review of current knowledge and consensus recommendation. *J Perinat Med* 2008;36:5–14.
- Schaeffer L, Gohlke H, Muller M, Heid IM, Palmer LJ, Kompauer E, Demmelmeir H, Illig T, Koletzko B, Heinrich J. Common genetic variants of the *FADS1 FADS2* gene cluster and their reconstructed haplotypes are associated with the fatty acid composition in phospholipids. *Hum Mol Genet* 2006;15:1745–56.
- Xie L, Innis SM. Genetic variants of the *FADS1 FADS2* gene cluster are associated with altered (n-6) and (n-3) essential fatty acids in plasma and erythrocyte phospholipids in women during pregnancy and in breast milk during lactation. *J Nutr* 2008;138:2222–8.
- Molto-Puigmarti C, Plat J, Mensink RP, Muller A, Jansen E, Zeegers MP, Thijs C. *FADS1 FADS2* gene variants modify the association between fish intake and the docosahexaenoic acid proportion in human milk. *Am J Clin Nutr* 2010;91:1368–010.
- Lattka E, Rzehak P, Szabo E, Jakobik V, Weck M, Weyermann M, Grallert H, Rothebacher D, Heinrich J, Brenner H, et al. Genetic variants in the *FADS* gene cluster are associated with arachidonic acid concentrations of human breast milk at 1.5 and 6 mo postpartum and influence the course of milk dodecanoic, tetraacosenoic, and trans-9-octadecenoic acid concentrations over the duration of lactation. *Am J Clin Nutr* 2011;93:382–91.
- Koletzko B, Lattka E, Zeilinger S, Illig T, Steer C. Genetic variants of the fatty acid desaturase gene cluster predict amounts of red blood cell docosahexaenoic and other polyunsaturated fatty acids in pregnant women: findings from the Avon Longitudinal Study of Parents and Children. *Am J Clin Nutr* 2011;93:211–9.
- Brenna JT, Salem N, Jr, Sinclair AJ, Cunnane SC.  $\alpha$ -Linolenic acid supplementation and conversion to n-3 long-chain polyunsaturated fatty acids in humans. *Prostaglandins Leukot Essent Fatty Acids* 2009;80:85–91.
- Tanaka T, Shen J, Abecasis GR, Kisialiou A, Ordovas JM, Guralnik JM, Singleton A, Bandinelli S, Cherubini A, Arnett D, et al. Genome-wide associations study of plasma polyunsaturated fatty acids in the InCHIANTI Study. *PLoS Genet* 2009;5:e1000338.
- Illig T, Gieger C, Zhai G, Romisch-Margl W, Wang-Sattler R, Prehn C, Altmaier E, Kastenmuller G, Kato B, Mewes HS, et al. A Genome-wide perspective of genetic variation in human metabolism. *Nat Genet* 2010;42:137–41.
- Tanjung C, Rzehak P, Mansyur M, Munasir Z, Sudoyo H, Immanuel S, Irawan R, Reischl E, Demmelmeir H, Koletzko B, et al. Study protocol to investigate the environmental and genetic aetiology of atopic dermatitis: the Indonesian Prospective Study of Atopic Dermatitis in Infants (ISADI). *BMJ Open* 2017;7:e012475.
- Glaser C, Rzehak P, Demmelmeir H, Klopp N, Heinrich J, Koletzko B. Influence of FADS polymorphisms on tracking of serum glycerophospholipid fatty acid concentrations and percentage composition in children. *PLoS ONE* 2011;6(7):e21933.
- Lattka E, Koletzko B, Zeilinger S, Hibbeln JR, Klopp N, Ring SM, Steer CD. Umbilical cord PUFA are determined by maternal and child fatty acid desaturase (*FADS*) genetic variants in the Avon Longitudinal Study of Parents and Children (ALSPAC). *Br J Nutr* 2013;109:1196–210.

21. Hosmer DW, Lemeshow S. Applied logistic regression. New York: John Wiley & Sons; 1989.
22. Gonzalez-Casanova I, Rzehak P, Stein AD, Feregrino RG, Dommarco JA, Barraza-Villareal A, Demmelmair H, Romieu I, Villalpando S, Martorell R, et al. Maternal single nucleotide polymorphisms in the fatty acid desaturase 1 and 2 coding regions modify the impact of prenatal supplementation with DHA on birth weight. *Am J Clin Nutr* 2016;103:1171–8.
23. Bokor S, Dumont J, Spinneker A, Gonzales-Gross M, Nova E, Widhalm K, Moschonis G, Stehle P, Amouvel P, De Henauw S, et al. Single nucleotide polymorphisms in the FADS gene cluster are associated with delta-5 and delta-6 desaturase activities estimated by serum fatty acid ratios. *J Lipid Res* 2010;51:2325–33.
24. Kohlboeck G, Glaser C, Tiesler C, Demmelmair H, Standl M, Romanos M, Koletzko B, Lehmann I, Heinrich J, LISAPlus study Group. Effect of fatty acid status in cord blood serum on children's behavioral difficulties at 10 y of age: results from LISAPlus study. *Am J Clin Nutr* 2011;94:1592–9.
25. Prieto-Sánchez MT, Ruiz-Palacios M, Blanco-Carnero JE, Pagan A, Hellmuth C, Uhl O, Peissner W, Ruiz-Alcaraz AJ, Parrilla JJ, Koletzko B, Larqué E. Placental MFSD2a transporter is related to decreased DHA in cord blood of women with treated gestational diabetes. *Clin Nutr* 2017;36:513–21.
26. Larqué E, Pagán A, Prieto MT, Blanco JE, Gil-Sánchez A, Zornoza-Moreno M, Ruiz-Palacios M, Gázquez A, Demmelmair H, Parrilla JJ, Koletzko B. Placental fatty acid transfer: a key factor in fetal growth. *Ann Nutr Metab* 2014;64:247–53.
27. Barman M, Nilsson S, Nalwai AT, Sandin A, Wold AE, Sandberg A-S. Single nucleotide polymorphisms in the FADS gene cluster but not the ELOVL2 gene are associated with serum polyunsaturated fatty acid composition and development of allergy (in a Swedish birth cohort). *Nutrients* 2015;7:10100–15.
28. Nielsen R. Molecular signatures of natural selection. *Annu Rev Genet* 2005;39:197–218.
29. Fumagalli M, Moltke I, Grarup N, Racimo F, Bjerregaard P, Jørgensen ME, Komeliussen TS, Gerbault P, Skotte L, Linneberg A, et al. Greenlandic Inuit show genetic signatures of diet and climate adaptation. *Science* 2015;349:1343–7.
30. Buckley MT, Racimo F, Allentoft ME, Jensen MK, Jonsson A, Huang H, Hormozdiari F, Sikora M, Marnetto D, Eskin E, et al. Selection in Europeans on fatty acid desaturases associated with dietary changes. *Mol Biol Evol* 2017;34:1307–18.
31. Klein RG. Darwin and the recent African origin of modern humans. *Proc Natl Acad Sci U S A* 2009;106:16007–9.
32. Karafet TM, Hallmark B, Cox MP, Sudoyo H, Downey S, Lansing JS, Hammer MF. Major east-west division underlies Y chromosome stratification across Indonesia. *Mol Biol Evol* 2010;27:1833–44.
33. Moreno-Estrada A, Gignoux CR, Fernández-López JC, Zakharia F, Sikora M, Contreras AV, Acuna-Alonzo V, Sandoval K, Eng C, Romero-Hidalgo S, et al. The genetics of Mexico recapitulates Native American substructure and affects biomedical traits. *Science* 2014;344:1280–5.

# The association of fatty acid desaturase gene polymorphisms on long-chain polyunsaturated fatty acid composition in Indonesian infants

## ORIGINALITY REPORT

**23%**  
SIMILARITY INDEX

**15%**  
INTERNET SOURCES

**22%**  
PUBLICATIONS

**0%**  
STUDENT PAPERS

## PRIMARY SOURCES

**1** [www.veneca.nl](http://www.veneca.nl) 1%  
Internet Source

**2** [novapublishers.com](http://novapublishers.com) 1%  
Internet Source

**3** [ihpmagazine.com](http://ihpmagazine.com) 1%  
Internet Source

**4** [diposit.ub.edu](http://diposit.ub.edu) 1%  
Internet Source

**5** [www.circumpolarhealthjournal.net](http://www.circumpolarhealthjournal.net) 1%  
Internet Source

**6** [www.efsa.europa.eu](http://www.efsa.europa.eu) 1%  
Internet Source

**7** [academic.oup.com](http://academic.oup.com) <1%  
Internet Source

**8** Hans Demmelmair, Talat Bashir Ahmed, Berthold Koletzko. "Content, variability, and" <1%



regulation of fatty acids in human milk",  
Elsevier BV, 2021

Publication

9

G. Kohlboeck. "Effect of fatty acid status in cord blood serum on children's behavioral difficulties at 10 y of age: results from the LISAPLUS Study", American Journal of Clinical Nutrition, 12/01/2011

Publication

<1 %

10

[www.theiem.org](http://www.theiem.org)

Internet Source

<1 %

11

Peter Rzehak. "Evidence for an association between genetic variants of the fatty acid desaturase 1 fatty acid desaturase 2 ( FADS1 FADS2) gene cluster and the fatty acid composition of erythrocyte membranes", British Journal Of Nutrition, 05/15/2008

Publication

<1 %

12

[2017.nutrition-growth.kenes.com](http://2017.nutrition-growth.kenes.com)

Internet Source

<1 %

13

Ding, Zhen, Guo-Liang Liu, Xiang Li, Xue-Yan Chen, Yi-Xia Wu, Can-Can Cui, Xi Zhang, Guang Yang, and Lin Xie. "Association of polyunsaturated fatty acids in breast milk with fatty acid desaturase gene polymorphisms among Chinese lactating mothers", Prostaglandins Leukotrienes and Essential Fatty Acids (PLEFA), 2016.

<1 %

14

Hans Demmelmair, Jannike Øyen, Tina Pickert, Astrid Rauh-Pfeiffer et al. "The effect of Atlantic salmon consumption on the cognitive performance of preschool children – A randomized controlled trial", Clinical Nutrition, 2018

Publication

---

<1 %

15

Cornelia Huth, Thomas Illig, Christian Herder, Christian Gieger et al. " Joint analysis of individual participants' data from 17 studies on the association of the variant -174G>C with circulating glucose levels, interleukin-6 levels, and body mass index ", Annals of Medicine, 2009

Publication

---

<1 %

16

Rasika A. Mathias, Vrindarani Pani, Floyd H. Chilton. "Genetic Variants in the FADS Gene: Implications for Dietary Recommendations for Fatty Acid Intake", Current Nutrition Reports, 2014

Publication

---

<1 %

17

[clinical-nutrition.imedpub.com](http://clinical-nutrition.imedpub.com)

Internet Source

---

<1 %

18

Gibson, Greg. "A Primer of Human Genetics", Oxford University Press

Publication

---

<1 %

19

Glaser, C.. "Role of FADS1 and FADS2 polymorphisms in polyunsaturated fatty acid metabolism", *Metabolism*, 201007

Publication

<1 %

20

[www.conferencecallsworld.com](http://www.conferencecallsworld.com)

Internet Source

<1 %

21

Juan Juan, Hongyan Huang, Xia Jiang, Andres V Ardisson Korat et al. "Joint effects of fatty acid desaturase 1 polymorphisms and dietary polyunsaturated fatty acid intake on circulating fatty acid proportions", *The American Journal of Clinical Nutrition*, 2018

Publication

<1 %

22

Kumar S. D. Kothapalli, , Kaixiong Ye, Maithili S. Gadgil et al. " Positive Selection on a Regulatory Insertion–Deletion Polymorphism in Influences Apparent Endogenous Synthesis of Arachidonic Acid ", *Molecular Biology and Evolution*, 2016

Publication

<1 %

23

Tejera, N., D. Vauzour, M. B. Betancor, O. Sayanova, S. Usher, M. Cochard, N. Rigby, N. Ruiz-Lopez, D. Menoyo, D. R. Tocher, J. A. Napier, and A. M. Minihane. "A Transgenic *Camelina sativa* Seed Oil Effectively Replaces Fish Oil as a Dietary Source of Eicosapentaenoic Acid in Mice", *Journal of Nutrition*, 2016.

<1 %

24

Hiroshi M Ueno, Satoshi Higurashi, Yuzuka Shimomura, Ryota Wakui et al. "Association of DHA Concentration in Human Breast Milk with Maternal Diet and Use of Supplements: A Cross-Sectional Analysis of Data from the Japanese Human Milk Study Cohort", Current Developments in Nutrition, 2020

Publication

<1 %

---

25

[nutritionandmetabolism.biomedcentral.com](https://nutritionandmetabolism.biomedcentral.com)

Internet Source

<1 %

---

26

[ehp.niehs.nih.gov](https://ehp.niehs.nih.gov)

Internet Source

<1 %

---

27

Bénédicte Renaville, Francesca Tulli, Massimiliano Bruno, Emilio Tibaldi, Maria Messina. " Fatty acid desaturase 2 ( ) insertion/deletion polymorphism impact on muscle fatty acid profile in European grayling ( ) ", British Journal of Nutrition, 2013

Publication

<1 %

---

28

Erlina Burhan, Ari Fahrial Syam, Ahmad Jabir Rahyussalim, Prasenohadi et al. "The emergence of COVID-19 in Indonesia: analysis of predictors of infection and mortality using independent and clustered data approaches", Cold Spring Harbor Laboratory, 2020

Publication

<1 %

---



29

Eva Lattka, Thomas Illig, Joachim Heinrich, Berthold Koletzko. "FADS Gene Cluster Polymorphisms: Important Modulators of Fatty Acid Levels and Their Impact on Atopic Diseases", *Journal of Nutrigenetics and Nutrigenomics*, 2009

Publication

&lt;1 %

30

Matthew T. Buckley, Fernando Racimo, Morten E. Allentoft, Majken K. Jensen et al. "Selection on the FADS region in Europeans", *Cold Spring Harbor Laboratory*, 2016

Publication

&lt;1 %

31

Rasika A. Mathias, Candelaria Vergara, Li Gao, Nicholas Rafaels et al. "FADS genetic variants and  $\omega$ -6 polyunsaturated fatty acid metabolism in a homogeneous island population", *Journal of Lipid Research*, 2010

Publication

&lt;1 %

32

[scholar.ui.ac.id](http://scholar.ui.ac.id)

Internet Source

&lt;1 %

33

"IUNS. 21st International Congress of Nutrition. Buenos Aires, Argentina, October 15-20, 2017: Abstracts", *Annals of Nutrition and Metabolism*, 2017

Publication

&lt;1 %

34

Mario Klingler, Hans Demmelmair, Berthold Koletzko, Claudia Glaser. "Fatty Acid Status

&lt;1 %

Determination by Cheek Cell Sampling  
Combined with Methanol-Based Ultrasound  
Extraction of Glycerophospholipids", Lipids,  
2011

Publication

---

35

[www.thieme-connect.com](http://www.thieme-connect.com)

Internet Source

<1 %

---

36

Giulia Cisbani, Alex Koppel, Dunja Knezevic,  
Ivonne Suridjan, Romina Mizrahi, Richard P.  
Bazinet. "Peripheral cytokine and fatty acid  
associations with neuroinflammation in AD  
and aMCI patients: An exploratory study",  
Brain, Behavior, and Immunity, 2020

Publication

<1 %

---

37

Agostoni, C.. "Reduced Docosahexaenoic Acid  
Synthesis May Contribute to Growth  
Restriction in Infants Born to Mothers Who  
Smoke", The Journal of Pediatrics, 200512

Publication

<1 %

---

38

Lahner, Edith, Giovanna Gentile, Flaminia  
Purchiaroni, Barbara Mora, Maurizio  
Simmaco, and Bruno Annibale. "Single  
nucleotide polymorphisms related to vitamin  
B12 serum levels in autoimmune gastritis  
patients with or without pernicious anaemia",  
Digestive and Liver Disease, 2015.

Publication

---

<1 %

39

Tanima De, Rita Christopher, Dindagur Nagaraja. "Influence of CYP2C9 polymorphism and phenytoin co-administration on acenocoumarol dose in patients with cerebral venous thrombosis", *Thrombosis Research*, 2014

Publication

---

<1 %

40

Tri Juli Edi Tarigan, Erni Hernawati Purwaningsih, Yusra Yusra, Murdani Abdullah et al. "Sambiloto (*Andrographis Paniculata*) Increases Glucagon-Like Peptide 1 Concentration in Prediabetes: A Double-Blind, Randomized, Placebo-Controlled Trial", *Research Square*, 2020

Publication

---

<1 %

41

Voruganti, V. Saroja, Paul B. Higgins, Sven O. E. Ebbesson, John Kennish, Harald H. H. GÅrning, Karin Haack, Sandra Laston, Eugene Drigalenko, Charlotte R. Wenger, William S. Harris, Richard R. Fabsitz, Richard B. Devereux, Jean W. MacCluer, Joanne E. Curr. "Variants in CPT1A, FADS1, and FADS2 are Associated with Higher Levels of Estimated Plasma and Erythrocyte Delta-5 Desaturases in Alaskan Eskimos", *Frontiers in Genetics*, 2012.

Publication

---

<1 %

42

Heidi AR Jensen, Laurine BS Harsløf, Maria S Nielsen, Line B Christensen et al. "FADS single-nucleotide polymorphisms are associated with behavioral outcomes in children, and the effect varies between sexes and is dependent on PPAR genotype", *The American Journal of Clinical Nutrition*, 2014

Publication

&lt;1 %

43

Matthew T. Buckley, Fernando Racimo, Morten E. Allentoft, Majken K. Jensen et al. "Selection in Europeans on Fatty Acid Desaturases Associated with Dietary Changes", *Molecular Biology and Evolution*, 2017

Publication

&lt;1 %

44

Niilo Färkkilä, Saku Torvinen, Harri Sintonen, Tiina Saarto, Heikki Järvinen, Juha Hänninen, Kimmo Taari, Risto P. Roine. "Costs of colorectal cancer in different states of the disease", *Acta Oncologica*, 2015

Publication

&lt;1 %

45

Solakivi, Tiina, Tarja Kunnas, Olli Jaakkola, Jaana Renko, Terho Lehtimäki, and Seppo T Nikkari. "Delta-6-desaturase gene polymorphism is associated with lipoprotein oxidation in vitro", *Lipids in Health and Disease*, 2013.

Publication

&lt;1 %



46

T. M. Karafet. "Major East-West Division Underlies Y Chromosome Stratification across Indonesia", *Molecular Biology and Evolution*, 08/01/2010

Publication

&lt;1 %

47

[jn.nutrition.org](http://jn.nutrition.org)

Internet Source

&lt;1 %

48

[pureadmin.qub.ac.uk](http://pureadmin.qub.ac.uk)

Internet Source

&lt;1 %

49

Yu, Howe-Ming, Marnie Newell, Kalpana Subedi, Randall J. Weselake, Vera Mazurak, and Catherine J. Field. "Effects of intermediates in the desaturation and elongation pathway of n-3 and n-6 fatty acids on the fatty acid composition of human breast cancer cells : Effects of polyunsaturated fatty acids on human breast cancer cells", *European Journal of Lipid Science and Technology*, 2015.

Publication

&lt;1 %

50

[genomemedicine.biomedcentral.com](http://genomemedicine.biomedcentral.com)

Internet Source

&lt;1 %

51

[helda.helsinki.fi](http://helda.helsinki.fi)

Internet Source

&lt;1 %

52

[pubs.sciepub.com](http://pubs.sciepub.com)

Internet Source

&lt;1 %

53

[www.nestlenutrition-institute.org](http://www.nestlenutrition-institute.org)

Internet Source

<1 %

---

54

"Diet, Nutrition, and Fetal Programming",  
Springer Science and Business Media LLC,  
2017

Publication

<1 %

---

55

Guilherme Eustáquio Furtado, Matheus Uba  
Chupel, Luciele Minuzzi, Miguel Patrício et al.  
"Exploring the potential of salivary and blood  
immune biomarkers to elucidate physical  
frailty in institutionalized older women",  
Experimental Gerontology, 2020

Publication

<1 %

---

56

Hellstrand, S., U. Ericson, B. Gullberg, B.  
Hedblad, M. Orho-Melander, and E.  
Sonestedt. "Genetic Variation in FADS Has  
Little Effect on the Association between  
Dietary PUFA Intake and Cardiovascular  
Disease", Journal of Nutrition, 2014.

Publication

<1 %

---

57

Markus Takkunen, Jyrki Ågren, Johanna  
Kuusisto, Markku Laakso, Matti Uusitupa,  
Ursula Schwab. "Dietary Fat in Relation to  
Erythrocyte Fatty Acid Composition in Men",  
Lipids, 2013

Publication

<1 %

---

58

Szilvia Bokor, Julie Dumont, Andre Spinneker, Marcela Gonzalez-Gross et al. "Single nucleotide polymorphisms in the FADS gene cluster are associated with delta-5 and delta-6 desaturase activities estimated by serum fatty acid ratios", Journal of Lipid Research, 2010

Publication

&lt;1 %

59

[bmcpregnancychildbirth.biomedcentral.com](http://bmcpregnancychildbirth.biomedcentral.com)

Internet Source

&lt;1 %

60

[coek.info](http://coek.info)

Internet Source

&lt;1 %

61

[doc.rero.ch](http://doc.rero.ch)

Internet Source

&lt;1 %

62

[id.123dok.com](http://id.123dok.com)

Internet Source

&lt;1 %

63

[tel.archives-ouvertes.fr](http://tel.archives-ouvertes.fr)

Internet Source

&lt;1 %

64

[www.nature.com](http://www.nature.com)

Internet Source

&lt;1 %

65

Karen Best, Maria Makrides. "Possible protective effect of prenatal omega-3 long-chain polyunsaturated fatty acids supplementation on persistent wheeze and asthma in early childhood", Evidence Based Medicine, 2017

Publication

&lt;1 %

|    |  |      |
|----|--|------|
| 66 | <a href="https://downloads.hindawi.com">downloads.hindawi.com</a><br>Internet Source   | <1 % |
| 67 | <a href="https://efsa.onlinelibrary.wiley.com">efsa.onlinelibrary.wiley.com</a><br>Internet Source   | <1 % |
| 68 | <a href="https://link.springer.com">link.springer.com</a><br>Internet Source   | <1 % |
| 69 | <a href="https://uowvivo.uow.edu.au">uowvivo.uow.edu.au</a><br>Internet Source   | <1 % |
| 70 | "Role of Long-Chain Omega-3 Fatty Acids in Cognitive and Emotional Development", Nutrition for Brain Health and Cognitive Performance, 2015.<br>Publication  | <1 % |
| 71 | Alicia Chemaly, Louis Arnould, Alassane Seydou, Pierre-Henry Gabrielle, Florian Baudin, Niyazi Acar, Catherine Creuzot-Garcher. "Plasma Fatty Acids and Primary Open-angle Glaucoma in the Elderly: the Montrachet Population-based Study", Research Square, 2021<br>Publication | <1 % |
| 72 | Beth T. Kao. "Mice raised on milk transgenically enriched with n-3 PUFA have increased brain docosahexaenoic acid", Lipids, 06/2006<br>Publication   | <1 % |

73

Dominique Brémond-Gignac. "Visual development in infants: physiological and pathological mechanisms :", Current Opinion in Ophthalmology, 04/2011

Publication

---

<1 %

74

G.M. Hon, S. Abel, C.M. Smuts, P. van, M.S. Hassan, S.J. van, R.T. Erasmus, T. Matsh. "Chapter 9 Gas Chromatography Results Interpretation: Absolute Amounts Versus Relative Percentages", IntechOpen, 2012

Publication

---

<1 %

75

Ning Ning Chung, Guy S. Jacobs, Herawati Sudoyo, Safarina G. Malik, Lock Yue Chew, J. Stephen Lansing, Murray P. Cox. "Sex-linked genetic diversity originates from persistent sociocultural processes at microgeographic scales", Royal Society Open Science, 2019

Publication

---

<1 %

76

Usha Ramakrishnan, Amanda Stinger, Ann M. DiGirolamo, Reynaldo Martorell, Lynnette M. Neufeld, Juan A. Rivera, Lourdes Schnaas, Aryeh D. Stein, Meng Wang. "Prenatal Docosahexaenoic Acid Supplementation and Offspring Development at 18 Months: Randomized Controlled Trial", PLOS ONE, 2015

Publication

---

<1 %



77

Usha Ramakrishnan, Ines Gonzalez-Casanova,  
Lourdes Schnaas, Ann DiGirolamo et al.

"Prenatal supplementation with DHA  
improves attention at 5 y of age: a  
randomized controlled trial", The American  
Journal of Clinical Nutrition, 2016

Publication

<1 %

78

[eprints.gla.ac.uk](http://eprints.gla.ac.uk)

Internet Source

<1 %

79

[hal.archives-ouvertes.fr](http://hal.archives-ouvertes.fr)

Internet Source

<1 %

80

[ods.od.nih.gov](http://ods.od.nih.gov)

Internet Source

<1 %

81

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<1 %

82

Agatha, G.. "Conjugated linoleic acid  
modulation of cell membrane in leukemia  
cells", Cancer Letters, 200406

Publication

<1 %

83

Antonio Gázquez, María Ruíz-Palacios, Elvira  
Larqué. "DHA supplementation during  
pregnancy as phospholipids or TAG produces  
different placental uptake but similar fetal  
brain accretion in neonatal piglets", British  
Journal of Nutrition, 2017

Publication

<1 %

84

Frédéric Destailats. "Differential effect of maternal diet supplementation with  $\alpha$ -Linolenic acid or n-3 long-chain polyunsaturated fatty acids on glial cell phosphatidylethanolamine and phosphatidylserine fatty acid profile in neonate rat brains", Nutrition & Metabolism, 2010

Publication

---

<1 %

85

James H. Cross, Ousman Jarjou, Nuredin Ibrahim Mohammed, Santiago Rayment Gomez et al. "Early postnatal hypoferrremia in low birthweight and preterm babies: A prospective cohort study in hospital-delivered Gambian neonates", EBioMedicine, 2020

Publication

---

<1 %

86

Michela Codini, Carmela Tringaniello, Lina Cossignani, Antonio Boccuto et al. "Relationship between Fatty Acids Composition/Antioxidant Potential of Breast Milk and Maternal Diet: Comparison with Infant Formulas", Molecules, 2020

Publication

---

<1 %

87

S. Cerolini, P.F. Surai, B.K. Speake, N.H.C. Sparks. "Dietary fish and evening primrose oil with vitamin E effects on semen variables in cockerels", British Poultry Science, 2010

Publication

---

<1 %

88

Schnittger, S, C Eder, S Jeromin, T Alpermann, A Fasan, V Grossmann, A Kohlmann, T Illig, N Klopp, H-E Wichmann, K-A Kreuzer, C Schmid, P Staib, R Peceny, N Schmitz, W Kern, C Haferlach, and T Haferlach. "ASXL1 exon 12 mutations are frequent in AML with intermediate risk karyotype and are independently associated with an adverse outcome", *Leukemia*, 2012.

Publication

&lt;1 %

89

Steen B. Haugaard. "Dietary intervention increases n-3 long-chain polyunsaturated fatty acids in skeletal muscle membrane phospholipids of obese subjects. Implications for insulin sensitivity", *Clinical Endocrinology*, 2/2006

Publication

&lt;1 %

90

[biomed.papers.upol.cz](http://biomed.papers.upol.cz)

Internet Source

&lt;1 %

91

[bmccancer.biomedcentral.com](http://bmccancer.biomedcentral.com)

Internet Source

&lt;1 %

92

[nutrition.highwire.org](http://nutrition.highwire.org)

Internet Source

&lt;1 %

93

[www.coursehero.com](http://www.coursehero.com)

Internet Source

&lt;1 %

94

[www.dzd-ev.de](http://www.dzd-ev.de)

Internet Source

&lt;1 %

95

[www.researchsquare.com](http://www.researchsquare.com)

Internet Source

&lt;1 %

96

"12th European Nutrition Conference (FENS), Berlin, Germany, October 20-23, 2015: Abstracts", *Annals of Nutrition and Metabolism*, 2015

Publication

&lt;1 %

97

Beverly Dosso, Charlotte Mae K Waits, Kelli N Simms, Susan Sergeant et al. "Impact of rs174537 on Critically Ill Patients with Acute Lung Injury: A Secondary Analysis of the OMEGA Randomized Clinical Trial", *Current Developments in Nutrition*, 2020

Publication

&lt;1 %

98

C. Molto-Puigmarti. "FADS1 FADS2 gene variants modify the association between fish intake and the docosahexaenoic acid proportions in human milk", *American Journal of Clinical Nutrition*, 05/01/2010

Publication

&lt;1 %

99

Chris R. Gignoux, Dara G. Torgerson, Maria Pino-Yanes, Lawrence H. Uricchio et al. "An admixture mapping meta-analysis implicates genetic variation at 18q21 with asthma susceptibility in Latinos", *Journal of Allergy and Clinical Immunology*, 2018

Publication

&lt;1 %

100 Fadilah Fadilah, Retnosari Andrajati, Ade Arsianti, Rafika Indah Paramita, Linda Erlina, Arry Yanuar. "Synthesis and in vitro Activity of Eugenyl Benzoate Derivatives as BCL-2 Inhibitor in Colorectal Cancer with QSAR and Molecular Docking Approach", Cold Spring Harbor Laboratory, 2020

Publication

<1 %

---

101 Graaff, Jolien Steenweg-de, Henning Tiemeier, Maartje Basten, Jolien Rijlaarsdam, Hans Demmelmair, Berthold Koletzko, Albert Hofman, Vincent W.V. Jaddoe, Frank C. Verhulst, and Sabine J. Roza. "Maternal LC-PUFA status during pregnancy and child problem behavior: the Generation R Study", Pediatric Research, 2014.

Publication

<1 %

---

102 Janice L.Y. Mong. "Associations of the Growth Hormone Receptor (*GHR*) Gene Polymorphisms with Adiposity and IGF-I Activity in Adolescents", Clinical Endocrinology, 01/2010

Publication

<1 %

---

103 M. Standl. "FADS gene cluster modulates the effect of breastfeeding on asthma. Results from the GINIplus and LISApplus studies : FADS genes, breastfeeding and asthma", Allergy, 01/2012

<1 %



104 Maria Teresa Segura, Hans Demmelmair, Susanne Krauss-Etschmann, Petra Nathan et al. "Maternal BMI and gestational diabetes alter placental lipid transporters and fatty acid composition", *Placenta*, 2017

Publication

---

105 Marie C. Conway, Emeir M. McSorley, Maria S. Mulhern, Toni Spence et al. "Maternal and child fatty acid desaturase genotype as determinants of cord blood long-chain PUFA (LCPUFA) concentrations in the Seychelles Child Development Study", *British Journal of Nutrition*, 2021

Publication

---

106 Shinichi Goto, Akira Sasaki, Ippei Takahashi, Yuri Mitsuhashi, Shigeyuki Nakaji, Atsushi Matsubara. "Relationship between cognitive function and balance in a community-dwelling population in Japan", *Acta Oto-Laryngologica*, 2017

Publication

---

107 Si-Wei Li, Kun Lin, Pei Ma, Zhen-Lu Zhang, Yi-Dan Zhou, Shuang-Yan Lu, Xin Zhou, Song-Mei Liu. "FADS Gene Polymorphisms Confer the Risk of Coronary Artery Disease in a Chinese Han Population through the Altered

Desaturase Activities: Based on High-Resolution Melting Analysis", PLoS ONE, 2013

Publication

- 
- 108 Suzanne J Meldrum, Michael A Smith, Susan L Prescott, Kathryn Hird, Karen Simmer. "Achieving definitive results in long-chain polyunsaturated fatty acid supplementation trials of term infants: factors for consideration", Nutrition Reviews, 2011  
Publication <1 %
- 
- 109 [genesandnutrition.biomedcentral.com](http://genesandnutrition.biomedcentral.com)  
Internet Source <1 %
- 
- 110 [pcwww.liv.ac.uk](http://pcwww.liv.ac.uk)  
Internet Source <1 %
- 
- 111 [sites.sph.harvard.edu](http://sites.sph.harvard.edu)  
Internet Source <1 %
- 
- 112 [www.biorxiv.org](http://www.biorxiv.org)  
Internet Source <1 %
- 
- 113 [www.ejast.org](http://www.ejast.org)  
Internet Source <1 %
- 
- 114 Ester MM Klaassen, Brenda EJT Thönissen, Guillaume van Eys, Edward Dompeling, Quirijn Jöbsis. "A systematic review of CD14 and toll-like receptors in relation to asthma in Caucasian children", Allergy, Asthma & Clinical Immunology, 2013  
Publication <1 %
-

115 Giovanni Malerba, Linda Schaeffer, Luciano Xumerle, Norman Klopp et al. "SNPs of the FADS Gene Cluster are Associated with Polyunsaturated Fatty Acids in a Cohort of Patients with Cardiovascular Disease", *Lipids*, 2008

Publication

<1 %

---

116 Rachele De Giuseppe, Carla Roggi, Hellas Cena. "n-3 LC-PUFA supplementation: effects on infant and maternal outcomes", *European Journal of Nutrition*, 2014

Publication

<1 %

---

117 Vera Zietemann, Janine Kröger, Cornelia Enzenbach, Eugene Jansen et al. " Genetic variation of the gene cluster and -6 PUFA composition in erythrocyte membranes in the European Prospective Investigation into Cancer and Nutrition-Potsdam study ", *British Journal of Nutrition*, 2010

Publication

<1 %

---

118 Antonio Gázquez, Olaf Uhl, María Ruíz-Palacios, Carolyn Gill et al. "Placental lipid droplet composition: Effect of a lifestyle intervention (UPBEAT) in obese pregnant women", *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*, 2018

Publication

<1 %

---

119

Isabella Pali-Schöll, Jennifer Namazy, Erika Jensen-Jarolim. "Allergic diseases and asthma in pregnancy, a secondary publication", World Allergy Organization Journal, 2017

Publication

<1 %

120

Millstein, Roberta L.. "Thinking about populations and races in time", Studies in History and Philosophy of Science Part C Studies in History and Philosophy of Biological and Biomedical Sciences, 2015.

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

# The association of fatty acid desaturase gene polymorphisms on long-chain polyunsaturated fatty acid composition in Indonesian infants

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---