

# Nucleotide mutation analyses of isolated lentogenic newcastle disease virus in live bird market

*by* Jola Rahmahani

---

**Submission date:** 26-Jul-2021 11:00AM (UTC+0800)

**Submission ID:** 1624048021

**File name:** ucleotide\_mutation\_analyses\_of\_isolated\_lentogenic\_newcastle.pdf (231.03K)

**Word count:** 4095

**Character count:** 22433

## Nucleotide mutation analyses of isolated lentogenic newcastle disease virus in live bird market

Jola Rahmahani<sup>1</sup>, Aisyah Nikmatuz Zahro<sup>2</sup>, Indah Laili Rahmawati<sup>3</sup>, Nurvita Putih<sup>3</sup>, Innah Wulandari<sup>3</sup>, Fedik Abdul-Rantam<sup>1,4,\*</sup>

- 1) Laboratory of Virology and Immunology, Department of Microbiology, Faculty of Veterinary Medicine, Airlangga University, Indonesia, 60115
- 2) Bachelor of Veterinary Science, Faculty of Veterinary Medicine, Airlangga University, Indonesia, 60115
- 3) Master of Vaccinology, Faculty of Veterinary Medicine, Airlangga University, Indonesia, 60115
- 4) Stem Cell Research and Development Center, Airlangga University, Indonesia, 60115

### ABSTRACT

Newcastle Disease (ND) is a major viral disease in Indonesia. It is an RNA virus belongs to Paramyxovirinae. It is well known that RNA virus is easily to mutate. In some cases, this mutation could generate virulence alteration. It is noted that mutation of NDV which has avirulent amino acid sequence on cleavage site, could mutate to be virulent Newcastle Disease Virus (NDV). It is needed to analyze the nucleotide and amino acid mutations and the effect of those to its virulence. The aim of this study was to analyze nucleotide and amino acid mutations of original isolated Lentogenic Newcastle Disease Virus (NDV). Samples were collected from cloacal swab of native chicken (*Gallus gallus domesticus*) suspected to be infected by Lentogenic NDV from live bird markets. They were inoculated into embryonated eggs, to isolate the virus. HA and HI assays were conducted to confirm that they were NDV. Positive samples were processed into serial passages in embryonated egg to observe their death time. Samples caused mortality of the embryonated eggs more than 90 hours post infection were suspected as Lentogenic NDV. They were processed to RT-PCR then sequenced. Lentogenic NDV confirmation was done comparing amino acid at Fusion protein cleavage site of the samples to Lasota/JF950510. Nucleotide and amino acid mutations were analyzed. The result showed that some nucleotide mutations were capable to change sequences of amino acid but the virulence of the samples remained the same to the reference sequence.

**Keywords:** NDV; RNA virus; mutations; virulence

### INTRODUCTION

*Avian Paramyxovirus serotype 1* which known as Newcastle Disease Virus (NDV) is negative single stranded RNA virus belongs to Genus Avulavirus, family Paramyxovirinae [1]. The genome of NDV has length 15,186 bp encoding six major structural protein 3-N-P-M-F-

\*Corresponding Author: Laboratory of Virology and Immunology, Department of Microbiology, Faculty of Veterinary Medicine, Airlangga University, Indonesia, 60115

Phone: +6231 5992785; Fax: +6231 5993015

E-mail: fedik-a-r@fkh.unair.ac.id And fedik.ar@gmail.com

HN-L-5 [2]. NDV is classified into three major strains based on clinical signs; Velogenic, Mesogenic, and Lentogenic [3]. Transcription of NDV is conducted in 3' to 5' direction [4]. Nucleoprotein (N) is firstly transcribed during replication [5]. Its length is 1,754 bp [2]. It protects genome of NDV and its positive sense from nucleases by the host cell [6]. Phosphoprotein (P) has length 1,451 bp [2]. It takes part during transcription [6] and prevents formation of unexpected nucleocapsid during replication [7]. RNA editing during P protein transcription produces V and W proteins [4]. Both proteins are virulent factor for NDV pathogenesis [6]. Matrix protein (M) has length 1,241 bp [2]. It controls RNA synthesis, assembling and budding of NDV virion. M protein is conserved protein among Paramyxovirinae family [5]. Sequence of M protein encoding gene has 20% homology compared to other Paramyxoviruses M protein [8].

NDV has two major glycoproteins that involve during NDV infection; Fusion (F) and Haemagglutinin-neuraminidase (HN) proteins [9]. F protein is type I integral protein which has length 1,729 bp that vary among Newcastle Disease. It is usually analyzed for genotyping NDV [10]. Activation of F protein is not dependent low acidity in endosomal compartment. NDV enter the host cell by direct fusion [11]. HN protein is type II integral protein which has length 2,002 bp. It has several functions such as virion-receptor binding [12], elusion of progeny virus from its host cell receptor, and Fusion protein activation [13]. The last transcribed gene is Polymerase protein (L). It has length 2,204 bp [2]. L, N, and P proteins form Ribonucleo-Protein complex for RNA synthesis [6]. Each gene is separated by intergenic regions (IGs) vary in length (1-47 nt) [2]. Sequence of IGs are not same among NDV but it usually ended with adenosine [14]. NDV is RNA virus [2]. Error that occurred during its replication causes RNA virus easily to mutate. It makes RNA virus is evolved quickly [15]. The mutation of amino acid in cleavage site of Fusion Protein of Newcastle Disease Virus caused the virulence of the virus is changed. It was proofed by outbreak in Australia during 1998-2000 [16]. This rapid mutation also forms a viral quasispecies. Viral quasispecies is dynamic distribution of virus that have same lineage but not identic as the result of genetic variation process, competition, and selection [2]. This study aims to analyze mutation that occurred in Lentogenic NDV.

## MATERIALS AND METHODS

**Samples collection and virus isolation:** Samples were collected from cloaca swab of native chicken (*Gallus gallus domesticus*) that suspected infected by Lentogenic NDV from five live bird markets in Surabaya (Bratang, Rungkut, Keputran, Kapasan, and Wonokromo). Transport medium was consisted of 105,000 IU penicillin (0.35 ml), 105,000 IU Streptomycin (0.525 ml) and 0.5 ml of sodium chloride. Samples were stored in 4°C refrigerator before inoculated into SPF embryonated eggs. All suspected samples were inoculated in 8 days old SPF embryonated eggs then incubated for five days in incubator (37°C). Virus isolation was conducted according to manual instruction from OIE [3]. Candling observation was done once daily. During incubation period, death embryonated eggs were recorded then chilled (4°C). In the end of incubation, embryonated eggs contained both death and live embryos were chilled (4°C). They were processed into confirmation test.

**HA and HI assays:** All samples were processed into Haemagglutination assay (HA) to detect the presence of Haemagglutinin protein in isolated samples. Both assays were done used micro technic method in V bottom 96 well microplate. HA assay was conducted using 50 µl of 0.5% chicken erythrocyte [3]. All positive HA samples were processed into Haemagglutination Inhibition assay (HI) using 2<sup>6</sup> NDV antiserum titre to confirm that isolated samples contained NDV. Each sample was diluted into 4HAU before processed into HI assay. HI assay was conducted using β-method [3]. Positive NDV samples were processed into serial passages in SPF embryonated eggs incubated for five days. The passages were done for five times to observe the death time of infected embryo. Embryos death at <60 hours post infection (hpi), 60-

90 hpi, and >90 hpi were suspected as Velogenic, Mesogenic and Lentogenic NDVs respectively [17]. Suspected lentogenic NDV were processed into RNA extraction.

**RNA extraction:** Suspected Lentogenic NDV samples processed into RNA extraction according to manufacturer protocol of R&A-BLUE™ iNTRON (BIOTECHNOLOGY, 17501). Extracted RNA of each samples were stored in -80°C.

**One Step RT-PCR:** One step RT-PCR was conducted according to manufacturer protocol of SuperScript® III One-Step RT-PCR System with Platinum®Taq DNA Polymerase (Invitrogen, 12574-026). Primer that used for one step RT-PCR was degenerated primer. PCR amplification was performed using following primers: forward primer 5'-GAG GTT ACC TCY ACY AAG CTR GAG-3' and reverse primer 5'-AGT CGG AGG ATG TTG GCA GC-3'. It was designed to amplify nucleotide number 4319-5005 of whole NDV genome (6865). 24 µl of RT-PCR master mix was added into each sample. The master mix was consisted of 12.5 µl of 2X RT Mix, 1 µl, respectively of primers, 8.5 µl NFW (DNase & RNase FREE, MP Biomedicals, 821932), and 1 µl SuperScript® III RT/Platinum®Taq Mix. One Step RT-PCR was run using thermo cycler machine (MJ Mini Personal Thermal Cycler, 1148). RT-PCR was arranged at 95°C for *Lid on*, 94°C predenaturation for 10 minutes, 94°C of denaturation for a minute, 56°C of annealing for a minute, 72°C of extension for 2 minutes and 72°C of extension prolong stage for 10 minutes. The cycles were repeated for 40 times. The product of PCR was stored in 4°C. Product of RT-PCR was visualized using 1.5% agarose gel (UltraPure™ Agarose, Invitrogen, 15510-019). DNA marker for electrophoresis was 100-1,000 bp (OneMark DNA Ladder). Agarose gel electrophoresis machine (PowerPac Basic, BIO-RAD) was run at 100 V, 400 mA for 40 minutes. Agarose gel then observed under UV illuminator then purified for sequencing according to protocol of QIAquick® PCR Purification Kit (Qiagen, 28104).

**Sequencing and molecular analyses:** RT-PCR product was sequenced using ABI 310 xL GENETIC ANALYZER (Applied Biosystem Inc.). RT-PCR product was labeled using BigDye® Terminator Sequencing RR-100. The sequencer machine was arranged as; first rapid thermal ramp 96°C for 10 seconds; second rapid thermal ramp 50°C for 5 seconds, third thermal ramp 60°C for 4 minutes. These cycles were repeated for 25 times. The samples were cooled in 4°C (fourth thermal ramp) until the samples had been ready to be precipitated. The samples were precipitated using 100 µl of absolute alcohol concentration. The sequence of nucleotides are symbolized as capital letters which are determined by International Union of Pure and Applied Chemistry (IUPAC). The sequence of nucleotides were analyzed using Biological Sequence Alignment Editor (BioEdit) version 7.0.5.3 and Molecular Evolutionary Genetics Analysis (MEGA) version 6.06. Isolated NDV were confirmed by comparing amino acid sequence in fusion protein cleavage site to Lentogenic NDV Lasota.

## RESULTS AND DISCUSSION

Thirty seven cloacal swabs were collected from native chicken (*Gallus gallus domesticus*) that show clinical signs such as mild diarrhea, subclinical respiratory infection, and prostration [18]. There were 37 samples were collected. All samples and Lentogenic NDV positive control (Lasota) were inoculated into allantoic fluid of 8 days old SPF embryonated eggs. They were incubated and observed for five days. Allantoic fluid was harvested then processed into HA and HI assays. Positive HA assay was showed by agglutination of 0.5% chicken erythrocyte in the V bottom microplate. Ten suspected samples were showed positive HA assay (Table1).

All positive HA assay samples and positive control were processed into HI assay using 2<sup>6</sup> of NDV antiserum. Positive HI assay was showed by the absence of agglutinated erythrocyte in the bottom of the microplate until well number six. Five of ten of suspected samples showed positive HI assay (Table 2). Cross protection is occurred among Paramyxoviruses. It can be

mistaken with Avian Paramyxovirus serotype 3 (APMV3) and APMV7. But APMV3 and APMV7 never been isolated from chicken before [17] and they are not isolated from Indonesia. Commonly Indonesian raising chicken such broiler, layer, and native chicken. NDV firstly isolated in Indonesia around 1926 [26]. Since its first emerging, NDV or so called 'tetelo' become endemic disease in Indonesia and cause annual outbreak [20, 26]. The samples were collected from native chicken (*Gallus gallus domesticus*) sold in bird market. Native chicken is traditionally reared in rural area which vaccination program is not conducted. This condition increases the possibility of natural infection of NDV. The aims of analysing lentogenic NDV is to know their mutation pattern as Lentogenic NDV has possibility to mutate into the virulence one. Thus it is very important to be observed.


**Table 1:** Titer of HA test of suspected Lentogenic NDV samples. Samples showed various titre

Samples	Titre of HA Test
NDVL1	2 <sup>6</sup>
NDVL2	2 <sup>6</sup>
NDVL3	2 <sup>9</sup>
NDVL5	2 <sup>4</sup>
NDVL6	2 <sup>6</sup>
NDVL7	2 <sup>9</sup>
NDVL8	2 <sup>5</sup>
NDVL9	2 <sup>9</sup>
NDVL10	2 <sup>8</sup>
NDVL11	2 <sup>9</sup>
LaSota	2 <sup>9</sup>

**Table 2:** Positive HI test samples means the isolated virus is NDV


Samples	HI titer
NDVL1	2 <sup>6</sup>
NDVL3	2 <sup>6</sup>
NDVL9	2 <sup>6</sup>
NDVL10	2 <sup>6</sup>
NDVL11	2 <sup>6</sup>
LaSota	2 <sup>6</sup>

Positive samples were observed in serial passages to select samples suspected Lentogenic NDV based on death time. Lentogenic NDV causes death of embryonated egg after 90 hours incubation [17]. All positive samples caused death of embryonated eggs after 90 hours post infection). It was supported the suggestion that isolated virus was Lentogenic NDV. But this suggestion compulsory to be confirmed by analysing amino acid at Fusion protein cleavage site of all samples. Suspected samples were processed into one step RT-PCR. RT-PCR product was visualized in 1.5% agarose gel, then it was observed under UV illuminator (Fig. 1).



**Figure 1:** Agarose gel electrophoresis of suspected Lentogenic NDV samples. The band of all samples were located between marker for 600 bp and 700 bp. It means that primer amplified targeted region of whole NDV genome. M: 100 bp DNA ladder marker. Lane I: NDVL1. Lane II : NDVL3. Lane III : NDVL9. Lane IV : NDVL10. Lane V : NDVL11. Lane VI : LaSota (positive control).

RT-PCR product was processed into sequencing. Nucleotide sequence of the samples were analyzed using BioEdit version 7.0.5.3 and Molecular Evolutionary Genetics Analysis (MEGA) version 6.06. to confirm that isolated virus was Lentogenic. They were analyzed in region of fusion protein cleavage site. Amino acid sequence at fusion protein cleavage site of NDV was located at amino acid number 112-117. Lentogenic NDV has only single basic amino acid at amino acid number 112 and 115, and Leucine at amino acid number 117 [19]. Isolated samples showed identical sequence compared to Lasota/JF950510.1 (Fig. 2). It can be seen that amino acid sequence of the samples, marked with 'dot' and colored as same as amino acid of the reference sequence.



**Figure 2:** Confirmation of isolated samples by comparing their amino acid sequence of Fusion protein to Lasota/JF950510.1.

After confirming that isolated samples were Lentogenic NDV, product of RT-PCR that had been sequenced was aligned to Lasota/JF950510.1 (Genebank). The Nucleotide mutations were classified according to the type and their alteration of amino acid mutation. It was done to understand the effect of nucleotide mutation to amino acid sequence.

Sixteen nucleotide mutations occurred in NDV1. They were occurred only on Fusion protein encoding gene. They were consisted of nine transition mutations and seven transversion mutations. These caused some of amino acid alteration (Table 3).

**Table 3:** Nucleotide and amino acid mutations of NDV1 only occurred on Fusion protein encoding gene

No.	Nucleotide mutation	Amino acid mutation	Type of mutation
1	A586G	I196V	Transition
2	G595A	G199S	Transition
3	G598A	I200V	Transition
4	G600C	-	Transversion
5	T618A	-	Transversion
6	C621G	-	Transversion
7	C623A	A208E	Transversion
8	A627G	-	Transition
9	A633T	-	Transversion
10	G636A	-	Transition
11	C639T	-	Transition
12	A642G	-	Transition
13	C644G	A215G	Transversion
14	T645C	-	Transition
15	G648A	-	Transition
16	A654T	Q218H	Transversion

Thirty six nucleotide mutations were occurred in NDV3. They were consisted of twenty two transition mutations and fourteen transversion mutations. Six mutations were occurred at genome junction, while the rest of mutations were occurred at gene encoding Fusion protein (Table 4). Eleven mutations were occurred in NDV9. They were consisted of eight transition mutations and three transversion mutations. All mutations were occurred at gene encoding Fusion protein (Table 5). Nucleotide and amino acid mutations which found on NDV10 and NDV11 were particularly same. They were consisted only of transition mutations. One mutation occurred at gene encoding Matrix protein while four mutations were occurred at gene encoding Fusion protein (Table 6).

According to the observation, 89.2% mutation occurred on Fusion protein encoding gene. Fusion protein is the major determinant protein in genotyping and virulence classification of

Newcastle Disease Virus. It is noted that Fusion protein is immunogenic protein and target of antibody-antigen neutralization [16]. Among samples, nucleotide and amino acid mutations on genome junction occurred only on NDVL3 (Table 5). Intergenic region of NDV extend between each protein encoding gene. The modification of its domain is capable to increase the virulence of NDV [24].

**Table 4:** Nucleotide and amino acid mutations of NDVL3 occurred on genome junction (red color) and Fusion protein encoding gene (black color)

No.	Nucleotide mutation	Amino acid mutation	Type of mutation
1	C102G	-	Transversion
2	A118C	K4Q	Transversion
3	T134C	L45P	Transition
4	A140T	Y47S	Transversion
5	G168A	-	Transition
6	T205C	-	Transition
7	C213T	-	Transition
8	C215T	S72k	Transition
9	C216T	-	Transition
10	C234T	-	Transition
11	A274G	I92V	Transition
12	T282C	-	Transition
13	G299C	C100S	Transversion
14	A339G	-	Transition
15	T342C	-	Transition
16	T357A	-	Transversion
17	C372A	-	Transversion
18	C387T	-	Transition
19	A405G	-	Transition
20	T414G	-	Transversion
21	G426A	-	Transition
22	T453C	-	Transition
23	G456A	-	Transition
24	C504T	-	Transition
25	T582C	-	Transition
26	G595A	G199S	Transition
27	G600C	-	Transversion
28	T618A	-	Transversion
29	C621G	-	Transversion
30	A627G	-	Transition
31	A633T	-	Transversion
32	G636T	-	Transversion
33	C639A	-	Transversion
34	C646G	L216V	Transversion
35	G648A	-	Transition
36	A663G	-	Transition

**Table 5:** Nucleotide and amino acid mutations of NDVL9 only occurred in Fusion protein encoding gene

No.	Nucleotide mutation	Amino acid mutation	Type of Mutation
1	T582C	-	Transition
2	G595A	G199S	Transition
3	G598A	V200I	Transition
4	G600C	-	Transversion
5	T618A	-	Transversion
6	C621G	-	Transversion
7	A627G	-	Transition
8	A633T	-	Transversion
9	G636A	-	Transition
10	C639T	-	Transition
11	A641G	-	Transition
12	G648A	-	Transition

**Table 6:** Nucleotide and amino acid mutations of NDVL10 and NDVL11 occurred on gene encoding Matrix protein (red) and Fusion protein encoding gene

No.	Nucleotide mutation	Amino acid mutation	Type of mutation
1	A23G	E8G	Transition
2	T582C	-	Transition
3	C621T	-	Transition
4	G636A	-	Transition
5	A642G	-	Transition

NDV is endemic viral disease in many regions include Indonesia [20]. It may be caused by the rise of poultry density but unfollowed by good biosecurity management [21]. Lentogenic NDV can be isolated from native chicken (*Gallus gallus domesticus*) in live bird markets. To define isolated NDV is Lentogenic strain several methods must be done such as observation of death time and analyzing sequence of amino acid at cleavage site of Fusion protein. RNA viruses are noted easily to mutate [2].

Some of nucleotide mutations that occurred caused amino acid mutations, but some of them are not. Mutations did not occurred in cleavage site of Fusion protein (nt 558-576) means the virulence of samples compared to reference sequence are same. The mutations that occurred may not effect the virulence of the samples. Even though it did not occur in this research, certain mutation in cleavage site of fusion protein transform the Lentogenic NDV into the virulence one [16]. Polymerase protein of RNA virus is characteristically for its low fidelity which lead RNA viruses are easily to mutate. These mutations may form quasispecies Newcastle Disease Virus [22]. Quasispecies is viruses genetically linked as the result of genetic variation process, competition, and selection. They are non identic but belongs to same lineage [2]. Each mutant in quasispecies is genetically linked [23]. It needs further research to understand concept of quasi species in Newcastle disease virus.

NDVL10 and NDVL11 are suspected to be one species since it is isolated from same live bird market. Among the samples, NDVL10 and NDVL11 have the least nucleotide and amino acid mutations. Furthermore, the position of the mutations are same. According to this finding it is suggested that these samples are 'Lentogenic NDV like vaccine. It could be found in wild birds too [25].

**Acknowledgment:** Thanks to member Newcastle Disease Research Team for the support. Thanks to Universitas Airlangga and Dean of Faculty of Veterinary Medicine for the support through Mandate Grant Fund 2018. Thanks for Magister and Bachelor students for helping the author checking the manuscript.

**Conflict of Interest:** The authors declare there is no conflict of interest regarding the research process and writing process. This manuscript never been submitted into any other publisher before.

## REFERENCES

1. Fenner. 2011. Fenner's Veterinary Virology 4<sup>th</sup> edition: Paramyxoviridae. Elsevier.
2. Knipe DM, Howley PM, Cohen JI, Griffin DE, Lamb RA, Martin MA, Racaniello VR, Roisman B. Fields Virology 2013; LIPPINCOT WILLIAMS & WILKINS.
3. OIE. 2012. OIE Manual of Diagnostic Test and Vaccines for Terrestrial Animals in Newcastle Diseases (Infection with Newcastle Diseases Virus).
4. Alamares JG. Newcastle disease virus virulence: Mechanism of the interferon antagonistic activity of the V protein and characterization of a putative virulence-specific antibody to the attachment protein. 2008. Any disertation. Medical School. University of Massachusetts.
5. Seal BS, King DJ, Sellers HS. The avian response to Newcastle diseases Virus. Dev Comp Immunol 2000;24:257-268.



6. Ganar K, Das M, Sinha S, Kumar S. Newcastle diseases virus: Current status and our understanding. *Virus Res* 2014;184:71-81.
7. Errington W, Emmerson PT. Assembly of recombinant Newcastle disease virus nucleocapsid protein into nucleocapsid-like structures is inhibited by the phosphoprotein. *J Gen Virol* 1997; 78:2335-2339.
8. Moran IS, Castano SC, Barroso IM, Kostetsky EY, Zhadan G, Gomez J, Shnyrov VL, Villar E. Thermal stability of matrix protein from Newcastle diseases virus. *Inter J Biol Macromol* 2013;61:390-395.
9. Samal S. Role of fusion protein in Newcastle disease virus pathogenesis [Disertation]. Faculty of Graduated School. University of Maryland 2012.
10. Miller PJ, Haddas R, Simanov L, Lublin A, Rehmani SF, Wajid A, Bibi T, Khan TA, Yaquob T, Setiyaningsih S, Afonso CL. Identification of new sub-genotypes of virulent Newcastle disease virus with potential panzootic features. *Infect Genet Evol* 2015;29:216-229.
11. Horvarth CM, Lamb RA. Studies on the fusion peptide of a Paramyxovirus fusion glycoprotein : Roles of conserved residues in cell fusion. *J Virol* 1992;66:2443-2445.
12. Yuan P, Swanson KW, Leser GP, Paterson RG, Lamb RA, Jardetzky TS. Structure of the Newcastle disease virus Hemagglutinin-Neuraminidase (HN) ectodomain reveals a four-helix bundle stalk. *Proc Natl Acad Sci USA* 2011;108:14920-14925.
13. Porotto M, Salah ZW, Gui L, DeVito I, Jurgens EM, Lu H, Yokoyama CC, Palermo LM, Lee KK, Moscona A. Regulation of Paramyxovirus fusion activation: the hemagglutinin-neuraminidase protein stabilizes the fusion protein in a pretriggered state. *J Virol* 2012;86: 12838-12848.
14. Krishnamurthy S, Samal SK. Nucleotide sequences of the trailer, Nucleocapsid protein gene and intergenic regions of Newcastle disease virus strain Beaudette C and completion of the entire genome sequence. *J Gen Virol* 1998;79:2419-2424.
15. Jenkins GM, Rambaut A, Pybus OG, Holmes EC. Rates of molecular evolution in RNA viruses : A quantitative phylogenetic analysis. *J Mol Evol* 2002;54:156-165.
16. Miller PJ, Kim LM, Ip HS, Afonso CL. Evolutionary dynamics of Newcastle disease virus. *Virology* 2009;391:64-72.
17. Alexander DJ. Newcastle disease, other avian Paramyxoviruses and pneumovirus infection. *Rev Sci Tech* 2000;19:443-462.
18. Alexander DJ, Bell JG, Alders RG. A Technology Review : Newcastle disease with special emphasis on its effect on village chickens. In : *FAO Animal Production and Health*. 2004.
19. Aldous EW, Alexander DJ. Detection and differentiation of Newcastle disease virus (avian Paramyxovirus type 1). *Avian Pathol* 2001;30:117-128.
20. Dharmayanti NI, Hartawan R, Hewajuli DA, Indriani R. Phylogenic analysis of genotype VII of Newcastle diseases virus in Indonesia. *Afri J Microbiol Res* 2014;8:1368-1374.
21. Chukwudi OE, Chukwuemeka ED, Mary U. Newcastle disease virus shedding among healthy commercial chickens and its epidemiological importance. *Pak Vet J* 2012;32:354-356.
22. Meng C, Qiu X, Yu S, Li C, Sun Y, Chen Z, Liu K, Zhang X, Tan L, Song C, Liu G, Ding C. Evolution of Newcastle disease virus quasispecies diversity and enhanced virulence after passage through chicken air sacs. *J Virol* 2016;90:2052-2063.
23. Lauring AS, Andino R. Quasispecies theory and the behaviour of RNA virus. *PLoS Pathog* 2010;6:e1001005.
24. Yan Y, Samal SK. Role of intergenic sequences in Newcastle disease virus RNA transcription and pathogenesis. *J Virol* 2008;82:1323-1331.
25. Ayala AJ, Dimitrov KM, Becker CR, Goraichuk IV, Arns CW, Bolotin VI, Ferreira HL, Gerilovych AP, Goujgoulova GV, Martini MC, Muzyka DV, Orsi MA, Scagion GP, Silva RK, Solodiankin OS, Stegnyy BT, Miller PJ, Afonso CL. Presence of vaccine-derived Newcastle disease viruses in wild birds. *PLoS One* 2016;11:e0162484.
26. Sa X, Paldural A, Nayak B, Samuel A, Bharoto EE, Prajitno TY, Collins PL, Samal SK. Complete genome sequence of Newcastle disease virus strains circulating in chicken populations of Indonesia. *J Virol* 2012;5969-5970.

# Nucleotide mutation analyses of isolated lentogenic newcastle disease virus in live bird market

## ORIGINALITY REPORT

12%

SIMILARITY INDEX

9%

INTERNET SOURCES

10%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Yu-Pin LIU, Chiu-Yen CHANG, Fan LEE, Chwei-Jang CHIOU, Hsiang-Jung TSAI. "Phylogenetic analysis of avian paramyxoviruses 1 isolated in Taiwan from 2010 to 2018 and evidence for their intercontinental dispersal by migratory birds", *Journal of Veterinary Medical Science*, 2020  
Publication 1%
- 2 [www.scipub.org](http://www.scipub.org)  
Internet Source 1%
- 3 "International Society for Animal Clinical Pathology (ISACP) Musashino, Tokyo, Japan, August 4–8, 2018", *Veterinary Clinical Pathology*, 2019  
Publication 1%
- 4 [doaj.org](http://doaj.org)  
Internet Source 1%
- 5 [link.springer.com](http://link.springer.com)  
Internet Source 1%

6	<a href="http://hdl.handle.net">hdl.handle.net</a> Internet Source	1 %
7	<a href="http://khartoumspace.uofk.edu">khartoumspace.uofk.edu</a> Internet Source	<1 %
8	<a href="http://www.e-sciencecentral.org">www.e-sciencecentral.org</a> Internet Source	<1 %
9	Desi Sandra Sari, Ernie Maduratna, Fourier Dzar Eljabbar Latief, Alexander Patera Nugraha, Ketut Suidiana, Fedik Abdul Rantam. "Osteogenic Differentiation and Biocompatibility of Bovine Teeth Scaffold with Rat Adipose-derived Mesenchymal Stem Cells", European Journal of Dentistry, 2019 Publication	<1 %
10	Xue-Di Cheng, Hua-Feng Xu, Xue-Mei Wei, Hai-Zhou Zhou. "Variation analysis of E1 and E2 in HCV subtypes", Archives of Virology, 2015 Publication	<1 %
11	<a href="http://escholarship.org">escholarship.org</a> Internet Source	<1 %
12	<a href="http://espace.library.uq.edu.au">espace.library.uq.edu.au</a> Internet Source	<1 %
13	<a href="http://journals.plos.org">journals.plos.org</a> Internet Source	<1 %
14	<a href="http://www.tsijournals.com">www.tsijournals.com</a> Internet Source	<1 %

15

[researchonline.jcu.edu.au](http://researchonline.jcu.edu.au)

Internet Source

&lt;1 %

16

Rather, Irfan Ahmad, Kwang-Ho Choi, Vivek K. Bajpai, and Yong-Ha Park. "Antiviral mode of action of *Lactobacillus plantarum* YML009 on Influenza virus H1N1", *Bangladesh Journal of Pharmacology*, 2015.

Publication

&lt;1 %

17

Youn Jeong Lee, Haan Woo Sung, Jun Gu Choi, Jae Hong Kim, Chang Seon Song. "Molecular epidemiology of Newcastle disease viruses isolated in South Korea using sequencing of the fusion protein cleavage site region and phylogenetic relationships", *Avian Pathology*, 2004

Publication

&lt;1 %

18

Lucy Harrier, Linda Paterson. "Inducibility studies with the arbuscular mycorrhizal fungus *Glomus mosseae* 3-phosphoglycerate kinase ( PGK ) gene promoter", *Current Genetics*, 2002

Publication

&lt;1 %

19

Vasudevan Gowthaman, Venkateswaran Ganesan, Thippicettipalayam Ramasamy Gopala Krishna Murthy, Sowmya Nair et al. "Molecular phylogenetics of Newcastle disease viruses isolated from vaccinated flocks during outbreaks in Southern India

&lt;1 %

reveals circulation of a novel sub - genotype",  
Transboundary and Emerging Diseases, 2018

Publication

20

[ijlr.org](http://ijlr.org)

Internet Source

<1 %

21

[nexusacademicpublishers.com](http://nexusacademicpublishers.com)

Internet Source

<1 %

22

[www.frontiersin.org](http://www.frontiersin.org)

Internet Source

<1 %

23

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<1 %

24

Bruce S. Seal. "Analysis of matrix protein gene nucleotide sequence diversity among Newcastle disease virus isolates demonstrates that recent disease outbreaks are caused by viruses of psittacine origin",  
Virus Genes, 06/1995

Publication

<1 %

25

Lilik Maslachah, Rahmi Sugihartuti. "Increase in neutrophil count after repeated exposure of Plasmodium berghei-infected mice to artemisinin",  
Universa Medicina, 2017

Publication

<1 %

26

Sachin Kumar, Flavia Militino Dias, Baibaswata Nayak, Peter L. Collins, Siba K. Samal.  
"Experimental avian paramyxovirus serotype-

<1 %

3 infection in chickens and turkeys",  
Veterinary Research, 2010

Publication

---

27

Esin Guven, Hamza Avcioglu, Ahmet Deniz, İbrahim Balkaya, Ugur Abay, Şevki Yavuz, Muzaffer Akyüz. "Prevalence and molecular characterization of Theileria equi and Babesia caballi in jereed horses in Erzurum, Turkey", Acta Parasitologica, 2017

Publication

---

<1 %

28

Martha Purnami Wulanjati, Nastiti Wijayanti, Aris Haryanto. "Phylogenetic Analysis of Newcastle Disease Virus from Indonesian Isolates Based on DNA-Sequence of Fusion Protein-Encoding Gene", Biotechnology(Faisalabad), 2018

Publication

---

<1 %

29

Momena Habib, Tahir Yaqub, Jawad Nazir, Wasim Shehzad et al. "Genomic and biological characterization of Newcastle disease viruses isolated from migratory mallards (Anas platyrhynchos)", Archives of Virology, 2018

Publication

---

<1 %

30

Perumal Arumugam Desingu, Shambhu Dayal Singh, Kuldeep Dhama, Obli Rajendran Vinodhkumar, Yashpal Singh Malik. "A sensitive haemadsorption technique based RT-PCR for concentration and detection of

<1 %

## Newcastle disease virus from clinical samples and allantoic fluid", VirusDisease, 2016

Publication

---

31

Shabbir, Muhammad Zubair, Siamak Zohari, Tahir Yaqub, Jawad Nazir, Muhammad Abu Shabbir, Nadia Mukhtar, Muhammad Shafee, Muhammad Sajid, Muhammad Anees, Muhammad Abbas, Muhammad Tanveer Khan, Asad Amanat Ali, Aamir Ghafoor, Abdul Ahad, Aijaz Ali Channa, Aftab Ahmad Anjum, Muhammad Nazir Kalhoro, Arfan Ahmad, Mohsan Ullah Goraya, Zahid Iqbal, Sohail Ahmad Khan, Hassan bin Aslam, Kiran Zehra, Muhammad Umer Sohail, Waseem Yaqub, Nisar Ahmad, Mikael Berg, and Muhammad Munir. "Genetic diversity of Newcastle disease virus in Pakistan: a countrywide perspective", *Virology Journal*, 2013.

<1 %

Publication

---

Exclude quotes  Off

Exclude matches  Off

Exclude bibliography  On