

A Novel Pleiotropic Effect of Beta-Blockers Useful or Not?

by Tara Mandiricha

Submission date: 28-Jul-2021 12:49PM (UTC+0800)

Submission ID: 1624945337

File name: A_Novel_Pleiotropic_Effect_of_Beta-Blockers_Useful_or_Not.pdf (1.08M)

Word count: 7928

Character count: 44777

A Novel Pleiotropic Effect of Beta-Blockers: Useful or Not?

Tara Mandiricha¹, Maftuchah Rochmanti^{2,*}

¹Master Student, Basic Medical Science of Pharmacology, Faculty of Medicine, Universitas Airlangga, Surabaya East Java Indonesia. (mandiricha103@gmail.com)

²Departement of Pharmacology, Faculty of Medicine, Universitas Airlangga, Surabaya East Java Indonesia, (maftuchah-r@fk.unair.ac.id)

*Corresponding author: Maftuchah Rochmanti Email: maftuchah-r@fk.unair.ac.id

ABSTRACT

β -adrenergic blockers or β -blockers (BBs) have been the antihypertensive treatment for the past 50 years. Several studies were developed to further explore the therapeutic effects of BB, ranging from use in liver cirrhosis with esophageal varices to involvement in cancer treatment. The Autonomic nervous system (ANS) dysfunction has been shown to cause several of these diseases. Immune suppression, cardiovascular dysfunction, hypertension, and even worse prognosis in cancer patients are due to chronic activation of the sympathetic nervous system (SNS). Based on its mechanism of action, the use of BB drugs has different therapeutic targets, and each has different advantages and side effects. Since the discovery of a new third-generation BB drug that has a complete combination of action, several diseases have hopes of being treated with this agent. Improving survival, hospital discharge, and quality of life affecting patients is the goal of the new therapeutic approach.

Keywords: β -blockers, antihypertensive, cardiovascular, esophageal varices, cancer

Correspondence:

Maftuchah Rochmanti
Department of Pharmacology, Faculty of Medicine, Universitas Airlangga,
Surabaya East Java Indonesia
Email: maftuchah-r@fk.unair.ac.id

INTRODUCTION

The involvement of the autonomic nervous system (ANS) has a considerable influence on the pathophysiology of cardiovascular disease (CVD). Activation of the sympathetic nervous system (SNS) is a short term compensatory response to hemodynamic changes resulting from abnormal cardiac function. This is also influenced by withdrawal activity by the Parasympathetic Nervous System (PNS). An imbalance of SNS and PNS activities in the long term will cause anatomic abnormalities and cardiac function. Myocardial remodeling is a sign of anatomical abnormalities that have an impact on cardiac dysfunction that often occurs in heart failure (HF). The etiology of HF varies, but the presence of autonomic dysfunction is characteristic of this disease. A worsening ANS imbalance can lead to a high risk of death. Therefore one of the medical approaches to improve the prognosis of chronic HF patients is to modulate the PNS and weaken SNS activation through blockade of the β -adrenergic system. Treatment of hypertension with BB has been used in clinical practice for the past 3 decades, but now new indications have been found from several studies conducted. It is known that BB can affect the immune system in the body, even being considered as a cancer treatment. Based on this, we are interested in reviewing the pleiotropic and new therapeutic effects of BB use both as a short-term and long-term treatment.

PHARMACOLOGY OF β -BLOCKER

β -blockers are one of the most widely used antihypertensives in clinical practice because of their benefits in the treatment of HF or acute myocardial infarction. This agent binds to the cardiac β -adrenergic receptors, thereby blocking the binding of endogenous catecholamines adrenaline and noradrenaline causing a decrease in the force contractile and heart rate. Beta-blockers (BB) are competitive antagonists at adrenergic receptors, where drug selectivity can be distinguished

according to the receptors. β -adrenoceptors are divided into 3 identified subtypes, β_1 , β_2 , and β_3 . Meanwhile, β_1 - and β_2 adrenoceptors are the strongest physiological mechanisms for increasing acute cardiac performance. β_1 -receptors are mainly expressed in the heart and induce positive chronotropic, dromotropic, bathmotropic, and inotropic effects. Meanwhile β_2 -receptors are mainly expressed in various types of smooth muscle cells located in blood vessels, skeletal muscle cells, and hollow visceral organs, such as bronchi or uterus which cause relaxation. β_3 -receptors are expressed in several tissues such as adipose tissue, heart, ANS, pancreas, colon, uterus, bladder, and gut. Moreover, these receptor differences include the presence of intrinsic sympathomimetic activity (ISA), lipid solubility that affects penetration into the central nervous system (CNS), as an inverse agonist, capacity to induce vasodilation, and pharmacokinetic profile. β -blockers have 3 generations that have been developed and applied in clinical practice. The first generation includes non-selective subtypes, for example propranolol which has the same affinity for β_1 and β_2 -receptors. The second generation includes drugs that have an affinity for β_1 rather than β_2 (selective), such as atenolol, bisoprolol, and metoprolol. The third generation contains selective and non-selective β_1 subtypes which have additional properties of α -1 receptor antagonists or the ability to induce the production of nitric oxide (NO) which causes vasodilation.

Several therapeutic effect of BB have been identified including decreased cardiac output (CO) and inhibition of renin release, thereby reducing myocardial workload and oxygen demand to reduce symptoms of angina and the risk of acute coronary syndrome (ACS). The mechanism of action of BB against norepinephrine and epinephrine at beta receptors is to prevent the cyclic adenosine monophosphate (cAMP) accumulation, protein kinase A (PKA) activation, and subsequent changes in intracellular calcium that mediate adrenergic responses. BB which is more lipophilic is usually associated with central nervous

system (CNS) side effects because it is able to cross the blood-brain barrier and is well absorbed from the intestine but undergoes extensive first pass metabolism in the liver. Meanwhile, hydrophilic BB (eg atenolol) is not fully absorbed from the intestine and does not undergo hepatic metabolism, has a longer half-life than lipophilic BB agents^{19,20}. Meanwhile, BB agents with intrinsic sympathomimetic activity (ISA) may be less prone to cause bradycardia¹⁹. Although the use of BB has been considered to be the primary safe therapy for CVD, it is also used in the treatment of migraine²¹, glaucoma²², hyperthyroidism²³, hepatic cirrhosis²⁴, chronic obstructive pulmonary disease (COPD)^{25,26}, sepsis^{27,28}, and currently considered cancer²⁹.

β-BLOCKER AND CANCER

β-adrenergic receptors can be detected in cancer cells and immune cells by influencing the proliferation process and multiple signaling pathways involved in cancer invasion, cancer-related inflammation, angiogenesis, and lymphangiogenesis³⁰. Deviated β2 expression is associated with oncogenic properties in breast cancer and increased metastases to axillary lymph nodes. This process is mediated by β2 via activation of the cAMP-calcium feed-forward loop and accumulation of mineral ion channels that increase breast cancer cell invasion. Meanwhile, β1 has been shown to be associated with increased lipolysis in cachexia cancer and β3 mutations contribute to breast cancer-related obesity³¹. Some evidence suggests that β-adrenergic receptor signaling can induce SSS activation and increase circulating catecholamine levels by the adrenal glands and locally from the term postganglionic sympathetic nerve fibers³². Activation of adrenergic receptors can regulate cancer-related signaling pathways including in the tumor microenvironment such as endothelial cells and macrophages which can lead to the development and spread of cancer metastases³³. In addition, activation of tumor beta adrenergic receptors is reported to increase the production of several factors that promote metastasis such as vascular endothelial growth factor (VEGF), matrix metalloproteinase 2 (MMP-2), MMP-9, IL-6, and IL-8³⁴. Therefore, blockade of β-adrenergic receptor signaling is expected to provide a protective role in cancer patients by changing the tumor microenvironment, preventing cancer development and metastasis through decreasing tumor cell viability and inhibiting the formation of vascular remodeling.^{35,36} The use of BB has also been reported to modulate the immune response and regulation of cytokines involved in cancer development³⁷. This is evidenced by the use of propranolol which can reduce mitogen-activated protein kinase (MAPK), 70-kDa heat shock protein (HSP70), iNOS activity, increases IL-10 and activates receptor activator of nuclear factor-κB ligand (RANKL). The MAPK pathway associated with cell proliferation, growth, transport, death, and many vital factors is very important in human tumors, while HSP70 is upregulated across a wide variety of cancers and is involved in tumor growth, invasion, migration and resistance to anti-cancer therapy, and RANKL which are frequently detected in the tumor microenvironment and participate in every step of cancer development³⁸⁻⁴⁰. Natural Killer cell (NK) is known to play an important role in innate immunity, killing virus-infected cells and tumors. At the cellular level, activation of β-adrenoceptors can decrease the cytotoxicity of NK cells against MADB106 mammary adenocarcinoma cells accompanied by pulmonary metastases in vitro, while at the systemic level

activation of local β-adrenoceptors can reduce the number of NK cells in the lungs⁴¹. Other evidence shows the results of in vitro studies on SK-BR-3 cells with 24-hour propranolol exposure that show decreased phosphorylation of multiple mitogenic active protein kinase (MAPK) and cAMP responsive element binding protein (CREB), as well as increased phosphorylation of protein kinase B (PKB), glycogen synthase kinase 3 (GSK3) and p53. In addition to breast cancer, propranolol can also inhibit damage due to catecholamine stimulation of adrenoceptors in people with pancreatic cancer⁴².

β-BLOCKER AND MIGRAINE

In the general population the prevalence of migraine cases is estimated to be 16% and more in women than men (ratio 3:1). Some people have episodic migraines and receive prophylactic therapy, but only 3% -13% of patients are reported to be free from post-control symptoms^{43,44}. Migraine is a neurovascular disorder that is common in about 10-15% of the general population⁴³. This neurovascular disorder is associated with a neurogenic inflammatory process characterized by the release of potent vasoactive neuropeptides such as calcitonin gene-related peptide (CGRP)⁴⁵, substance P (SP)⁴⁶, and neurokinin A⁴⁷. Meanwhile, migraine headaches that are preceded by an aura phase of about 20% -30% may be triggered directly by cortical spreading depression (CSD), a slow depolarization wave that spreads throughout the cortex^{48,49}. The involvement of CSD causes complex molecular changes in upregulation of cortical genes involved in inflammatory processing such as cyclooxygenase-2, TNF-α, IL-1b, or matrix metalloproteinase (MMP). CSD waves have also been reported to induce activation of trigeminovascular pathways derived from meningeal nociceptors^{48,50}. β-blockers are widely used as prophylactic therapy for migraines and effective in reducing the frequency of attacks by 50%⁴⁹. Abortive and prophylactic treatment for migraine may be needed in patients with acute symptoms. Prophylactic therapy can reduce the frequency, duration, or severity of attacks⁵¹. Inhibition of adrenoceptor β1 such as propranolol is reported to be able to modulate negative trigeminovascular nociception in thalamocortical neurons in migraine. Abnormalities in sensory processing of the thalamus and cortical can cause some migraine cardinal symptoms such as photophobia and phonophobia so this thalamic neuron may be able to be targeted as a preventive in migraine⁵². Whereas the activation of MMP causes an increase in blood brain barrier permeability and activation of trigeminal nerve afferents. These neurons act to supply sensory innervation to large blood vessels in the cerebrovascular and meningeal systems. Migraine headaches may occur as a result of a cascade; pannexin-1 megachannels open, which activates caspase-1 and triggers the release of proinflammatory agents thereby activating the astrocytic NF-κB and finally transduction of the inflammatory response to the sensory trigeminal nerves that innervate pial vasculature⁵³. In addition, propranolol also inhibits the release of nitric oxide (NO) by blocking inducible Nitric Oxide Synthase (iNOS) which has the potential to reduce the regulation of activation of trigeminovascular complexes^{54,55}. BB drugs such as metoprolol and propranolol are known to have a better therapeutic response because they can cross the blood-brain barrier (BBB), very lipophilic, and modify nerve stimulation. β1 receptor antagonists can modulate the processing of cortical information expressed as changes in visual evoked potentials (VEP) mediated

through adrenaline and noradrenaline, contingent negative variation (CNV), and auditory evoked potentials (AEP), which are usually abnormal in migraine patients. The use of metoprolol and propranolol has been proven to be able to normalize high CNV and decrease VEP amplitude, which shows that BB has a significant effect on visual system stimulation in migraine patients⁴⁹. Furthermore, the ventrobasal complex is innervated by noradrenergic fibers which are mostly found in the ipsilateral coeruleus locus. Noradrenaline is found in the thalamus and its release is triggered by the activation of the locus coeruleus neurons. Adrenergic receptors α and β are found in somatosensory thalamus. β receptors are found in ventrobasal complexes, especially the receptor subtype β_1 ⁵². Recent findings suggest that propranolol can interfere with the process of chronic sensitization in the rostral ventromedial medulla (RVM) and locus coeruleus (LC)⁵⁶. Propranolol and timolol have a high affinity to serotonergic system by blocking 5-HT_{2C} and 5-HT_{2B} receptors because 5-HT (serotonin) receptors which have a significant role in migraine pathophysiology⁵⁷.

β -BLOCKER AND COPD

Chronic obstructive pulmonary disease (COPD) is a chronic inflammatory lung disease, which is characterized by progressive limitation of air flow and is associated with an abnormal lung inflammatory response to harmful particles or gases^{58,59}. The systemic inflammatory process in COPD involves several cells including macrophages, epithelial cells, dendritic cells, neutrophils, eosinophils, and T lymphocytes and B lymphocytes. These cells release many inflammatory mediators that play a role in the pathophysiology of COPD such as lipids, cytokines, chemokines, and growth factors⁶⁰. The β -agonist drug class is a treatment that is often used in chronic obstructive pulmonary disease (COPD) where the use of this drug also has the potential for cardiovascular side effects in obstructive pulmonary disease patients. Therefore it is recommended the use of β -agonists must be careful in patients with pulmonary disease or severe cardiovascular disease and who use BB⁶¹. β_2 -adrenoceptor stimulation in the lungs can cause bronchodilation and has been recommended as a short-term treatment and long-term maintenance therapy for patients with asthma bronchial and chronic obstruction pulmonary disease (COPD) or better known as COPD⁶². According to *The Global Initiative for Chronic Obstructive Lung Disease (GOLD)* and *The European Society of Cardiology (ESC) Guidelines*, BB has been recommended as a treatment for HF with reduced ejection fraction (HFrEF) accompanied by COPD. In this case, the BB that must be chosen is a selective β_1 blocker (eg bisoprolol, metoprolol, or nebivolol) while non-selective is not recommended^{63,64}. The use of BB (β_1 selective) can be considered in several ways: First, reducing the risk of COPD exacerbations and heart disease by reducing the influence of adrenergic and inflammatory effects. Second, during COPD exacerbations, there is an increase in catecholamines in circulation, which will increase the risk of myocardial ischemia, tachycardia, worsening HF, and hypertension including increasing the risk of stroke. All cardiovascular risks involving endogenous exacerbations can be corrected by BB administration. Third, the administration of a β -agonist (salbutamol) in patients with COPD exacerbations causes tachycardia. Salbutamol is a bad selective ligand and causes β_1 -adrenoceptor activation in the heart. This shows that the administration of β_2 -agonists has been

shown to increase the risk of CVD, therefore BB administration is needed to prevent cardiovascular risk induced by β_2 -agonists [51,54]^{62,65}.

Propranolol can inhibit G protein-dependent signaling and reduce MAPK activation through β -dependent signaling, which results in reduced MUC5AC expression and mucosal hypersecretion induced by cigarette smoke. Chronic administration of β -inverse agonists such as propranolol to animal models of obstructive respiratory disease and clinical trials shows decreased lung infiltrates, increased respiratory smooth muscle response to β -agonists, restores β_2 -adrenoceptor (β_2 -AR) expression, decreased norepinephrine in serum, decreased cytokines (TNF- α , IL-8), and decreased mucus production and eosinophilic inflammation. Meanwhile, acute administration of β_2 -AR inverse agonists such as nadolol or carvedilol does not affect the airway response, but after 28 days of treatment, the inverse agonist can significantly reduce the airway response to antigens⁶⁶.

According to a cohort study that the risk of COPD patients is reduced in the group receiving BB among patients with comorbidities such as HF, coronary artery disease (CAD), ischemic heart disease, hypertension, cardiac arrhythmias, and pulmonary circulation disease (pulmonary embolism and cor pulmonale)⁶⁷. This is also supported by a prospective COPD Gene cohort study proving that BB has an acceptable safety profile in patients with GOLD II to IV criteria, including severe COPD that uses oxygen at home, and shows a decrease in exacerbations in the population⁶⁸. Based on this analysis, several meta-analyses from cohort trials mention the use of BB, especially selective β_1 including metoprolol, bisoprolol, and nebivolol can be considered in reducing COPD exacerbations with clear heart disease (eg. HFrEF or post-MI), but use BB combined with other drugs (ex. CCB or vasodilator drugs) need to be monitored because they can cause adverse effects⁶⁹.

β -BLOCKER AND HEPATIC CIRRHOSIS

In hepatic cirrhosis, splanchnic vasodilation occurs accompanied by increased blood flow to the splanchnic organs and portal system. This causes systemic vasodilation and will subsequently trigger plasma volume expansion and increase CO. If this hyperkinetic circulation continues to increase blood flow to the splanchnic organs, then an increase in portal pressure occurs⁷⁰. As a result of splanchnic vasodilation causes a progressive decrease in circulating volume thereby triggering activation of the neurohumoral system (SNS, renin-angiotensin-aldosterone system [RAAS], and arginine vasopressin [AVP] system). If the body cannot compensate to compensate for the vasodilation, it will lead to refractory ascites, hyponatremia, and even hepatorenal syndrome (HRS)⁷¹.

NSBB is a treatment that is often used for portal hypertension in hepatic cirrhosis. According to some RCTs, it has been proven that non selective BB (NSBB) can prevent initial variceal bleeding and subsequent rebleeding. Other evidence reports that NSBB prevents liver decompensation in patients with compensated cirrhosis. Therefore, the introduction of a new NSBB, carvedilol, has a better therapeutic effect than the previous NSBB⁷². Carvedilol has an intrinsic anti- α_1 adrenergic activity that causes intrahepatic vasodilation so that it can reduce portal pressure⁷¹. Carvedilol is reported to be more effective in decreasing hepatic venous pressure gradient (HVPG) than propranolol or nadolol⁷³. Moreover, carvedilol tends not to cause hypotension but reduces portal pressure significantly beyond propranolol. This

explains that carvedilol has been better tolerated than propranolol⁷⁰. In addition, carvedilol has a cytoprotective, antioxidant effect by scavenging and suppressing reactive oxygen species (ROS), anti-inflammatory, anti-fibrotic effect, increase insulin sensitivity and improve mitochondrial function^{74,75}. NSBB has been reported to reduce intestinal permeability and LPS-binding protein (LBP-dissolved protein acute phase response) and IL-6 which were found to be high related to the risk of varicose bleeding⁷⁵.

The β -adrenergic blocker has an effect on decreasing hepatic venous pressure gradient (HVPG) in patients with clinically significant portal hypertension (CSPH), in which patients with splanchnic vasodilation and hyperdynamic circulation occur. This explains the use of NSBB might be able to prevent the development of varicose veins in CSPH patients. NSBB mechanically acts by lowering the heart rate (HR) and inhibits splenic vasodilatation, thus the hemodynamic effect of NSBB depends on the severity of the hyperdynamic condition. This shows that the use of NSBB has advantages in splanchnic hemodynamics and vice versa disadvantages in systemic hemodynamics⁷⁶. The use of NSBB in hepatic cirrhosis is proven to effectively reduce portal pressure which has an impact on reducing the risk of bleeding and rebleeding of varicose veins. Based on Baveno IV and *The American Association for the Study of Liver Diseases (AASLD) Guidelines* recommend NSBB as primary and secondary prophylaxis (combination with endoscopic band ligation (EBL)) bleeding varicose veins in patients with cirrhosis accompanied by esophageal varices⁷⁶. However, not all NSBBs have the same mechanism, because the presence of additional anti- α 1-adrenergic activity in carvedilol is more effective in reducing portal pressure, but can cause a clear decrease in systemic arterial pressure⁷⁶. Carvedilol has also been shown to prevent rebleeding (secondary prophylaxis) of varicose veins in cirrhosis than other NSBBs⁷⁷. Overall, carvedilol is a treatment of portal hypertension in cirrhosis patients with safe and good efficacy of esophageal varices⁷⁸.

Another beneficial effect of NSBB through reduced portal hypertension and its sympatholytic action can increase intestinal congestion and edema by normalizing intestinal transit so as to prevent bacterial translocation. In animal models of portal hypertension, propranolol can increase intestinal motility and reduce the overgrowth of enteric bacterial flora, migration of microbiota into the systemic circulation and the development of spontaneous bacterial peritonitis (SBP). This is supported by a meta-analysis study showing the protective effect of NSBB on the development of SBP⁷⁴. Meanwhile, current guidelines recommend NSBB or EBL as an equivalent therapy for primary prophylaxis, but there is new evidence reporting that NSBB use in compensated cirrhosis patients accompanied by CSPH can prevent the first decompensation, ie ascites⁷⁹.

β -BLOCKER AND HYPERTHYROIDISM

Hyperthyroidism is a metabolic disorder in which the thyroid gland produces too much thyroid hormone. This situation can be controlled by administering drugs, but some people fall into worsening conditions such as thyroid storm and thyrotoxicosis⁸⁰. Thyroid storm is a life-threatening condition that requires prompt diagnosis and prompt treatment. This situation has manifestations of decompensation in various organs including loss of consciousness, high fever, HF, diarrhea, and jaundice⁸¹. Whereas thyrotoxicosis refers to the clinical state that is

produced by the activity of thyroid hormone that is too high in the tissue and is generally caused by inappropriate levels of tissue thyroid hormone. While hyperthyroidism is included in the form of thyrotoxicosis due to improper synthesis and secretion of thyroid hormone⁸². As a result, there is an increased risk of atrial fibrillation, embolic events, coronary events, and heart failure, especially in elderly patients and in those with underlying heart disease⁸³. In overcoming such conditions it is important especially to deal with cardiac dysfunction which if not treated immediately will continue to worsen⁸⁴. The most common medication given to thyrotoxicosis patients to improve symptoms is non-cardioselective BB⁸². The advantage of using BB is necessary because the cardiac manifestations of hyperthyroidism are caused by an increase in the action of catecholamines. In the thyrotoxicosis condition there is an increase in tissue sensitivity to catecholamines and an increase in the number of β 1-adrenergic receptors and protein binding protein guanosine triphosphate. This supports the fact that the administration of NSBB in hyperthyroid patients is especially useful for managing heart problems⁸⁵.

NSBB such as propranolol (20-40 mg every 6 hours) or who work longer hours (eg atenolol/bisoprolol) can control adrenergic symptoms such as palpitations and tremors, especially during the initial stages before antithyroid drugs. High doses of propranolol (40 mg qid) can inhibit peripheral T4 to T3 conversion by inhibiting the enzymatic activity of Dio1 (iodothyronine deiodinases). Specifically, thyroid hormone activation is mediated by the conversion of Dio1 and Dio2 from T4 to T3^{86,87}. Whereas cardioselective BB with high cardioprotective effects and good prevention of atrial fibrillation are alternative options, especially for patients with asthma. The use of NSBB is needed to control hyperthyroidism especially before thyroidectomy therapy⁸⁶. NSBB (eg propranolol) is a standard treatment in thyroid storm and thyrotoxicosis to ward off hyperadrenergic states that can trigger a cardiovascular collapse in pre-existing cardiac dysfunction⁸⁸. Meanwhile, tachycardia and atrial fibrillation that occur in thyroid storm are also recommended to get intravenous β 1-selective BB (landiolol, esmolol) or oral (bisoprolol) as treatment⁸¹.

β -BLOCKER AND SEPSIS

Sepsis is a complex condition characterized by simultaneous activation of inflammation and coagulation in response to microbes. Sepsis has manifestations as a systemic inflammatory response syndrome or symptoms of sepsis through the release of proinflammatory cytokines, procoagulants, and adhesion molecules from damaged immune cells and/or endothelium. At present, sepsis is a severe multisystem disease with a high mortality rate⁸⁹. In general, first-line management for sepsis is a combination of alpha and beta-agonist, generally noradrenaline and dobutamine. However, there are several potential mechanisms underlying the protective effect of BB during sepsis, both cardiac and non-cardiac⁹⁰.

High levels of endogenous and exogenous catecholamines are most likely due to excessive sympathetic stimulation in sepsis, despite adequate fluid resuscitation. This causes diastolic dysfunction, the dominant phenotype in myocardial dysfunction associated with sepsis so that BB can be indicated in this condition. BBs such as esmolol are reported to reduce HR which allows better diastolic filling and weakens the toxic effects of catecholamine. It is

characterized by inflammation, oxidative stress and abnormal intracellular calcium trafficking that can cause apoptosis and even necrosis⁹⁰. Increased catecholamine levels in sepsis cause catecholamine-induced cardiomyopathy and cardiac damage due to excess calcium, which causes cardiomyocyte necrosis. Sepsis also causes a decrease in myocardial beta-adrenergic receptor density and transduction of the disturbed b-adrenergic stimulant signal. Therefore, prevention of further cardiomyocyte damage due to excessive SNS stimulation is the key to managing sepsis⁹¹.

In septic shock, BB can stabilize circulation and increase myocardial damage. BB has a role in suppressing catecholamine storms, controlling ventricular velocity and reducing mortality in acute myocardial infarction (AMI) patients and chronic heart failure. Propranolol has been reported to reduce ventricular progress by 15% in children with severe burns, inhibit heart stress caused by burns, reduce heart work, and have no significant effect on MAP. Esmolol has been widely used as a study in critical patients, because of its high selectivity to β_1 receptor and its short and fast acting characteristics. A study also showed that β_1 -blockers do not increase oxygen consumption in septic patients and do not affect blood flow in the liver and lower extremities or peripheral vascular resistance therefore tissue perfusion can be maintained. The beneficial effect of BB in sepsis patients is supported by a systematic review and meta-analysis that esmolol is safe and effective in improving 28-day mortality and controlling the ventricular rate in sepsis patients after fluid resuscitation and does not have a significant adverse effect on tissue perfusion⁹².

β -BLOCKER AND GLAUCOMA

Glaucoma is one of the main causes of irreversible blindness. The most common type of glaucoma is primary open-angle glaucoma. Increased intraocular pressure (IOP) is often associated with primary open-angle glaucoma due to chronic progressive resistance to drainage of aqueous humor through the trabecular meshwork in the anterior chamber. Increased IOP is a major risk factor in the onset and development of glaucoma and is the only risk factor that can be modified. Common topical treatments used to reduce IOP include selective or non-selective BB⁹³.

The mechanism of action of BB for the treatment of glaucoma by reducing IOP through the production of suppressed aqueous humor. In general, it is believed that the mechanism is based on interactions with the β_2 -adrenergic receptor in the ciliary epithelium because β -receptors are found in the ciliary eye and are mostly β_2 subtypes. BB acts to induce vasoconstriction in the ciliary arteries so that it can reduce the production of aqueous humor. BB can reach the posterior eye segment after topical application can cause vasoconstriction and negatively affect ocular blood flow. BB is available as a solution for the eyes such as timolol, betaxolol, carteolol, levobunolol, and metipranolol⁹⁴. There is evidence that shows the use of topical BB as glaucoma therapy for patients with respiratory and cardiovascular complications leading to adverse outcomes⁹³. The first BB available for eye solution is timolol. Previously timolol was often used as a treatment for glaucoma. But now the use of timolol has decreased since the presence of a new agent namely prostaglandin analogs because of its ability to reduce IOP better and minimal side effects than BB^{95,96}.

CONCLUSION

Several studies have shown the efficacy of BB for several diseases with or without other comorbidities, which is evidenced by the significant decrease in mortality and morbidity in some cases, especially CVD. Its use, known as antihypertension, has been widely debated because of the findings of several BB drugs and their pleiotropic effects. However, the presence of hypertension induced by SNS hyperactivation, proves that BB placement is very plausible as a first line treatment. After more than 50 years of its discovery, new findings have emerged that show a protective effect outside the cardiovascular system, especially in the field of oncology. In a retrospective clinical study, the clinical benefits of using BB as a cancer therapy have long been suspected, but this is still debated given there is no explanation that explains the optimal use of BB as adjuvant therapy or regarding its clinical justification. This new concept will continue to be carried out research and evaluation to improve the prognosis of cancer patients.

ACKNOWLEDGEMENTS

The authors would like to express my special thanks of gratitude to the Head of Master Program in Basic Medical Science, Faculty of Medicine Universitas Airlangga, Prof. Dr. Kuntaman, dr, MS, Sp.MK(K) for providing support and encouraging me throughout my study.

REFERENCES

1. F. Triposkiadis, G. Karayannis, G. Giamouzis, J. Skoularigis, G. Louridas, and J. Butler, "The Sympathetic Nervous System in Heart Failure. Physiology, Pathophysiology, and Clinical Implications," *J. Am. Coll. Cardiol.*, vol. 54, no. 19, pp. 1747-1762, 2009, doi: 10.1016/j.jacc.2009.05.015.
2. M. van Bilsen *et al.*, "The autonomic nervous system as a therapeutic target in heart failure: a scientific position statement from the Translational Research Committee of the Heart Failure Association of the European Society of Cardiology," *Eur. J. Heart Fail.*, vol. 19, no. 11, pp. 1361-1378, 2017, doi: 10.1002/ejhf.921.
3. P. Hanna, K. Shivkumar, and J. L. Ardell, "Calming the Nervous Heart: Autonomic Therapies in Heart Failure," *Card. Fail. Rev.*, vol. 4, no. 2, p. 92, 2018, doi: 10.15420/cfr.2018.20.2.
4. C. Fumagalli, N. Maurizi, N. Marchionni, and D. Fornasari, " β -blockers: Their new life from hypertension to cancer and migraine," *Pharmacol. Res.*, vol. 151, no. September 2019, p. 104587, 2020, doi: 10.1016/j.phrs.2019.104587.
5. C. Wiysonge, H. Bradley, J. Volmink, B. Mayosi, and L. Opie, "Beta-blockers for hypertension. Cochrane Database of Systematic Reviews," *Cochrane Database Syst. Rev.*, vol. 1, no. 1, pp. 1-74, 2017, doi: 10.1002/14651858.CD002003.pub5.www.cochranelibrary.com.
6. J. G. Baker, "The selectivity of β -adrenoceptor antagonists at the human β_1 , β_2 and β_3 adrenoceptors," *Br. J. Pharmacol.*, vol. 144, no. 3, pp. 317-322, 2005, doi: 10.1038/sj.bjp.0706048.
7. O. E. Brodde, H. Bruck, and K. Leineweber, "Cardiac adrenoceptors: Physiological and pathophysiological relevance," *J. Pharmacol. Sci.*, vol. 100, no. 5, pp. 323-337, 2006, doi: 10.1254/jphs.CRJ06001X.
8. R. Gordan, J. K. Gwathmey, and L.-H. Xie, "Autonomic and endocrine control of cardiovascular function," *World J. Cardiol.*, vol. 7, no. 4, p. 204, 2015, doi:

- 10.4330/wjcv7.i4.204.
9. N. S. Rehsia and N. S. Dhalla, "Mechanisms of the beneficial effects of beta-adrenoceptor antagonists in congestive heart failure," *Exp. Clin. Cardiol.*, vol. 15, no. 4, 2010.
 10. M. G. Perrone and A. Scilimati, *β 3 -Adrenoceptor agonists and (Antagonists as) inverse agonists. History, perspective, constitutive activity, and stereospecific binding*, 1st ed., vol. 484, no. C. Elsevier Inc., 2010.
 11. G. Pucci, M. G. Ranalli, F. Battista, and G. Schillaci, "Effects of β -Blockers with and Without Vasodilating Properties on Central Blood Pressure: Systematic Review and Meta-Analysis of Randomized Trials in Hypertension," *Hypertension*, vol. 67, no. 2, pp. 316–324, 2016, doi: [10.1161/HYPERTENSIONAHA.115.06467](https://doi.org/10.1161/HYPERTENSIONAHA.115.06467).
 12. D. E. Becker, "Cardiovascular Drugs : Implications for Dental Practice," *Anesth Prog*, vol. 3006, no. 07, pp. 178–186, 2007.
 13. R. Khatib and F. Wilson, *Pharmacology of Medications Used in the Treatment of Atherosclerotic Cardiovascular Disease*. Elsevier Inc., 2018.
 14. C. Ham et al., "Beta-blocker use and fall risk in older individuals: Original results from two studies with meta-analysis," *Br. J. Clin. Pharmacol.*, vol. 83, no. 10, pp. 2292–2302, 2017, doi: [10.1111/bcp.13328](https://doi.org/10.1111/bcp.13328).
 15. Bain, "Beta-blocker use in cardiovascular disease," *Br. J. Card. Nurs.*, vol. 13, no. 10, pp. 491–497, 2018, doi: [10.12968/bjca.2018.13.10.491](https://doi.org/10.12968/bjca.2018.13.10.491).
 16. Kravets, "Hyperthyroidism: Diagnosis," *Am. Fam. Physician*, vol. 93, no. 5, pp. 364–370, 2016, doi: [10.1007/BF01579317](https://doi.org/10.1007/BF01579317).
 17. R. J. Groszmann et al., "Beta-blockers to prevent gastroesophageal varices in patients with cirrhosis," *N. Engl. J. Med.*, vol. 353, no. 21, pp. 2254–2261, 2005, doi: [10.1056/NEJMoa044456](https://doi.org/10.1056/NEJMoa044456).
 18. Albouaini, M. Andron, A. Alahmar, and M. Egred, "Beta-blockers use in patients with chronic obstructive pulmonary disease and concomitant cardiovascular conditions," *Int J. COPD*, vol. 2, no. 4, pp. 535–540, 2007.
 19. S. Duffy et al., "Effect of beta-blockers on exacerbation rate and lung function in chronic obstructive pulmonary disease (COPD)," *Respir. Res.*, vol. 18, no. 1, pp. 4–10, 2017, doi: [10.1186/s12931-017-0609-7](https://doi.org/10.1186/s12931-017-0609-7).
 20. Y. R. Lee, M. S. Seth, D. Soney, and H. Dai, "Benefits of Beta-Blockade in Sepsis and Septic Shock: A Systematic Review," *Clin. Drug Investig.*, no. 0123456789, 2019, doi: [10.1007/s40261-019-00762-z](https://doi.org/10.1007/s40261-019-00762-z).
 21. Tan, M. Harazim, B. Tang, A. McLean, and M. Nalos, "The association between pre-morbid beta blocker exposure and mortality in sepsis - A systematic review," *Crit. Care*, vol. 23, no. 1, pp. 1–12, 2019, doi: [10.1186/s13054-019-2562-y](https://doi.org/10.1186/s13054-019-2562-y).
 22. Z. Na et al., "The effects of beta-blocker use on cancer prognosis: A meta-analysis based on 319,006 patients," *Onco. Targets. Ther.*, vol. 11, pp. 4913–4944, 2018, doi: [10.2147/OTT.S167422](https://doi.org/10.2147/OTT.S167422).
 23. Montoya et al., "Use of non-selective β -blockers is associated with decreased tumor proliferative indices in early stage breast cancer," *Oncotarget*, vol. 8, no. 4, pp. 6446–6460, 2017, doi: [10.18632/oncotarget.14119](https://doi.org/10.18632/oncotarget.14119).
 24. R. Udumyan et al., "Beta-blocker drug use and survival among patients with pancreatic adenocarcinoma," *Cancer Res.*, vol. 77, no. 13, pp. 3700–3707, 2017, doi: [10.1158/0008-5472.CAN-17-0108](https://doi.org/10.1158/0008-5472.CAN-17-0108).
 25. O. Fiala et al., "Incidental use of beta-blockers is associated with outcome of metastatic colorectal cancer patients treated with bevacizumab-based therapy: A single-institution retrospective analysis of 514 patients," *Cancers (Basel)*, vol. 11, no. 12, pp. 1–13, 2019, doi: [10.3390/cancers11121856](https://doi.org/10.3390/cancers11121856).
 26. H. Choi et al., "Meta-analysis of the effects of beta blocker on survival time in cancer patients," *J. Cancer Res. Clin. Oncol.*, vol. 140, no. 7, pp. 1179–1188, 2014, doi: [10.1007/s00432-014-1658-7](https://doi.org/10.1007/s00432-014-1658-7).
 27. N. Amaya et al., "Non-selective beta blockers inhibit angiosarcoma cell viability and increase progression free- and overall-survival in patients diagnosed with metastatic angiosarcoma," *Oncoscience*, vol. 5, no. March, p. 124, 2018, doi: [10.18632/oncoscience.413](https://doi.org/10.18632/oncoscience.413).
 28. P. Le et al., "Chronic stress in mice remodels lymph vasculature to promote tumour cell dissemination," *Nat. Commun.*, vol. 7, 2016, doi: [10.1038/ncomms10634](https://doi.org/10.1038/ncomms10634).
 29. Juhasz et al., "The complex function of Hsp70 in metastatic cancer," *Cancers (Basel)*, vol. 6, no. 1, pp. 42–66, 2014, doi: [10.3390/cancers6010042](https://doi.org/10.3390/cancers6010042).
 30. Y. Y. Lei, W. J. Wang, J. H. Mei, and C. L. Wang, "Mitogen-activated protein kinase signal transduction in solid tumors," *Asian Pacific J. Cancer Prev.*, vol. 15, no. 20, pp. 8539–8548, 2014, doi: [10.7314/APJCP.2014.15.20.8539](https://doi.org/10.7314/APJCP.2014.15.20.8539).
 31. Renema, B. Navet, M. F. Heymann, F. Lezot, and D. Heymann, "RANK-cad signalling in cancer," *Biosci. Rep.*, vol. 36, no. 4, pp. 1–17, 2016, doi: [10.1042/BSR20160150](https://doi.org/10.1042/BSR20160150).
 32. J. F. Chung, S. J. Lee, and A. K. Sood, "Immunological and pleiotropic effects of individual β -blockers and their relevance in cancer therapies," *Expert Opin. Investig. Drugs*, vol. 25, no. 5, pp. 501–505, 2016, doi: [10.1517/13543784.2016.1164141](https://doi.org/10.1517/13543784.2016.1164141).
 33. R. Peixoto, M. de L. Pereira, and M. Oliveira, "Beta-blockers and cancer: Where are we?," *Pharmaceuticals*, vol. 13, no. 6, 2020, doi: [10.3390/ph13060105](https://doi.org/10.3390/ph13060105).
 34. T. A. Smitherman, R. Burch, H. Sheikh, and E. Loder, "The prevalence, impact, and treatment of migraine and severe headaches in the United States: A review of statistics from national surveillance studies," *Headache*, vol. 53, no. 3, pp. 427–436, 2013, doi: [10.1111/head.12074](https://doi.org/10.1111/head.12074).
 35. R. B. Lipton, M. E. Bigal, M. Diamond, F. Freitag, and M. L. Reed, "Migraine prevalence, disease burden, and the need for preventive therapy Herein we describe the epidemiology of migraine," *Neurology*, vol. 68, no. 5, pp. 343–349, 2007.
 36. P. L. Durham, "CGRP-Receptor Antagonists - A Fresh Approach to Migraine Therapy?," *N. Engl. J. Med.*, vol. 350, no. 11, pp. 1073–1075, 2004, doi: [10.1056/NEJMp048016](https://doi.org/10.1056/NEJMp048016).
 37. C. Rosa and R. Fantozzi, "The role of histamine in neurogenic inflammation," *Br. J. Pharmacol.*, vol. 170, no. 1, pp. 38–45, 2013, doi: [10.1111/bph.12266](https://doi.org/10.1111/bph.12266).
 38. Samsam, R. Covenas, R. Ahangari, J. Yajeya, and J. Narvaez, "Magnetic Nanoparticles From Fabrication to Clinical Applications," *Drug Dev. Res.*, vol. 64, no. April, pp. 294–314, 2007, doi: [10.1002/ddr](https://doi.org/10.1002/ddr).
 39. Borsook, N. Maleki, and R. Burstein, "Migraine," *Neurobiol. Brain Disord. Biol. Basis Neurol. Psychiatr. Disord.*, pp. 693–708, 2015, doi: [10.1016/B978-0-12-398270-4.00042-2](https://doi.org/10.1016/B978-0-12-398270-4.00042-2).
 40. T. Sprenger, M. Viana, and C. Tassorelli, "Current prophylactic medications for migraine and their

- potential mechanisms of action," *Neurotherapeutics*, vol. 15, no. 2, pp. 313–323, 2018, doi: [10.1007/s13311-018-0621-8](https://doi.org/10.1007/s13311-018-0621-8).
41. X. Zhang, D. Levy, V. Kainz, R. Noseda, M. Jakubowski, and R. Burstein, "Activation of central trigeminovascular neurons by cortical spreading depression," *Ann. Neurol.*, vol. 69, no. 5, pp. 855–865, 2011, doi: [10.1002/ana.22329](https://doi.org/10.1002/ana.22329).
 42. S. D. Silberstein, "Preventive migraine treatment," *Contin. Lifelong Learn. Neurol.*, vol. 21, no. 4, pp. 973–989, 2015, doi: [10.1212/CON.0000000000000199](https://doi.org/10.1212/CON.0000000000000199).
 43. K. G. Shields and P. J. Goadsby, "Propranolol modulates trigeminovascular responses in thalamic ventroposteromedial nucleus: A role in migraine?," *Brain*, vol. 128, no. 1, pp. 86–97, 2005, doi: [10.1093/brain/awh298](https://doi.org/10.1093/brain/awh298).
 44. Urits et al., "CGRP Antagonists for the Treatment of Chronic Migraines: a Comprehensive Review," *Curr. Pain Headache Rep.*, vol. 23, no. 5, pp. 1–10, 2019, doi: [10.1007/s11916-019-0768-y](https://doi.org/10.1007/s11916-019-0768-y).
 45. M. Ramadan, "Prophylactic migraine therapy: Mechanisms and evidence," *Curr. Pain Headache Rep.*, vol. 8, no. 2, pp. 91–95, 2004, doi: [10.1007/s11916-004-0022-z](https://doi.org/10.1007/s11916-004-0022-z).
 46. A. Pradhan, Z. Bertels, and S. Akerman, "Targeted Nitric Oxide Synthase Inhibitors for Migraine," *Neurotherapeutics*, vol. 15, no. 2, pp. 391–401, 2018, doi: [10.1007/s13311-018-0614-7](https://doi.org/10.1007/s13311-018-0614-7).
 47. GOLD 2017, "Global Initiative for Chronic Obstructive Lung Disease: Pocket Guide To COPD Diagnosis, Management, and Prevention, A Guide for Health Care Professionals," *Gold*, pp. 1–33, 2017, doi: [10.1097/00008483-200207000-00004](https://doi.org/10.1097/00008483-200207000-00004).
 48. R. Bracke and G. G. Brusselle, *Chronic Obstructive Pulmonary Disease*, Fourth Edi., vol. 2–2. Elsevier, 2015.
 49. J. Barnes, "Cellular and molecular mechanisms of chronic obstructive pulmonary disease," *Clin. Chest Med.*, vol. 35, no. 1, pp. 71–86, 2014, doi: [10.1016/j.ccm.2013.10.004](https://doi.org/10.1016/j.ccm.2013.10.004).
 50. Maltais et al., "β-Blockers in COPD: A Cohort Study From the TONADO Research Program," *Chest*, vol. 153, no. 6, pp. 1315–1325, 2018, doi: [10.1016/j.chest.2018.01.008](https://doi.org/10.1016/j.chest.2018.01.008).
 51. G. Baker and R. G. Wilcox, "β-Blockers, heart disease and COPD: Current controversies and uncertainties," *Thorax*, vol. 72, no. 3, pp. 271–276, 2017, doi: [10.1136/thoraxjnl-2016-208412](https://doi.org/10.1136/thoraxjnl-2016-208412).
 52. J. Vestbo et al., "Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease GOLD executive summary," *Am. J. Respir. Crit. Care Med.*, vol. 187, no. 4, pp. 347–365, 2013, doi: [10.1164/rccm.201204-0596PP](https://doi.org/10.1164/rccm.201204-0596PP).
 53. Ponikowski et al., "2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure," *Eur. Heart J.*, vol. 37, no. 27, pp. 2129–2200m, 2016, doi: [10.1093/eurheartj/ehw128](https://doi.org/10.1093/eurheartj/ehw128).
 54. W. MacNee, "Beta-blockers in COPD — A controversy resolved?," *N. Engl. J. Med.*, vol. 381, no. 24, pp. 2367–2368, 2019, doi: [10.1056/NEJMe1912664](https://doi.org/10.1056/NEJMe1912664).
 55. D. W. Tan, J. L. Wong, S. T. Tie, J. A. Abisheganaden, A. Y. Lim, and W. F. Wong, "β 2 -adrenoceptor in obstructive airway diseases: Agonism, antagonism or both?," *World J. Respiriol.*, vol. 5, no. 3, pp. 199–206, 2014, doi: [10.5320/wjr.v5.i3.199](https://doi.org/10.5320/wjr.v5.i3.199).
 56. O. Nielsen, L. Pedersen, B. F. Sode, and M. Dahl, "β-Blocker Therapy and Risk of Chronic Obstructive Pulmonary Disease – A Danish Nationwide Study of 1.3 Million Individuals," *EclinicalMedicine*, vol. 7, pp. 21–26, 2019, doi: [10.1016/j.eclinm.2019.01.004](https://doi.org/10.1016/j.eclinm.2019.01.004).
 57. Lipworth, J. Wedzicha, G. Devereux, J. Vestbo, and M. T. Dransfield, "Beta-blockers in COPD: Time for reappraisal," *Eur. Respir. J.*, vol. 48, no. 3, pp. 880–888, 2016, doi: [10.1183/13993003.01847.2015](https://doi.org/10.1183/13993003.01847.2015).
 58. G. Rodrigues, Y. P. Mendoza, and J. Bosch, "Beta-blockers in cirrhosis: Evidence-based indications and limitations," *JHEP Reports*, vol. 2, no. 1, p. 100063, 2020, doi: [10.1016/j.jhepr.2019.12.001](https://doi.org/10.1016/j.jhepr.2019.12.001).
 59. Moctezuma-Velazquez, S. Kalainy, and J. G. Abraldes, "Beta-blockers in patients with advanced liver disease: Has the dust settled?," *Liver Transplant.*, vol. 23, no. 8, pp. 1058–1069, 2017, doi: [10.1002/lt.24794](https://doi.org/10.1002/lt.24794).
 60. V. Giannelli, B. Lattanzi, U. Thalheimer, and M. Merli, "Beta-blockers in liver cirrhosis," *Ann. Gastroenterol.*, vol. 27, no. 1, pp. 20–26, 2014.
 61. N. Rajoriya and D. Tripathi, "Non-selective beta-blockers in cirrhosis: Current concepts and controversies," *World J. Pharmacol.*, vol. 5, no. 1, pp. 15–31, 2016, doi: [10.5497/wjp.v5.i1.15](https://doi.org/10.5497/wjp.v5.i1.15).
 62. Reiberger and M. Mandorfer, "Beta adrenergic blockade and decompensated cirrhosis," *J. Hepatol.*, vol. 66, no. 4, pp. 849–859, 2017, doi: [10.1016/j.jhep.2016.11.001](https://doi.org/10.1016/j.jhep.2016.11.001).
 63. J. Yang, K. Ge, L. Chen, and J. Le Yang, "The efficacy comparison of carvedilol plus endoscopic variceal ligation and traditional, nonselective β-blockers plus endoscopic variceal ligation in cirrhosis patients for the prevention of variceal rebleeding: A meta-analysis," *Eur. J. Gastroenterol. Hepatol.*, vol. 31, no. 12, pp. 1518–1526, 2019, doi: [10.1097/MEG.0000000000001442](https://doi.org/10.1097/MEG.0000000000001442).
 64. P. Zacharias, R. Jeyaraj, L. Hobolth, F. Bendtsen, L. L. Gluud, and M. Y. Morgan, "Carvedilol versus traditional, non-selective beta-blockers for adults with cirrhosis and gastroesophageal varices," *Cochrane Database Syst. Rev.*, vol. 2018, no. 10, 2018, doi: [10.1002/14651858.CD011510.pub2](https://doi.org/10.1002/14651858.CD011510.pub2).
 65. Villanueva et al., "β blockers to prevent decompensation of cirrhosis in patients with clinically significant portal hypertension (PREDESCI): a randomised, double-blind, placebo-controlled, multicentre trial," *Lancet*, vol. 393, no. 10181, pp. 1597–1608, 2019, doi: [10.1016/S0140-6736\(18\)31875-0](https://doi.org/10.1016/S0140-6736(18)31875-0).
 66. N. Ferretti and J. Yee, "UC Irvine," *J. Educ. Teach. Emerg. Med.*, vol. 4, no. 3, pp. 1–24, 2019.
 67. Satoh et al., "2016 Guidelines for the management of thyroid storm from The Japan Thyroid Association and Japan Endocrine Society (First edition)," *Endocr. J.*, vol. 63, no. 12, pp. 1025–1064, 2016.
 68. D. S. Ross et al., "2016 American Thyroid Association Guidelines for Diagnosis and Management of Hyperthyroidism and Other Causes of Thyrotoxicosis," *Thyroid*, vol. 26, no. 10, pp. 1343–1421, 2016, doi: [10.1089/thy.2016.0229](https://doi.org/10.1089/thy.2016.0229).
 69. B. Biondi and G. J. Kahaly, "Cardiovascular involvement in patients with different causes of hyperthyroidism," *Nat. Rev. Endocrinol.*, vol. 6, no. 8, pp. 431–443, 2010, doi: [10.1038/nrendo.2010.105](https://doi.org/10.1038/nrendo.2010.105).
 70. Genev, M. D. Lundholm, M. A. Emanuele, E. McGee, and V. Mathew, "Thyrotoxicosis-induced cardiomyopathy treated with venoarterial extracorporeal membrane oxygenation," *Hear. Lung*, vol. 49, no. 2, pp. 165–166, 2020, doi: [10.1016/j.hrtng.2019.06.006](https://doi.org/10.1016/j.hrtng.2019.06.006).
 71. P. M. Osuna, M. Udovicic, and M. D. Sharma,

- "Hyperthyroidism and the Heart," *Methodist Debakey Cardiovasc. J.*, vol. 13, no. 2, pp. 60–63, 2017, doi: [10.14797/mdcj-13-2-60](https://doi.org/10.14797/mdcj-13-2-60).
72. J. Kahaly, L. Bartalena, L. Hegedüs, L. Leenhardt, K. Poppe, and S. H. Pearce, "2018 European thyroid association guideline for the management of graves' hyperthyroidism," *Eur. Thyroid J.*, vol. 7, no. 4, pp. 167–186, 2018, doi: [10.1159/000490384](https://doi.org/10.1159/000490384).
73. B. K. Singh and P. M. Yen, "A clinician's guide to understanding resistance to thyroid hormone due to receptor mutations in the TR α and TR β isoforms," *Clin. Diabetes Endocrinol.*, vol. 3, no. 1, pp. 1–11, 2017, doi: [10.1186/s40842-017-0046-z](https://doi.org/10.1186/s40842-017-0046-z).
74. Abubakar, V. Singh, A. Arora, and S. Alsunaid, "Propranolol-Induced Circulatory Collapse in a Patient With Thyroid Crisis and Underlying Thyrocardiac Disease: A Word of Caution," *J. Investig. Med. High Impact Case Reports*, vol. 5, no. 4, 2017, doi: [10.1177/2324709617747903](https://doi.org/10.1177/2324709617747903).
75. G. Polat, R. A. Ugan, E. Cadirci, and Z. Halici, "Sepsis ve septik şok: Mevcut tedavi stratejileri ve yeni yaklaşımlar," *Eurasian J. Med.*, vol. 49, no. 1, pp. 53–58, 2017, doi: [10.5152/eurasianjmed.2017.17062](https://doi.org/10.5152/eurasianjmed.2017.17062).
76. E. C. Boerma and M. Singer, "Beta-blockers in sepsis: Time to reconsider current constraints?," *Br. J. Anaesth.*, vol. 119, no. 4, pp. 560–561, 2017, doi: [10.1093/bja/aex266](https://doi.org/10.1093/bja/aex266).
77. Suzuki *et al.*, "Sepsis-induced cardiac dysfunction and β -adrenergic blockade therapy for sepsis," *J. Intensive Care*, vol. 5, no. 1, pp. 1–10, 2017, doi: [10.1186/s40560-017-0215-2](https://doi.org/10.1186/s40560-017-0215-2).
78. Li, W. Sun, Y. Guo, Y. Ren, Y. Li, and Z. Yang, "Prognosis of β -adrenergic blockade therapy on septic shock and sepsis: A systematic review and meta-analysis of randomized controlled studies," *Cytokine*, vol. 126, no. July 2019, 2020, doi: [10.1016/j.cyto.2019.154916](https://doi.org/10.1016/j.cyto.2019.154916).
79. Viswanathan *et al.*, "Resource Utilization Among Glaucoma Patients in the UK Treated with Beta-Blocker and Non-Beta-Blocker Adjunctive Therapy: A Retrospective Cohort Analysis," *Adv. Ther.*, vol. 34, no. 7, pp. 1695–1706, 2017, doi: [10.1007/s12325-017-0541-8](https://doi.org/10.1007/s12325-017-0541-8).
80. M. Hoste, "Beta-Blockers," *Glaucoma Second Ed.*, vol. 1, pp. 548–558, 2015, doi: [10.1016/B978-0-7020-5193-7.00053-4](https://doi.org/10.1016/B978-0-7020-5193-7.00053-4).
81. H. Heo, "Comparison of Prostaglandin Analog Treatment Patterns," *J. Manag. Care Spec. Pharm.*, vol. 25, no. 9, pp. 1001–1010, 2019.
82. T. Li *et al.*, "Comparative Effectiveness of First Line Meds in POAG," vol. 123, no. 1, pp. 129–140, 2017, doi: [10.1016/j.ophtha.2015.09.005.Comparative](https://doi.org/10.1016/j.ophtha.2015.09.005.Comparative).
83. P. Andell *et al.*, "B-Blocker use and mortality in copd patients after myocardial infarction: A swedish nationwide observational study," *J. Am. Heart Assoc.*, vol. 4, no. 4, pp. 1–8, 2015, doi: [10.1161/JAHA.114.001611](https://doi.org/10.1161/JAHA.114.001611).
84. T. Tankeu *et al.*, "Effect of propranolol on heart rate variability in hyperthyroidism NCT03393728 NCT," *BMC Res. Notes*, vol. 11, no. 1, pp. 1–4, 2018, doi: [10.1186/s13104-018-3224-x](https://doi.org/10.1186/s13104-018-3224-x).
85. Gao *et al.*, "Effect of propranolol on atrial fibrillation in hyperthyroidism," *Int J. Clin. Exp. Med.*, vol. 9, no. 2, pp. 1245–1253, 2016.

Table 1. Pleiotropic Effects of Beta-Blockers

Disease	BB Recommendation	Mechanism of Action	References
Vascular System			
Cancer	Propranolol	<ul style="list-style-type: none"> ▪ Anti-apoptotic ▪ Anti-proliferative ▪ Anti-angiogenesis 	[4,23,27] [30,33]
Hepatic cirrhosis is accompanied by: <ul style="list-style-type: none"> ▪ First bleeding variceal ▪ Variceal rebleeding 	Carvedilol Propranolol	<ul style="list-style-type: none"> ▪ Vasculoprotective ▪ Cytoprotective ▪ Antioxidant effect ▪ Anti-inflammatory ▪ Anti-fibrotic effect 	[58,59,60] [61,62,63]
Glaucoma	Timolol Betaxolol Carteolol Levobunolol Metirapranolol	<ul style="list-style-type: none"> ▪ Vasculoprotective ▪ Suppress aqueous humor production 	⁹⁴
Neurogenic System			
Migraine with aura (<i>classic migraine</i>)	Propranolol Metoprolol Timolol	<ul style="list-style-type: none"> ▪ Neuroprotective ▪ Vasculoprotective 	[4,40,43]
Cardiac System			
<ul style="list-style-type: none"> ▪ COPD with HFrEF ▪ COPD with post-MI 	Bisoprolol Metoprolol Nebivolol	<ul style="list-style-type: none"> ▪ Cardioprotective ▪ Vasculoprotective ▪ Anti-inflammatory effects ▪ Antioxidant ▪ Anti-arrhythmic 	[4,57,83]
Sepsis	Esmolol Propranolol	Cardioprotective	[76,78]
Endocrine System			
<ul style="list-style-type: none"> ▪ Hyperthyroidism ▪ Thyrotoxicosis ▪ Thyroid storm 	Propranolol	<ul style="list-style-type: none"> ▪ Anti-hyperthyroid ▪ Anti-apoptotic ▪ Anti-arrhythmic 	[70,72] [84,85]

BB: Beta-blocker; COPD: Chronic obstructive pulmonary disease; HfrEF: Heart failure with reduced ejection fraction; MI: Myocardial infarction

A Novel Pleiotropic Effect of Beta-Blockers Useful or Not?

ORIGINALITY REPORT

20%

SIMILARITY INDEX

12%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Zainab Zuhair Ali, Fatimah J. Al-Hasani. "Evaluation of Surface Roughness of Some Biomedical Titanium Alloys by Pack Cementation Coating", Key Engineering Materials, 2021
Publication 1%
- 2** "Liver Diseases", Springer Science and Business Media LLC, 2020
Publication 1%
- 3** www.ncbi.nlm.nih.gov
Internet Source 1%
- 4** "Comprehensive Cardiovascular Medicine in the Primary Care Setting", Springer Science and Business Media LLC, 2019
Publication 1%
- 5** George J. Kahaly. "Chapter 1 Diagnosis and Endocrine Management of Graves' Disease", Springer Science and Business Media LLC, 2015
Publication <1%

6

Han Qiao, Zhaowen Cui, Shengbing Yang, Dingkun Ji et al. "Targeting Osteocytes to Attenuate Early Breast Cancer Bone Metastasis by Theranostic Upconversion Nanoparticles with Responsive Plumbagin Release", ACS Nano, 2017

Publication

<1 %

7

www.heanoti.com

Internet Source

<1 %

8

onlinelibrary.wiley.com

Internet Source

<1 %

9

www.mysciencework.com

Internet Source

<1 %

10

Tetsurou Satoh, Osamu Isozaki, Atsushi Suzuki, Shu Wakino et al. "2016 Guidelines for the management of thyroid storm from The Japan Thyroid Association and Japan Endocrine Society (First edition)", Endocrine Journal, 2016

Publication

<1 %

11

jintensiveware.biomedcentral.com

Internet Source

<1 %

12

Amanda C. Bultas, Besu Teshome, Sara K. Richter, Stephen Schafers, Emily Cooke, William B. Call. "Use of Nonselective β -Blockers in Patients With End-Stage Liver

<1 %

Disease and Select Complications", Annals of Pharmacotherapy, 2019

Publication

13

annalsofian.org

Internet Source

<1 %

14

docplayer.net

Internet Source

<1 %

15

clindiabetesendo.biomedcentral.com

Internet Source

<1 %

16

event.ners.unair.ac.id

Internet Source

<1 %

17

mts.intechopen.com

Internet Source

<1 %

18

"Proceedings: 15th Asia-Pacific Federation for Clinical Biochemistry and Laboratory Medicine (APFCB) Congress 2019 from 17-20 November, 2019, Jaipur, India", Indian Journal of Clinical Biochemistry, 2019

Publication

<1 %

19

Sanchit Sharma, Samagra Agarwal, Deepak Gunjan, Kanav Kaushal, Abhinav Anand, Srikant Mohta, Shalimar, Anoop Saraya.
"Long-term Outcomes with Carvedilol versus Propranolol in Patients with Index Variceal Bleed: 6-year Follow-up Study", Journal of Clinical and Experimental Hepatology, 2021

Publication

<1 %

20	brain.oxfordjournals.org Internet Source	<1 %
21	pt.scribd.com Internet Source	<1 %
22	trialsjournal.biomedcentral.com Internet Source	<1 %
23	David Borsook, Nasim Maleki, Rami Burstein. "Migraine", Elsevier BV, 2015 Publication	<1 %
24	Mohammed I.Y. Elmallah, Marine Cordonnier, Valentin Vautrot, Gaëtan Chanteloup, Carmen Garrido, Jessica Gobbo. "Membrane-anchored heat-shock protein 70 (Hsp70) in cancer", Cancer Letters, 2020 Publication	<1 %
25	Richard Gordan, Judith K Gwathmey, Lai-Hua Xie. "Autonomic and endocrine control of cardiovascular function", World Journal of Cardiology, 2015 Publication	<1 %
26	svmi.web.ve Internet Source	<1 %
27	www.dovepress.com Internet Source	<1 %
28	Ann M Hoste. "Beta-Blockers", Elsevier BV, 2015	<1 %

29

Rerdin Julario, Ricardo Adrian Nugraha, Bagus Putra Dharma Khrisna, Tony Santoso Putra et al. "Simple Laboratory and Clinical Parameters as Predictor for Electrocardiographic Abnormalities Among Hospitalized Patients with Chronic Kidney Disease: A Cross-Sectional Study", Cold Spring Harbor Laboratory, 2021

Publication

<1 %

30

Alexa Montoya, Clarissa N. Amaya, Andres Belmont, Nabih Diab et al. "Use of non-selective β -blockers is associated with decreased tumor proliferative indices in early stage breast cancer", Oncotarget, 2016

Publication

<1 %

31

Ivan Urits, Kyle Gress, Karina Charipova, Alec M. Zamarripa et al. "Pharmacological options for the treatment of chronic migraine pain", Best Practice & Research Clinical Anaesthesiology, 2020

Publication

<1 %

32

K. G. Shields. "Propranolol modulates trigeminovascular responses in thalamic ventroposteromedial nucleus: a role in migraine?", Brain, 11/17/2004

Publication

<1 %

33 Neil Rajoriya, Dhiraj Tripathi. "Non-selective beta-blockers in cirrhosis: Current concepts and controversies", World Journal of Pharmacology, 2016

Publication

<1 %

34 pesquisa.bvsalud.org

Internet Source

<1 %

35 Omar Sabra, Maria Muhammad Ali, Maha Al Zayer, Saleh Altuwaijri. "Frequency of Migraine as a Chief Complaint in Otolaryngology Outpatient Practice", 'Hindawi Limited', 2015

Internet Source

<1 %

36 globenewswire.com

Internet Source

<1 %

37 www.pubmedcentral.nih.gov

Internet Source

<1 %

38 www.science.gov

Internet Source

<1 %

39 Christian J. Steib, Alexander L. Gerbes. "Secondary prophylaxis for variceal bleeding: carvedilol vs. propranolol", Hepatology International, 2017

Publication

<1 %

40 François Maltais, Roland Buhl, Andrea Koch, Valeria C. Amatto et al. "β-Blockers in COPD",

<1 %

41 Xiao-Fang Li, Yi-Min Mao. "Beta-blockers in COPD: A systematic review based on recent research", Life Sciences, 2020 <1 %
Publication

42 core.ac.uk <1 %
Internet Source

43 utmb.influent.utsystem.edu <1 %
Internet Source

44 www.liebertpub.com <1 %
Internet Source

45 www.nature.com <1 %
Internet Source

46 www.thieme-connect.de <1 %
Internet Source

47 Wolfram Doehner, Dilek Ural, Karl Georg Haeusler, Jelena Čelutkienė et al. "Heart and brain interaction in patients with heart failure: overview and proposal for a taxonomy. A position paper from the Study Group on Heart and Brain Interaction of the Heart Failure Association", European Journal of Heart Failure, 2018 <1 %
Publication

48 repository.unair.ac.id
Internet Source

<1 %

49

thorax.bmj.com

Internet Source

<1 %

50

Carlos Moctezuma-Velazquez, Sylvia Kalainy, Juan G. Abrales. "Beta-blockers in patients with advanced liver disease: Has the dust settled?", Liver Transplantation, 2017

Publication

<1 %

51

Dell'Era, A.. "Acute variceal bleeding: Pharmacological treatment and primary/secondary prophylaxis", Best Practice & Research Clinical Gastroenterology, 200804

Publication

<1 %

52

Jacopo Agrimi, Danilo Menicucci, Marco Laurino, Chelsea D Mackey et al. "Cardiac-generated sympathetic stress alters heart-brain communication, reduces EEG-theta activity, and increases locomotor behavior", Cold Spring Harbor Laboratory, 2021

Publication

<1 %

53

Maxime Caru, Denis Corbin, Delphine Périé, Valérie Lemay et al. "Doxorubicin treatments induce significant changes on the cardiac autonomic nervous system in childhood acute lymphoblastic leukemia long-term survivors", Clinical Research in Cardiology, 2019

Publication

<1 %

54

Stephen D. Silberstein. "Current Preventive Therapy: Preventive Treatment Mechanisms", Headache Currents, 9/2006

Publication

<1 %

55

www.hindawi.com

Internet Source

<1 %

56

www.labome.org

Internet Source

<1 %

57

www.tandfonline.com

Internet Source

<1 %

58

David Kockerling, Rooshi Nathwani, Roberta Forlano, Pinelopi Manousou, Benjamin H Mullish, Ameet Dhar. "Current and future pharmacological therapies for managing cirrhosis and its complications", World Journal of Gastroenterology, 2019

Publication

<1 %

59

Stefania Paolillo, Simona Dell'Aversana, Immacolata Esposito, Alessandra Poccia, Pasquale Perrone Filardi. "The use of β -blockers in patients with heart failure and comorbidities: Doubts, certainties and unsolved issues", European Journal of Internal Medicine, 2021

Publication

<1 %

60

Yoshiaki Kubota, Wan Ting Tay, Tiew - Hwa Katherine Teng, Kuniya Asai et al. "Impact of

<1 %

beta - blocker use on the long - term outcomes of heart failure patients with chronic obstructive pulmonary disease", ESC Heart Failure, 2021

Publication

61	academic.oup.com Internet Source	<1 %
62	archive.org Internet Source	<1 %
63	dokumen.pub Internet Source	<1 %
64	e-journal.unair.ac.id Internet Source	<1 %
65	epdf.pub Internet Source	<1 %
66	heart.bmj.com Internet Source	<1 %
67	journals.physiology.org Internet Source	<1 %
68	orca.cf.ac.uk Internet Source	<1 %
69	res.mdpi.com Internet Source	<1 %
70	www.frontiersin.org Internet Source	<1 %

71

"Cardiac Considerations in Chronic Lung Disease", Springer Science and Business Media LLC, 2020

Publication

<1 %

72

"Diagnostic Methods for Cirrhosis and Portal Hypertension", Springer Science and Business Media LLC, 2018

Publication

<1 %

73

"Variceal Bleeding in Liver Cirrhosis", Springer Science and Business Media LLC, 2021

Publication

<1 %

74

Sergi Guixé-Muntet, Chang-Peng Zhu, Wei-Fen Xie, Jordi Gracia-Sancho. "Novel therapeutics for portal hypertension and fibrosis in chronic liver disease", Pharmacology & Therapeutics, 2020

Publication

<1 %

75

Sofia Simona Jakab, Guadalupe Garcia-Tsao. "Evaluation and Management of Esophageal and Gastric Varices in Patients with Cirrhosis", Clinics in Liver Disease, 2020

Publication

<1 %

76

Chel Hun Choi, Taejong Song, Tae Hyun Kim, Jun Kuk Choi et al. "Meta-analysis of the effects of beta blocker on survival time in cancer patients", Journal of Cancer Research and Clinical Oncology, 2014

Publication

<1 %

77 Chun Chen, Jing Zhang, Zemei Zhou. "Benefits of esmolol in sepsis and septic shock in adults: a meta-analysis of randomized controlled trials", Research Square, 2021 <1 %
Publication

78 Wolfgang Kreisel, Denise Schaffner, Adhara Lazaro, Jonel Trebicka, Irmgard Merfort, Annette Schmitt-Graeff, Peter Deibert. "Phosphodiesterases in the Liver as Potential Therapeutic Targets of Cirrhotic Portal Hypertension", International Journal of Molecular Sciences, 2020 <1 %
Publication

79 imed.pub <1 %
Internet Source

Exclude quotes On

Exclude matches Off

Exclude bibliography On

A Novel Pleiotropic Effect of Beta-Blockers Useful or Not?

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
