

# Benzalkonium chloride effectiveness as a disinfectant against hospital-associated methicillin-resistant Staphylococcus aureus (HA- MRSA)

*by* Koyuki Atifa Rahmi

---

**Submission date:** 28-Jul-2021 12:51PM (UTC+0800)

**Submission ID:** 1624945903

**File name:** ociated\_methicillin-resistant\_Staphylococcus\_aureus\_HA-MRSA.pdf (397.39K)

**Word count:** 4193

**Character count:** 22986



9  
**Benzalkonium chloride effectiveness as a disinfectant against hospital-associated methicillin-resistant *Staphylococcus aureus* (HA-MRSA)**

Koyuki Atifa Rahmi<sup>1</sup>, Priyo Budi Purwono<sup>2</sup> and Maftuchah Rochmanti<sup>3\*</sup>

<sup>1</sup>Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

<sup>2</sup>Department of Medical Microbiology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

<sup>3</sup>Department of Pharmacology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

Email: [maftuchah-r@fk.unair.ac.id](mailto:maftuchah-r@fk.unair.ac.id)

25  
Received 21 January 2018; Received in revised form 13 November 2018; Accepted 23 November 2018

**ABSTRACT**

**Aims:** Benzalkonium chloride is used to disinfect hospital instruments to prevent nosocomial infection caused by microorganisms, such as Methicillin Resistant *Staphylococcus aureus* (MRSA). There are strains of MRSA isolated from hospitals that were found to be resistant towards benzalkonium chloride. This research was aimed to compare the affectivity of different concentrations of benzalkonium chloride to inhibit the growth of Hospital-Associated MRSA (HA-MRSA) and determine the Minimum Inhibitory Concentration (MIC) of benzalkonium chloride against HA-MRSA.

**Methodology and results:** The samples were five HA-MRSA isolates obtained from Dr. Soetomo Hospital Surabaya. It was identified by amplification of *SCCmec* genes. The HA-MRSA with *SCCmec* type III was divided into six flasks based on the concentration of benzalkonium chloride in their inoculation media (0 µg/mL, 0.625 µg/mL, 1.25 µg/mL, 2.5 µg/mL, 5 µg/mL, and 10 µg/mL). The growth of HA-MRSA in media was determined by the number of colonies after treatment. The result showed that the MIC of benzalkonium chloride for HA-MRSA was 5 µg/mL, where no growth of bacterial was observed. There was significant difference in MRSA colony count between different groups of benzalkonium chloride concentrations ( $p = 0.001$ ), and there was negative correlation between benzalkonium chloride concentration and HA-MRSA growth ( $p = 0.0001$  and  $r = -0.880$ ).

**Conclusion, significance and impact of study:** The concentration of benzalkonium chloride influences the growth of HA-MRSA. The higher the concentration, the fewer HA-MRSA growth. Application of benzalkonium chloride according to MIC will prevent HA-MRSA resistance towards benzalkonium chloride.

**Keywords:** HA-MRSA, benzalkonium chloride, concentration, MIC

**INTRODUCTION**

Benzalkonium chloride is a disinfectant used to prevent the spread of nosocomial infections through hospital instruments (McDonnel and Russell, 1999; Fazlara and Ekhtelat, 2012). Benzalkonium chloride is a Quaternary Ammonium Compound (QAC) that works as a disinfectant by damaging the phospholipid bilayer of the bacterial cell, then entering into the cell and denaturing the essential proteins and inactivating the metabolic enzymes required by the cell. The destruction of cell and enzyme proteins then causes cell death (Rutala and Weber, 2008).

Hospital-Associated Methicillin Resistant *Staphylococcus aureus* (HA-MRSA) is one of the bacteria that cause nosocomial infections (Gordon and Lowy, 2008). The source of MRSA transmission can be from carrier individuals or contaminated instruments (Otto, 2012; Watkins *et al.*, 2012; Bellows *et al.*, 2013). MRSA is

43  
a *S. aureus* strain that is resistant to methicillin and other  $\beta$ -lactam class antibiotics (e.g. penicillin and cephalosporin) (Todar *et al.*, 2000). MRSA is resistant to methicillin because it has a *mecA* gene which is located in a large chromosome element called *SCCmec*. The mode of resistance of MRSA to antibiotics is a change in Penicillin Binding Protein (PBP) so that it is resistant to beta-lactam class antibiotics (Watkins *et al.*, 2012). The most common HA-MRSA causing nosocomial infection is Staphylococcal Cassette Chromosome *mecA* (*SCCmec*) type III, which composed 94.4% of all the HA-MRSA (Ashgar, 2014; Santosaningsih *et al.*, 2014), and 98% in Dr. Soetomo Hospital Surabaya (Lelitawati, 2016). Because of its high prevalence, the HA-MRSA type which is used in this research is HA-MRSA with *SCCmec* type III.

Benzalkonium chloride is generally used as a disinfectant for hospital instruments to rid them of

\*Corresponding author

microorganisms, such as MRSA (Al-Masaudi *et al.*, 1988). However, there have been reported MRSA strains resistant to benzalkonium chloride which were isolated from Kyushu University Hospital, Japan (Akimitsu *et al.*, 1999). Meanwhile, there is lack of study concerning the presence of MRSA strains, especially HA-MRSA (including type III), which are resistant to benzalkonium chloride in Indonesia. This study determined the effectiveness of benzalkonium chloride in various concentrations as disinfectant against HA-MRSA with SCC*mec* type III.

## MATERIALS AND METHODS

### Ethical clearance

Ethical clearance for this research was obtained from Ethical Committee of Faculty of Medicine Universitas Airlangga with the ethical clearance certificate number 183/EC/KEPK/FKUA/2016.

### Bacterial isolates

Five samples of MRSA bacterial isolates were obtained from the isolates taken from the patients who were admitted to Dr. Soetomo Hospital Surabaya with MRSA infection and then stored in the Laboratory of Clinical Microbiology (No. 183/EC/KEPK/FKUA/2016).

### Determination of SCC*mec* type

To identify HA-MRSA with SCC*mec* type III, a confirmatory test was performed using multiplex PCR method (Milheirico *et al.*, 2007). Five samples of MRSA with isolate number 448, 575, 646, 686 and 700 were cultured on slant agar medium, then sent to Microbiology Laboratory of Medical Faculty of Brawijaya University for DNA isolation and Polymerase Chain Reaction (PCR).

There were 3 primer pairs used in this experiment, (1) *mecA* primer pair for detecting *mecA* gene found in MRSA with 162 base pairs of amplicon, with *mecA* P4: TCCAGATTACAACCTTCACCAGG and *mecA* P7: CCACTTCATATCTTGTAACG; (2) *mecI* primer pair for detecting *mec* complex found in HA-MRSA SCC*mec* type II and III with 209 base pairs of amplicon, with *mecI* P2: ATCAAGACTTGCATTCAGGC and *mecI* P3: GCGGTTTCAATTCACCTTGTC; and (3) SCC*mec* III primer pair for detecting J1 region found in HA-MRSA SCC*mec* type III with 243 base pairs of amplicon, with SCC*mec* III J1 F: CATTGTGAAACACAGTACG and SCC*mec* III J1 R: GTTATTGAGACTCCTAAAGC.

The HA-MRSA SCC*mec* type III identification method above was based on methods used in Murakami *et al.* (1991), Zhang *et al.* (2005) and Milheirico *et al.* (2007). The reagent used for DNA isolation was obtained from iNtRON Biotechnology, Korea. PCR was performed using GoTaq Green Master Mix reagent (Promega, United States of America) and processed with a thermocycler from Bio-Rad, United States of America.

### Preparation of medium containing benzalkonium chloride

A 0.1 % solution of benzalkonium chloride was prepared by mixing 10  $\mu$ L benzalkonium chloride (80%) from Brataco, Indonesia and 7990  $\mu$ L aquadest into the test tube, then the solution was mixed using micropipette. The amount of 0.1% benzalkonium chloride used for each concentration is 0  $\mu$ L for 0  $\mu$ g/mL as a positive control (group 1), 12.5  $\mu$ L for 0.625  $\mu$ g/mL (group 2), 25  $\mu$ L for 1.25  $\mu$ g/mL (group 3), 50  $\mu$ L for two media, which are 2.5  $\mu$ g/mL (group 4) and as a negative control (group 7), 100  $\mu$ L for 5  $\mu$ g/mL (group 5), 200  $\mu$ L for 10  $\mu$ g/mL (group 6). Six groups were inoculated with MRSA (groups 1-6) and one group is not (group 7). All media were prepared by mixing 0.1% benzalkonium chloride according to the required amount with previously sterilized liquid nutrient agar in the Petri dish with the final volume of medium 20 mL. The Petri dish was then placed in room temperature (25 °C) for 15 min until the nutrient agar was solidified and then incubated in 37 °C for 24 h (Akimitsu *et al.*, 1999).

### Preparation of bacterial suspension

The colony of HA-MRSA isolate number 646 was taken from agar plate with a sterile loop needle and suspended into a nutrient broth medium in a test tube until it reached 0.5 McFarland turbidity, then the suspension was mixed using vortex. The homogeneous suspension was further diluted six times by 4 mL dilution in nutrient broth medium (final concentration of bacteria: Approx.  $10^6$ - $10^7$  CFU/mL) (Garcia, 2010).

### Inoculation of bacteria

Fifty microliter suspension of diluted bacteria was dripped using micropipette onto the agar medium that has been solidified in a Petri dish (except the medium for negative control). The bacterial suspension was then spread using a sterile spreader. The bacterial-inoculated medium was kept at room temperature until the bacterial suspension on the medium dried, then incubated at 37 °C for 24 h (Akimitsu *et al.*, 1999). Four replications were performed on each concentration of benzalkonium chloride that was tested.

### Counting colonies

Colony counting was done using BZG 40 colony counter from Hoskin Scientific, Canada. Due to the non-scaling calculations (in some of the Petri dish, the bacteria formed a fog-like growth, so the number of colonies could not be counted), the results of the colony count were converted to ordinal data, with 0 representing 0 colonies from 50  $\mu$ L of suspension (0 CFU/mL), +1 representing 10-100 colonies from 50  $\mu$ L of suspension (200-2,000 CFU/mL), +2 representing 100-1,000 colonies from 50  $\mu$ L of suspension (2,000-20,000 CFU/mL), and +3 representing more than 1,000 colonies from 50  $\mu$ L of

suspension (>20,000 CFU/mL) or fog-like growth (Brugger *et al.*, 2012).

#### MIC determination

The effectiveness of benzalkonium chloride as a disinfectant against HA-MRSA was calculated in the number of colonies grown on each agar nutrient medium. The Minimum Inhibitory Concentration (MIC) was determined by the result of the colony counting. MIC of benzalkonium chloride against MRSA was the concentration of benzalkonium chloride in the medium with zero colonies of MRSA growth (Akimitsu *et al.*, 1999).

#### Data analysis


Data which was obtained from seven groups was processed using IBM SPSS Statistics 23 with Kruskal-

Wallis comparative test to determine the difference of HA-MRSA growth on media with different benzalkonium chloride concentration. Spearman correlation test was used to find out the correlation between benzalkonium chloride concentration and HA-MRSA growth.

## RESULTS


### HA-MRSA identification

There were five isolates tested, isolates numbered 448, 575, 646, 686, and 700. Results of PCR of those 5 MRSA isolates are shown in Figure 1. *mecA* gene was positive in isolates 448, 575, 646, 686, and 700, confirming that all of those isolates are MRSA. *SCCmec* type II and III was positive in isolate number 646. PCR HA-MRSA with a target DNA fragment with a length of 243 base pairs (detection of *SCCmec* type III) was positive in isolate number 646.



**Figure 1:** Results of HA-MRSA *SCCmec* type III PCR visualization with gel electrophoresis. (a) PCR HA-MRSA with a target DNA fragment with a length of 162 base pairs (detection of *mecA* gene), positive in isolates 448, 575, 646, 686, and 700. (b) PCR HA-MRSA with target DNA fragment with length of 209 base pairs (detection of *SCCmec* type II and III), positive in isolate number 646. (c) PCR HA-MRSA with a target DNA fragment with a length of 243 base pairs (detection of *SCCmec* type III), positive in isolate number 646. (d) Resume of the PCR result from each electrophoresis shown in a, b, and c. The isolate that gave positive band in all PCR test, thus identified as HA-MRSA type III, was isolate 646.





d)	0 µg/mL or Control (+)	0.625 µg/mL	1.25 µg/mL	2.5 µg/mL	5 µg/mL	10 µg/mL	Control (-)
Replication 1	+3	+3	+2	+2	0	0	0
Replication 2	+3	+3	+2	+2	0	0	0
Replication 3	+3	+3	0	0	0	0	0
Replication 4	+3	+3	+2	0	0	0	0

**Figure 2:** Results of HA-MRSA growth on nutrient agar medium containing different concentrations of benzalkonium chloride. (a) Medium showing bacterial growth of +3 [observed at positive control (group 1) and 0.625 µg/mL (group 2) of benzalkonium chloride]. (b) Medium showing bacterial growth of +2 [observed at 1.25 µg/mL (group 3) and 2.5 µg/mL (group 4) of benzalkonium chloride]. (c) Medium showing no bacterial growth [observed at 5 µg/mL (group 5) and 10 µg/mL (group 6) of benzalkonium chloride, also negative control (group 7)]. (d) Resume of the bacterial colony growth after converted to ordinal data that shows the MIC of benzalkonium chloride against HA-MRSA, which is 5 µg/mL.

**Table 1:** Log of number of bacterial colonies which grew on media with different concentrations of benzalkonium chloride.

		Log of number of colonies				
Benzalkonium chloride concentration		1 <sup>st</sup> replication	2 <sup>nd</sup> replication	3 <sup>rd</sup> replication	4 <sup>th</sup> replication	Mean value
0 µg/mL (Control (+))	log(CFU/50 µL)	3.258397804	3.32056168	3.548757829	3.638788667	3.469674773
	log(CFU/mL)	4.5594278	4.621591676	4.849787824	4.939818663	4.770704768
0.625 µg/mL	log(CFU/50 µL)	3.384532615	Undetermined*	3.46834733	3.56937391	3.480725379
	log(CFU/mL)	4.681964459	Undetermined*	4.769377326	4.870403905	4.780749231
1.25 µg/mL	log(CFU/50 µL)	2.930949031	2.557507202	0	2.909556029	2.705007959
	log(CFU/mL)	4.231979027	3.858537198	0	4.210586025	4.005609445
2.5 µg/mL	log(CFU/50 µL)	2.195899652	2.471291711	0	0	2.053078443
	log(CFU/mL)	3.496929648	3.772321707	0	0	3.355068206
5 µg/mL	log(CFU/50 µL)	0	0	0	0	0
	log(CFU/mL)	0	0	0	0	0
10 µg/mL	log(CFU/50 µL)	0	0	0	0	0
	log(CFU/mL)	0	0	0	0	0
Control (-)	log(CFU/50 µL)	0	0	0	0	0
	log(CFU/mL)	0	0	0	0	0


### Benzalkonium chloride sensitivity test

Isolate number 646 was then tested for its sensitivity to benzalkonium chloride. The results of HA-MRSA growth on nutrient agar medium containing six different concentrations of benzalkonium chloride is shown in Figure 2.


The number of HA-MRSA colonies which grew on each nutrient agar medium is shown in Table 1. The

number of colonies was then converted to ordinal data and is shown in Figure 2. The graph showing the mean values of the number of HA-MRSA colonies which grew on each nutrient agar medium is shown in Figure 3.

Figure 2, Table 1, and Figure 3 show that the MIC of benzalkonium chloride for HA-MRSA with SCC<sub>mec</sub> type III was 5 µg/mL because benzalkonium chloride in that concentration was effective to kill all bacteria inoculated in the media.



**Figure 3:** Mean value of the log of number of bacterial colonies (in CFU/50  $\mu$ L) which grew on each medium containing different benzalkonium chloride concentration. Dot at 5  $\mu$ g/mL indicated the MIC of benzalkonium chloride against HA-MRSA which was tested.



**Figure 4:** Log of number of bacterial colonies (in CFU/50  $\mu$ L) which grew on each medium containing different benzalkonium chloride concentration. Linear trendline (black) shows negative correlation between concentration of benzalkonium chloride and the growth of HA-MRSA.

There is significant difference of the number of HA-MRSA growth on the media with different concentrations of benzalkonium chloride, evidenced by the significance value of 0.001 ( $p < 0.05$ ) by Kruskal-Wallis test.

There is a significant relationship between the concentration of benzalkonium chloride with the growth of HA-MRSA when analyzed by Spearman correlation test. It showed the value of significance at 0.0001 ( $p < 0.05$ ). The correlation coefficient obtained was -0.880, which means the negative correlation between the two variables was strong ( $> 0.75$  or  $< -0.75$ ). The higher concentration of benzalkonium chloride will decrease the growth of HA-MRSA. The negative correlation is shown in Figure 4.

## DISCUSSION

49

Minimum Inhibitory Concentration (MIC) of benzalkonium chloride against HA-MRSA isolate with SCCmec type III obtained from Dr. Soetomo Hospital Surabaya was 5  $\mu$ g/mL. This MIC was similar to the MIC of benzalkonium chloride against benzalkonium chloride-sensitive MRSA in a research conducted in Japan by Akimitsu in 1999. According to the research, HA-MRSA with SCCmec type III used in this study had no resistance to benzalkonium chloride but has definitely been resistant to antibiotic methicillin because it has *mecA* gene that encodes resistance to methicillin in all MRSAs.

Previous tests of HA-MRSA isolated from various hospitals in Rome, Italy, resulted in benzalkonium chloride having different MICs on the various HA-MRSA strains tested. The MICs were 2  $\mu$ g/mL, 4  $\mu$ g/mL, 8  $\mu$ g/mL, and 16  $\mu$ g/mL (Raggi and Filippini, 2013). These results suggested that different gene mutations between different HA-MRSA strains may also affect the resistance of HA-MRSA to benzalkonium chloride. This is the reason why this research is important to be conducted.

Research on the effectiveness of benzalkonium chloride as a disinfectant against MRSA has been done before. From these studies, it can be observed that MRSA resistance to benzalkonium chloride is associated with MRSA resistance to certain antibiotics (Fatholahzadeh *et al.*, 2008; Ahmad *et al.*, 2013). This happens because both have the same resistance mechanism, that is through the efflux pump encoded by the same gene (Ubarretxena-Belandia, 2003; Weinstein *et al.*, 2005; Quistgaard *et al.*, 2016). Mutation of genes that cause MRSA resistance to benzalkonium chloride is a mutation of the *femA* gene that also causes resistance to methicillin and oxacillin (Akimitsu *et al.*, 1999), as well as *qac* gene mutations that also lead to erythromycin, penicillin and ampicillin resistance (He *et al.*, 2014). From the studies it can be concluded that MRSA resistance mechanisms to various types of antibiotics and disinfectants, especially benzalkonium chloride, are not only caused by mutations of one particular gene, but there are several gene mutations that play a role (Reiter *et al.*, 2010). Gene mutations in MRSA that cause resistance towards antibiotics are influenced by environmental condition and frequent use of disinfectants (He *et al.*, 2014). The MIC of benzalkonium chloride against HA-MRSA might be different at places other than health care institutions, because the MIC is also influenced by previous benzalkonium chloride exposure towards the microorganism (He *et al.*, 2014). Thus, the MIC of benzalkonium chloride against MRSA at places other than health care institutions are probably different than those at health care institutions which use benzalkonium chloride frequently as disinfectant.

Different types of MRSA and MRSA growing in different environments are likely to have different gene mutations and patterns of resistance to different antibiotics and disinfectants. This is the reason why researches of effectivity of benzalkonium chloride against HA-MRSA has been done in various countries and

environments using many different types of MRSA, including in Indonesia using HA-MRSA with SCCmec type III as was done in this research.

Contamination of hospital equipment by HA-MRSA will cause nosocomial infection if those contaminated equipment are exposed to the patients. Patients with nosocomial infection caused by HA-MRSA will be difficult to treat using antibiotics, because HA-MRSA has been proven resistant towards numerous antibiotics, such as methicillin. Thus, the application of benzalkonium chloride as disinfectant is important to prevent the nosocomial infection caused by methicillin resistant bacteria (Mollema *et al.*, 2010; Otto, 2012).

There are some limitations in this study. First, this research only used one strain of MRSA, which is HA-MRSA with SCCmec type III, and second, this study was only conducted at Dr. Soetomo Hospital, Surabaya. However, the result of this research can be used as initial data for further studies concerning MRSA resistance towards benzalkonium chloride in Indonesia.

## CONCLUSION

In conclusion, HA-MRSA isolated from Dr. Soetomo Hospital Surabaya is still sensitive to benzalkonium chloride with concentration 5 µg/mL. This concentration can be implemented not only at Dr. Soetomo Hospital Surabaya, but also at another health care institutions in and around Surabaya.

## ACKNOWLEDGEMENTS

We thank Dr. Dewi Santosaningsih for assistance with multiplex PCR for confirmatory test to identify HA-MRSA with SCCmec type III.

## REFERENCES

- Ahmad, B., Urbas, F., Johar, J., Ahmed, J. and Bashir, S. (2013). Biocides susceptibility pattern and phenotypic detection of efflux pump in *Staphylococcus aureus* Isolates from two tertiary hospitals in Pakistan. *African Journal of Microbiology Research* 7(25), 3171-3178.
- Akimitsu, K., Akimitsu, N., Hamamoto, H., Inoue, R., Shoji, M., Akamine, A., Takemori, K. Hamasaki, N. and Sekimizu, K. (1999). Increase in resistance of methicillin-resistant *Staphylococcus aureus* to  $\beta$ -lactams caused by mutations conferring resistance to benzalkonium chloride, a disinfectant widely used in hospitals. *Antimicrobial Agents and Chemotherapy* 43(12), 3042-3043.
- Al-Masaudi, S. B., Day, M. J. and Russell, A. D. (1988). Sensitivity of methicillin-resistant *Staphylococcus aureus* strains to some antibiotics, antiseptics and disinfectants. *Journal of Applied Bacteriology* 65, 329-337.
- Ashgar, A. (2014). Molecular characterization of methicillin-resistant *Staphylococcus aureus* isolated from tertiary care hospitals. *Pakistan Journal of Medical Science* 30(4), 689-702.
- Bellows, C., Smith, A., Wheeler, J. and Morici, L. (2013). Nasal carriage of methicillin-resistant *Staphylococcus aureus* among students at a Louisiana Medical University. *The Brazilian Journal of Infectious Diseases* 17(1), 118-119.
- Brugger, S., Baumberger, C., Jost, M., Jenni, W., Brugger, U. and Mühlemann, K. (2012). Automated counting of bacterial colony forming units on agar plates. *PLoS ONE* 7(3), 33695.
- Fatholahzadeh, B., Emaneini, M., Gilbert, G., Udo, E., Aligholi, M., Modarressi, M., Nouri, K., Sedaghat, H. and Feizabadi, M. (2008). Staphylococcal Cassette Chromosomemec (SCCmec) analysis and antimicrobial susceptibility patterns of methicillin-resistant *Staphylococcus aureus* (MRSA) isolates in Tehran, Iran. *Microbial Drug Resistance* 14(3), 217-220.
- Fazlara, A. and Ekhtelat, M. (2012). The disinfectant effects of benzalkonium chloride on some important foodborne pathogens. *American-Eurasian Journal of Agricultural and Environmental Science* 12(1), 23-29.
- Garcia, L. (2010). Preparation of routine media and reagents used in antimicrobial susceptibility testing. In: *Clinical Microbiology Procedures Handbook*. 3rd Edn. ASM Press, Washington, DC. pp. 181-201.
- Gordon, R. and Lowy, F. (2008). Pathogenesis of methicillin-resistant *Staphylococcus aureus* infection. *Clinical Infectious Diseases* 46(S5), S350-S359.
- He, G., Chen, H., Thorpe, C., Walsh, D., Varela, M., Pan, H. and Landry, M. (2014). Detection of benzalkonium chloride resistance in community environmental isolates of staphylococci. *Journal of Medical Microbiology* 63(5), 735-741.
- Lelitawati, M. (2016). Distribusi gen SCCmec tipe III, IV, dan V pada isolat methicillin resistant *Staphylococcus aureus* (MRSA) pasien rawat inap bedah RSUD. Dr. Soetomo Surabaya. PhD Thesis. Universitas Airlangga, Indonesia.
- McDonnel, G. and Russell, A. (1999). Antiseptics and disinfectants: Activity, action, and resistance. *Clinical Microbiology Reviews* 12(1), 147-179.
- Milheirico, C., Oliveira, D. and de Lencastre, H. (2007). Update to the multiplex PCR strategy for assignment of *mec* element types in *Staphylococcus aureus*. *Antimicrobial Agents and Chemotherapy* 51(9), 3374-3377.
- Mollema, F., Severin, J., Nouwen, J., Ott, A., Verbrugh, H. and Vos, C. (2010). Successful treatment for carriage of methicillin resistant *Staphylococcus aureus* and importance to follow up. *Antimicrobial Agents and Chemotherapy* 54(9), 4020-4025.
- Murakami, K., Minamide, W., Wada, K., Nakamura, E., Teraoka, H. and Watanabe, S. (1991). Identification of methicillin-resistant strains of Staphylococci by polymerase chain reaction. *Journal of Clinical Microbiology* 29(10), 2240-2244.
- Otto, M. (2012). MRSA virulence and spread. *Cell Microbiol* 14(10), 1513-1521.

- Quistgaard, E., Löw, C., Guettou, F. and Nordlund, P. (2016).** Understanding transport by the major facilitator superfamily (MFS): Structures pave the way. *Nature Reviews Molecular Cell Biology* **17(2)**, 123-132.
- Raggi, C. and Filippini, P. (2013).** Methicillin resistance, biofilm formation and resistance to benzalkonium chloride in *Staphylococcus aureus* clinical isolates. *Clinical Microbiology* **2**, 121.
- Reiter, K., Machado, A., Freitas, A. and Barth, A. (2010).** High prevalence of methicillin-resistant *Staphylococcus aureus* with SCCmec type III in cystic fibrosis patients in southern, Brazil. *Revista da Sociedade Brasileira de Medicina Tropical* **43(4)**, 377-381.
- Rutala, W. and Weber, D. (2008).** Guideline for disinfection and sterilization in healthcare facilities. Central for Disease Control. North Carolina.
- Santosningsih, D., Santoso, S., Budayanti, N., Kuntaman, Lestari, E., Farida, H., Hapsari, R., Hadi, P., Winarto, Milheirico, C., Maquelin, K., Willemse-Erix, D., van Belkum, A., Severin, J. and Verbrugh, H. (2014).** Epidemiology of *Staphylococcus aureus* harboring the *mecA* or Panton-Valentine Leukocidin genes in Hospitals in Java and Bali, Indonesia. *American Journal of Tropical Medicine and Hygiene*. **90(4)**, 728-734.
- Todar, K. (2000).** *Todar's Online Textbook of Bacteriology*. Kenneth Todar. Wisconsin. pp. 1-3.
- Ubarretxena-Belandia, I. (2003).** Three-dimensional structure of the bacterial multidrug transporter EmrE shows it is an asymmetric homodimer. *The EMBO Journal* **22(23)**, 6175-6181.
- Watkins, R., David, M. and Salata, R. (2012).** Current concepts on the virulence mechanisms of methicillin-resistant *Staphylococcus aureus*. *Journal of Medical Microbiology* **61(Pt\_9)**, 1179-1193.
- Weinstein, Robert A. and Hooper, David C. (2005).** Efflux pumps and nosocomial antibiotic resistance: A primer for Hospital Epidemiologists. *Clinical Infectious Diseases* **40(12)**, 1811-1817.
- Zhang, K., McClure, J., Elsayed, S., Louie, T., and Conly, J. (2005).** Novel multiplex PCR assay for characterization and concomitant subtyping of Staphylococcal cassette chromosome *mec* Types I to V in methicillin-resistant *Staphylococcus aureus*. *Journal of Clinical Microbiology* **43(10)**, 5026-33.



# Benzalkonium chloride effectiveness as a disinfectant against hospital-associated methicillin-resistant *Staphylococcus aureus* (HA-MRSA)

---

## ORIGINALITY REPORT

---

**21** %  
SIMILARITY INDEX

**16** %  
INTERNET SOURCES

**18** %  
PUBLICATIONS

**0** %  
STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** Rachel MOYEN, Gabriel AHOMBO, Etienne NGUIMBI, Nina Esther ONTSIRA, Rock Fabien NIAMA, Gatsiel Claudette YALA, Delphin LOUEMBE. "Activity of beta-lactam antibiotics and production of beta-lactamases in bacteria isolated from wound infections in Brazzaville, Congo", *African Journal of Microbiology Research*, 2014  
Publication **1** %
- 2** P. H. McCay. "The effect of sub-inhibitory concentrations of benzalkonium chloride on the competitiveness of *Pseudomonas aeruginosa* grown in continuous culture", *Microbiology*, 10/08/2009  
Publication **1** %
- 3** Aurora Fernández-Astorga. "Disinfectant tolerance and antibiotic resistance in psychrotrophic Gram-negative bacteria isolated from vegetables", *Letters in Applied Microbiology*, 5/1995 **1** %

4

Paolo Andreini, Simone Bonechi, Monica Bianchini, Alessandro Mecocci, Vincenzo Di Massa. "Chapter 57 Automatic Image Analysis and Classification for Urinary Bacteria Infection Screening", Springer Science and Business Media LLC, 2015

Publication

---

1 %

5

Rathi, D-N., Persson, E. J., Maurer, F. H. J., Sudesh, K.. "Biosynthesis of P(3HB-co-3HV-co-3HHp) terpolymer by *Cupriavidus necator* PHB-4 transformant harboring the highly active PHA synthase gene of *Chromobacterium* sp. USM2", *Malaysian Journal of Microbiology*, 2013

Publication

---

1 %

6

[annalsmicrobiology.biomedcentral.com](http://annalsmicrobiology.biomedcentral.com)

Internet Source

---

1 %

7

[mountainscholar.org](http://mountainscholar.org)

Internet Source

---

1 %

8

Wenjing Geng, Yonghong Yang, Dejing Wu, Guoying Huang et al. " Molecular characteristics of community-acquired, methicillinresistant isolated from Chinese children ", *FEMS Immunology & Medical Microbiology*, 2010

Publication

---

1 %

9

[www.lib.unair.ac.id](http://www.lib.unair.ac.id)

Internet Source

1 %

10

Megan C. Jennings, Kevin P. C. Minbiole, William M. Wuest. "Quaternary Ammonium Compounds: An Antimicrobial Mainstay and Platform for Innovation to Address Bacterial Resistance", ACS Infectious Diseases, 2015

Publication

1 %

11

Zaini Mohd Zain, Muhammad Fikri Johari, Nurul Shahirah Mohd Husin, Nurul Syamimi Rozman et al. "Staphylococcus aureus Nasal Carriage in Medical Students of Universiti Teknologi MARA", Journal of Clinical and Health Sciences, 2017

Publication

1 %

12

Michael Jonatan, Ricardo Adrian Nugraha, Tan Nicko Octora, Firas Farisi Alkaff et al. "31 Influence of Night Shift Clinical Rotation Towards Diurnal Variation of Blood Pressure of The Clinical Clerks Compared with Preclinical Students", Journal of Hypertension, 2017

Publication

1 %

13

[cdn.intechopen.com](http://cdn.intechopen.com)

Internet Source

1 %

14

[scholarscompass.vcu.edu](http://scholarscompass.vcu.edu)

Internet Source

1 %

- |    |   |      |
|----|---|------|
| 15 | Lim, Eun Seob, Jang Eun Lee, Joo-Sung Kim, and Ok Kyung Koo. "Isolation of indigenous bacteria from a cafeteria kitchen and their biofilm formation and disinfectant susceptibility", LWT - Food Science and Technology, 2017.<br>Publication | 1 %  |
| 16 | Núria Rodriguez, Sònia Roig, Núria Olmedo, José G. Lorén, Núria Rius. "A new bioassay method for quantitative analysis of tetracyclines", Journal of Industrial Microbiology & Biotechnology, 2004<br>Publication                             | 1 %  |
| 17 | <a href="http://mafiadoc.com">mafiadoc.com</a><br>Internet Source   | 1 %  |
| 18 | <a href="http://ujcontent.uj.ac.za">ujcontent.uj.ac.za</a><br>Internet Source   | 1 %  |
| 19 | <a href="http://www.researchgate.net">www.researchgate.net</a><br>Internet Source   | <1 % |
| 20 | AG Bell, GW Coombs, B Cater, C Douglass. "First report of a -positive multidrug-resistant isolated from a dog in New Zealand ", New Zealand Veterinary Journal, 2016<br>Publication   | <1 % |
| 21 | Dhanesha Nanayakkara, Dimanthi Jayatilake, Suranga Kodithuwakku. "Development of a single nucleotide polymorphism-based DNA   | <1 % |



marker for fall armyworm (Lepidoptera: Noctuidae) biotyping: a case study from the fall armyworm outbreak in Sri Lanka", The Canadian Entomologist, 2020

Publication

22

[journals.plos.org](https://journals.plos.org)

Internet Source

<1 %

23

[www.dovepress.com](http://www.dovepress.com)

Internet Source

<1 %

24

Eva Medina, Dietmar Helmut Pieper. "Chapter 492 Tackling Threats and Future Problems of Multidrug-Resistant Bacteria", Springer Science and Business Media LLC, 2016

Publication

<1 %

25

[www.eoriwyoming.org](http://www.eoriwyoming.org)

Internet Source

<1 %

26

Shipra Bijpuria, Rakesh Sharma, Bhupesh Taneja. " Deletion of RsmE 16S rRNA methyltransferase leads to low level increase in aminoglycoside resistance in ", Cold Spring Harbor Laboratory, 2020

Publication

<1 %

27

[academic.oup.com](http://academic.oup.com)

Internet Source

<1 %

28

Muhammad Zafrullah Arifin, Yulius Hermanto, Agung Budi Sutiono. "Quality of life in patients with traumatic brain injury: A single tertiary

<1 %

center experience", Research Square Platform  
LLC, 2021

Publication

29

[aip.scitation.org](http://aip.scitation.org)

Internet Source

<1 %

30

[sites.kowsarpub.com](http://sites.kowsarpub.com)

Internet Source

<1 %

31

Fateh Rahimi, Leili Shokoohizadeh.  
"Characterization of methicillin resistant  
Staphylococcus aureus strains among  
inpatients and outpatients in a referral  
hospital in Tehran, Iran", Microbial  
Pathogenesis, 2016

Publication

<1 %

32

Hadi Sedigh Ebrahim-Saraie, Mohammad  
Motamedifar, Jamal Sarvari, Seyedeh Mahsan  
Hoseini Alfatemi. "Emergence of SCCmec  
Type I Obtained from Clinical Samples in  
Shiraz Teaching Hospitals, South-West of  
Iran", Jundishapur Journal of Microbiology,  
2015

Publication

<1 %

33

Mangamuri, U. K., Vijayalakshmi Muvva, V.,  
Poda, S., Kamma, S.. "Isolation, Identification  
and Molecular Characterization of Rare  
Actinomycetes from Mangrove Ecosystem of  
Nizampatnam", Malaysian Journal of  
Microbiology, 2012

<1 %

34

Murilo Batista Abud, Ricardo Noguera Louzada, David Leonardo Cruvinel Isaac, Leonardo Gomes Souza et al. "In vivo and in vitro toxicity evaluation of liposome-encapsulated sirolimus", International Journal of Retina and Vitreous, 2019

Publication

---

35

[etd.auburn.edu](http://etd.auburn.edu)

Internet Source

---

<1 %

36

[www.pnas.org](http://www.pnas.org)

Internet Source

---

<1 %

37

"OS18: Interstitial Lung Disease 1",  
Respirology, 2013

Publication

---

<1 %

38

Geoffrey Omuse, Kristien Nel Van Zyl, Kim Hoek, Shima Abdulgader, Samuel Kariuki, Andrew Whitelaw, Gunturu Revathi.  
"Molecular characterization of Staphylococcus aureus isolates from various healthcare institutions in Nairobi, Kenya: a cross sectional study", Annals of Clinical Microbiology and Antimicrobials, 2016

Publication

---

<1 %

39

Ian F. Parney. "Neuroradiographic changes following convection-enhanced delivery of the recombinant cytotoxin interleukin 13—

<1 %

PE38QQR for recurrent malignant glioma",  
Journal of Neurosurgery, 02/2005

Publication

---

40

Ihsan Edan Abdulkareem AlSaimary.  
"Prevalence of  $\beta$ -lactamase producing and non-producing Staphylococcus aureus associated with patients in intensive care unit", International Journal of Medicine and Medical Sciences, 2012

Publication

---

<1 %

41

Ji-Hyeon Kim, Junfei Ma, Sungjin Jo, Seunghun Lee, Chang Su Kim. "Enhancement of Antibacterial Properties of a Silver Nanowire Film via Electron Beam Irradiation", ACS Applied Bio Materials, 2020

Publication

---

<1 %

42

Parhizgari, Najmeh, Seyed Sajjad Khoramrooz, Seyed Ali Asghar Malek Hosseini, Masoud Marashifard, Mahboobeh Yazdanpanah, Mohammad Emaneini, Farzaneh Gharibpour, Mehdi Mirzaii, Davood Darban-Sarokhalil, Masoud Moein, and Mahmood Naraki. "High frequency of multidrug-resistant Staphylococcus aureus with SCCmec type III and Spa types t037 and t631 isolated from burn patients in southwest of Iran", Apmis, 2015.

Publication

---

<1 %



43 Zhang, Jiuyang, Yung Pin Chen, Kristen P. Miller et al. "Antimicrobial Metallopolymers and Their Bioconjugates with Conventional Antibiotics against Multidrug-Resistant Bacteria", Journal of the American Chemical Society  
Publication <1 %

---

44 [cds.ismrm.org](https://cds.ismrm.org)  
Internet Source <1 %

---

45 [link.springer.com](https://link.springer.com)  
Internet Source <1 %

---

46 [new.seceij.net](https://new.seceij.net)  
Internet Source <1 %

---

47 [www.aimst.edu.my](http://www.aimst.edu.my)  
Internet Source <1 %

---

48 [www.frontiersin.org](http://www.frontiersin.org)  
Internet Source <1 %

---

49 Azizollah Ebrahimi, Zohreh Arvaneh, Mohamadreza Mahzounieh, Sharareh Lotfalian. "Antibiotic Resistance Induction by Benzalkonium Chloride Exposure in Nosocomial Pathogens", International Journal of Infection, 2016  
Publication <1 %

---

50 Marc G. Ghany, Doris B. Strader, David L. Thomas, Leonard B. Seeff. "Diagnosis, <1 %

# management, and treatment of hepatitis C: An update", Hepatology, 2009

Publication

51

[www.omicsonline.org](http://www.omicsonline.org)

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

# Benzalkonium chloride effectiveness as a disinfectant against hospital-associated methicillin-resistant Staphylococcus aureus (HA-MRSA)

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---