

# Cefotaxime vs ceftriaxone for the prolongation of latency period in preterm premature rupture of membranes

*by* Shinta Dewi Rasti

---

**Submission date:** 28-Jul-2021 01:01PM (UTC+0800)

**Submission ID:** 1624949088

**File name:** of\_latency\_period\_in\_preterm\_premature\_rupture\_of\_membranes.pdf (316.08K)

**Word count:** 5208

**Character count:** 27214

# Cefotaxime vs ceftriaxone for the prolongation of latency period in preterm premature rupture of membranes

Shinta Dewi Rasti<sup>1</sup>,  
Maftuchah Rochmanti<sup>2</sup>,  
Relly Yanuari  
Primariawan<sup>3</sup>

## Abstract

**Introduction:** Antibiotics are recommended as the main therapy for preterm premature rupture of membranes (PPROM). But the research on antibiotics other than the recommended macrolides regimens is still lacking. This research aims to evaluate whether there are effects differences of cefotaxime and ceftriaxone given on pregnancy with PPRM by comparing the duration of the latency period and the infants outcomes.

**Material and Methods:** Data was taken through medical records retrospectively at Dr. Soetomo Surabaya General Hospital, Indonesia, during the period of January-December 2017. The inclusion criteria were a history of PPRM in pregnancy <37 weeks, given cefotaxime or ceftriaxone therapy, and have labor data. The analysis was performed by the Mann-Whitney comparison test for the latency period and Fisher's exact test for infant outcomes.

**Results:** There were 52 samples obtained. The antibiotics used were cefotaxime 3 grams once and ceftriaxone 2 grams once. The results of the analysis showed that there were no significant differences between the types of antibiotics with the length of the latency period, with a value of  $p = 0.601$  ( $p > 0.05$ ), where cefotaxime had a median of 52,67 hours and ceftriaxone was 34,17 hours. Cefotaxime was found to be more able to extend the latency period for >48 hours with a percentage of 57.8%, whereas in ceftriaxone only 42.9%. There are no significant differences in infant outcomes; infant birth weight and Apgar score among the two therapies used.

**Conclusion:** Cefotaxime was more preferably to be used in the Dr. Soetomo Surabaya General Hospital. Although there are no diffe-

13

**1** Faculty of Medicine, Universitas Airlangga.

**2** Department of Pharmacology, Faculty of Medicine, Universitas Airlangga.

**3** Department of Obstetric and Gynecology, Faculty of Medicine, Universitas Airlangga.

## Contact information:

**Dr. Maftuchah Rochmanti.**

**Tel:** +62 818315479.

✉ maftuchah-r@fk.unair.ac.id

Received 29-11-2019; Accepted 14-01-2020

rences in infant outcomes between the two antibiotics, cefotaxime appears to be more able to extend the latency period for more than 48 hours which gives better prospects for fetal lung maturation. Both cefotaxime and ceftriaxone have succeeded in preventing infections in women with PPROM.

#### Keywords

PPROM; Antibiotic; Cefotaxime; Ceftriaxone; Cephalosporin; Latency Period.

## 1 Introduction

Premature rupture of membranes (PROM) is the rupture of the membranes before the onset of labor. In many cases, this happens in term pregnancy, but when the rupture of membranes occurs before 37 weeks gestation, this is known as a preterm PROM (PPROM) [1, 2]. PPROM complicates about 3 percent of pregnancies and leads to one-third of preterm births. This increases the risk of prematurity and leads to some other perinatal and neonatal complications, including 1 to 2 percent risk of fetal death [1].

The incidence of PPROM is not only a problem when it causes preterm labor. Women with prolonged PPROM are at higher risk for developing chorioamnionitis, which can be obtained from an increase in bacterial colonization before membrane rupture (causing PPROM) or after membrane rupture (PPROM complications) [3, 4]. The risk of infection increases with the decreasing gestational age when membrane rupture and the increasing duration of the ruptured membrane [3].

The use of antibiotics in women with PPROM has been shown to increase the latency period, which is defined as the period between ruptured membranes and onset of labor [5-8]. There is high-quality evidence that antibiotics for PPROM can reduce the risk of complications due to prematurity and the risk of maternal and neonatal infection [7,

9]. This means that antibiotics can be used at the same time to prevent the occurrence of infections and help maintain pregnancy in the PPROM to the maximum possible time or prevent preterm labor. Erythromycin is the first line choice antibiotic used for PPROM worldwide, but the recommendation was made conditional and there is no consensus regarding the best choice to this day. Cephalosporin group is another non-famous therapeutic option. Some study has already suggested that amoxicillin, 3<sup>rd</sup> generation cephalosporins, and erythromycin in the association erythromycin-amoxicillin can be used and have been shown to improve neonatal outcome [10, 11]. But the research on cephalosporin and its regimen as a PPROM therapy is still lacking.

Therefore, this study aims to evaluate whether there are effects differences between the new alternative cephalosporin antibiotics used in Dr. Soetomo Surabaya General Hospital for PPROM; cefotaxime and ceftriaxone by comparing the duration of the latency period and the infant outcomes.

## 21 Methods

### Study Design and Ethics

This analytic study with a cross-sectional design took place from September 2018 - June 2019. This research was reviewed by the Health Research Ethics Committee of Dr. Soetomo Surabaya General

Hospital and has been declared "Eligible of Ethics" with certificate number 0648/KEPK/lx/2018 on September 21<sup>st</sup>, 2018. Maternal informed consent was obtained through the help of ethics committee. We used secondary data in the form of medical records taken from the Central Medical Record and VK IRD in Dr. Soetomo Surabaya General Hospital. The population of this study was all pregnant women diagnosed with preterm premature rupture of membrane (PPROM) in Dr. Soetomo Surabaya General Hospital for January-December 2017 period. The diagnosis criteria for PPRM is the rupture of membranes without the begin of initial labor signs after 1 hour later. We used the 2017 data because that is the latest medical records which are complete for a year.

### Statistical Analysis

Samples were obtained using a total sampling technique so that all parts of the population that met the inclusion criteria were taken. The inclusion criteria for this study sample are history of PPRM in pregnancies <37 weeks, given cefotaxime or ceftriaxone therapy, having delivery data, and medical records in good and complete condition. Whereas the exclusion criteria are patients who did not deliver at the Dr. Soetomo Surabaya General Hospital and medical records that are damaged and difficult to read. The independent variable in this study is the antibiotic used in pregnant women with PPRM in Dr. Soetomo Surabaya General Hospital. While the dependent variable is the length of the latent period and the outcome in infants. The time of antibiotic administration cannot be a control variable because antibiotics were given in different time frames after the rupture of membranes. These differences occur because Dr. Soetomo is a tertiary hospital so some patients came after being referred. It also depends on the time when patients decide to go to the hospital after the rupture of membranes.

The collected data was analyzed by computerization with the Statistical Package for Social Sci-

ces (SPSS) software. Data is presented in the form of frequency distribution tables and described in narrative form. Analysis of the two variables that are thought to have a relationship is done through the Mann-Whitney and Fisher's exact statistical tests that illustrate the comparison of influences between variables.

The Mann-Whitney test was conducted to determine whether there were differences in the effect of antibiotic therapy between groups used on the latency period. At first, a chi-square test was performed to determine whether there are differences in infant outcomes between the types of antibiotics used. After the test is done, it was found that there were > 20% of tables that have an expected value of less than 5. Therefore, the p-value seen is the p-value of Fisher's exact test.

## Results

### Demographic Data

A total of 242 patients with history of PPRM were found in the medical records of Dr. Soetomo Surabaya General Hospital during the period of January – December 2017. Out of the total, there were only 123 patients who gave birth in Dr. Soetomo Surabaya General Hospital. Among them, there were only 52 patients who met the inclusion criteria.

In general, pregnant women with a history of PPRM are mostly in the age group of 20-35 years, which is as much as 61.5%. It is more common in multigravida women, with a percentage of 48.3%. Most PPRM occurred in the 24-31 week gestational age group, which was 40.4% and the least frequency at gestational age <24 weeks as much as 3.8%. There were 54% of samples with an estimated birth weight (EBW) ≥ 1500 grams when PPRM occurs and 46% with EBW <1500 grams.

There were 76.9% of samples that gave birth through vaginal delivery and 23.1% through cesa-

rean section. About 51.9% of the labor happens after being induced or terminated and 48.1% occur spontaneously. There were 36.7% of babies born with low birth weight (LBW), 28.6% with very low birth weight (VLBW), 22.4% with normal weight, and 12.2% with extremely low birth weight (ELBW). The majority of the 1<sup>st</sup> minute Apgar score was low (<7) with the percentage of 55.3%, but 59.6% of the 5<sup>th</sup> minute Apgar score was more than 7. Approximately 5.8% baby died within the first 24 hours after labor. Also, we found that 100% of women with PPROM didn't suffer any infection after the rupture of membranes.

**Table 1** shows the demographic data based on each antibiotic therapy used. Some of the data like the gestational age might be a moderator or intervening variable, which is a variable that can affect the dependent variable in this study or the latency period.

### Antibiotic use


It was found that there were no other antibiotics used for PPROM patients at Dr. Soetomo Surabaya General Hospital apart from cefotaxime or ceftriaxone. There were only two regimens found to be used, which are cefotaxime given at a dose of 3 grams once and ceftriaxone given at a dose of 2 grams once dissolved in a 100cc intravenous NaCl 0,9% (normal saline/PZ) infusion.<sup>20</sup>

The latency period consists of the time between the rupture of membranes (ROM) and antibiotic administration and the time from antibiotic administration until labor. Instead of using mean, the median was used to represent data broadly in this study because of the non-normal distribution. **Figure 1** shows that the length between the ROM and antibiotic administration was shorter and the time until labor was longer in the cefotaxime group. It can be interpreted that antibiotic was administrated sooner in the cefotaxime group and the latency period in the cefotaxime group involved more antibiotics than in the ceftriaxone group.


**Table 1.** Demographic data.

Demographic Data	Cefotaxime	Ceftriaxone
	n = 45 (%)	n = 7 (%)
Age		
<20 years	15.6	0
20-35 years	57.8	85.7
>35 years	26.7	14.3
Primigravida	46.3	14.3
Multigravida	53.7	85.7
Gestational Age		
<24 weeks	4.4	0
24-31 weeks	37.8	57.1
32-33 weeks	20	0
34-36 weeks	37.8	42.9
Estimated Birth Weight (EBW)		
<1500 grams	41.9	71.4
≥1500 grams	58.1	28.6
Mode of Delivery		
Pervaginam	73.3	100
Sectio caesarea	26.7	0
Terminated	53.3	42.9
Spontaneous	46.7	57.1
Infant Birth Weight		
≥2500 grams (Normal)	21.4	28.6
1500-2499 grams (LBW)	40.5	14.3
1000-1499 grams (VLBW)	28.6	28.6
<1000 grams (ELBW)	9.5	28.6
1 <sup>st</sup> minute Apgar Score		
7-10	45	42.9
4-6	17.5	28.6
0-3	37.5	28.6
5 <sup>th</sup> minutes Apgar Score		
7-10	62.5	42.9
4-6	12.5	28.6
0-3	25	28.6
Infant Condition		
Survive	93.3	100
Death within 24 hours	6.7	0

**Figure 1:** Time Mapping between The Ruptures of Membranes (ROM), Antibiotic Administration, and Labor



**Figure 2:** Percentage of Latency Period Length



**Table 2.** Latency period based on gestational age and Mann-Whitney test.

Gestational Age	Median of Latency Period (hours)						P value
	Cefotaxime			Ceftriaxone			
Weeks	Median	Min	Max	Med	Min	Max	
<24	35,92	17,83	54	there are no samples in			0.601
24-31	74,17	7,3	200,5	127,48	34,17	193,18	
32-33	64,08	19,75	320,5	there are no samples in			
34-36	33,67	8,28	85,58	26,45	24,78	29,37	
Total	52,67	7,3	320,5	34,17	24,78	193,18	

**Table 3.** Infant outcomes.

Infant Outcomes	Cefotaxime		Ceftriaxone		P value
	N	%	N	%	
	Mean	Notes	Mean	Notes	
Birth Weight (grams)	1821,43	LBW	1471,43	VLBW	0.339
APGAR Score 1 <sup>st</sup> minutes	5	Moderately depressed	5	Moderately depressed	0.711
APGAR Score 5 <sup>th</sup> minutes	6	Moderately depressed	6	Moderately depressed	0.131

It was found that there were 57,8% of cefotaxime group samples which have latency period of >48 hours, whereas in ceftriaxone there were only 42,9% (Figure 2)

### Analysis Results

The Mann-Whitney test shown a P-value of 0.601 (p >0.05), which means that there were no statistical differences. The duration of the latency period was illustrated through the median because of the non-normal distribution. If seen as a whole regard-

less of a certain gestational age group, the median latency period of cefotaxime group is 52 hours and 40 minutes and ceftriaxone group is 34 hours and 10 minutes. In Table 2 & 3 there are no differences in all aspect of infant outcomes between the two antibiotics.

### Discussion

The most frequently used antibiotic is cefotaxime 3 grams once given until labor. This therapeutic choi-

ce does not match the recommendations of Indonesian Association for Obstetrics and Gynaecology (POGI) (2016) and WHO (2015), which states that erythromycin is the first choice for prophylaxis after PPROM. The difference in this use of antibiotics can be due to the administration of antibiotics adjusted to their susceptibility to the bacteria that cause infection. In the previous research, cephalosporin as first-line therapy for the care of mothers and newborns in PPROM is indicated because of the high susceptibility of *Escherichia coli* to antibiotics [12]. *Escherichia coli* was one of the most common microorganism isolated from women with PPROM along with *Staphylococcus* or *Streptococcus* species [14-16]. In addition, the most common bacteria which cause urinary tract infection (UTI) is *Escherichia coli*, where UTI is one of the causes of preterm labor in PPROM [17, 18].

The cephalosporins are divided into 3 generations according to their spectrum of activity, where activity against gram-negative bacilli increases from first- to third-generation drugs [19, 20]. Third-generation cephalosporins are preferred in many clinical situations because of their proven record of clinical efficacy, favorable pharmacokinetics and low frequency of adverse effect [21]. Cefotaxime, ceftriaxone and other cephalosporins are safe to be used during pregnancy and considered as category B by FDA [22]. Cefotaxime was found to have the best gram-positive coverage of the 3<sup>rd</sup> generation agents, but ceftriaxone was as safe and as effective as cefotaxime in treating serious bacterial infections [21, 23-25]. A study conducted in Malang, East Java, Indonesia showed that cefotaxime generally was more cost-effective than ceftriaxone which made cefotaxime more preferable [26].

It was found that cefotaxime group was able to extend the latent period longer, with a median of approximately 2 days and maximum 13 days. As seen in **Figure 1**, antibiotic in the cefotaxime group was used more effectively since the sooner administration of antibiotic after ROM occurred was as-

sociated with a better outcome [7]. Also, as seen in **Figure 2**, the median and the majority of latency period in the cefotaxime group was more than 48 hours, where 48 is the time required until the corticosteroid given has worked to stimulate pulmonary maturation [27]. This result can be caused by the majority of pregnant women treated with cefotaxime being in the 24-31 week gestational age group whereas those treated with ceftriaxone didn't have any sample in that age group as shown in Table 1 and 2. According to Melamed *et al.* [8], the duration of the latency period ranges between 0 and 59 days and is inversely proportional to the gestational age at admission. Women with a short latent period (<48 hours) are characterized by a higher rate of cervical dilatation and a higher gestational age at entry and are more likely to be nulliparous [8]. However, no other studies are comparing these 2 antibiotic regimens.

A study that also compared the latency period between antibiotics from the cephalosporin group by Fortunato *et al.* [28], stated that there was no difference in the latency period between patients treated with ceftizoxime and those given other antibiotic therapy (cefoxitin, cefazolin, ampicillin). Other studies by Lee *et al.* [29], stated that a new combination of antibiotics consisting of ceftriaxone, clarithromycin, and metronidazole was able to extend the latency period longer with a median value of 23 days, compared to the therapeutic regimen of ampicillin and/or cephalosporin alone which had a median value of 12 days.

Another comparison to be considered is the worldwide recommended erythromycin which is still recommended because there have been many studies on its effectiveness for PPROM [30]. According to WHO (2015), the choice of erythromycin was based on the findings of ORACLE I study with over 2000 women, which showed that erythromycin decreases the risk of necrotizing enterocolitis (NEC) in the newborn compared to co-amoxiclav [7]. Another study that supports the use of erythromycin sta-

ted that erythromycin was found to be associated with a decrease in the primary composite outcome (neonatal death, chronic lung disease or major cerebral abnormality on ultrasound;  $p = 0.08$ ) and in single adverse neonatal outcomes ( $p = 0.02$ ) when compared to placebo [31]. However, a recent study mentioned that erythromycin alone is insufficient to control the growth of Gram-positive and Gram-negative bacteria in patients with PPROM [15]. In particular, *Escherichia coli* and group B *Streptococcus* isolates showed high rates of resistance to erythromycin [15, 32].

Seelbach-Goebel [33], summarized the results of the ORACLE I trials which state that 60.9% of pregnant women given erythromycin 4x/day orally give birth within 7 days after the rupture of membranes. There were no differences in latency to delivery, incidence of chorioamnionitis, or neonatal outcomes when erythromycin was compared with other drugs from one class of macrolides, azithromycin [34, 35]. The latency period after erythromycin administration was 65,6 hours and azithromycin 61,7 hours [36].

Other regimens, amoxicillin/clavulanic acid 325 mg 4x/day orally or erythromycin 250 mg + amoxicillin/clavulanic acid 325 mg, which all mentioned in ORACLE I trial were also associated with a significantly lower rate of deliveries within 48 hours compared to placebo [33]. But a fourfold higher rate of NEC was found in both groups treated with amoxicillin/clavulanic acid [33].

Several protocols recommend other different choices of antibiotic. Some Latin American countries recommend the use of ampicillin or amoxicillin [12]. Those antibiotics also showed a significant increase in the latency period and a lower incidence of clinical amnionitis [12, 37]. But the use of ampicillin and amoxicillin was related to adverse symptoms, such as nausea, vomiting and abdominal pain [37, 38]. Thus, amoxicillin and ampicillin are often replaced by a cephalosporin in clinical practice [38].

Based on the infant outcomes, both antibiotic groups have the same low birth weight output, even though cefotaxime group has a slightly higher value. In the Fisher's exact test, p-value in lower case = 0.339 was obtained, which means there was no difference in birth weight of babies between antibiotic therapy A and B. In accordance with the ORACLE Trial I study conducted between 1994 and 2000, the use of antibiotics was not directly related to birth weight [30, 33]. But a randomized controlled trial using oral ampicillin 1 g ( $n = 59$ ) vs placebo ( $n = 51$ ), showed that the mean birth weight was significantly higher in those received ampicillin (2885 vs. 2336 g;  $P < 0.05$ ) [39].

The Apgar score provides an accepted and convenient method for reporting the status of the newborn infant immediately after birth and the response to resuscitation if needed [40]. The average Apgar score both in the first minute and the fifth minute were identical between the two groups. In line with the Fisher's exact test which showed that there were no differences in the Apgar score between the two types of antibiotics used. The Apgar score is influenced by the length of the latency period. Salan [41], found that there was a correlation between the length of the latency period and the infant Apgar results with  $P < 0.001$  and the Odds ratio was 8.5, which means that the length of the latent period affected the infant Apgar score with an 8.5-fold increased risk.

## Conclusion

Based on this study, the most commonly used therapeutic regimen is cefotaxime 3 grams once given until labor. No differences were found in latency period and infant outcomes in terms of birth weight and Apgar scores between the use of cefotaxime and ceftriaxone. But cefotaxime appears to be more able to extend the latency period for more than 48 hours which gives better prospects for fetal lung maturation and more cost-effective than ceftriaxo-



ne. There was no incidence of infection after the rupture of membranes with the use of these two antibiotics, which mean that cephalosporin has succeeded in preventing infections in women with PPROM. This research, however, is subject to several limitations since the sample of the ceftriaxone group was small.

## Acknowledgment

The author would like to thank Atika, S.Sc., M.Kes, who provided guidance and assistance during this research, assisting data processing, methodology and statistics. And the authors would like to give appreciation to all colleagues in the obstetrics and gynecology department who have provided support and inspiration during the work.

## References

1. Medina TM, Hill DA. Preterm premature rupture of membranes: diagnosis and management. *Am Fam Physician* 2006; 73(4):659-664.
2. Caughey AB, Robinson JN, Norwitz ER. Contemporary diagnosis and management of preterm premature rupture of membranes. *Rev Obstet Gynecol* 2008; 1(1):11-22.
3. Mercer BM. Preterm premature rupture of the membranes: current approaches to evaluation and management. *Obstet Gynecol Clin North Am* 2005; 32(3):411-28.
4. Stock O, Gordon L, Kapoor J, et al. Chorioamnionitis Occurring in Women With Preterm Rupture of the Fetal Membranes Is Associated With a Dynamic Increase in mRNAs Coding Cytokines in the Maternal Circulation. *Reprod Sci* 2015; 22(7):852-859.
5. Bowes WA Jr. The role of antibiotics in the prevention of preterm birth. *F1000 Med Rep* 2009; 1. pii: 22. doi: 10.3410/M1-22
6. Johnston MM, Sanchez-Ramos L, Vaughn AJ, Todd MW, Benrubi GI. Antibiotic therapy in preterm premature rupture of membranes: a randomized, prospective, double-blind trial. *Am J Obstet Gynecol* 1990; 163(3):743-747.
7. Kenyon S, Boulvain M, Neilson J. Antibiotics for preterm premature rupture of membranes. *Cochrane Database Syst Rev* 2001; (4):CD001058.
8. Melamed N, Hadar E, Ben-Haroush A, Kaplan B, Yogev Y. Factors affecting the duration of the latency period in preterm premature rupture of membranes. *J Matern Fetal Neonatal Med* 2009; 22(11):1051-1056.
9. Cousens S, Blencowe H, Gravett M, Lawn JE. Antibiotics for pre-term pre-labour rupture of membranes: prevention of neonatal deaths due to complications of pre-term birth and infection. *Int J Epidemiol* 2010; 39 Suppl 1(Suppl 1):134-143.
10. Schmitz T, Sentilhes L, Lorthe E, Gallot D, Madar H, Doret-Dion M, Beucher G, Charlier C, Cazanave C, Delorme P, Garabedian C, Azria É, Tessier V, Senat MV, Kayem G. Preterm premature rupture of membranes: CNGOF Guidelines for clinical practice - Short version. *Gynecol Obstet Fertil Senol* 2018; 46(12):998-1003.
11. Tchirikov M, Schlalbritz-Loutsevitch N, Maher J, Buchmann J, Naberezhnev Y, Winarno AS, Seliger G. Mid-trimester preterm premature rupture of membranes (PPROM): etiology, diagnosis, classification, international recommendations of treatment options and outcome. *J Perinat Med* 2018; 46(5):465-488.
12. Machado DCS, Dias JdM, Rosado LEP. Antibiotic prophylaxis in preterm rupture of membranes. *Hum Reprod Arch* 2018; 32(3):e000217.
13. Saghafi N, Pourali L, Ghazvini K, Maleki A, Ghavidel M, Karbalaieizadeh Babaki M. Cervical bacterial colonization in women with preterm premature rupture of membrane and pregnancy outcomes: A cohort study. *Int J Reprod Biomed (Yazd)* 2018; 16(5):341-348.
14. Zeng L, Zhang L, Shi J, Gu L, Grogan W, Gargano MM, Chen C. The primary microbial pathogens associated with premature rupture of the membranes in China: A systematic review. *Taiwanese Journal of Obstetric and Gynecology* 2014; 53(4):443-451.
15. Li YY, Kong CW, To WWK. Pathogens in preterm prelabour rupture of membranes and erythromycin for antibiotic prophylaxis: a retrospective analysis. *Hong Kong Med J* 2019; 25(4):287-94.
16. Adewumi OA, Olofinbiyi BA, Oyekale OT, Loto OM, Abu SH, Sotunsa JO. Microbiological Pattern in Preterm Prelabour Rupture of the Fetal Membranes in South-Western Nigeria. *Obstet Gynecol Int J* 2017; 6(4):00215
17. Majdawati A. Pola Kepekaan Kuman Penyebab Infeksi Saluran Kemih dan Resistensinya. *Mutiara Medika* 2001; 1:55-63.
18. Behzadi P, Behzadi E, Yazdanbod H, Aghapour R, Akbari Cheshmeh M, Salehian Omran D. A survey on urinary tract infections associated with the three most common uropathogenic bacteria. *Maedica (Buchar)* 2010; 5(2):111-115.
19. Thompson RL, Wright AJ. Cephalosporin antibiotics. *Mayo Clinic Proceedings* 1983; 58(2):79-87.
20. Kalman D, Barriere SL. Review of the pharmacology, pharmacokinetics, and clinical use of cephalosporins. *Tex Heart Inst J* 1990; 17(3):203-215.
21. Klein NC, Cunha BA. Third-generation cephalosporins. *Med Clin North Am* 1995; 79(4):705-719.
22. Sá Del Fiol F, Gerenutti M, Groppo FC. Antibiotics and pregnancy. *Pharmazie* 2005; 60:483-493.

23. Mandell LA, Bergeron MG, Ronald AR, Vega C, Harding G, Saginur R, Feld R, Duperval R, Landis SJ, Miedzinski LJ, et al. Once-daily therapy with ceftriaxone compared with daily multiple-dose therapy with cefotaxime for serious bacterial infections: a randomized, double-blind study. *J Infect Dis* 1989; 160(3):433-441.
24. Smith CR, Petty BG, Hendrix CW, Kernan WN, Garver PL, Fox K, Beamer A, Carbone K, Threlkeld M, Lietman PS. Ceftriaxone compared with cefotaxime for serious bacterial infections. *J Infect Dis* 1989; 160(3):442-447.
25. Simmons BP, Gelfand MS, Grogan J, Craft B. Cefotaxime twice daily versus ceftriaxone once daily. A randomized controlled study in patients with serious infections. *Diagn Microbiol Infect Dis* 1995; 22(1-2):155-157.
26. Illahi RK, Pramestutie HR, Susilaningtyas WS. Cost-Effectiveness Analysis of Cefotaxime, Ceftriaxone, and Levofloxacin in the Treatment of Typhoid Fever: A Study From Kanjuruhan Hospital, Malang, Indonesia. *Value in Health* 2018; 21:S65.
27. Bonanno C, Wapner RJ. Antenatal corticosteroids in the management of preterm birth: are we back where we started?. *Obstet Gynecol Clin North Am* 2012; 39(1):47-63.
28. Fortunato SJ, Welt SI, Eggleston M, Cole J, Bryant EC, Dodson MG. Prolongation of the latency period in preterm premature rupture of the membranes using prophylactic antibiotics and tocolysis. *J Perinatol* 1990; 10(3):252-6.
29. Lee J, Romero R, Kim SM, Chaemsaitong P, Yoon BH. A new antibiotic regimen treats and prevents intra-amniotic inflammation/infection in patients with preterm PROM. *J Matern Fetal Neonatal Med* 2016; 29(17):2727-37.
30. Kenyon S, Taylor DJ, Tarnow-Mordi WO; ORACLE Collaborative Group. ORACLE--antibiotics for preterm prelabour rupture of the membranes: short-term and long-term outcomes. *Acta Paediatr Suppl* 2002; 91(437):12-5.
31. Buitendijk SE. Erythromycin for premature rupture of membranes is beneficial for infant. *Ned Tijdschr Geneesk* 2001; 145(38):1828-31.
32. Kibret M, Abera B. Antimicrobial susceptibility patterns of *E. coli* from clinical sources in northeast Ethiopia. *Afr Health Sci* 2011; 11 Suppl 1(Suppl 1):S40-S45.
33. Seelbach-Goebel B. Antibiotic Therapy for Premature Rupture of Membranes and Preterm Labor and Effect on Fetal Outcome. *Geburtshilfe Frauenheilkd* 2013; 73(12):1218-1227.
34. Navathe R, Schoen CN, Heidari P, Bachilova S, Ward A, Tepper J, Visintainer P, Hoffman MK, Smith S, Berghella V, Roman A. Azithromycin vs erythromycin for the management of preterm premature rupture of membranes. *Am J Obstet Gynecol* 2019; 221(2):144.e1-144.e8.
35. Pierson RC, Gordon SS, Haas DM. A retrospective comparison of antibiotic regimens for preterm premature rupture of membranes. *Obstet Gynecol* 2014; 124(3):515-519.
36. Gelber S, Brent E, Varrey A, Fridman B, Sapra K, Frayer W. Equivalence of erythromycin and azithromycin for treatment of PPROM. *Am J Obstet Gynecol* 2013; 208(1):S291.
37. Mercer BM, Miodovnik M, Thurnau GR, Goldenberg RL, Das AF, Ramsey RD, et al. Antibiotic therapy for reduction of infant morbidity after preterm premature rupture of the membranes: a randomized controlled trial. *JAMA* 1997; 278(12):989-995.
38. Desai SV, Tank P. Handbook on Preterm Prelabor Rupture of Membranes in a Low Resource Setting. *JAYPEE* 2012; 35-36.
39. Nadisauskiene R, Bergström S, Kilda A. Ampicillin in the treatment of preterm labor: a randomised, placebo-controlled study. *Gynecol Obstet Invest* 1996; 41(2):89-92.
40. American Academy Of Pediatrics Committee On Fetus And Newborn; American College Of Obstetricians And Gynecologists Committee On Obstetric Practice. The Apgar Score. *Pediatrics* 2015; 136(4):819-822.
41. Salan YDC. Effect of Length of Premature Rupture of Membranes Period toward Apgar Score of The Infant and Maternal Leukocyte Serum in Ulin General Hospital Banjarmasin. *Berkala Kedokteran* 2017; 13:199-204.

**Publish in The International  
Arabic Journal of Antimicrobial Agents**

The Journal is an open access peer-reviewed journal that publishes scientific papers about all aspects of antimicrobials. The journal will publish original research articles, reviews, brief reports and case reports dealing with basic and clinical antibacterial agents, antiviral, antiprotozoals, antituberculous, antifungal and antihelminthes agents. All manuscripts must be prepared in English, and are subject to a rigorous and fair peer-review process. Accepted papers will immediately appear online. The journal aims to advance the knowledge, attitude and the research of chemotherapy in the Arabic world in cooperation with international, national scientific and public societies as well as research centers with similar aims and objectives.

# Cefotaxime vs ceftriaxone for the prolongation of latency period in preterm premature rupture of membranes

## ORIGINALITY REPORT

17%

SIMILARITY INDEX

12%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://scialert.net">scialert.net</a> Internet Source	2%
2	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet Source	1%
3	Nir Melamed. "Factors affecting the duration of the latency period in preterm premature rupture of membranes", The Journal of Maternal-Fetal & Neonatal Medicine, 2009 Publication	1%
4	<a href="http://thieme-connect.com">thieme-connect.com</a> Internet Source	1%
5	<a href="http://www.hkmj.org">www.hkmj.org</a> Internet Source	1%
6	<a href="http://repository.tudelft.nl">repository.tudelft.nl</a> Internet Source	1%
7	<a href="http://media.neliti.com">media.neliti.com</a> Internet Source	1%
8	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	

1 %

9

"Committee Opinion No. 644 Summary",  
Obstetrics & Gynecology, 2015

Publication

1 %

10

Daniela Cristina Schroff Machado, Joice de  
Moura Dias, Luiza Emylce Pelá Rosado.  
"Antibiotic prophylaxis in preterm rupture of  
membranes", Human Reproduction Archives,  
2017

Publication

1 %

11

Brian M. Mercer. "Preterm Premature  
Rupture of the Membranes: Current  
Approaches to Evaluation and Management",  
Obstetrics and Gynecology Clinics of North  
America, 2005

Publication

1 %

12

[www.mysciencework.com](http://www.mysciencework.com)

Internet Source

1 %

13

[repository.unair.ac.id](http://repository.unair.ac.id)

Internet Source

1 %

14

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

<1 %

15

Reshama Navathe, Corina N. Schoen, Paniz  
Heidari, Sophia Bachilova et al. "Azithromycin  
vs erythromycin for the management of

<1 %

preterm premature rupture of membranes",  
American Journal of Obstetrics and  
Gynecology, 2019

Publication

---

16

[www.psmid.org.ph](http://www.psmid.org.ph)

Internet Source

<1 %

---

17

Ramesh Devkota, G. M. Khan, Kadir Alam,  
Amisha Regmi, Binaya Sapkota. "Medication  
utilization pattern for management of  
pregnancy complications: a study in Western  
Nepal", BMC Pregnancy and Childbirth, 2016

Publication

<1 %

---

18

Sarah Dotters-Katz. "Antibiotics for  
Prophylaxis in the Setting of Preterm Prelabor  
Rupture of Membranes", Obstetrics and  
Gynecology Clinics of North America, 2020

Publication

<1 %

---

19

"OS18: Interstitial Lung Disease 1",  
Respirology, 2013

Publication

<1 %

---

20

Ramakanta Sahoo, Sushreesmita Mohanty.  
"STUDY ON AETIOLOGY & RISK FACTORS IN  
SPONTANEOUS LATE PRETERM PREMATURE  
RUPTURE OF MEMBRANES", Journal of  
Evolution of Medical and Dental Sciences,  
2019

Publication

---

<1 %

21 Birjna A. Hirani, Bariki L. Mchome, Nicholas S. Mazuguni, Michael J. Mahande. "The decision delivery interval in emergency caesarean section and its associated maternal and fetal outcomes at a referral hospital in northern Tanzania: a cross-sectional study", BMC Pregnancy and Childbirth, 2017  
Publication <1 %

---

22 Gidon Test, Amalia Levy, Arnon Wiznitzer, Moshe Mazor, Gershon Holcberg, Alexander Zlotnik, Eyal Sheiner. "Factors affecting the latency period in patients with preterm premature rupture of membranes", Archives of Gynecology and Obstetrics, 2010  
Publication <1 %

---

23 docplayer.net  
Internet Source <1 %

---

24 journal.unair.ac.id  
Internet Source <1 %

---

25 mafiadoc.com  
Internet Source <1 %

---

26 nepjol.info  
Internet Source <1 %

---

27 www.humanreproductionarchives.com  
Internet Source <1 %

---

28 www.karger.com

<1 %

29

B. Seelbach-Goebel. "Antibiotic Therapy for Premature Rupture of Membranes and Preterm Labor and Effect on Fetal Outcome", *Geburtshilfe und Frauenheilkunde*, 2013

Publication

<1 %

30

Faisal O. Al-Qurashi, Abdullah A. Yousef, Bassam H. Awary. "Epidemiological aspects of prematurity in the Eastern region of Saudi Arabia", *Saudi Medical Journal*, 2016

Publication

<1 %

31

H. Lode, J. Eller, M. Raffenberg, H. Mauch. "Role of long-acting cephalosporins in the treatment of lower respiratory tract infections", *Clinical Microbiology and Infection*, 2000

Publication

<1 %

32

M. Guckert, E. Clouqueur, E. Drumez, C. Petit, V. Houfflin-Debarge, D. Subtil, C. Garabedian. "Is homecare management associated with longer latency in preterm premature rupture of membranes?", *Archives of Gynecology and Obstetrics*, 2019

Publication

<1 %

33

[docplayer.es](https://www.docplayer.es)  
Internet Source

<1 %

34

"The Role of HFCs in Mitigating 21st Century Climate Change", Climate Policy, 2016.

Publication

<1 %

35

KÅRE S. OLAFSEN, JOHN A. RØNNING, LAURITZ BREDRUP DAHL, STEIN ERIK ULVUND, BJØRN HELGE HANDEGÅRD, PER IVAR KAARESEN. "Infant responsiveness and maternal confidence in the neonatal period", Scandinavian Journal of Psychology, 2007

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On



# Cefotaxime vs ceftriaxone for the prolongation of latency period in preterm premature rupture of membranes

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---