

High intensity interval training for improving maximum aerobic capacity in women with sedentary lifestyle a systematic review and meta analysis

by Fajar Syamsudin

Submission date: 19-Oct-2021 10:41AM (UTC+0800)

Submission ID: 1677685157

File name: th_sedentary_lifestyle_a_systematic_review_and_meta_analysis.pdf (491.54K)

Word count: 6425

Character count: 36013

53
High-intensity interval training for improving maximum aerobic capacity in women with sedentary lifestyle: a systematic review and meta-analysis

FAJAR SYAMSUDIN¹, CITRAWATI DYAH KENCONO WUNGU², EMA QURNIANINGSIH³, LILIK HERAWATI⁴

^{1,4}Master Program of Sport Health Science, Faculty of Medicine, Universitas Airlangga, INDONESIA

^{2,3}Department of Biochemistry, Faculty of Medicine, Universitas Airlangga, INDONESIA

⁴Department of Physiology, Faculty of Medicine, Universitas Airlangga, INDONESIA

Published online: June 30, 2021

(Accepted for publication June 15, 2021)

DOI:10.7752/jpes.2021.04226

Abstract

33
Introduction: Sedentary lifestyle behavior increases the risk of non-communicable diseases such as diabetes mellitus, cardiovascular disease, and cancer. High-intensity interval training (HIIT) is an effective and efficient exercise to overcome this problem. It has been proven that HIIT effectively increases the maximal aerobic capacity (VO2Max) as an indicator of cardiovascular health. *Objective:* The purpose of this study was to analyze the effect of HIIT on women with sedentary lifestyle and to evaluate the difference in VO2max using the Pre and Post group design. *Methods:* Three electronic databases were searched (i.e., PubMed, ScienceDirect, and ProQuest) for original research articles. The search was performed using the following search term – ‘high intensity interval training’ and ‘maximum aerobic capacity’ and ‘sedentary’. Inclusion criteria were controlled trials, healthy 18–55-year-old women, training duration >2 weeks, and VO2max assessed during pre and post training. The RevMan 5.4 software was used for analysis in fixed/random effect models. *Result:* A total of 7 studies met the inclusion criteria and were included in the meta-analysis. Meta-analysis was divided into 2 analysis groups, i.e., 6- and 12-week training. The standardized mean difference (SMD) values of 6- and 12-week-training were 1.31 (85% CI 0.42–2.21) and 1.13 (0% CI 0.70–1.57), respectively. *Conclusion:* HIIT with a minimum of 4 weeks of exercise training (3 times per week, 18-30 minutes per session) is an effective training protocol for increasing maximal aerobic capacity in women with a sedentary lifestyle.

Key Words : HIIT, female, lifestyle, VO2max, fitness, health

Introduction

52
Sedentary style (lack of physical activity or sedentary lifestyle) increases along with the development of times (M. Lee et al., 2021; Ozemek, Lavie, & Rognum, 2019). In addition, there are a lot of evidence showing that sedentary lifestyle increases various risks that can disrupt health conditions (I. M. Lee et al., 2012), including including can increase the risk of diabetes by 112%, increase the risk of cardiovascular diseases such as heart attacks and strokes by 147%, increase the risk of death from cardiovascular disease by 90% and increase the risk of death from various causes by 49% (Rahayu, Aprilawati, Mahmud, Purwanto, & Herawati, 2019; Wilmot et al., 2012).

4
The World Health Organization (WHO) (World Health Organization, 2020) has recommended at least 150-300 minutes of moderate physical activity (40-60% HRM) or 75-150 minutes of high-intensity/ vigorous physical activity (60-85% HRM) per week to maintain and improve the stamina of healthy adults (18-64 years old). However, 31.1% of adults worldwide fail to meet these minimum physical activity guidelines, and inactivity increases with age (Hallal et al., 2012). Lack of time, low motivation, and non-compliance to established guidelines are frequently mentioned as the challenge to engage in physical activity (Hoare, Stavreski, Jennings, & Kingwell, 2017; Rech et al., 2016).

To carry out physical activity, the body needs oxygen to burn calories into energy (Guyton & Hall E, 2016). However, the oxygen capacity of each person is different (Hoeger, Hoeger, Fawson, & Hoeger, 2019). A higher oxygen capacity is an indicator of the level of cardiovascular health and enables body to perform greater physical activity (Saghiv, Sagiv, Saghiv, & Sagiv, 2020; Scott, 1998). The parameter of oxygen coverage is called the maximal aerobic capacity, in short known as VO2Max (Maximal Oxygen Volume), stated in units of mL·kg⁻¹·min⁻¹. VO2max is an indicator of cardiovascular health due to its vital role in the body (Batacan, Duncan, Dalbo, Tucker, & Fenning, 2017). VO2max can be increased by routine and programmed exercise (Arboleda-Sema, Feito, Patiño-Villada, Vargas-Romero, & Arango-Vélez, 2019; Astorino et al., 2017).

HIIT is one of the exercises recommended for someone who doesn't have much leisure time. HIIT is less time-consuming compared to over low or moderate intensity or continuous training, making it more preferable to provide similar or greater health benefits than the recommended physical activity (Norton, Norton, & Sadgrove, 2010; Ramos, Dalleck, Tjonna, Beetham, & Coombes, 2015). HIIT can overcome the challenge of

1788

Corresponding Author: LILIK HERAWATI, E-mail: lilik_heraw@fk.unair.ac.id

lack of time to perform physical (Reichert, Barros, Domingues, & Hallal, 2007; Syamsudin, Herawati, Qurnianingsih, & Wungu, 2021), making HIIT the third most popular sport in the world (Thompson, 2018). It was also reported that HIIT is good to improve fitness, cardiovascular function, insulin sensitivity, and reduce atherosclerosis (Kong et al., 2016; Lanzi et al., 2015).

As such, HIIT is in great demand and beneficial method for improving fitness and cardiovascular health levels (Batacan et al., 2017). However, the current systematic review of HIIT remains limited to subjects that are mixed between men and women (Kriel, Askew, & Solomon, 2019; Viana et al., 2019). It has not focused on female subjects although the prevalence of obesity is more dominant in women (World Health Organization, 2016) and problems associated with ovulatory dysfunction are mostly found in obese females (Broughton & Moley, 2017). In addition, there is no systematic review that is specifically focused on sedentary lifestyle or physical inactivity. Therefore, this systematic review was aimed to analyze the effect of HIIT on maximum aerobic capacity in female subjects whose sedentary lifestyle and physical inactivity in the normal, overweight, and obese categories based on published studies.

Materials & methods

This systematic review article followed the guidelines for Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) (Moher et al., 2016), and was registered in the International Prospective Register of Systematic Review (PROSPERO) (Chien, Khan, & Siassakos, 2012) (CRD42020218218).

Search Strategy. Electronic databases were searched in PubMed, Science Direct, and ProQuest. The keywords used in the search process were completed with MeSH Terms, Titles / Abstracts, and asterisks (*) for words that are likely to have various types of subsequent words. "High Intensity Interval Training OR High Intensity Interval Exercise OR High Intensity Intermittent Exercise OR Interval Training OR Interval Exercise OR HIIT OR Sprint Interval Training OR Sprint Training" AND "Sedentary OR Sedentary* OR Sedentary Lifestyle OR Sedentary Behaviour* OR Inactive OR Inactive* OR Physical Inactivity" AND "VO2Max OR VO2 Max OR VO2Max OR Maximum Oxygen Volume OR Maximum Oxygen Capacity OR Maximal Oxygen Uptake" were used as keywords. In addition to the obtained articles, several articles beside the unpublished database were used.

Inclusion and Exclusion criteria : Participants, Interventions, Comparisons and Outcomes. Studies on female subjects whose a sedentary lifestyle between the ages of 18-65 years, subjects were given high-intensity training with intervals, training included weight-bearing or non-weight bearing. Studies were not limited to treatment comparisons, may be compared to any exercise treatment such as MICT (Moderate Intensity Continuous Training), circuit training, resistance training or control, and each study should have a result on VO2Max (Maximum Oxygen Volume).

Data Extraction. The characteristics of the study must meet data on gender, age, VO2max, filtering sedentary questionnaires, tools used in the intervention, bouts/active and rest/passive phases of training, and data in each group must have Pre and Post Test data. For any incomplete data, the author of the article was contacted to provide complete data or clarification of the data needed. However, if there was no response, it was included or excluded with several conditions.

Assessment of Bias. Two investigators independently performed the electronic search and retrieved the articles that matched with our searched terms. Any disagreement was settled by discussion and consensus with all the authors. Final decision was merely based on the agreements of all authors.

Study Quality Assessment. The quality of each study was assessed using the Jadad Score. Jadad Score consists of three items related to the description of randomization (0–2 points), blinding (0–2 points), and drop-out (0–1 points) with a total of five scores. Higher scores indicate better quality. High-quality trials were defined as trials that score more than 2. A low-quality trial was defined as trials that score 2 or less.

Data Analysis. Analysis was conducted using Review Manager 5.3 software (The Cochrane Collaboration, UK). Hardy Weinberg Equilibrium was examined by Chi Square test when the raw information was not provided. The standardized mean difference (SMD) data were evaluated in all obtained studies. Heterogeneity assumption was assessed with Cochrane Q statistic and I² statistic. The pool estimated ORs was calculated with either fixed or random effects model assumptions. If Q test showed significant result ($p < 0.05$), a random effects model was used. Otherwise, if Q test showed insignificant result ($p > 0.05$), a fixed effect model was used. 95% confidence interval (CI) of pool estimated SMD was also calculated. Inverted funnel plots were used to find any publication bias.

Result

The search strategy from electronic and non-electronic database resulted in 2321 and 4 articles, respectively. After checking for duplication using Mendeley, the number of articles decreased by 34, leading to exclusion of 2264 articles through the screening process by reading titles and abstracts. The remaining 24 articles were selected in full text; however, after understanding them in full text, 16 articles were published for the following reasons: (1) Five data were mixed with men; (2) Two single articles were bout; (3) Six articles did not provide VO2Max data; (4) One article was not sedentary; (5) One article involved youth subjects; (6) One HIIT

article was conducted simultaneously with Resistance Training; and (7) One subject article had disease. In the end, 7 articles provided sufficient data was analyzed in a systematic review. 6 articles were then used for meta-analysis and divided into 2 categories such as 6-week and 12-week meta-analysis.

Study Quality. Results of the quality of all studies were analyzed (Astorino et al., 2013; Bhati, Bansal, & Moiz, 2017; Clark, Rosa, DeRevere, & Astorino, 2019; Metcalfe, Tardif, Thompson, & Vollaard, 2016; Nie et al., 2018; Trilk, Singhal, Bigelman, & Cureton, 2011) using the jasad score. A total of six studies (Astorino et al., 2013; Bhati et al., 2017; Clark et al., 2019; Nie et al., 2018; Trilk et al., 2011) were high quality, and only one study (Metcalfe et al., 2016) was low quality.




Fig. 1 Prisma Flow diagram illustrating the study selection process.

Characteristics of Participant. The characteristics of participants were summarized in Table 2. All 130 participants were included in the systematic review. Study (Astorino et al., 2013; Bhati et al., 2017; Nie et al., 2018; Zhang et al., 2015) used subjects with an average age of 17-30 years, while study (Clark et al., 2019; Metcalfe et al., 2016; Trilk et al., 2011) used subjects with an average age of 30-45 years. The number of participants in each study was different, ranging from 8-18 people in each group. Studies by (Bhati et al., 2017; Metcalfe et al., 2016) involved normal BMI subjects, two studies by (Astorino et al., 2013; Nie et al., 2018) involved the overweight category, while a study by Trilk et al. (2011) involved subjects with class 1 obesity and a study by Clark et al. (2019) involved subjects with class 2 obesity.

Characteristics of Intervention. High-intensity interval training (HIIT) describes physical exercise that is characterized by brief, intermittent bursts of vigorous activity, interspersed by periods of rest or low-intensity exercise (Gibala, Little, Macdonald, & Hawley, 2012; MacInnis & Gibala, 2017). The HIIT programs were summarized in Table 1. Five studies (Astorino et al., 2013; Clark et al., 2019; Metcalfe et al., 2016; Nie et al., 2018; Trilk et al., 2011) used a cycle ergometer as an intervention tool, while two studies (Bhati et al., 2017; Zhang et al., 2015) used treadmills.

Duration of training varied, intervention in three studies (Astorino et al., 2013; Nie et al., 2018; Zhang et al., 2015) were given over twelve weeks, four studies (Bhati et al., 2017; Clark et al., 2019; Metcalfe et al., 2016) were given for six weeks, one study (Trilk et al., 2011) was given for four weeks. For one-week intervention, nearly all studies (Astorino et al., 2013; Bhati et al., 2017; Clark et al., 2019; Metcalfe et al., 2016;

Triik et al., 2011) were performed 3 times, and one study Nie et al., (2018) was performed 3-4 times progressively.

Volume of training provided varied, studies (Metcalf et al., 2016; Zhang et al., 2015) took no more than 15 minutes, a fixed volume of training from the beginning of the meeting to the end of the meeting was carried out in the study by (2 groups) Bhati et al., (2017) and study by Clark et al., (2019), with volume of 22 minutes, 43 minutes and 25 minutes. In addition, study by (Astorino et al., 2013; Triik et al., 2011) had a progressive volume, while a study by Nie et al. (2018) used energy expenditure benchmarks.

Intensity of all studies was high, but had different measuring instruments, four studies (Astorino et al., 2013; Bhati et al., 2017; Clark et al., 2019; Zhang et al., 2015) used 70-100% HRmax, one study (Nie et al., 2018) used 90 - 95% VO2max, and two studies (Metcalf et al., 2016; Triik et al., 2011) asked subjects to do maximum sprint / all-out.

Table 1. Quality Assessment of Articles

No	Study	Randomization	Blinding	Drop-Out	Total of	Concealment
1.	Metcalf et al., (2016)	0	1	1	2	Low quality
2.	Bhati et al., (2017)	2	0	1	3	High quality
3.	Triik et al., (2011)	2	0	1	3	High quality
4.	Clark et al., (2019)	1	1	1	3	High quality
5.	Astorino et al., (2013)	1	1	1	3	High quality
6.	Nie et al., (2018)	2	1	1	4	High quality
7.	Zhang et al., (2015)	2	0	1	3	High quality

Note: Jadad Score consists of three items related to the description of randomization (0–2 points), blinding (0–2 points), and drop-out (0–1 points) with a total of five scores. Higher scores indicated better quality. High-quality trials were defined as trials that scored more than 2. A low-quality trial was defined as trials that scored 2 or less.

Sedentary Lifestyle. All subjects were categorized as healthy, four studies (Bhati et al., 2017; Clark et al., 2019; Metcalf et al., 2016; Triik et al., 2011) used the PAR-Q (Physical Activity Readiness Questionnaire) to assess the health quality of subjects, while four other studies did not use the PAR-Q questionnaire but provided information that subjects were healthy and did not have a disease that could harm the subject itself. To determine sedentary lifestyle subjects, two studies by Astorino and Nie (Astorino et al., 2013; Nie et al., 2018) used PYTPAQ (Past Year Total Physical Activity Questionnaire), a study by Metcalf et al. (2016) used IPAQ (International Physical Activity Questionnaire), while another study did not use a questionnaire but confirmed that the subjects had sedentary lifestyle, with different criteria in each study. More details can be seen in Table 2.

Table 2. Characteristics of study population and studies

No	Author	Country	Sedentary Questionnaire	Age & BMI	Group Size	Interventions	Pre VO2Max (mL·kg ⁻¹ ·min ⁻¹)	Post VO2Max (mL·kg ⁻¹ ·min ⁻¹)	P-value of Pre Post Test
1.	Metcalf et al., (2016)	United Kingdom	IPAQ, PAR-Q Body mass stable and no conscious change in diet or physical activity patterns over the preceding 6 months	A : 36 ± 9 B : 24.1 ± 3.5 (overweight)	Size (35) Male (17) Female (18)	T : Cycle Ergometer D : 6 weeks, 3x per week V : 10 min per training I : 10 – 40 s sprint, 3-4.50 min passive, ariation and progression	31.7 ± 4.6	34.7±5.2	< 0.001
2.	Bhati et al., (2017)	India	PAR-Q Not engaging in any leisure time physical activity or	A : 22.0 ± 0.3 B : 21.3 ± 0.3 (normal)	Size 28 LVHIIT (17) HVHIIT (15)	T : Treadmill D : 6 weeks, 3 x per week V : 22 min (LVHIIT) I : 4 min active – 3 min recovery,	20.4 ± 0.6	23.1 ± 0.6	< 0.001

No	Author	Country	Sedentary Questionnaire	Age & BMI	Group Size	Interventions	Pre VO ₂ Max (mL·kg ⁻¹ ·min ⁻¹)	Post VO ₂ Max (mL·kg ⁻¹ ·min ⁻¹)	P-value of Pre Post Test
			there was only one instance of low-intensity exercise or sports activity for less than 20 min in a week No participation in any aerobic or resistance training in the last 6 months	A : 22.0 ± 0.3 B : 21.3 ± 0.3 (normal)	Size 28 LVHIIT (17) HVHIIT (15)	active : 85–95% HRmax, Passive : 70% HRmax T : Treadmill D : 6 weeks, 3 x per week V : 43 min (HVHIIT) I : 4 min active – 3 min recovery, active : 85–95% HRM, passive : 70% HRM	20.9 ± 0.6	22.6 ± 0.8	< 0.04
3.	Trilk et al., (2011)	USA	Sedentary Exercise < 1 day per week	A : 35.7 ± 6.3 B : 30.1 ± 6.8 (class 1 obesity)	Size 28 SIT (14) Control (14)	T : Cycle Ergometer D : 4 weeks, 3 x per week V : 18 – 31.30 menit I : 30 s active, 4 min passive, 4-7 sets, progressive, Active : All-out (sprint), Passive : Active recovery	21.6 ± 1.1	24.5 ± 1.1	< 0.05
4.	Clark et al., (2019)	USA	PAR-Q Sedentary < 2 hour of moderate-to-vigorous activity per week in the previous 12 months	A : 39.1 ± 4.3 B : 37.5 ± 10.5 (class 2 obesity)	Size 17 HIIT Tradit (9) HIIT Period (8)	T : Cycle Ergometer D : 6 weeks, 3x per week V : 25 min I : WU – 5 min, Active 60 s – passive 60 s, 10 sets, active : 80–95% HRM, passive : -	21.5 ± 3.2	22.5 ± 3.2	< 0.44
				A : 39.1 ± 4.3 B : 37.5 ± 10.5 (class 2 obesity)	Size 17 HIIT Tradit (9) HIIT Period (8)	T : Cycle Ergometer D : 6 weeks, 3x per week V : 25 min I : WU – 5 min, variation active and passive, variation sets, active : 80–95% HRM, passive : -	17.3 ± 2.4	18.0 ± 2.2	< 0.49
5.	Astorino et al.,	USA	PYT-PAQ	A : 23.1 ± 5.6	Size 27 HLHIIT	T : Cycle Ergometer	29.4 ± 5.9	6 week 34.8 ±	< 0.05

No	Author	Country	Sedentary Questionnaire	Age & BMI	Group Size	Interventions	Pre VO2Max (mL·kg ⁻¹ ·min ⁻¹)	Post VO2Max (mL·kg ⁻¹ ·min ⁻¹)	P-value of Pre Post Test
	(2013)			B : 25.3 ± 4.3 (overweight)	(10) LOHIIT (10) Control (7)	D : 12 weeks, 3x per week V : 12-23 min I : 6-10 s bouts, 75 s recovery, 6-10 sets, active : 85-100 % HRM, passive: -		5.7 12 weeks 35.8 ± 5.7	
				A : 24.7 ± 7.6 B : 22.5 ± 4.5 (overweight)	Size 27 HI.HIIT (10) LO.HIIT (10) Control (7)	T : Cycle Ergometer D : 12 weeks, 3x per week V : 12-23 min I : 6-10 s bouts, 75 s recovery, 6-10 sets, active : 75-95 % HRM, passive : -	29.4 ± 5.9	6 weeks 34.1 ± 4.9 12 weeks 36.4 ± 4.4	< 0.05
6.	Nie et al., (2018)	China	PYT-PAQ Body fat percentage ≥35% Body weight remained constant (±2 kg) during the past 3 months No regular physical activities or exercise training	A : 21.0 ± 1.1 B : 26.3 ± 3.6 (overweight)	Size 48 HIIT (17) MICT (15) Control (16)	T : Cycle Ergometer D : 12 weeks, 3-4x per week, progressive V : The duration of training was based on the use of energy, which was 200-300kJ I : 4 min bouts 90% VO2max, 3 min passive	30.2 ± 4.4	34.3 ± 4.6	< 0.05
7	Zhang et al., (2015)	China	No regular physical training Body weight remained constant (±2 kg) during the past 3 month	A : 21.0 ± 1.0 B : 25.8 ± 2.7 (overweight)	Size 35 HIIT (12) MICT (12) Control (11)	T : Treadmill D : 12 weeks, 4x per week V : 15 min I : 4 min 85-95% HRM, 3 min 50-60% HRM, and 7 min active rest	33.1 ± 3.0	37.7 ± 3.0	< 0.05

Questionnaire: PAR-Q, Physical Activity Readiness Questionnaire; IPAQ, International Physical Activity Questionnaire; PYT-PAQ, Past Year Total Physical Activity Questionnaire. **Age & BMI :** A, Age; B, BMI. **Interventions:** T, Treatment; D, Duration; V, Volume; I, Intensity. **Group :** HIIT, High Intensity Interval Training; LVHIIT, Low Volume High Intensity Interval Training; HVHIIT, High Volume High Intensity Interval Training; SIT, Sprint Interval Training; HI.HIIT, High Intensity-High Intensity Interval Training; LO.HIIT, Lower Intensity-High Intensity Interval Training.

Seven research groups with 87 subjects were included to prove the effects of HIIT for six weeks. Because heterogeneity was found in the statistical analysis, a sensitivity analysis was performed to evaluate the

source of the heterogeneity. It was found that the heterogeneity between the studies was mainly due to Bhati's et al., (2017) study with the 2 research groups of HV and LV, because no significant heterogeneity was found after this study was excluded. The heterogeneity between studies for all models (12) was 85% and the P value for heterogeneity was 0.004. Random -Effect Model was applied. The results of the analysis showed a significant increase in VO2max with a p value of 0.04 (<0.05) and an SMD value of 1.31 with a percentage of 85% with a minimum and maximum CI limit of 0.42 and 2.21, respectively. Therefore, it was concluded that 6-week training led to 1.31 times increase in VO2max from pre-test to post test.




Fig. 2. Forest plot of 6-week training

Four research groups consisting of 49 subjects were included to prove effect of HIIT for twelve weeks. In this meta-analysis, there was no significant change in SMD by excluding any study, indicating that there was no single study that affected the statistical significance in whole results.

Results of analysis showed a significant increase of VO2max with p-value 0.00001 ($p < 0.05$), SMD value of 1.45, percentage of 81%, minimum and maximum CI limit of 0.80 and 2.11, respectively. Therefore, it was concluded that 12-week training led to increased VO2max from pre-test to post-test.




Fig. 3. Forest plot of 12-week training

23

The funnel plot image showed an asymmetrical distribution of the study, indicating an imbalance in left and right side of center line boundary. This might be due to the heterogeneity of the study by Bhati. Therefore, it was concluded that publication bias affected the relationship between 6-week HIIT training and VO2max.

The funnel plot image showed a symmetrical distribution of study, indicating a balanced distribution to the left and right side of the center line boundary. Therefore, it was concluded that publication bias did not affect the relationship between 12-week HIIT training and VO2max.




Fig. 4. Funnel Plot of 6-week training




Fig. 5. Funnel Plot of 12-week training

Discussion

The aim of this systematic review and meta-analysis was to review some references analyzing HIIT in sedentary women and determine its potential effects on cardiovascular health. In addition, this meta-analysis aimed to identify the HIIT protocol in relation to the comparison of the pre and post test results of VO₂max. Seven studies with 10 training groups were evaluated for evidence showing that HIIT can increase VO₂max in sedentary women. The main finding from the meta-analysis was that HIIT training significantly increased VO₂max.

In the Pre Test data, two studies by (Metcalf et al., 2016; Zhang et al., 2015) showed average VO₂max levels, 5 studies (Amaro-Gahete et al., 2019; Astorino et al., 2013; Clark et al., 2019; Nie et al., 2018; Trilk et al., 2011) showed that the subjects had fair VO₂max levels, while study by Bhati (2) and Clark showed poor VO₂max levels in 3 research groups. After HIIT training, increased VO₂max were found in all studies, except three studies by (Amaro-Gahete et al., 2019; Astorino et al., 2013; Nie et al., 2018) which managed to increase the VO₂max to an average level from the previously fair level. However, this proved that people with a sedentary lifestyle have a low level of cardiovascular fitness, highlighting the important role of HIIT to improve cardiovascular fitness in people with sedentary lifestyle.

The study by Bhati et al. (2017) explained that there was no significant difference in post test results of VO₂max between HIIT volume training with a duration of 43 minutes compared to that of 22 minutes which was carried out with the same number of meetings and durations. This result indicated that training in the same number of sessions but different training volumes led to similar or slightly different results.

A study by Metcalfe et al. (2016) proved that time-saving was indeed one advantage of HIIT. It was shown this study that training in one meeting which lasted only 10 minutes and a total of only 30 minutes for one week provided significant results between pre and post.

Studies by (Clark et al., 2019; Trilk et al., 2011) which involved subjects with class 1 and class 2 obesity showed that exercise is very important for overweight people because their fitness level is very low. The fitness level was determined based on Hoeger et al. (2019) criteria suggesting that fair level is for women aged 30-39 years with VO₂max <20 - 27,9 mL·kg⁻¹·min⁻¹ and poor level if they have VO₂max <19. mL·kg⁻¹·min⁻¹. In the study by Trilk et al. (2011) the VO₂max level was only 21.6 ± 1.1 - 24.5 ± 1.1 mL·kg⁻¹·min⁻¹, while in the study by Clark et al. (2019) the VO₂max level was only 17.3 ± 2.4 - 18.0 ± 2.2 mL·kg⁻¹·min⁻¹. Clark et al. (2019) also compared HIIT with the same exercise program from the first meeting to the last, with HIIT training programs that varied in active volume (sprint), passive (recovery) and intensity. The results showed that a variety of exercise programs led to better results than a monotonous training program.

It was shown in a study by Astorino et al. (2013) which compared HI.HIIT and LO.HIIT that a longer meeting in HIIT training led to better VO₂max results. It was shown that in week 3, 6, 9, and 12, there were increased VO₂max levels.

The forest plot of 6-week training showed that the intervention led to 1.31 times increase of VO₂max from pre-test to post test, with a minimum limit of increase of 0.42 and a maximum limit of increase of 2.21. Furthermore, in forest plot Fig. 3, the 12-week training showed that the intervention led to 1.13 times increase of VO₂max with a minimum limit of increase of 0.70 and a maximum limit of increase of 1.57. Both forest plots Fig. 2. and Fig. 3. used a confidence level of 95% and an error rate of 5%.

Publication bias is a problem in meta-analysis because it only takes published research (without unpublished research) and only combines research results without critical identification, leading to an overly optimistic meta-analysis model. Funnel plot is a technique used for identification of publication bias in meta-analysis. The following is a funnel plot on the effect of HIIT on VO₂max in women with sedentary lifestyle, which was divided based on the duration of treatment.

Funnel plot Fig. 4. shows asymmetrical distribution of the research, because there was a study which showed a great result effect causing no compatibility between the average data and the maximum data from the effect data of other studies. Therefore, it was suggested that the research group of funnel plot had a high probability of bias. Furthermore, funnel plot Fig. 5. shows the symmetrical distribution of research because there was no great difference of results effect from one another. Therefore, it was suggested that the research group of funnel plot had a small possibility of bias.

Conclusion

The findings of this systematic review indicated that HIIT is an effective training protocol to increase the maximal aerobic capacity in female subjects with a sedentary lifestyle. A minimum of 4 weeks of exercise with a frequency of 3 times per week, a training duration of 18-30 minutes per session, and various types of HIIT exercises such as circuit training, treadmill, and cycle ergometer must be considered to achieve an increased maximal aerobic capacity. However, it should be noted that in order to achieve the expected results, this exercise must achieve 85-95% HRM.

Conflict Of Interest

No potential conflict of interest relevant to this article was reported.

References

- Amaro-Gahete, F. J., De-La-O, A., Jurado-Fasoli, L., Dote-Montero, M., Gutiérrez, Á., Ruiz, J. R., & Castillo, M. J. (2019). Changes in physical fitness after 12 weeks of structured concurrent exercise training, high intensity interval training, or whole-body electromyostimulation training in sedentary middle-aged adults: A randomized controlled trial. *Frontiers in Physiology*, 10(APR), 451. <https://doi.org/10.3389/fphys.2019.00451>
- Arboleda-Serna, V. H., Feito, Y., Patiño-Villada, F. A., Vargas-Romero, A. V., & Arango-Vélez, E. F. (2019). Effects of a high-intensity interval training compared to moderate-intensity continuous training on maximal oxygen consumption and blood pressure in healthy men: A randomized controlled trial. *Biomedica*, 39(3), 524–536. <https://doi.org/10.7705/biomedica.v39i3.4451>
- Astorino, T. A., Edmunds, R. M., Clark, A., King, L., Gallant, R. A., Namm, S., Wood, K. M. (2017). High-Intensity Interval Training Increases Cardiac Output and V-O₂max. *Medicine and Science in Sports and Exercise*, 49(2), 265–273. <https://doi.org/10.1249/MSS.0000000000001099>
- Astorino, T. A., Schubert, M. M., Palumbo, E., Stirling, D., McMillan, D. W., Cooper, C., ... Gallant, R. (2013). Magnitude and time course of changes in maximal oxygen uptake in response to distinct regimens of chronic interval training in sedentary women. *European Journal of Applied Physiology*, 113(9), 2361–2369. <https://doi.org/10.1007/s00421-013-2672-1>
- Batacan, R. B., Duncan, M. J., Dalbo, V. J., Tucker, P. S., & Fenning, A. S. (2017). Effects of high-intensity interval training on cardiometabolic health: A systematic review and meta-analysis of intervention studies. *British Journal of Sports Medicine*. <https://doi.org/10.1136/bjsports-2015-095841>
- Bhati, P., Bansal, V., & Moiz, J. A. (2017). Comparison of different volumes of high intensity interval training on cardiac autonomic function in sedentary young women. *International Journal of Adolescent Medicine and Health*, 31(6), 1–13. <https://doi.org/10.1515/ijamh-2017-0073>
- Broughton, D. E., & Moley, K. H. (2017). Obesity and female infertility: potential mediators of obesity's impact. *Fertility and Sterility*. <https://doi.org/10.1016/j.fertnstert.2017.01.017>
- Chien, P. F. W., Khan, K. S., & Siassakos, D. (2012). Registration of systematic reviews: PROSPERO. *BJOG: An International Journal of Obstetrics and Gynaecology*. <https://doi.org/10.1111/j.1471-0528.2011.03242.x>
- Clark, A., Rosa, A. B. D. La, DeRevere, J. L., & Astorino, T. A. (2019). Effects of various interval training regimes on changes in maximal oxygen uptake, body composition, and muscular strength in sedentary women with obesity. *European Journal of Applied Physiology*, 119(4), 879–888. <https://doi.org/http://dx.doi.org/10.1007/s00421-019-04077-x>
- Gibala, M. J., Little, J. P., Macdonald, M. J., & Hawley, J. A. (2012). Physiological adaptations to low-volume, high-intensity interval training in health and disease. *Journal of Physiology*, 590(5):1077-84. <https://doi.org/10.1113/jphysiol.2011.224725>
- Guyton, A. C., & Hall E, J. (2016). Guyton and Hall Textbook of Medical Physiology, Thirteenth Edition. Elsevier. [https://doi.org/10.1016/S0140-6736\(02\)79504-4](https://doi.org/10.1016/S0140-6736(02)79504-4)
- Hallal, P. C., Andersen, L. B., Bull, F. C., Guthold, R., Haskell, W., Ekelund, U., Wells, J. C. (2012). Global physical activity levels: Surveillance progress, pitfalls, and prospects. *The Lancet*. [https://doi.org/10.1016/S0140-6736\(12\)60646-1](https://doi.org/10.1016/S0140-6736(12)60646-1)
- Hoare, E., Stavreski, B., Jennings, G. L., & Kingwell, B. A. (2017). Exploring Motivation and Barriers to Physical Activity among Active and Inactive Australian Adults. *Sports*, 5(3), 47. <https://doi.org/http://dx.doi.org/10.3390/sports5030047>
- Hoeger, W. W. K., Hoeger, S. A., Fawson, A. L., & Hoeger, C. I. (2019). Principles and labs for fitness and wellness, 606.
- Kong, Z., Fan, X., Sun, S., Song, L., Shi, Q., & Nie, J. (2016). Comparison of high-intensity interval training and moderate-to-vigorous continuous training for cardiometabolic health and exercise enjoyment in obese young women: A randomized controlled trial. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0158589>
- Kriel, Y., Askew, C. D., & Solomon, C. (2019). Sprint interval exercise versus continuous moderate intensity exercise: acute effects on tissue oxygenation, blood pressure and enjoyment in 18–30 year old inactive men. *PeerJ*. <https://doi.org/http://dx.doi.org/10.7717/peerj.7077>
- Lanzi, S., Codecasa, F., Cornacchia, M., Maestrini, S., Capodaglio, P., Brunani, A., Malatesta, D. (2015). Short-term HIIT and Fatmax training increase aerobic and metabolic fitness in men with class II and III obesity. *Obesity*. <https://doi.org/10.1002/oby.21206>
- Lee, I. M., Shiroma, E. J., Lobelo, F., Puska, P., Blair, S. N., Katzmarzyk, P. T., Wells, J. C. (2012). Effect of physical inactivity on major non-communicable diseases worldwide: An analysis of burden of disease and life expectancy. *The Lancet*, 380(9838), 219–229. [https://doi.org/10.1016/S0140-6736\(12\)61031-9](https://doi.org/10.1016/S0140-6736(12)61031-9)
- Lee, M., Lee, S., Robinson, S. R., Parrott, K. R., Lee, M., Lee, S., Parrott, K. R. (2021). Relationship between physical activity levels and perceived residential environment for rural older adults. *JPES*, 21(May), 1317–1324. <https://doi.org/10.7752/jpes.2021.03167>
- MacInnis, M. J., & Gibala, M. J. (2017). Physiological adaptations to interval training and the role of exercise

- intensity. *Journal of Physiology*. <https://doi.org/10.1113/JP273196>
- Metcalfe, R. S., Tardif, N., Thompson, D., & Vollaard, N. B. J. (2016). Changes in aerobic capacity and glycaemic control in response to reduced-exertion high-intensity interval training (REHIT) are not different between sedentary men and women. *Applied Physiology, Nutrition and Metabolism*, 41(11), 1117–1123. <https://doi.org/10.1139/apnm-2016-0253>
- Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., Whitlock, E. (2016). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Revista Espanola de Nutricion Humana y Dietetica*. <https://doi.org/10.1186/2046-4053-4-1>
- Nie, J., Zhang, H., Kong, Z., George, K., Little, J. P., Tong, T. K., Shi, Q. (2018). Impact of high-intensity interval training and moderate-intensity continuous training on resting and postexercise cardiac troponin T concentration. *Experimental Physiology*, 103(3), 370–380. <https://doi.org/10.1113/EP086767>
- Norton, K., Norton, L., & Sadgrove, D. (2010). Position statement on physical activity and exercise intensity terminology. *Journal of Science and Medicine in Sport*. <https://doi.org/10.1016/j.jsams.2009.09.008>
- Ozemek, C., Lavie, C. J., & Rognmo, Ø. (2019). Global physical activity levels - Need for intervention. *Progress in Cardiovascular Diseases*. <https://doi.org/10.1016/j.pcad.2019.02.004>
- Rahayu, T., Aprilawati, D. W. I., Mahmud, J., Purwanto, B., & Herawati, L. (2019). Both Upper and Lower Extremity-Only Video Game-Based Exercises (Exergaming) Affect Blood Glucose Serum Levels and Heart Rates but not Oxygen Saturation in Teenagers. *Journal of Physical Education and Sport*, 19(3), 802–807. <https://doi.org/10.7752/jpes.2019.s3114>
- Ramos, J. S., Dalleck, L. C., Tjonna, A. E., Beetham, K. S., & Coombes, J. S. (2015). The Impact of High-Intensity Interval Training Versus Moderate-Intensity Continuous Training on Vascular Function: a Systematic Review and Meta-Analysis. *Sports Medicine*. <https://doi.org/10.1007/s40279-015-0321-z>
- Rech, C., Camargo, E., Almeida, M., Bronoski, R., Okuno, N., & Reis, R. (2016). Barriers for physical activity in overweight adults. *Revista Brasileira de Atividade Física & Saúde*, 21(3), 272–279. <https://doi.org/10.12820/rbafs.v.21n3p272-279>
- Reichert, F. F., Barros, A. J. D., Domingues, M. R., & Hallal, P. C. (2007). The role of perceived personal barriers to engagement in leisure-time physical activity. *American Journal of Public Health*. <https://doi.org/10.2105/AJPH.2005.070144>
- Saghiv, M. S., Sagiv, M. S., Saghiv, M. S., & Sagiv, M. S. (2020). Introduction to Exercise Physiology. In *Basic Exercise Physiology*. https://doi.org/10.1007/978-3-030-48806-2_1
- Scott, J. R. (1998). Fox's Physiological Basis for Exercise and Sport, 6th Edition. *Medicine & Science in Sports & Exercise*. <https://doi.org/10.1097/00005768-199812000-00016>
- Syamsudin, F., Herawati, L., Qurnianingsih, E., & Wungu, D. C. K. (2021). HIIT for Improving Maximal Aerobic Capacity in Adults Sedentary Lifestyle: Review Article. *Halaman Olahraga Nusantara (Jurnal Ilmu Keolahragaan)*, 4(1), 1–11.
- Thompson, W. R. (2018). Worldwide survey of fitness trends for 2019. *Acsm's health and fitness journal*. <https://doi.org/10.1249/FIT.0000000000000438>
- Trilk, J. L., Singhal, A., Bigelman, K. A., & Cureton, K. J. (2011). Effect of sprint interval training on circulatory function during exercise in sedentary, overweight/obese women. *European Journal of Applied Physiology*, 111(8), 1591–1597. <https://doi.org/10.1007/s00421-010-1777-z>
- Viana, R. B., Naves, J. P. A., Coswig, V. S., De Lira, C. A. B., Steele, J., Fisher, J. P., & Gentil, P. (2019). Is interval training the magic bullet for fat loss? A systematic review and meta-analysis comparing moderate-intensity continuous training with high-intensity interval training (HIIT). *British Journal of Sports Medicine*. <https://doi.org/10.1136/bjsports-2018-099928>
- Wilmot, E. G., Edwardson, C. L., Achana, F. A., Davies, M. J., Gorely, T., Gray, L. J., Biddle, S. J. H. (2012). Sedentary time in adults and the association with diabetes, cardiovascular disease and death: Systematic review and meta-analysis. *Diabetologia*, 55(11), 2895–2905. <https://doi.org/10.1007/s00125-012-2677-z>
- World Health Organization. (2016). Global Health Observatory (GHO) data Overweight and obesity.
- World Health Organization. (2020). WHO Guidelines on physical activity, sedentary behaviour. World Health Organization.
- Zhang, H., Tong, T. K., Qiu, W., Wang, J., Nie, J., & He, Y. (2015). Effect of high-intensity interval training protocol on abdominal fat reduction in overweight Chinese women: A randomized controlled trial. *Kinesiology*, 47(1), 57–66.

High intensity interval training for improving maximum aerobic capacity in women with sedentary lifestyle a systematic review and meta analysis

ORIGINALITY REPORT

16%

SIMILARITY INDEX

11%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	www.slideshare.net Internet Source	1%
2	www.thieme-connect.com Internet Source	<1%
3	open.library.ubc.ca Internet Source	<1%
4	repository.nwu.ac.za Internet Source	<1%
5	M. Wewege, R. van den Berg, R. E. Ward, A. Keech. "The effects of high-intensity interval training vs. moderate-intensity continuous training on body composition in overweight and obese adults: a systematic review and meta-analysis", Obesity Reviews, 2017 Publication	<1%
6	Regina Márcia Ferreira Silva, Carolina Rodrigues Mendonça, Matias Noll. "Barriers to high school and university students' physical	<1%

activity: A systematic review protocol",
International Journal of Educational Research,
2021

Publication

7

Leandro Sant'Ana, Jeferson Vianna, Aline Ribeiro, Natália Reis et al. "Chronic Effects of Different Intensities of Interval Training on Hemodynamic, Autonomic and Cardiorespiratory Variables of Physically Active Elderly People", Research Square, 2020

Publication

<1 %

8

scholarworks.boisestate.edu

Internet Source

<1 %

9

Nicole Korman, Michael Armour, Justin Chapman, Simon Rosenbaum, Steve Kisely, Shuichi Suetani, Joseph Firth, Dan Siskind. "High Intensity Interval training (HIIT) for people with severe mental illness: A systematic review & meta-analysis of intervention studies– considering diverse approaches for mental and physical recovery", Psychiatry Research, 2020

Publication

<1 %

10

TOMOAKI MATSUO, KOUSAKU SAOTOME, SATOSHI SEINO, NOBUTAKE SHIMOJO et al. "Effects of a Low-Volume Aerobic-Type Interval Exercise on V'O2max and Cardiac

<1 %

Mass", Medicine & Science in Sports & Exercise, 2014

Publication

11

cronfa.swan.ac.uk

Internet Source

<1 %

12

shura.shu.ac.uk

Internet Source

<1 %

13

Jinlei Nie, Haifeng Zhang, Zhaowei Kong, Keith George, Jonathan P. Little, Tomas K. Tong, Feifei Li, Qingde Shi. "Impact of high-intensity interval training and moderate-intensity continuous training on resting and postexercise cardiac troponin T concentration", *Experimental Physiology*, 2018

Publication

<1 %

14

M. Hajishafiee, G. Askari, B. Iranj, R. Ghiasvand, N. Bellissimo, J. Totosy de Zepetnek, A. Salehi-Abargouei. "The Effect of n-3 Polyunsaturated Fatty Acid Supplementation on Androgen Status in Patients with Polycystic Ovary Syndrome: A Systematic Review and Meta-Analysis of Clinical Trials", *Hormone and Metabolic Research*, 2016

Publication

<1 %

15

Nana Wu, Shannon Bredin, Veronica Jamnik, Michael Koehle, Yanfei Guan, Erin Shellington, Yongfeng Li, Jun Li, Darren Warburton.

<1 %

"Association Between Physical Activity Level and Cardiovascular Risk Factors in Adolescents Living with Type 1 Diabetes: A Cross-sectional Study", Research Square, 2020

Publication

16

dspace.stir.ac.uk

Internet Source

<1 %

17

Vinícius Mallmann Schneider, Paula Frank, Sandra C. Fuchs, Rodrigo Ferrari. "Effects of recreational sports and combined training on blood pressure and glycosylated hemoglobin in middle-aged and older adults: a systematic review and meta-analysis", Cold Spring Harbor Laboratory, 2021

Publication

<1 %

18

www.tandfonline.com

Internet Source

<1 %

19

Lora-Pozo, Lucena-Anton, Salazar, Galán-Mercant, Moral-Munoz. "Anthropometric, Cardiopulmonary and Metabolic Benefits of the High-Intensity Interval Training Versus Moderate, Low-Intensity or Control for Type 2 Diabetes: Systematic Review and Meta-Analysis", International Journal of Environmental Research and Public Health, 2019

Publication

<1 %

20 Hongyan Zhao, Shan Hu, Jidong Rong. "Gene polymorphism associated with Angiotensinogen(M235T), Endothelial lipase (584C/T) and susceptibility to coronary artery disease: A meta-analysis", Cold Spring Harbor Laboratory, 2020
Publication

21 Todd A. Astorino, Ross M. Edmunds, Amy Clark, Rachael Gallant et al. "Change in maximal fat oxidation in response to different regimes of periodized high-intensity interval training (HIIT)", European Journal of Applied Physiology, 2017
Publication

22 erpub.chnpu.edu.ua:8080
Internet Source

23 Ming Yu Claudia Wong, Pak-Kwong Chung, Ka-Man Leung. "The Relationship Between Physical Activity and Self-Compassion: a Systematic Review and Meta-analysis", Mindfulness, 2020
Publication

24 mafiadoc.com
Internet Source

25 Yayi Zhao, Li Liu, Helen Yue - lai Chan. "Dementia care education interventions on healthcare providers' outcomes in the nursing

home setting: A systematic review", Research
in Nursing & Health, 2021

Publication

26

P. Bhati, D. Singla, M.E. Hussain. "Resistance training and modulation of cardiac autonomic control in animal models: a systematic review", Comparative Exercise Physiology, 2019

Publication

<1 %

27

www.antrocom.net

Internet Source

<1 %

28

Alicen A. Whitaker, Mohammed Alwatban, Andrea Freemyer, Jaime Perales-Puchalt, Sandra A. Billinger. "Effects of High Intensity Interval Exercise on Cerebrovascular Function: A Systematic Review", Cold Spring Harbor Laboratory, 2020

Publication

<1 %

29

Tomas K. Tong, Haifeng Zhang, Hongru Shi, Yang Liu, Jingwen Ai, Jinlei Nie, Zhaowei Kong. "Comparing Time Efficiency of Sprint vs. High-Intensity Interval Training in Reducing Abdominal Visceral Fat in Obese Young Women: A Randomized, Controlled Trial", Frontiers in Physiology, 2018

Publication

<1 %

30

ir.nrf.ac.za

Internet Source

<1 %

31	Hassane Zouhal, Abderraouf Ben Abderrahman, Arash Khodamoradi, Ayoub Saeidi et al. "Effects of physical training on anthropometrics, physical and physiological capacities in individuals with obesity: A systematic review", Obesity Reviews, 2020 Publication	<1 %
32	aaltodoc.aalto.fi Internet Source	<1 %
33	edepositireland.ie Internet Source	<1 %
34	recil.grupolusofona.pt Internet Source	<1 %
35	ugspace.ug.edu.gh Internet Source	<1 %
36	www.mdpi.com Internet Source	<1 %
37	www.nrcresearchpress.com Internet Source	<1 %
38	www.scielo.org.co Internet Source	<1 %
39	Anna Paula Azevedo de Carvalho, Carlos Adam Conte Junior. "Green strategies for active food packagings: A systematic review on active properties of graphene-based nanomaterials and biodegradable polymers",	<1 %

40 James A. Timmons, Steen Knudsen, Tuomo Rankinen, Lauren G. Koch et al. "Using molecular classification to predict gains in maximal aerobic capacity following endurance exercise training in humans", *Journal of Applied Physiology*, 2010

Publication

41 Malin Knutsen Glette, Siri Wiig. "The role of organizational factors in how efficiency-thoroughness trade-offs potentially affect clinical quality dimensions – a review of the literature", *International Journal of Health Governance*, 2021

Publication

42 Marine Dupuit, Florie Maillard, Bruno Pereira, Luis Marquezi Marcelo, Herbert Lancha Antonio, Nathalie Boisseau. "Effect of high intensity interval training on body composition in women before and after menopause: A meta - analysis", *Experimental Physiology*, 2020

Publication

43 Shubham Misra, Manabesh Nath, Vijay Hadda, Deepti Vibha. "Effect of various treatment modalities on the novel coronavirus (nCOV-2019) infection in humans:

a systematic review & meta-analysis", Cold Spring Harbor Laboratory, 2020

Publication

44

Xi Li, Tianyu Han, Xu Zou, Han Zhang, Wenpin Feng, Han Wang, Yulin Shen, Li Zhang, Guoliang Fang. "Long-term high-intensity interval training increases serum neurotrophic factors in elderly overweight and obese Chinese adults", European Journal of Applied Physiology, 2021

Publication

<1 %

45

Y. Türk, W. Theel, M. J. Kasteleyn, F. M. E. Franssen, P. S. Hiemstra, A. Rudolphus, C. Taube, G. J. Braunstahl. "High intensity training in obesity: a Meta-analysis", Obesity Science & Practice, 2017

Publication

<1 %

46

bdtd.ucb.br:8443

Internet Source

<1 %

47

booksc.org

Internet Source

<1 %

48

jssm.org

Internet Source

<1 %

49

www.fisiologiadelejercicio.com

Internet Source

<1 %

50

www.researchsquare.com

Internet Source

<1 %

51

Martin J. MacInnis, Martin J. Gibala.
"Physiological adaptations to interval training
and the role of exercise intensity", *The Journal
of Physiology*, 2017

Publication

<1 %

52

Miguel Angel Martínez-González, Constanza
López-Fontana, José Javier Varo, Almudena
Sánchez-Villegas, J Alfredo Martinez.
"Validation of the Spanish version of the
physical activity questionnaire used in the
Nurses' Health Study and the Health
Professionals' Follow-up Study", *Public Health
Nutrition*, 2007

Publication

<1 %

53

Zhi-Jian Wu, Zhu-Ying Wang, Hao-En Gao,
Xian-Feng Zhou, Fang-Hui Li. "Impact of high-
intensity interval training on cardiorespiratory
fitness, body composition, physical fitness,
and metabolic parameters in older adults: A
meta-analysis of randomized controlled
trials", *Experimental Gerontology*, 2021

Publication

<1 %

54

Amy S. Ha, Johan Y. Y. Ng, Rosa S. Wong.
"Examining the relationship between
children's health - related quality of life and
their perception of parental support toward
physical activity: A longitudinal study", *Applied
Psychology: Health and Well-Being*, 2021

Publication

<1 %

55

Elisa Marin-Couture, Louis Pérusse, Angelo Tremblay. "The fit-active profile to better reflect the benefits of a lifelong vigorous physical activity participation: mini-review of literature and population data.", *Applied Physiology, Nutrition, and Metabolism*, 2021

Publication

<1 %

56

Laura W.M. Vergoossen, Jacobus F.A. Jansen, Walter H. Backes, Miranda T. Schram.

"Cardiometabolic determinants of early and advanced brain alterations: Insights from conventional and novel MRI techniques", *Neuroscience & Biobehavioral Reviews*, 2020

Publication

<1 %

57

Tucker Stuart, Le Cai, Alex Burton, Philipp Gutruf. "Wireless and battery-free platforms for collection of biosignals", *Biosensors and Bioelectronics*, 2021

Publication

<1 %

58

Xinyue Zhang, Dongmei Xu, Guozhen Sun, Zhixin Jiang, Jinping Tian, Qijun Shan. "Effects of high - intensity interval training in patients with coronary artery disease after percutaneous coronary intervention: A systematic review and meta - analysis", *Nursing Open*, 2021

Publication

<1 %

59

Internet Source

<1 %

60

eprints.whiterose.ac.uk
Internet Source

<1 %

61

repository.upi.edu
Internet Source

<1 %

62

researchonline.ljmu.ac.uk
Internet Source

<1 %

Exclude quotes On
Exclude bibliography On

Exclude matches Off

High intensity interval training for improving maximum aerobic capacity in women with sedentary lifestyle a systematic review and meta analysis

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
