

# The Relationship Of Insulin-Like Growth Factor 1 With Bone Mass In Obese Female

*by* Adek Ardiansyah

---

**Submission date:** 19-Oct-2021 11:23AM (UTC+0800)

**Submission ID:** 1677732096

**File name:** Insulin\_Like\_Growth\_Factor\_1\_With\_Bone\_Mass\_In\_Obese\_Female.pdf (271.74K)

**Word count:** 3124

**Character count:** 16655

## The Relationship Of Insulin-Like Growth Factor 1 With Bone Mass In Obese Female

Adek Ardiansyah<sup>1</sup>, Lilik Herawati<sup>1,2</sup>, Damayanti Tinduh<sup>3\*</sup>

<sup>1</sup>Sport Health Science, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

<sup>2</sup>Department of Physiology, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

<sup>3</sup>Physical Medicine and Rehabilitation Department, Faculty of Medicine Universitas Airlangga/Dr. Soetomo General Hospital, Surabaya, Indonesia

\* [damayanti.tinduh@fk.unair.ac.id](mailto:damayanti.tinduh@fk.unair.ac.id)

### ABSTRACT

Obesity causes peripheral insulin resistance and hyperinsulinaemia, which causes a decrease in the secretion of Growth Hormone (GH) and Insulin-like Growth Factors 1 (IGF-1). Decreased GH and IGF-1 secretion decreases bone mineral density (BMD) and increases the risk of fractures. This study aimed to analyze the relationship of insulin-like growth factor 1 with bone mass in obese female. This study used a cross sectional study method using 30 obese female aged 19-23 years, body mass index (BMI) 25-35 kg/m<sup>2</sup>, normal blood pressure, normal resting heart rate (RHR), normal hemoglobin (Hb) and fasting blood glucose (FBG) < 100 mg/dL. Measurement of IGF-1 levels used the Enzym Link Immunosorbent Assay (ELISA) method. Measurement of bone mass using TANITA (Body Composition Analyzer DC3607601 (2) -1604 FA, TANITA Corporation of America, Inc., USA). The data analysis technique used the Pearson product-moment test with Statistical Package for Social Science (SPSS). The results showed that mean levels of IGF-1 (1.17±0.10) ng/mL and bone mass (2.49±0.06) kg ( $r = 0.712$ ,  $p \leq 0.001$ ). Our findings suggest that there was a positive correlation between IGF-1 levels and bone mass in obese female.

**Keywords :** IGF-1 levels, Bone Mass, Obesity

Received August 7, 2020; Revised August 30, 2020; Accepted September 23, 2020



STRADA Jurnal Ilmiah Kesehatan, its website, and the articles published there in are licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.

### BACKGROUND

Obesity is a condition that exceeds a person's relative body weight as a result of accumulation of nutrients, especially carbohydrates, fats and proteins. Obesity is currently a worldwide problem (Ayu and Handayani, 2016). Therefore, obesity is a dangerous condition. This is supported by the prevalence of obesity of ~ 20% (Pasquali *et al.*, 2020). According to the World Health Organization (WHO) in 2014, the prevalence of overweight at over 18 years of age reached 39%, where 15% were female. This prevalence increased 2 times greater than the prevalence of obesity in 1980. According to WHO (2016), more than 1.9 billion people aged over 18 years are overweight and 600 million of them are obese. The high prevalence makes obesity a problem in the world.

Obesity is a risk factor for various health problems that are usually experienced by adults such as diabetes mellitus, hypertension and high cholesterol. The causes of obesity are very complex in the sense that there are many factors that cause obesity, such as

environmental, genetic, psychological, health, drugs, development and physical activity. The cause of obesity is multi-factorial, thus the food intake factor is one of many factors. Influential food intake is one that contains fat and is high in calories. Some of the factors that cause obesity include a history of obesity in the elderly, lack of consumption of vegetables and fruit and smoking habits (Dewi, 2015).

Obesity experiences hyperinsulinemia condition due to peripheral insulin resistance which causes reduced growth hormone (GH) secretion. The resulting decrease in GH can inhibit the liver from secreting IGF-1 (Lewitt *et al.*, 2014). Growth hormone (GH) and its physiological mediator, insulin-like growth factor-1 (IGF-1), have a major role in bone growth. In general, the effect of GH promoting linear growth appears to depend on the production of IGF-1 and possibly other IGF peptides. IGF-1 acts on cartilage, stimulates cell proliferation and synthesis of Deoxyribo Nucleic Acid (DNA), Ribonucleic Acid (RNA), proteins and proteoglycans. IGF-1 is a major component of the organic skeleton matrix and is the most important differentiating factor for osteoblasts. If there is a decrease in the secretion of GH and IGF-1, it can reduce bone mineral density (BMD) and increase the risk of fracture (Ekbote *et al.*, 2015). According to research conducted by Chalista *et al.* (2017) that there is a relationship between IGF-1 and low bone mineral density in the elderly. However, the relationship between IGF-1 and bone mineral density in obesity is unclear. On the basis of the above background, the researchers wanted to find out whether there was a relationship between IGF-1 and bone mass in obese female.

## METHODS

This study used a cross sectional study method using 30 obese female aged 19-23 years, body mass index (BMI) 25-35 kg/m<sup>2</sup>, normal blood pressure, normal resting heart rate (RHR), normal hemoglobin (Hb) and fasting blood glucose (FBG) < 100 mg/dL. All of these research procedures have been approved by the Health Research Ethics Commission of the Faculty of Medicine, Airlangga University, Surabaya number 81/EC/KEPK/FKUA/2020.

Measurement of height using a stadiometer (SECA, Chino, CA). Measurement of body weight, BMI, percentage body fat (PBF), fat mass (FM), free fat mass (FFM) and bone mass (BM) using TANITA (Body Composition Analyzer DC3607601 (2) -1604 FA, TANITA Corporation of America, Inc., USA). Blood pressure was measured twice at a 5 min interval on the right arm using a standard mercury sphygmomanometer (Baumanometer; Baum, Copiague, NY, USA) and recorded as an average of the two readings. Measurement of FBG using ACCU-CHEK (ACCU-CHEK® Performance, Mannheim, Germany) with mg/dL concentration units. Measurement of Hb using Mission Hemoglobin (Mission® Hb Test Strips (Whole Blood), ACON Laboratories, Inc. 10125 Mesa Rim Road, San Diego, CA 92121, USA) with a unit of concentration of g/dL.

Collecting blood was from cubital veins with 4ml after a 12-hour overnight fasting. Blood was centrifuged for 15 minutes at 3000 rpm. Serum was separated and stored at -80 °C for analysis of IGF-1 levels the following day. Blood draws are conducted at 07.00 a.m. Measurement of serum IGF-1 levels using an Enzyme-Linked Immunosorbent Assay (ELISA) kit (Catalog No: E-EL-H0086; Lot No: 5KH3L8GCSC; Elabscience, China) with a standard curve range of 1.56-100 ng/mL and IGF-1 sensitivity level of 0.94 ng/mL.

Statistical analysis used packet statistical software for social science (SPSS) (Chicago, IL, USA). The normality test uses the Kolmogorov – Smirnov test. Correlation test using Pearson product-moment with a significant level ( $p \leq 0.05$ ). All data are presented as mean±SE.

## RESULTS

The basic profiles of the samples, including age, weight, height, body mass index, percentage body fat, fat mass, free fat mass, systolic blood pressure, diastolic blood pressure, resting heart rate, fasting blood glucose, hemoglobin, insuline-like growth factor-1, bone mass are displayed in Table 1 and Table 2.


**Table 1.** Characteristics of Research Subjects

Parameters	n	Mean	Std. Error
Age (years)	30	21.56	0.13
Weight (kg)	30	72.31	1.25
Height (m)	30	1.58	0.01
Body mass index (kg/m <sup>2</sup> )	30	28.98	0.29
Percentage body fat (%)	30	44.54	0.57
Fat mass (kg)	30	33.00	0.90
Free fat mass (kg)	30	40.77	0.61
Systolic blood pressure (mmHg)	30	112.67	0.82
Diastolic blood pressure (mmHg)	30	75.00	0.93
Resting heart rate (bpm)	30	79.37	1.81
Fasting Blood Glucose (mg/dL)	30	90.16	1.20
Hemoglobin (g/dL)	30	15.23	0.25

**Table 2.** Association between IGF-1 serum levels with bone

Parameter	n	Mean±SE	r	p-values
IGF-1 (ng/mL)	30	1.17±0.10		
Bone Mass (kg)	30	2.49±0.06	0.712**	0.000

\*\* Significant with  $p \leq 0.01$  by Pearson's product-moment



**Figure 1.** Correlations between IGF-1 levels and bone mass

Based on Table 2 and Figure 1, the results of the Pearson product-moment test show that there was a positive correlation between IGF-1 levels and bone mass ( $r = 0.712$ ,  $p \leq 0.001$ ).

## DISCUSSION

Obesity is a world health problem that must be considered because obesity increases the risk of various chronic diseases such as type 2 diabetes (T2D), hypertension, and cardiovascular disease (CVD) (Gadde *et al.*, 2018). In addition, the condition of obesity also occurs peripheral insulin resistance and hyperinsulinaemia, leading to decreased secretion of Growth Hormone (GH) (Lewitt *et al.*, 2014) and Insulin-like Growth Factors 1

(Rasmussen *et al.*, 1995). Decreased secretion of growth hormone (GH) and insulin-like growth factor-1 (IGF-1) reduces bone mineral density and increases the risk of fractures, such as fractures of the ankle, leg, humerus and vertebral column (Yakar *et al.*, 2018; Premaor *et al.*, 2014). The relationship of IGF-1 with bone mass or bone density has been studied in adults as well as elderly population. Some researchers have found a positive association of IGF-1 concentrations with bone density (Zhao *et al.*, 2008; Rucker *et al.*, 2004; Langlois *et al.*, 1998; Szulc *et al.*, 2004). The results showed that there was a positive correlation between IGF-1 levels and bone mass (Table 2). These results are in line with the results of research conducted by Chalista *et al.* (2017) that there is a relationship between IGF-1 levels and low bone mineral density (BMD). Likewise, the results of research conducted by Fornari *et al.* (2017) concluded that there was a correlation between IGF-1 levels and lumbar and femoral BMD in obese subjects.

Obesity is often associated with abnormalities in GH and IGF-1 secretion (Fornari *et al.*, 2017). GH secretion abnormalities can inhibit the liver from secreting IGF-1 (Lewitt *et al.*, 2014). GH and IGF-1 have a major role in bone growth and maintaining bone health (Fornari *et al.*, 2017). In general, the effects of GH promoting bone growth appear to be dependent on IGF-1 production. The results of research by Fornari *et al.* (2017) showed that increased IGF-1 levels were associated with increased BMD in obesity. IGF-1 acts on cartilage, stimulates cell proliferation and synthesis of Deoxyribo Nucleic Acid (DNA), Ribonucleic Acid (RNA), proteins and proteoglycans. IGF-1 is a major component of the organic skeleton matrix and is the most important differentiating factor for osteoblasts. If there is a decrease in the secretion of GH and IGF-1, it can reduce bone mineral density (BMD) and increase the risk of fracture (Ekbote *et al.*, 2015).

IGF-1 is also referred to as somatomedin C, mostly synthesized in the liver and regulated by GH which has a major role in cell growth and development (Guo *et al.*, 2020), energy metabolism (TaeHo Kim *et al.*, 2015). In addition, IGF-1 also plays a role in maintaining tissue homeostasis, anti-apoptosis, mitogenic, anti-inflammatory, antioxidant and metabolic properties, contributing to skeletal muscle plasticity, maintenance of muscle strength and muscle mass and increased bone mass (Guo *et al.*, 2020; Vitale *et al.*, 2019; Maggio *et al.*, 2013). IGF-1 can be synthesized by endocrine, paracrine and autocrine mechanisms (Stein *et al.*, 2018). IGF-1 synthesis is stimulated by nutrients and GH in the liver and other tissues (Lewitt *et al.*, 2014). GH, through IGF-1 stimulates the proliferation of epiphyseal cartilage so that more space is formed for bone formation and stimulates osteoblast activity (Sherwood, 2014). Osteoblast is a type of bone cell that produces organic matrix in bone. According to Sihombing *et al.* (2012) osteoblasts make, secrete and precipitate new bone matrix organic elements called osteoid. Osteoblasts contain alkaline phosphatase enzymes which indicate that these cells are not only associated with matrix creation, but also mineralization. When osteoblast activity deposits new bone on the external surface, other cells in the bone, osteoclasts (bone destroyers), dissolve bone tissue on the inner surface beside the marrow cavity (Sherwood, 2014). Osteoclasts are large multinuclear cells located along the surface of the bone where bone resorption, remodeling and repair occur (Sihombing *et al.*, 2012).

**CONCLUSIONS**

Based on the results of the study, it can be concluded that there is a positive correlation between IGF-1 levels and bone mass in obese female. Based on the results of the study, it is recommended that further research be carried out by adding other variables such as growth hormone (GH), insulin, glucose, Insulin-Like Growth Factor Binding Protein 3 (IGFBP3), serum glutamic pyruvic transaminase (SGPT) and serum glutamic oxaloacetic transaminase (SGOT).

**REFERENCES**

- Ayu, D.S. and Handayani, O.W.K. 2016. Diary Teratas (Terapi Anak Obesitas) dalam Perubahan Perilaku Gizi Siswa Sekolah Dasar. *Unnes Journal of Public Health*, 5(2): 167-175.
- Chalista, S.I., Chong, L.T., Wajda, M. et al. 2017. Association of Insulin-like Growth Factor 1, Bone Mass and Inflammation to Low-Energy Distal Radius Fractures and Fractures Healing in Elderly Women Attending Emergency Care. *Orthopedic Surgery*, 9(4): 380-385. <https://doi.org/10.1111/os.12358>.
- Dewi, M.C. 2015. Faktor-Faktor yang Menyebabkan Obesitas pada Anak. *Majority*, 4(8): 53-56.
- Ekbote, V.H., Khadilkar, V.V., Shilvant, D.S., et al. 2015. Relation of Insulin-like growth factor 1 and bone parameters in 7-15 years old apparently, healthy Indian children. *Indian Journal Endocrinol Metab*, 19(6): 770-774.
- Fornari, R., Marocco, C., Francomano, D., Fittipaldi, S., Lubrano, C., Bimonte, V.M., Donini, L.M., Nicolai, E., Aversa, A., Lenzi, A., Greco, E.A., and Migliacci, S. 2017. Insulin growth factor-1 correlates with higher bone mineral density and lower inflammation status in obese adult subjects. *Eating and Weight Disorders (EWD)*, 23(3): 375-381. <https://doi.org/10.1007/s40519-017-0362-4>.
- Gadde, K.M., Martin, C.K., Berthoud, H.R. and Heymsfield, S.B. (2018) 'Obesity: Pathophysiology and Management', *Journal of the American College of Cardiology*, 71(1): 69-84. <https://doi.org/10.1016/j.jacc.2017.11.011>.
- Guo, J., Xie, J., Zhou, B., Gaman, M-A., Kord-Varkaneh, H., Clark, C.C.T., Salehi-Sahlabadi, A., Li, Y, Han, X., Hao, Y. and Liang, Y. 2020. The influence of zinc supplementation on IGF-1 levels in humans: A systematic review and meta-analysis. *Journal of King Saud University - Science*, <https://doi.org/10.1016/j.jksus.2020.01.018>.
- Lewitt, M., Dent, M. and Hall, K. 2014. The Insulin-Like Growth Factor System in Obesity, Insulin Resistance and Type 2 Diabetes Mellitus. *Journal of Clinical Medicine*, 3(4): 1561-1574. <https://doi.org/10.3390/jcm3041561>.
- Langlois JA, Rosen CJ, Visser M, Hannan MT, Harris T, Wilson PW, et al. 1998. Association between insulin-like growth factor I and bone mineral density in older women and men: The Framingham Heart Study. *J Clin Endocrinol Metab*, 83(12): 4257-4262. <https://doi.org/10.1210/jcem.83.12.5308>.
- Maggio, M., De Vita, F., Lauretani, F., Butto, V., Bondi, G., Cattabiani, C. and Ceda, G.P. 2013. IGF-1, the cross road of the nutritional, inflammatory and hormonal pathways to frailty. *Nutrients* 5(10): 4184-4205. <https://doi.org/10.3390/nu5104184>.
- Pasquali, R., Casanueva, F., Haluzik, M. Huisteijn, L.V., Ledoux, S., Monteiro, M.P, Salvador, J., Santini, F., Toplak, H., Dekkers., O.M. 2020. European Society of

- Endocrinology Clinical Practice Guideline: Endocrine Work-Up in Obesity. *European Journal of Endocrinology*, 182(1): G1-G32. <https://doi.org/10.1530/EJE19-0893>.
- Premaor, M.O., Comim, F.V., and Compston, J.E. 2014. Obesity and fractures. *Arq Bras Endocrinol Metab*, 58(5): 470-477. <https://doi.org/10.1590/0004-2730000003274>.
- Rasmussen, M.H.; Hvidberg, A.; Juul, A.; Main, K.M.; Gotfredsen, A.; Skakkebae, N.E.; Hilsted, J. 1995. Massive weight loss restores 24-h growth hormone release profiles and serum insulin-like growth factor-I levels in obese subjects. *J. Clin. Endocrinol. Metab*, 80: 1407-1415.
- Rucker D, Ezzat S, Diamandi A, Khosravi J, Hanley DA. 2004. IGF-I and testosterone levels as predictors of bone mineral density in healthy, community-dwelling men. *Clin Endocrinol (Oxf)*, 60(4): 491-499. <https://doi.org/10.1111/j.1365-2265.2004.02006.x>.
- Sherwood, L. 2014. *Fisiologi Manusia: dari Sel ke Sistem, Edisi 8*. Jakarta: Penerbit Buku kedokteran EGC.
- Sihombing, I., Wangko, S., Kalangi, S. 2012. Peran Esterogen Pada Remodeling Tulang. *Jurnal Biomedik*, 4(3): 18-28.
- Stein, A.M., Silva, T.M.V., Coelho, F.G.M., Arantes, F.J., Costa, J.L.S., Teodoro, L. and Santos-Galduróz, R.F. 2018. Physical exercise, IGF-1 and cognition A systematic review of experimental studies in the elderly. *Dement Neuropsychol*, 12(2): 114-122. <http://dx.doi.org/10.1590/1980-57642018dn12-020003>.
- Szulc P, Joly-Pharaboz MO, Marchand F, Delmas PD. 2004. Insulin-like growth factor I is a determinant of hip bone mineral density in men less than 60 years of age: MINOS study. *Calcif Tissue Int*, 74(4): 322-329. <https://doi.org/10.1007/s00223-003-0090-9>.
- TaeHo Kim, Chang, J.S., Kim, H., Lee, K.H. and Kong, I.D. 2015. Intense Walking Exercise Affects Serum IGF-1 and IGFBP3. *Journal of Lifestyle Medicine*, 5(1): 21-25. <http://dx.doi.org/10.15280/jlm.2015.5.1.21>.
- Vitale, G., Pellegrino, G., Vollery, M. and Hofland, L.J., 2019. ROLE of IGF-1 System in the Modulation of Longevity: Controversies and New Insights From a Centenarians' Perspective. *Frontiers in Endocrinology*, 10(27): 1-11. <https://doi.org/10.3389/fendo.2019.00027>.
- World Health Organization (WHO). 2014. *Global Status Report on Noncommunicable Diseases 2014*. Switzerland: WHO Press.
- World Health Organization (WHO). *Obesity and overweight*. 2016. Switzerland: WHO Press Available at: <http://www.who.int/mediacentre/factsheets/fs311/en/>.
- Yakar, S., Werner, H. and Rosen, C.J. 2018. Insulin-like growth factors: actions on the skeleton. *Journal of Molecular Endocrinology*. 61(1): T115-T137. <https://doi.org/10.1530/JME-17-0298>.
- Zhao HY, Liu JM, Ning G, Zhao YJ, Chen Y, Sun LH, et al. 2008. Relationships between insulin-like growth factor-I (IGF-I) and OPG, RANKL, bone mineral density in healthy Chinese women. *Osteoporos Int*, 19(2): 221-226. <https://doi.org/10.1007/s00198-007-0440-y>.

# The Relationship Of Insulin-Like Growth Factor 1 With Bone Mass In Obese Female

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

16%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.ebmonline.org">www.ebmonline.org</a> Internet Source	1 %
2	<a href="http://www.chinesehepatology.net.cn">www.chinesehepatology.net.cn</a> Internet Source	1 %
3	<a href="http://www.researchsquare.com">www.researchsquare.com</a> Internet Source	1 %
4	<a href="http://www.ruvival.de">www.ruvival.de</a> Internet Source	1 %
5	Vassilios Perifanis, Timoleon Vyzantiadis, Konstantinos Tziomalos, Sofia Vakalopoulou et al. "Effect of zoledronic acid on markers of bone turnover and mineral density in osteoporotic patients with beta-thalassaemia", Annals of Hematology, 2006 Publication	1 %
6	<a href="http://journal.unnes.ac.id">journal.unnes.ac.id</a> Internet Source	1 %
7	<a href="http://www.pagepress.org">www.pagepress.org</a> Internet Source	1 %



8	<a href="http://biomedpharmajournal.org">biomedpharmajournal.org</a> Internet Source	<1 %
9	<a href="http://core.ac.uk">core.ac.uk</a> Internet Source	<1 %
10	<a href="http://www.scielo.br">www.scielo.br</a> Internet Source	<1 %
11	Alexander Törpel, Fabian Herold, Dennis Hamacher, Notger Müller, Lutz Schega. "Strengthening the Brain—Is Resistance Training with Blood Flow Restriction an Effective Strategy for Cognitive Improvement?", <i>Journal of Clinical Medicine</i> , 2018 Publication	<1 %
12	Cláudia V. Bender, Heraldo L. D. Silveira, Natália S. Santos, Juliano Cavagni et al. "Oral, dental, and craniofacial features in chronic acid sphingomyelinase deficiency", <i>American Journal of Medical Genetics Part A</i> , 2020 Publication	<1 %
13	<a href="http://eprints.nottingham.ac.uk">eprints.nottingham.ac.uk</a> Internet Source	<1 %
14	<a href="http://jonuns.com">jonuns.com</a> Internet Source	<1 %
15	<a href="http://scholar.sun.ac.za">scholar.sun.ac.za</a> Internet Source	<1 %

- |    |   |      |
|----|---|------|
| 16 | <a href="http://www.nationalchildrensstudy.gov">www.nationalchildrensstudy.gov</a><br>Internet Source   | <1 % |
| 17 | <a href="http://www.turkosteoporozdergisi.org">www.turkosteoporozdergisi.org</a><br>Internet Source   | <1 % |
| 18 | Research and Perspectives in Endocrine Interactions, 2004.<br>Publication   | <1 % |
| 19 | Stephen R. Pye, Bader Almusalam, Steven Boonen, Dirk Vanderschueren et al. "Influence of Insulin-Like Growth Factor Binding Protein (IGFBP)-1 and IGFBP-3 on Bone Health: Results from the European Male Ageing Study", Calcified Tissue International, 2011<br>Publication | <1 % |
| 20 | <a href="http://academic.oup.com">academic.oup.com</a><br>Internet Source   | <1 % |
| 21 | <a href="http://eprints.lib.hokudai.ac.jp">eprints.lib.hokudai.ac.jp</a><br>Internet Source   | <1 % |
| 22 | <a href="http://researchspace.ukzn.ac.za">researchspace.ukzn.ac.za</a><br>Internet Source   | <1 % |
| 23 | <a href="http://scholarsrepository.llu.edu">scholarsrepository.llu.edu</a><br>Internet Source   | <1 % |
| 24 | <a href="http://www.iabpi.com">www.iabpi.com</a><br>Internet Source   | <1 % |
| 25 | <a href="http://www.researcherslinks.com">www.researcherslinks.com</a><br>Internet Source   | <1 % |

<1 %

26

H.-Y. Zhao. "Relationships between insulin-like growth factor-I (IGF-I) and OPG, RANKL, bone mineral density in healthy Chinese women", Osteoporosis International, 02/2008

Publication

<1 %

27

Shreyasee Amin, B Lawrence Riggs, L Joseph Melton, Sara J Achenbach, Elizabeth J Atkinson, Sundeep Khosla. "High Serum IGFBP-2 Is Predictive of Increased Bone Turnover in Aging Men and Women", Journal of Bone and Mineral Research, 2007

Publication

<1 %

28

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<1 %

29

[alzres.biomedcentral.com](http://alzres.biomedcentral.com)

Internet Source

<1 %

30

[app.trdizin.gov.tr](http://app.trdizin.gov.tr)

Internet Source

<1 %

31

[coek.info](http://coek.info)

Internet Source

<1 %

32

[ir-library.ku.ac.ke](http://ir-library.ku.ac.ke)

Internet Source

<1 %

33

[jme.bioscientifica.com](http://jme.bioscientifica.com)

Internet Source

<1 %

34	<a href="https://journals.plos.org">journals.plos.org</a> Internet Source	<1 %
35	<a href="https://nation.africa">nation.africa</a> Internet Source	<1 %
36	<a href="https://nutritionandmetabolism.biomedcentral.com">nutritionandmetabolism.biomedcentral.com</a> Internet Source	<1 %
37	<a href="https://pure.coventry.ac.uk">pure.coventry.ac.uk</a> Internet Source	<1 %
38	<a href="https://www.eje-online.org">www.eje-online.org</a> Internet Source	<1 %
39	<a href="https://www.ijcp.elmerpress.com">www.ijcp.elmerpress.com</a> Internet Source	<1 %
40	<a href="https://www.kbb-forum.net">www.kbb-forum.net</a> Internet Source	<1 %
41	E.I. Barengolts. "Effects of progesterone on serum levels of IGF-1 and on femur IGF-1 mRNA in ovariectomized rats", <i>Journal of Bone and Mineral Research</i> , 10/1996 Publication	<1 %
42	Jaime Guevara-Aguirre, Carlos Torres, Gabriela Peña, María Palacios et al. "IGF-I deficiency and enhanced insulin sensitivity due to a mutated growth hormone receptor gene in humans", <i>Molecular and Cellular Endocrinology</i> , 2021 Publication	<1 %

43

Derek LeRoith, Jeff M.P. Holly, Briony E. Forbes. "Insulin-like growth factors: Ligands, binding proteins, and receptors", *Molecular Metabolism*, 2021

Publication

---

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On

# The Relationship Of Insulin-Like Growth Factor 1 With Bone Mass In Obese Female

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---