

Treadmill and Ergo cycle Exercises Increase Insulin-Like Growth Factor-1 Levels in Obese Female

by Adi Pranoto

Submission date: 16-Jun-2021 12:23PM (UTC+0800)

Submission ID: 1607299135

File name: Increase_Insulin-Like_Growth_Factor-1_Levels_in_Obese_Female.pdf (715.72K)

Word count: 7791

Character count: 34183

Treadmill and Ergo cycle Exercises Increase Insulin-Like Growth Factor-1 Levels in Obese Female

Adi Pranoto,¹ Zakaria Sedy Wardana,² Misbakhul Munir,^{2,3} Hayuris Kinandita,^{2,3} Desiana Merawati,⁴ Sugiharto,⁴ Purwo Sri Rejeki^{2,3*}

¹Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

²Sport Health Science, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

³Department of Physiology, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

⁴Sport Science Department, Faculty of Sport Science State University of Malang, Malang, Indonesia

Abstract: Obesity increases the risk incidence of metabolic disorders, including insulin resistance, causing an increase in blood glucose. Physical exercises, mediated by Insulin-Like Growth Factor-1 (IGF-1), contribute to improving insulin sensitivity and lowering blood glucose. IGF-1 increases insulin sensitivity and maintains glucose homeostasis by activating the Phosphoinositide 3-Kinase (PI3K) pathway and translocation of Glucose Transporter Protein-4 (GLUT-4). The purpose of the study was to analyze the effect of a moderate-intensity treadmill and ergo cycle exercises on the increasing IGF-1 levels in the obese female. This research was a true experimental study with the randomized pretest-posttest control group design. Subjects were 27 obese females aged 18-22. Subjects were randomly divided into three groups, that is CONT (n=9, group without intervention), ERGO-Exercise (n=9, Ergocycle Exercise), and TREAD-Exercise (n=9, Treadmill Exercise). Interventions for TREAD-Exercise and ERGO-Exercise were conducted continuously for 30 minutes, from 07.00-10.00 a.m. Blood samples were taken pre-exercise and 15 minutes post-exercise. IGF-1 levels were examined by the Enzyme-Linked Immunosorbent Assay (ELISA) method. Data were analyzed using ANOVA test and LSD post hoc test using Statistic Package for Social Science (SPSS). The research results showed that one session of a moderate-intensity treadmill and ergo cycle exercise for 30 minutes increases IGF-1 levels compared to the control. It is necessary to conduct advanced research to analyze chronic intervention (training) for increasing IGF-1 levels in obese females.

Keywords: IGF-1 levels, treadmill, ergo cycle, obese female

跑步机和人体工学循环运动可增加肥胖女性的胰岛素样生长因子-1 水平

摘要: 肥胖会增加包括胰岛素抵抗在内的代谢异常的风险发生率, 从而导致血糖升高。由胰岛素样生长因子-1 (IGF-1) 介导的体育锻炼有助于提高胰岛素敏感性和降低血糖。IGF-1 通过激活磷酸肌醇 3-激酶 (PI3K) 途径和葡萄糖转运蛋白 4 (GLUT-4) 的易位来提高胰岛素敏感性并维持葡萄糖稳态。这项研究的目的是分析中等强度的跑步机和人体工学循环运动对肥胖女性中 IGF-1 水平升高的影响。这项研究是一项真正的实验研究, 采用了随机的前测后测对照组设计。受试者是 27 名 18-22 岁的肥胖女性。将受试者随机分为三组, 即 CONT (n = 9, 无干预组), ERGO 锻炼 (n = 9, Ergocycle 锻炼) 和 TREAD-锻炼 (n = 9, 跑步机锻炼)。从上午 07.00-10.00 开始连续 30 分钟进行 TREAD 运动和 ERGO 运动的干预, 在运动前和运动后 15 分钟采

Received (date): 3 November 2020

About the authors: Adi Pranoto, Medicine, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia; Zakaria Sedy Wardana, Sport Health Science, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia; Misbakhul Munir, Hayuris Kinandita, Purwo Sri Rejeki, Sport Health Science, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia; Department of Physiology, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia; Desiana Merawati, Sugiharto, Sport Science Department, Faculty of Sport Science State University of Malang, Malang, Indonesia.

Corresponding author Purwo Sri Rejeki, purwo-s-r@fk.unair.ac.id; purwo_faal@yahoo.com

集血样。通过酶联免疫吸附测定 (ELISA) 方法检查 IGF-1 水平。使用 ANOVA 检验对数据进行分析, 并使用社会科学统计软件包 (SPSS) 对 LSD 事后检验进行分析。研究表明, 与对照组相比, 在中等强度的跑步机上进行一次持续 30 分钟的健身操, 可以增加 IGF-1 的水平。有必要进行高级研究以分析肥胖女性中增加 IGF-1 水平的慢性干预 (培训)。

关键词: IGF-1 水平, 跑步机, 人为周期, 肥胖女性

1. Introduction

Obesity is a metabolic disease that has reached epidemic proportions [1]. Obesity is considered an epidemic, and recently, it has been a pandemic [2] and syndemic [3]. That is because obesity prevalence keeps rising from year to year[4]. It is estimated that 1.9 billion people above 18 years old suffer from obesity, 650 million are obese, 11% male, and 15% female [5]. Globally, more than a third of adult people suffer from obesity [6]. It is estimated that in 2025, obesity prevalence will become 18% male and 21% female [7]. According to [8], obesity prevalence for people above 18 years old in Indonesia was 21.8%. This number was higher than the obesity prevalence in 2013 (14.8%) and 2007 (10.5%). The high increase in obesity prevalence becomes a serious problem that will threaten human resource quality [9] and the health problem of countries worldwide [10].

Obesity is a disease that has a high risk in the incidence of a very serious health problem that will threaten the world's public health [11, 12]. It is because obesity is one of many causes of disability and early death [6, 13], not only in adult people but also in children and teenagers around the world [1, 14]. Obesity triggers various health problems, such as increased incidence of cardiovascular diseases [15], type 2 diabetes mellitus, hypertension [12], stroke, some cancers, gallstone, osteoarthritis [16], respiratory disorders [1], muscle dysfunction [17], rheumatic diseases, metabolic syndrome [18] and metabolic disorders, including insulin resistance [19]. The cause of obesity is multifactorial. However, a general factor contributing to bodyweight increase is an imbalance between energy intake and energy expenditure [20–22]. Lifestyle modification is recommended for basic management in obesity treatment [12]. Lifestyle modification with an exercise-based non-pharmacological approach is the right strategy [23, 24]. Physical exercises are considered an effective and efficient method in preventing the increase in obesity prevalence [25, 26] because they increase energy expenditure and maintain glucose homeostasis mediated by Insulin-Like Growth Factor-1 (IGF-1) [10, 17].

IGF-1, also called somatomedin C, is mostly synthesized by the liver. It is regulated by Growth Hormone (GH), which plays the main role in cell growth, cell development, and energy homeostasis [17, 27]. Physical exercise was proven to increase IGF-1 levels in acute and chronic exercises [28–30]. It is because physical exercises stimulate the brain to activate the hypothalamus [31]. The hypothalamus secretes Growth Hormone Releasing Hormone (GhRH). This GHRH is transferred to the anterior pituitary [27, 32] and then stimulates GH secretion [29, 33]. GH modulates IGF-1 secretion from the liver to the blood circulation [17, 34]. IGF-1 in the circulation will bind to Insulin-Like Growth Factor Binding Protein 3 (IGFBP3) [17]. The binding of IGF-1 and IGFBP3 will affect energy balance, decrease body weight, and free fat mass [17]. It also affects muscle strength and muscle mass [27]. Previous research showed different results. As reported in [35], acute response from moderate-intensity exercise using ergo cycle for 20 minutes increased IGF-1 serum levels significantly in 20-29-year-old men. Nevertheless, exercise with an intensity above 80% of VO_{2max} decreased IGF-1 levels significantly [18]. It was concluded in [36] that acute exercise using an ergo cycle with moderate and high intensity for 15 minutes increased IGF-1 serum levels significantly in healthy women. Low-intensity aerobic training using an ergo cycle for 60 minutes decreased significantly by 9% of IGF-1 levels [37]. According to the previously mentioned research, it is still unclear whether exercise will decrease or increase IGF-1 levels. According to the background above, the objective of this research was to compare moderate-intensity treadmill and ergo cycle exercises to the increase of IGF-1 levels in obese female teenagers.

2. Materials and Methods

2.1. Experimental Design

This research was a true experimental study with the randomized pretest-post-test control group design. Subjects were 27 females, 18-22 year old, body mass index (BMI) 25.5-32.5 kg/m², percentage body fat (PBF) > 30%, fasting blood glucose (FBG) < 100 mg/dL,

normal hemoglobin (Hb), normal blood pressure, and normal resting heart rate. Subjects were randomly divided into three groups, that is CONT (n=9, group without intervention), ERGO-Exercise (n=9, Ergocycle Exercise), and TREAD-Exercise (n=9, Treadmill Exercise). All subjects received information verbally or in writing about the research. Subjects filled out and signed informed consent before participating in the study. All procedures in this research were approved by the Health Research Ethics Committee, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia (309/EC/KEPK/FKUA/2019).

2.2. Exercise Protocol

The exercise program was applied and supervised by professionals at the Fitness Center of the Malang City Health Department, East Java, Indonesia. Treadmill

exercise and ergo cycle exercise were conducted with moderate-intensity 60-70% of HR_{max} for 40 minutes. It was consist of 5 minutes warming up (50-60% HR_{max}), 30 minutes core exercise (60-70% HR_{max}), and 5 minutes cooling down (50-60 HR_{max}) [38, 39]. Treadmill exercises and ergo cycle exercises were done with the continuous method at 08.00 a.m. [40, 41]. Instruments used for this intervention were treadmill (Pulsar 4.0 HP Cosmos Sports & Medical, Nussdorf-Traunstein, Germany) and ergo cycle (Monark 828 E, Version 1010 Art. No: 7950-296, Vansbro, Sweden). Monitoring heart rate during exercise using a polar heart rate monitor (Polar H10 Heart Rate Sensor, Inc., USA). This study was carried at the Physical Freshness Laboratory with a room temperature of $26\pm 1^{\circ}C$ and humidity level in the room 50-70 %. Figure 1 shows the treadmill and ergo cycle exercise program in clearer detail.




Fig. 1 Treadmill and ergo cycle exercise program

2.3. Anthropometric Measurements

The body height of the subject was measured using a stadiometer (SECA, Chino, CA). Anthropometric measurements included body weight, BMI, PBF, fat mass (FM), free fat mass (FFM), muscle mass (MM), bone mass (BM). These measurements were conducted using TANITA (Body Composition Analyzer DC3607601(2)-1604 FA, TANITA Corporation of America, Inc).

2.4. Physiological Condition

Blood pressure was measured using an automated device (OMRON Model HEM-7130 L, Omron Co., JAPAN) at the non-dominant arm three times consecutively with a 1-2 minute interval between two measurements while the participants were in a seated position. Measurement of resting heart rate (RHR) using a Pulse Oximeter (PO 30 Pulse Oximeter, Beurer North America LP, 900 N Federal Highway, Suite 300, Hallandale Beach, FL 33009).

2.5. Blood Samples

Four milliliters of blood samples were taken from the cubital vein after 12-hour overnight fasting (pre-exercise) [36, 42]. At the time of drawing blood, the subject was in a sleeping position. Pre-exercise blood samples were taken 30 minutes after the intervention [42], whereas post-exercise blood samples were taken 15 minutes after the intervention [36]. Blood samples were centrifuged for 10 minutes at 3000 rpm [17]. Serum was collected and saved at $-80^{\circ}C$ to analyze IGF-1 levels on the next day [10].

2.6. IGF-1 Serum Levels Assessment

IGF-1 levels were examined using an Enzyme-Linked Immunosorbent Assay (ELISA) kit (Catalog No. E-EL-H0086; Elabscience, Inc., China, 2019) with a standard curve range 1.56–100 ng/mL and sensitivity up to 0.94 ng/mL. FBG was measured in mg/dL using ACCU-

CHEK (ACCU-CHEK® Performa, Mannheim, Germany). Hb was measured in g/dL using Easy Touch (Easy Touch GCHb, Taiwan).

2.7. Statistical Analysis

Data were analyzed by Statistic Package for Social Science (SPSS) Statistics for Windows, version 16 (SPSS Inc., Chicago, IL, USA). Normality was tested using the Shapiro-Wilk test, whereas Homogeneity was measured using the Levene test. Statistical difference was tested using the Paired Sample T-Test, ANOVA, and Least Significant Difference (LSD) post hoc test. All data were

presented in mean±SD. All statistical analysis was conducted using the level of significance ($P<0.05$).

3. Research Results

The basic profiles of the subjects, including age, body weight, body height, body mass index, percentage body fat, fat mass, free fat mass, muscle mass, bone mass, resting heart rate, systolic blood pressure, diastolic blood pressure, fasting blood glucose, and hemoglobin are displayed in Table 1.

Table 1 Characteristics of the subjects

Variable	CONT (n = 9)	ERGO-Exercise (n = 9)	TREAD-Exercise (n = 9)	ANOVA P-Value
Age (years)	20.78±0.97	20.33±1.00	20.89±0.78	0.415
Body Weight (kg)	73.49±7.96	68.21±7.88	71.15±7.43	0.367
Body Height (m)	1.60±0.05	1.55±0.05	1.57±0.04	0.173
Body mass index (kg/m ²)	28.49±1.57	28.244±2.39	28.61±1.67	0.918
Body fat percentage (%)	42.92±2.43	43.48±4.01	43.72±2.67	0.857
Fat mass (kg)	34.19±6.44	30.69±5.25	33.31±4.50	0.381
Free fat mass (kg)	40.67±3.78	41.08±4.40	40.99±3.20	0.972
Muscle mass (kg)	38.28±3.47	38.64±4.02	38.57±2.93	0.973
Bone mass (kg)	2.39±0.314	2.43±0.38	2.42±0.27	0.956
Resting heart rate (bpm)	75.33±7.81	74.67±6.93	78.78±7.99	0.479
Systolic blood pressure (mmHg)	114.44±5.27	115.55±5.27	112.22±4.41	0.370
Diastolic blood pressure (mmHg)	74.44±5.27	76.67±5.00	75.55±5.27	0.666
Fasting blood glucose (mg/dL)	90.22±7.90	90.67±4.21	88.22±8.23	0.737
Hemoglobin (g/dL)	15.03±0.93	14.42±1.32	14.89±1.46	0.567

Note: One way-ANOVA. Data are presented as mean±SD. CONT: Control group; ERGO-Exercise: Ergocycle exercise group; TREAD-Exercise: Treadmill exercise group.

Data in Table 1 show the means of the subjects' characteristics in each group. The ANOVA test results showed no difference in the subjects' characteristics on all variables of each group ($P>0.05$). Analytical results on pre-exercise and post-exercise IGF-1 levels are presented in Figure 2.

The level of both forms of IGF-1 was assessed pre-exercise and post-exercise. The results of the Paired Sample T-Test in the CONT group showed that there was no significant difference in the mean IGF-1 levels between pre-exercise and post-exercise (9.75±1.74 vs. 9.45±2.44 ng/mL, (p-value=0.710)) (Figure 1). However, the ERGO-Exercise group showed a significant difference in mean IGF-1 levels between pre-exercise and post-exercise (9.79±3.00 vs. 14.76±6.71 ng/mL, (p-value=0.045)) (Figure 2). Likewise, the TREAD-Exercise group showed a significant difference in mean IGF-1 levels between pre-exercise and post-exercise (9.75±2.63 vs. 14.98±2.84 ng/mL, (p-value=0.000))

(Figure 1). The results of the analysis of IGF-1 levels pre-exercise, post-exercise, and delta (Δ) post – pre in all groups can be seen in Figure 3.




Fig. 2 IGF-1 levels pre-exercise vs. post-exercise. CONT: Control group; ERGO-Exercise: Ergocycle exercise group; TREAD-Exercise: Treadmill exercise group. Data are presented as mean±SD. p-Value was obtained using Paired Sample T-Test to compare post-exercise and pre-exercise IGF-1 levels
*Significant vs pre-exercise ($p<0.05$)




Fig. 3 IGF-1 levels pre-exercise, post-exercise, and delta (Δ) post – pre. CONT: Control group; ERGO-Exercise: Ergocycle exercise group; TREAD-Exercise: Treadmill exercise group. Data are presented as mean \pm SD. p-Values were obtained using one way-ANOVA, followed by an LSD post hoc test compare pre-exercise, post-exercise, delta (Δ) post – pre IGF-1 level in the group *Significant vs control group (CONT) ($p < 0.05$).

Data in Figure 3 show no significant difference in pre-exercise IGF-1 levels in all groups ($P > 0.05$). However, post-exercise and delta (Δ) (post – pre) IGF-1 levels differ significantly in all groups ($P < 0.05$). LSD post hoc test showed that there was a significant difference between ERGO-Exercise and CONT IGF-1 levels ($p = 0.018$), as well as TREAD-Exercise and CONT ($p = 0.014$); whereas, ERGO-Exercise and TREAD-Exercise IGF-1 levels showed no significant difference ($p = 0.919$). Likewise, delta (Δ) (post – pre) showed a significant difference in mean IGF-1 levels between ERGO-Exercise and CONT ($p = 0.013$), TREAD-Exercise with CONT ($p = 0.010$). In contrast, ERGO-Exercise with TREAD-Exercise did not show a significant difference ($P > 0.05$).

4. Discussion

This research aimed to analyze the difference between treadmill and ergo cycle exercise to increase IGF-1 levels in obese females. The results showed no significant difference in pre-exercise IGF-1 levels among all groups ($p = 0.999$). Therefore, the three groups in this research had the same characteristics before the treadmill and ergo cycle intervention.

Results from this research showed that there was a significant difference in post-exercise IGF-1 levels between TREAD-Exercise and CONT ($p = 0.014$). Likewise, delta (Δ) (post – pre) showed a significant difference in mean IGF-1 levels between TREAD-Exercise with CONT ($p = 0.010$). It was similar to the result from the research conducted in [28] concluded that

treadmill exercise with intensity 60% of VO_{2max} increased IGF-1 mRNA levels significantly. As reported in [43], moderate-intensity exercise increased IGF-1 levels. The IGF-1 levels possibly increased in TREAD-Exercise because of the effect of exercise. Physical exercises stimulate the brain to activate the hypothalamus [31]. The hypothalamus secretes GhRH and, then, GhRH stimulates GH secretion [32, 33]. GH modulates IGF-1 secretion from the liver to the blood circulation [34].

There was a significant difference in post-exercise IGF-1 levels between ERGO-Exercise with CONT ($p = 0.018$). Likewise, delta (Δ) (post – pre) showed a significant difference in mean IGF-1 levels between ERGO-Exercise and CONT ($p = 0.013$). It was similar to the research results reported in [36], revealing that acute ergo cycle exercise with moderate intensity increased IGF-1 serum levels significantly. As concluded in [35], the acute response from moderate-intensity exercise increased IGF-1 serum levels significantly. The increase of IGF-1 in ERGO-Exercise was possible because of increased energy needs for muscle contraction for exercise. When exercising, there is an increase in energy needs and glucose uptake for muscle contraction, so that energy stored in muscle decreases. This decrease, in turn, increases IGF-1 release to the blood circulation. The increase of IGF-1 secretion to the circulation activates Phosphoinositide 3-Kinase (PI3K) pathway and Glucose Transporter Protein-4 (GLUT-4) translocation for maintaining glucose homeostasis and energy balance during exercise [44].

IGF-1, or somatomedin C, is mostly synthesized by the liver and regulated by GH, which plays the main role in cell growth, cell development, and energy metabolism [17, 27]. IGF-1 plays a role in body composition related to the alteration of lean body mass and fat mass [28]. IGF-1 also plays a role in increasing insulin sensitivity and maintaining glucose homeostasis [44]. Besides that, IGF-1 is significant for tissue homeostasis, anti-apoptotic, mitogenic, anti-inflammation, antioxidant, metabolism, skeletal muscle plasticity, muscle strength and mass maintenance, and neural and cardiovascular protection [27, 45, 46]. IGF-1 can be synthesized through endocrine, paracrine, and autocrine mechanism [29]. Acute and chronic exercise both can increase IGF-1 levels in plasma and serum [28, 29]. Exercises induce IGF-1 secretion through activation of the hypothalamus [31]. The hypothalamus secretes GhRH, which then should be transferred to the anterior pituitary through hypothalamic-hypophyseal portal vessels [32]. GhRH stimulates GH secretion [33]. GH modulates IGF-1

secretion from the liver to the blood circulation [34]. IGF-1 in the circulation will bind to IGFBP3 [17]. The binding of IGF-1 and IGFBP3 will affect energy balance, decrease body weight, free fat mass [17], and the maintenance of muscle strength and mass [27].

There was no significant difference in post-exercise IGF-1 levels between TREAD-Exercise with ERGO-Exercise ($p=0.999$). Likewise, delta (Δ) (post – pre) there was no significant difference in IGF-1 levels between TREAD-Exercise with ERGO-Exercise ($p=0.897$). The limitations of applying this research results cannot be applied to the elderly and people who have a high risk, such as hypertension, coronary heart disease (CHD), stroke, respiratory disorders, etc. Other than that, the limitation of this study is that we only analyzed one dependent variable, namely the IGF-1 level. The intervention given in this study was only acute exercise. It is necessary to conduct advanced research to analyzed chronic intervention (training) in obese female teenagers with the addition of several dependent variables, such as growth hormone (GH), Growth Hormone Releasing Hormone (GhRH), Insulin-Like Growth Factor Binding Protein 3 (IGFBP3), blood glucose, insulin, and homeostasis model assessment of insulin resistance (HOMA-IR). Hopefully, the results of these future research will benefit from perfecting optimal exercise programs for obesity treatment in the future.

6. Conclusion

This research showed that one session of the moderate-intensity treadmill and ergo cycle exercises for 30 minutes increases IGF-1 levels compared to the control.

98

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Acknowledgment

No financial support was provided for this project. We would like to express our gratitude to the Faculty of the Sports Science State University of Malang that has provided facilities in the screening process of a prospective research subject. We are thankful to the Fitness Center of the Malang City Health Department that has provided facilities. Also, we greatly appreciate and wish to thank Palang Merah Indonesia (PMI) Blood Transfusion Unit (UTD) Malang that has assisted the blood sampling and blood centrifuge processes. Our gratitude includes but is not limited to all the parties of to Physiology Laboratory Faculty of Medicine Universitas Brawijaya Malang, who have helped the analysis process

of IGF-1 level and all volunteers who have participated in this study.

References

- [1] AKTAR N., QURESHI N.K., and FERDOUS H.S. Obesity: A Review of Pathogenesis and Management Strategies in Adult. *Delta Medical College Journal*, 2017, 5(1): 35-48. <https://doi.org/10.3329/dmcj.v5i1.31436>.
- [2] WINDARTI N., HLAING S.W., and KAKINAKA M. Obesity Kuznets curve: international evidence. *Public Health*, 2019, 169: 26-35. <https://doi.org/10.1016/j.puhe.2019.01.004>.
- [3] SWINBURN B.A., KRAAK V.I., ALLENDER S., et al. The Global Syndemic of Obesity, Undernutrition, and Climate Change: The Lancet Commission report. *The Lancet*, 2019, 393(10173): 791-846. [https://doi.org/10.1016/S0140-6736\(18\)32822-8](https://doi.org/10.1016/S0140-6736(18)32822-8).
- [4] CHOOI Y.C., DING C., and MAGKOS F. The epidemiology of obesity. *Metabolism: Clinical and Experimental*, 2019, 92: 6-10. <https://doi.org/10.1016/j.metabol.2018.09.005>.
- [5] WORLD HEALTH ORGANIZATION (WHO). *Obesity and overweight*. 2016. <http://www.who.int/mediacentre/factsheets/fs311/en/>.
- [6] AL KIBRIA G.M. Prevalence and factors affecting underweight, overweight and obesity using Asian and World Health Organization cutoffs among adults in Nepal: Analysis of the Demographic and Health Survey 2016. *Obesity Research & Clinical Practice*, 2019, 13(2): 129-136. <https://doi.org/10.1016/j.orcp.2019.01.006>.
- [7] NCD Risk Factor Collaboration. Trends in adult body-mass index in 200 countries from 1975 to 2014: A pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *The Lancet*, 2016, 387(10026): 1377-1396. [https://doi.org/10.1016/S0140-6736\(16\)30054-X](https://doi.org/10.1016/S0140-6736(16)30054-X).
- [8] RISKESDAS. Laporan Nasional Riset Kesehatan Dasar. Jakarta: Kemenkes RI. 2018. <http://www.kesmas.kemkes.go.id>.
- [9] AKTER, S., RAHMAN, M.M., ABE, S.K., and SULTANA P. Prevalence of diabetes and prediabetes and their risk factors among Bangladeshi adults: a nationwide survey. *Bulletin of the World Health Organization*, 2014, 92(3): 204-213. <https://doi.org/10.2471/BLT.13.128371>.
- [10] TSUCHIYA Y., ANDO D., GOTO K., et al. High-intensity exercise causes greater irisin response compared with low-intensity exercise under similar energy consumption. *The Tohoku Journal of Experimental Medicine*, 2014, 233(2): 135-140. <https://doi.org/10.1620/tjem.233.135>.
- [11] WORLD HEALTH ORGANIZATION (WHO). *Noncommunicable Diseases Country Profiles 2018*, World Health Organization. 2018. <https://doi.org/10.1620/tjem.233.135>.
- [12] GADDE K.M., MARTIN C.K., BERTHOUD H.R., and HEYMSFIELD S.B. Obesity: Pathophysiology and Management. *Journal of the American College of*

- Cardiology*, 2018, 71(1): 69-84. <https://doi.org/10.1016/j.jacc.2017.11.011>.
- [13] ROSELLA L.C., KORNAS K., HUANG A., et al. Population risk and burden of health behavioral-related all-cause, premature and amenable deaths in Ontario, Canada: Canadian Community Health Survey-linked mortality files. *Annals of Epidemiology*, 2019, 32: 49-57. <https://doi.org/10.1016/j.annepidem.2019.01.009>.
- [14] KHAN S., and SHAH A.J. Causes and Complication of Obesity among the Children International Journal of Nutrition and Health Sciences Causes and Complication of Obesity among the Children. *Acta Scientific Nutritional Health*, 2017, 1(1): 47-50.
- [15] MARIE NG., FLEMING T., ROBINSON M., et al. Global regional and national prevalence of overweight and obesity in children and adults 1980-2013: A systematic analysis. *Lancet*, 2014, 384(9945): 766-781. [https://doi.org/10.1016/S0140-6736\(14\)60460-8](https://doi.org/10.1016/S0140-6736(14)60460-8). Global.
- [16] BALES C.W., and BUHR G. Is Obesity Bad for Older Persons? A Systematic Review of the Pros and Cons of Weight Reduction in Later Life. *Journal of the American Medical Directors Association*, 2008, 9(5): 302-312. <https://doi.org/10.1016/j.jamda.2008.01.006>.
- [17] TAE HO KIM, CHANG, J.S., KIM, H., et al. Intense Walking Exercise Affects Serum IGF-1 and IGF1BP3. *Journal of Lifestyle Medicine*, 2015, 5(1): 21-25. <http://dx.doi.org/10.15280/jlm.2015.5.1.21>.
- [18] MAJORCZYK M., and SMOLAĞ D. Effect of Physical Activity on IGF-1 and IGF1BP Levels in The Context of Civilization Diseases Prevention. *Rocz Panstw Zakl Hig*, 2016, 67(2): 105-111.
- [19] UMOH A.V., and ABAH G.M. Haemoglobin Genotypes: A Prevalence Study and Implications for Reproductive Health in Uyo, Nigeria. *Nigerian Journal of Medicine*, 2010, 19(1): 36-41.
- [20] FAN W., and EVANS R.M. Exercise Mimetics: Impact on Health and Performance. *Cell Metabolism*, 2017, 25(2): 242-247. <https://doi.org/10.1016/j.cmet.2016.10.022>.
- [21] RAMBHOJAN C., BOUAZIZ-AMAR E., LARIFLA L., et al. Ghrelin, adipokines, metabolic factors in relation with weight status in school-children and results of a 1-year lifestyle intervention program. *Nutrition & Metabolism*, 2015, 12(43): 1-10. <https://doi.org/10.1186/s12986-015-0039-9>.
- [22] NORHEIM F., LANGLEITE T.M., HJORTH M., et al. The effects of acute and chronic exercise on PGC-1 α , irisin and browning of subcutaneous adipose tissue in humans. *FEBS Journal*, 2014, 281(3): 739-749. <https://doi.org/10.1111/febs.12619>.
- [23] KOLAWOLE A.K., KEVIN N.U., OLUWOLE I., and ADEMOLA S. The Association of Socio-Demographic Factors with Overweight/Obesity among Students (Ages 18-35 Years) in Cavendish University, Uganda. *Epidemiology (Sunnyvale)*, 2017, 7(328): 1-6. <https://doi.org/10.4172/2161-1165.1000328>.
- [24] MURAWSKA-CIALOWICZ E., WOJNA J., and ZUWALA-JAGIELLO J. Crossfit training changes brain-derived neurotrophic factor and irisin levels at rest, after wingate and progressive tests and improves aerobic capacity and body composition of young physically active men and women. *Journal of Physiology and Pharmacology*, 2015, 66(6): 811-821.
- [25] HUH J.Y., MOUGIOS V., KABASAKALIS A., et al. Exercise-induced irisin secretion is independent of age or fitness level and increased irisin may directly modulate muscle metabolism through AMPK activation. *Journal of Clinical Endocrinology and Metabolism*, 2014, 99(11): E2154-E2161. <https://doi.org/10.1210/jc.2014-1437>.
- [26] BOSTRÖM P., WU J., JEDRYCHOWSKI M.P., et al. A PGC1 α -dependent myokine that drives browning of white fat and thermogenesis. *Nature*, 2012, 481(7382): 463-468. <https://doi.org/10.1038/nature10777>. A.
- [27] GUO J., XIE J., ZHOU B., et al. The influence of zinc supplementation on IGF-1 levels in humans: A systematic review and meta-analysis. *Journal of King Saud University-Science*, 2020, 32(3): 1824-1830. <https://doi.org/10.1016/j.jksus.2020.01.018>.
- [28] BERRY N.T., HUBAL M., and WIDEMAN L. The effects of an acute exercise bout on GH and IGF-1 in prediabetic and healthy African Americans: A pilot study investigating gene expression. *PLoS ONE*, 2018, 13(1): e0191331. <https://doi.org/10.1371/journal.pone.0191331>.
- [29] STEIN A.M., SILVA T.M.V., COELHO F.G.M., et al. Physical exercise, IGF-1 and cognition A systematic review of experimental studies in the elderly. *Dementia & Neuropsychologia*, 2018, 12(2): 114-122. <http://dx.doi.org/10.1590/1980-57642018dn12-020003>.
- [30] BERG U., and BANG P. Exercise and Circulating Insulin-Like Growth Factor I. *Hormone Research*, 2004, 62(1): 50-58. <https://doi.org/10.1159/000080759>.
- [31] WOOD C.J., CLOW A., HUCKLEBRIDGE F., et al. Physical fitness and prior physical activity are both associated with less cortisol secretion during psychosocial stress. *An International Journal Anxiety, Stress & Coping*, 2017, 31(2): 135-145. <https://doi.org/10.1080/10615806.2017.1390083>.
- [32] MARKS D.B., MARKS A.D., and MARKS C.M. *Biokimia Kedokteran Dasar: Sebuah Pendekatan Klinis*. Jakarta: EGC. 2014.
- [33] LARON Z. Insulin-like growth factor 1 (IGF-1): a growth hormone. *Journal of Clinical Pathology*, 2001, 54(5): 311-316. <https://doi.org/10.1136/mp.54.5.311>.
- [34] CLEMMONS D.R. The relative roles of growth hormone and IGF-1 in controlling insulin sensitivity. *The Journal of Clinical Investigation*, 2004, 113(1): 96-105.
- [35] TSAI C.-L., WANG C.-H., PAN C.-Y., et al. Executive function and endocrinological responses to acute resistance exercise. *Frontiers in Behavioral Neuroscience*, 2014, 8(262): 1-12. <https://doi.org/10.3389/fnbeh.2014.00262>.

- [36] MANNERKORPI K., LANDIN-WILHELMSSEN K., LARSSON A., et al. Acute effects of physical exercise on the serum insulin-like growth factor system in women with fibromyalgia. *BMC Musculoskeletal Disorders*, 2017, 18(37): 1-8. <https://doi.org/10.1186/s12891-017-1402-y>.03
- [37] NISHIDA Y., MATSUBARA T., TOBINA T., et al. Effect of Low-Intensity Aerobic Exercise on Insulin-Like Growth Factor-I and Insulin-Like Growth Factor-Binding Proteins in Healthy Men. *International Journal of Endocrinology*, 2010, 452820: 8 pages. <https://doi.org/10.1155/2010/452820>. 94
- [38] DIAS K.A., INGUL C.B., TJONNA A.E., et al. Effect of High-Intensity Interval Training on Fitness, Fat Mass and Cardiometabolic Biomarkers in Children with Obesity: A Randomised Controlled Trial. *Sports Medicine*, 2017, 48(3): 733-746. <https://doi.org/10.1007/s40279-017-0777-0>.
- [39] WEWEGE M., VAN DEN BERG R., WARD R.E., and KEECH A. The effects of high-intensity interval training vs. moderate-intensity continuous training on body composition in overweight and obese adults: a systematic review and meta-analysis. *Obesity Reviews*, 2017, 18(6): 635-646. <https://doi.org/10.1111/obr.12532>.
- [40] NYGAARD H., NYGAARD H., SLETTALOKKEN G., et al. Irisin in blood increases transiently after single sessions of intense endurance exercise and heavy strength training. *PLoS ONE*, 2015, 10(3): 1-12. <https://doi.org/10.1371/journal.pone.0121367>.
- [41] KRAEMER R.R., SHOCKETT P., WEBB N.D., et al. A transient elevated irisin blood concentration in response to prolonged, moderate aerobic exercise in young men and women. *Hormone and Metabolic Research*, 2014, 46(2): 150-154. <https://doi.org/10.1055/s-0033-1355381>.
- [42] DASKALOPOULOU S.S., COOKE A.B., GOMEZ Y.H., et al. Plasma irisin levels progressively increase in response to increasing exercise workloads in young, healthy, active subjects. *European Journal of Endocrinology*, 2014, 171(3): 343-352. <https://doi.org/10.1530/EJE-14-0204>.
- [43] YOON S.J., LEE M.J., LEE H.M., and LEE J.S. Effect of low-intensity resistance training with heat stress on the HSP72, anabolic hormones, muscle size, and strength in elderly women. *Aging Clinical and Experimental Research*, 2017, 29: 977-984.
- [44] HAYWOOD N.J., SLATER T.A., MATTHEWS C.J., and WHEATCROFT S.B. The insulin like growth factor and binding protein family: Novel therapeutic targets in obesity & diabetes. *Molecular Metabolism*, 2019, 19: 86-96. <https://doi.org/10.1016/j.molmet.2018.10.008>.
- [45] VITALE G., PELLEGRINO G., VOLLERY M., and HOFLAND L.J. Role of IGF-1 System in the Modulation of Longevity: Controversies and New Insights From a Centenarians' Perspective. *Frontiers in Endocrinology*, 2019, 10(27): 1-11. <https://doi.org/10.3389/fendo.2019.00027>.
- [46] MAGGIO M., DE VITA F., LAURETANI F., et al. IGF-1, the cross road of the nutritional, inflammatory and hormonal pathways to frailty. *Nutrients*, 2013, 5(10): 4184-4205. <https://doi.org/10.3390/nu5104184>.

参考文献:

- [1] AKTAR N., QURESHI N.K.和 FERDOUS H.S.肥胖：成人发病机理和治疗策略的回顾。三角洲医学院学报，2017，5（1）：35-48。
<https://doi.org/10.3329/dmcj.v5i1.31436>。
- [2] WINDARTI N., HLAING S.W.和 KAKINAKA M.肥胖症库兹涅茨曲线：国际证据。公共卫生，2019，169：26-35。
<https://doi.org/10.1016/j.puhe.2019.01.004>。
- [3] SWINBURN B.A., KRAAK V.I., ALLENDER S.等。《肥胖，营养不良与气候变化全球综合症：柳叶刀委员会报告》。柳叶刀，2019，393（10173）：791-846。
[https://doi.org/10.1016/S0140-6736\(18\)32822-8](https://doi.org/10.1016/S0140-6736(18)32822-8)。
- [4] CHOOI Y.C., DING C.和 MAGKOSF.肥胖的流行病学。代谢：临床与实验，2019，92：6-10。
<https://doi.org/10.1016/j.metabol.2018.09.005>。
- [5] 世界卫生组织（世卫组织）。肥胖和超重。2016。
<http://www.who.int/mediacentre/factsheets/fs311/en/>。
- [6] AL KIBRIA G.M.亚洲和世界卫生组织对尼泊尔成年人的体重过轻，超重和肥胖的发生率及其影响因素：对2016年人口与健康调查的分析。肥胖研究与临床实践，2019，13（2）：129-136。
<https://doi.org/10.1016/j.orcp.2019.01.006>。
- [7] NCD 风险因素协作。1975年至2014年间200个国家/地区的成年人体重指数趋势：对1698项基于人口的测量研究进行汇总分析，共有1920万参与者。柳叶刀，2016，387（10026）：1377-1396。
[https://doi.org/10.1016/S0140-6736\(16\)30054-X](https://doi.org/10.1016/S0140-6736(16)30054-X)。
- [8] RISKESDAS. 国家基本卫生研究报告。雅加达：卫生部 RI。2018。
<http://www.kesmas.kemkes.go.id>。
- [9] AKTER, S., RAHMAN, M.M., ABE, S.K. 和 SULTANAP. 孟加拉国成年人中糖尿病和前驱糖尿病的患病率及其危险因素：一项全国性调查。世界卫生组织通报，2014，92（3）：204-213。
<https://doi.org/10.2471/BLT.13.128371>。
- [10] TSUCHIYA Y., ANDO D., GOTO K.等。与在相同能量消耗下的低强度运动相比，高强度运动引起的虹膜反应更大。东北实验医学，2014，233（2）：135-140。
<https://doi.org/10.1620/tjem.233.135>。
- [11] 世界卫生组织（世卫组织）。世界卫生组织《2018年非传染性疾病预防国家概况》。2018。
<https://doi.org/10.16/j.jad.2010.09.007>。

- [12] GADDE K.M., MARTIN C.K., BERTHOUD H.R.和 HEYMSFIELD S.B.肥胖:病理生理学和管理。美国心脏病学院学报, 2018, 71 (1): 69-84。
<https://doi.org/10.1016/j.jacc.2017.11.011>。
- [13] ROSELLA L.C., KORNAS K., HUANG A.等。加拿大安大略省的人口风险和与健康行为相关的全因、早产和应得的死亡负担:加拿大社区健康调查所关联的死亡率档案。流行病学年刊, 2019, 32: 49-57。
<https://doi.org/10.1016/j.annepidem.2019.01.009>。
- [14] KHAN S.和 SHAH A.J.儿童肥胖的原因和并发症国际营养与健康科学杂志儿童肥胖的原因和并发症。科学营养健康学报, 2017, 1 (1): 47-50。
- [15] MARIE NG., FLEMING T., ROBINSON M.等。1980-2013年儿童和成人超重和肥胖的全球区域和国家患病率:系统分析。柳叶刀, 2014, 384 (9945): 766-781,
[https://doi.org/10.1016/S0140-6736\(14\)60460-8](https://doi.org/10.1016/S0140-6736(14)60460-8).Global。
- [16] BALES C.W.和 BUHR G.肥胖对老年人有害吗?对减轻体重的利弊的系统评价。美国医学指导协会杂志, 2008, 9 (5): 302-312。
<https://doi.org/10.1016/j.jamda.2008.01.006>。
- [17] TAE HO KIM, CHANG, J.S., KIM, H.等。剧烈的步行运动会影响血清 IGF-1 和 IGFBP3。生活方式医学杂志, 2015, 5 (1): 21-25。
<http://dx.doi.org/10.15280/jlm.2015.5.1.21>
- [18] MAJORCZYK M.和 SMOLA GD.在预防文明疾病的背景下, 体育锻炼对 IGF-1 和 IGFBP 水平的影响。Rocz Panstw Zakl Hig, 2016, 67 (2): 105-111。
- [19] UMOH A.V.和 ABAH G.M.血红蛋白基因型:在尼日利亚乌约的流行病学研究及其对生殖健康的影响。尼日利亚医学杂志, 2010, 19 (1): 36-41。
- [20] FAN W.和 EVANS R.M.模拟运动:对健康和表现的影响。细胞代谢, 2017, 25 (2): 242-247。
<https://doi.org/10.1016/j.cmet.2016.10.022>。
- [21] RAMBHOJAN C., BOUAZIZ-AMAR E., LARIFLA L.等。生长激素释放激素, 脂肪因子, 与学龄儿童体重状况相关的代谢因子以及一项为期 1 年的生活方式干预计划的结果。营养与代谢, 2015, 12 (43): 1-10。
<https://doi.org/10.1186/s12986-015-0039-9>。
- [22] NORHEIM F., LANGLEITE T.M., HJORTH M.等。急性运动对人皮下脂肪 PGC-1 α , 虹膜素和褐变的影响。FEBS Journal, 2014, 281 (3): 739-749。
<https://doi.org/10.1111/febs.12619>。
- [23] KOLAWOLE A.K., KEVIN N.U., OLUWOLE I.和 ADEMOLA S.乌干达卡文迪什大学社会人口统计学因素与超重/肥胖学生 (18-35 岁) 的联系。流行病学 (森尼韦尔), 2017, 7 (328): 1-6。
<https://doi.org/10.4172/2161-1165.1000328>。
- [24] MURAWSKA-CIALOWICZ E., WOJNA J. 和 ZUWALA-JAGIELLO J. Crossfit 训练在经过 Wingate 和进行性测试后, 改变了休息时脑源性神经营养因子和虹膜素的水平, 并改善了年轻运动活跃男性的有氧能力和身体成分和女人。生理与药理学报, 2015, 66 (6): 811-821。
- [25] HUH J.Y., MOUGIOS V., KABASAKALIS A.等。运动引起的虹膜素分泌与年龄或体适能水平无关, 虹膜素增加可能通过 AMPK 激活直接调节肌肉代谢。临床内分泌代谢杂志, 2014, 99 (11): E2154-E2161。
<https://doi.org/10.1210/jc.2014-1437>。
- [26] BOSTRÖMP., WU J., JEDRYCHOWSKI M.P.等。PGC1 α 依赖性肌动蛋白, 可导致白色脂肪褐变和生热。自然, 2012, 481 (7382): 463-468。
<https://doi.org/10.1038/nature10777.A>。
- [27] GUO J., XIE J., ZHOU B., 等。补锌对人体 IGF-1 水平的影响:系统评价和荟萃分析。沙特国王大学学报, 2020, 32 (3): 1824-1830。
<https://doi.org/10.1016/j.jksus.2020.01.018>。
- [28] BERRY N.T., HUBAL M.和 WIDEMAN L.急性运动发作对糖尿病前期和健康非裔美国人 GH 和 IGF-1 的影响:一项研究基因表达的初步研究。公共科学学报, 2018, 13 (1): e0191331。
<https://doi.org/10.1371/journal.pone.0191331>。
- [29] STEIN A.M., SILVA T.M.V., COELHO F.G.M.等。体育锻炼, IGF-1 和认知老年人实验研究的系统综述。痴呆症和神经心理疾病, 2018, 12 (2): 114-122。
<http://dx.doi.org/10.1590/1980-57642018dn12-020003>。
- [30] BERG U.和 BANG P.运动与循环中的胰岛素样生长因子 I 激素研究, 2004, 62 (1): 50-58。
<https://doi.org/10.1159/000080759>。
- [31] WOOD C.J., CLOW A., HUCKLEBRIDGE F.等。身体适应性和先前的体育锻炼都与社会心理压力期间皮质醇的分泌减少有关。国际期刊《焦虑, 压力与应对》, 2017, 31 (2): 135-145。
<https://doi.org/10.1080/10615806.2017.1390083>。

- [32] MARKS D.B., MARKS A.D.和 MARKS C.M.基础医学生物化学：一种临床方法。雅加达：EGC。2014。
- [33] LARON Z. 胰岛素样生长因子 1 (IGF-1)：一种生长激素。临床病理学杂志，2001，54 (5)：311-316。
<https://doi.org/10.1136/mp.54.5.311>。
- [34] CLEMMONS D.R.生长激素和 IGF-1 在控制胰岛素敏感性中的相对作用。临床研究杂志，2004，113 (1)：96-105。
- [35] TSAI C-L.， WANG C-H.， PAN C-Y.， 等。对急性抵抗运动的执行功能和内分泌反应。行为神经科学前沿，2014，8 (262)：1-12。
<https://doi.org/10.3389/fnbeh.2014.00262>。
- [36] MANNERKORPI K.， LANDIN-WILHELMSEN K.， LARSSON A.等。体育锻炼对纤维肌痛妇女血清胰岛素样生长因子系统的急性影响。BMC 肌肉骨骼疾病，2017，18 (37)：1-8。
<https://doi.org/10.1186/s12891-017-1402-y>。
- [37] NISHIDA Y.， MATSUBARA T.， TOBINA T.等。低强度有氧运动对健康男性中胰岛素样生长因子-I 和胰岛素样生长因子结合蛋白的影响。国际内分泌学杂志，2010，452820：8页。
<https://doi.org/10.1155/2010/452820>。
- [38] DIAS K.A.， INGUL C.B.， TJONNA A.E.等。高强度间歇训练对肥胖儿童体能，脂肪量和心脏代谢生物标志物的影响：一项随机对照试验。运动医学，2017，48 (3)：733-746。
<https://doi.org/10.1007/s40279-017-0777-0>。
- [39] WEWEGE M.， VAN DEN BERG R.， WARD R.E.和 KEECH A.高强度间歇训练与对超重和肥胖成年人进行身体强度中等强度连续训练：系统评价和荟萃分析。肥胖评论，2017，18 (6)：635-646。
<https://doi.org/10.1111/obr.12532>。
- [40] NYGAARD H.， NYGAARD H.， SLETTALOKKEN G.等。单次剧烈的耐力运动和沉重的力量训练后，血液中的鸢尾素会短暂增加。公共科学学报，2015，10 (3)：1-12。
<https://doi.org/10.1371/journal.pone.0121367>。
- [41] KRAEMER R.R.， SHOCKETT P.， WEBB N.D.等。男性和女性长期，中度有氧运动后短暂的虹膜血药浓度升高。激素与代谢研究，2014，46 (2)：150-154。
<https://doi.org/10.1055/s-0033-1355381>。
- [42] DASKALOPOULOU S.S.， COOKE A.B.， GOMEZ Y.H.等。在年轻，健康，活跃的受试者中，血浆虹膜素水平会随着运动量的增加而逐渐增加。欧洲内分泌学杂志，2014，171 (3)：343-352。
<https://doi.org/10.1530/EJE-14-0204>。
- [43] YOON S.J.， LEE M.J.， LEE H.M.和 LEE J.S. 低强度抵抗训练和热应激对老年妇女 HSP72，合成代谢激素，肌肉大小和力量的影响。老化临床与实验研究，2017，29：977-984。
- [44] HAYWOOD N.J.， SLATER T.A.， MATTHEWS C.J.和 WHEATCROFT S.B. 胰岛素样生长因子和结合蛋白家族：肥胖症和糖尿病的新型治疗靶标。分子代谢，2019，19：86-96。
<https://doi.org/10.1016/j.molmet.2018.10.008>。
- [45] VITALE G.， PELLEGRINO G.， VOLLERY M.和 HOFLAND L.J. IGF-1 系统在长寿调节中的作用：百岁老人的观点之争和新见解。内分泌前沿，2019，10 (27)：1-11。
<https://doi.org/10.3389/fendo.2019.00027>。
- [46] MAGGIO M.， DE VITA F.， LAURETANI F.等。IGF-1 是脆弱，营养，炎症和激素途径的交叉通道。营养素，2013，5 (10)：4184-4205。
<https://doi.org/10.3390/nu5104184>。

Treadmill and Ergo cycle Exercises Increase Insulin-Like Growth Factor-1 Levels in Obese Female

ORIGINALITY REPORT

19%

SIMILARITY INDEX

17%

INTERNET SOURCES

10%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Sunmi Park, Mark C Willingham, Jun Qi, Sheue-Yann Cheng. "Metformin and JQ1 synergistically inhibit obesity-activated thyroid cancer", *Endocrine-Related Cancer*, 2018
Publication <1 %
- 2 Lakshini Herat, Caroline Rudnicka, Yasunori Okada, Satsuki Mochizuki, Markus Schlaich, Vance Matthews. "The Metalloproteinase ADAM28 Promotes Metabolic Dysfunction in Mice", *International Journal of Molecular Sciences*, 2017
Publication <1 %
- 3 jme.bioscientifica.com
Internet Source <1 %
- 4 protocolexchange.researchsquare.com
Internet Source <1 %
- 5 eurosurveillance.org
Internet Source <1 %
- 6 www.assignmentfinish.com
Internet Source <1 %

7	www.banglajol.info Internet Source	<1 %
8	www.ncbi.nlm.nih.gov Internet Source	<1 %
9	www.springermedizin.de Internet Source	<1 %
10	Gul F. Yarim, Bugrahan B. Yagci, Murat Yarim, Mahmut Sozmen, Didem Pekmezci, Sena Cenesiz, Gokmen Z. Pekmezci, Efe Karaca. "Serum concentration and skin tissue expression of insulin-like growth factor 2 in canine generalized demodicosis", <i>Veterinary Dermatology</i> , 2015 Publication	<1 %
11	bib.irb.hr Internet Source	<1 %
12	cjbmb.bjmu.edu.cn Internet Source	<1 %
13	labtestsonline.org Internet Source	<1 %
14	www.berich.com.tw Internet Source	<1 %
15	www.intechopen.com Internet Source	<1 %

16

Internet Source

<1 %

17

www.medrxiv.org

Internet Source

<1 %

18

Agustí Noya, Guillermo Ripoll, Isabel Casasús, Albina Sanz. "Effects of immunocastration performed at two live weights on the growth physiology, temperament and testicular development of feral beef bulls", *Animal Science Journal*, 2019

Publication

<1 %

19

casedocs.omniagentsolutions.com

Internet Source

<1 %

20

cerdosmart.info

Internet Source

<1 %

21

cpb-us-e2.wpmucdn.com

Internet Source

<1 %

22

curis.ku.dk

Internet Source

<1 %

23

koreascience.or.kr

Internet Source

<1 %

24

lib.dr.iastate.edu

Internet Source

<1 %

25

pen.sagepub.com

Internet Source

<1 %

26	pure-oai.bham.ac.uk Internet Source	<1 %
27	researchers.mq.edu.au Internet Source	<1 %
28	www.biomedcentral.com Internet Source	<1 %
29	www.nutritionj.com Internet Source	<1 %
30	www.tandfonline.com Internet Source	<1 %
31	www.termedia.pl Internet Source	<1 %
32	Kuo-Chin Lin, Duan-Shin Lee, Shao-An Chien, Yueh-Hui Vanessa Chiang. "The effect of a badminton teaching system using Kinect on badminton skill performance", 體育學報, 2020 Publication	<1 %
33	Nonie K. Lesaux, Michael J. Kieffer, S. Elisabeth Faller, Joan G. Kelley. "The Effectiveness and Ease of Implementation of an Academic Vocabulary Intervention for Linguistically Diverse Students in Urban Middle Schools", Reading Research Quarterly, 2010 Publication	<1 %

alzres.biomedcentral.com

34

Internet Source

<1 %

35

cancerbiomedcentral.com

Internet Source

<1 %

36

docplayer.net

Internet Source

<1 %

37

e-journal.unair.ac.id

Internet Source

<1 %

38

jcst.ict.ac.cn:8080

Internet Source

<1 %

39

pesquisa.bvsalud.org

Internet Source

<1 %

40

pubmed.ncbi.nlm.nih.gov

Internet Source

<1 %

41

trialsjournal.biomedcentral.com

Internet Source

<1 %

42

www.chvm.net

Internet Source

<1 %

43

www.fcpc120.com

Internet Source

<1 %

44

www.gzlongteng.cn

Internet Source

<1 %

45

www.wjgnet.com

Internet Source

<1 %

46	zh.argospine.org Internet Source	<1 %
47	ClinicalTrials.gov Internet Source	<1 %
48	Min-Gul Kim, Baik-Hwan Cho, Soo-Wan Chae, Tae-Sun Park, Dal-Sik Kim. "What Do Very Low Plasma Concentrations of High-sensitivity C-reactive Protein (hs-CRP) Mean among Healthy Middle-aged Koreans?", <i>Journal of Lifestyle Medicine</i> , 2015 Publication	<1 %
49	Wen-Kang Gao, Yan-Yun Shu, Jin Ye, Xiao-Li Pan. "Circadian clock and liver energy metabolism", <i>World Chinese Journal of Digestology</i> , 2020 Publication	<1 %
50	bmcbiotechnol.biomedcentral.com Internet Source	<1 %
51	bmchealthservres.biomedcentral.com Internet Source	<1 %
52	fbj.springeropen.com Internet Source	<1 %
53	okayama.pure.elsevier.com Internet Source	<1 %
54	preview-etsmjournal.biomedcentral.com Internet Source	<1 %

- 55 www.chinagene.cn Internet Source <1 %
-
- 56 www.hdzxxx.com Internet Source <1 %
-
- 57 www.karger.com Internet Source <1 %
-
- 58 www.x-mol.com Internet Source <1 %
-
- 59 "42nd Annual Meeting Association for European Paediatric Cardiology, 16–19 May, 2007; Warsaw Medical Academy Congress Center", *Cardiology in the Young*, 2007
Publication <1 %
-
- 60 "OPENING LECTURE", *Gynecological Endocrinology*, 2009
Publication <1 %
-
- 61 J Ren. "Increases in insulin-like growth factor-1 level and peroxidative damage after gestational ethanol exposure in rats", *Pharmacological Research*, 2003
Publication <1 %
-
- 62 R Rabkin. "Work-induced changes in skeletal muscle IGF-1 and myostatin gene expression in uremia", *Kidney International*, 06/28/2006
Publication <1 %
-
- 63 dsnlab.web.unc.edu

Internet Source

<1 %

64

etds.ntut.edu.tw

Internet Source

<1 %

65

helda.helsinki.fi

Internet Source

<1 %

66

hsag.co.za

Internet Source

<1 %

67

journal.unnes.ac.id

Internet Source

<1 %

68

link.springer.com

Internet Source

<1 %

69

repository.ipb.ac.id:8080

Internet Source

<1 %

70

richtlijnendatabase.nl

Internet Source

<1 %

71

sph.umd.edu

Internet Source

<1 %

72

teachmephysiology.com

Internet Source

<1 %

73

www.biolreprod.org

Internet Source

<1 %

74

www.minervamedica.it

Internet Source

<1 %

75 www.novusbio.com Internet Source <1 %

76 www.scitepress.org Internet Source <1 %

77 H G L Rauch. "A signalling role for muscle glycogen in the regulation of pace during prolonged exercise", British Journal of Sports Medicine, 2005
Publication <1 %

78 Lovell, D.I.. "The hormonal response of older men to sub-maximum aerobic exercise: The effect of training and detraining", Steroids, 201204
Publication <1 %

79 Steven Riechman, Robert Zoeller, G. Balasekaran, Fredric Goss, Robert Robertson. "Prediction of 2000 m indoor rowing performance using a 30 s sprint and maximal oxygen uptake", Journal of Sports Sciences, 2001
Publication <1 %

80 alm.plos.org Internet Source <1 %

81 biokimiasatulima.wordpress.com Internet Source <1 %

82 caod.oriprobe.com Internet Source

<1 %

83

d-nb.info

Internet Source

<1 %

84

digitalcommons.psjhealth.org

Internet Source

<1 %

85

ecommons.luc.edu

Internet Source

<1 %

86

elifesciences.org

Internet Source

<1 %

87

eprints.nottingham.ac.uk

Internet Source

<1 %

88

jurnal.poltekkes-solo.ac.id

Internet Source

<1 %

89

libdigital.umac.mo

Internet Source

<1 %

90

pureportal.coventry.ac.uk

Internet Source

<1 %

91

researchwith.montclair.edu

Internet Source

<1 %

92

scholars.wlu.ca

Internet Source

<1 %

93

uk.pcmag.com

Internet Source

<1 %

94	www.analefefs.ro Internet Source	<1 %
95	www.caahs-acss.ca Internet Source	<1 %
96	www.cnxc001.cn Internet Source	<1 %
97	www.dwu2k.com Internet Source	<1 %
98	www.hindawi.com Internet Source	<1 %
99	www.koreascience.or.kr Internet Source	<1 %
100	E. Bognetti, F. Meschi, A. Pattarini, A. Zoja, G. Chiumello. "Post-exercise Albuminuria Does Not Predict Microalbuminuria in Type 1 Diabetic Patients", Diabetic Medicine, 1994 Publication	<1 %
101	J-G Scharf. "The IGF axis and hepatocarcinogenesis", Molecular Pathology, 6/1/2001 Publication	<1 %
102	Sindy Cornelia Nelwan, Ricardo Adrian Nugraha, Anang Endaryanto, Asti Meizarini et al. " Atopic Biomarker Changes after Exposure to Lipopolysaccharide: A Small Experimental	<1 %

Study in Wistar Rat ", Cold Spring Harbor
Laboratory, 2021

Publication

103 A R Weale. "Intestinal adaptation after
massive intestinal resection", Postgraduate
Medical Journal, 2005 $<1\%$

Publication

104 Corinne Malpuech-Brugère, Wilhelmine P.H.G.
Verboeket-van de Venne, Ronald P. Mensink,
Marie-Agnès Arnal et al. "Effects of Two
Conjugated Linoleic Acid Isomers on Body Fat
Mass in Overweight Humans", Obesity
Research, 2004 $<1\%$

Publication

105 Pontus Boström, Jun Wu, Mark P.
Jedrychowski, Anisha Korde et al. "A PGC1- α -
dependent myokine that drives brown-fat-like
development of white fat and
thermogenesis", Nature, 2012 $<1\%$

Publication

106 Sarah K. Wise, Sandra Y. Lin, Elina Toskala,
Richard R. Orlandi et al. "国际过敏与鼻科学共
识声明：变应性鼻炎", International Forum of
Allergy & Rhinology, 2019 $<1\%$

Publication

107 docsplayer.com $<1\%$

Internet Source

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Treadmill and Ergo cycle Exercises Increase Insulin-Like Growth Factor-1 Levels in Obese Female

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
