

DAFTAR PUSTAKA

- Kendrisic M., Petakovic S. 2013. Obstetric Anesthesia : Post Operative analgetik After Caesarean Section - The Simpler, The Better!. Fifth Annual Spring Scientific Symposium In Anaesthesiology and Intensive care. Faculty of Medicine, NIS; page 130-133.
- Gibbons L, Belizan JM, Lauer JA, Betrán AP, Merialdi M, Althabe F. 2010. The global numbers and costs of additionally needed and unnecessary caesarean sections performed per year: overuse as a barrier to universal coverage. World health report. Geneva, Switzerland. Volume 30; page 1-31.
- Kementerian Kesehatan Republik Indonesia. 2013. Riset Kesehatan Dasar. Jakarta. Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI.
- Bamigboye AA, Hofmeyr GJ. 2010. Caesarean section wound infiltration with local anaesthetic for post operative pain relief - any benefit?. SAMJ, Volume 100, No 5; page 313-319.
- Borges N de C, Pereira LV, Moura LA de, Silva TC, Pedroso CF. Predictors for Moderate to Severe Acute Postoperative Pain after Cesarean Section. Pain Research and Management. 2016;2016:1–6.
- Srivastava U, Verma S, Singh TK, Gupta A, Saxsena A, Jagar KD, dkk. Efficacy of trans abdominis plane block for post cesarean delivery analgetik: A double-blind, randomized trial. Saudi J Anaesth. 2015;9(3):298–302.
- McDonnell JG, Curley G, Carney J, Benton A, Costello J, Maharaj CH, dkk. The analgesic efficacy of transversus abdominis plane block after cesarean delivery: a randomized controlled trial. Anesth Analg. Januari 2008;106(1):186–91.
- Mieszkowski MM, Mayzner-Zawadzka E, Tuyakov B, Mieszkowska M, Żukowski M, Waśniewski T, dkk. Evaluation of the effectiveness of the Quadratus Lumborum Block type I using ropivacaine in postoperative analgetik after a cesarean section — a controlled clinical study. Ginekologia Polska. 2018;89(2):8.
- Blanco R, Ansari T, Riad W, Shetty N. Quadratus Lumborum Block Versus Transversus Abdominis Plane Block for Postoperative Pain After Cesarean Delivery: A Randomized Controlled Trial. Reg Anesth Pain Med. Desember 2016;41(6):757–62.
- Vejnović TR, Costa SD, Ignatov A. New Technique for Caesarean Section. Geburtshilfe Frauenheilkd. September 2012;72(9):840–5.
- Hofmeyr G.J, Mathai M, Shah A.N, Novikova N. Techniques for caesarean section. Chochrane database of systematc reviews. 2008, Issue 1. Art. No.:CD004662.
- Penn Z, Maghami Ghaem. Indications for caesarean section. Best pract res clin obstet gynaecol. 2001;15(1):1-15.

- Dodd JM, Anderson ER, Gates S, Grivell RM. Surgical techniques for uterine incision and uterine closure at the time of caesarian section. Cochrane database of systematic reviews. 2014, Issue 7. Art. No.:CD004732.
- Anderson G, Aldamluji N. Risks of aspiration in pregnant women undergoing non-obstetric procedures: intubate or not? *Br J Hosp Med (Lond)*. 2011 Apr;72(4):238.
- Chang BA MD. Physiologic changes of pregnancy. In: Chestnut H. David, editors. *Obstetric Anesthesia principles and practice* 3rd ed. Alabama: Elsevier; 2004. p. 15-32.
- Somboonviboon W, Kyokong O, Charuluxananan S, Narasethakamol A, Incidence and Risk Factors of Hypotension and Bradycardia after Spinal Anesthesia for Cesarean Section, *J Med Assoc Thai* 2008; 91 (2): 181-7
- Berges PU, Regional Anesthesia for Obstetrics. *Clinical Anesthesiology* 1969; 2 : 141-166.
- Samuel SIU, MD, Jack M. Colman, MD, Cardiovascular problems and pregnancy : An approach to management. *CCJM*. 2004: Dec;71(12):977-985.
- G. Edward Morgan, Jr. et.al, *Clinical anesthesiology, 3th: Obstetric Anesthesia*, the McGraw Hill Companies, Inc, New York, 2002, p: 804-847
- Check GT, Gutasche BB. Maternal physiologic alterations during pregnancy. In: Levinson Gershon. Shnider M Sol, editors. *Anesthesia for obstetrics* 2nd ed. Williams and Wilkins, Baltimore. 1987. p.3-10
- Mitra S, Ahuja V, Urman RD, Vadivelu N. Mechanisms of pain in Urman RD, Vadivelu N, Portenoy RK, eds. *Perioperative Pain Management*. 2013. Oxford. Oxford University Press, pp 11-18.
- Brunton LL. Goodman and Gilman's *The Pharmacological Basis of Therapeutics*, 12th ed. 2011. San Diego. McGraw-Hill.
- Kuthiala G, Chaudhary G. Ropivacaine: A review of its pharmacology and clinical use. *Indian J Anaesth* 2011;55:104-10.
- Transversus Abdominis Plane Block: Background, Indications, Contraindications. 20 September 2018 [dikutip 5 Agustus 2019]; Tersedia pada: <https://emedicine.medscape.com/article/2000944-overview#a1>
- Elsharkawy H, El-Boghdady K, Barrington M. Quadratus Lumborum Block: Anatomical Concepts, Mechanisms, and Techniques. *Anesthesiology*. Februari 2019;130(2):322–35.
- Ultrasound-Guided Transversus Abdominis Plane and Quadratus Lumborum Blocks [Internet]. NYSORA. 2018 [dikutip 11 Agustus 2019]. Tersedia pada: <https://www.nysora.com/regional-anesthesia-for-specific-surgical-procedures/abdomen/ultrasound-guided-transversus-abdominis-plane-quadratus-lumborum-blocks/>
- Peck T, Hill S. *Pharmacology for Anaesthesia and Intensive Care*, 4th ed. 2014. Cambridge. Cambridge University Press.

- Ituk, E., Habib, A.S., 2018. Enhanced recovery after cesarean delivery. *F1000 Research*, 1-11. <https://doi.org/10.12688/f1000research.13895>. 16 september 2019 (16.45)
- Hunt J, Fletcher H. Patient Survey (QoR – 40). <http://webcache.googleusercontent.com/search?q=cache:3rARjrBQSpQJ:links.lww.com/ALN/A104+&cd=3&hl=en&ct=clnk&gl=id>. 16 september 2019 (16.30)
- Ciechanowicz, S., Setty, T., Robson, E., Sathasivam, C., Chazapis, M., Dick, J., Carvalho, B., Sultan, P., 2019. Development and evaluation of an obstetric quality-of-recovery score (ObsQoR-11) after elective Caesarean delivery. *British Journal of Anaesthesia* 122, 69–78. <https://doi.org/10.1016/j.bja.2018.06.011>
- Howle, R., Ciechanowicz, S., Heppolette, C., Nakhjavani, B., Sultan, P., 2018. Validation of an obstetric quality-of-recovery scoring tool (ObsQoR-11) after emergency Caesarean delivery. *British Journal of Anaesthesia* 121, e19–e20. <https://doi.org/10.1016/j.bja.2018.05.020>
- Habib, A. S., & Ituk, U. (2018). Enhanced recovery after cesarean delivery [version 1; referees: 2 approved]. In *F1000Research* (Vol. 7). <https://doi.org/10.12688/f1000research.13895.1>
- Macones, G. A., Caughey, A. B., Wood, S. L., Wrench, I. J., Huang, J., Norman, M., Pettersson, K., Fawcett, W. J., Shalabi, M. M., Metcalfe, A., Gramlich, L., Nelson, G., & Wilson, R. D. (2019). Guidelines for postoperative care in cesarean delivery: Enhanced Recovery After Surgery (ERAS) Society recommendations (part 3). *American Journal of Obstetrics and Gynecology*, 221(3), 247.e1-247.e9. <https://doi.org/10.1016/j.ajog.2019.04.012>
- Mallan, D., Sharan, S., Saxena, S., Singh, T. K., & Faisal. (2019). Anesthetic techniques: Focus on transversus abdominis plane (TAP) blocks. *Local and Regional Anesthesia*. <https://doi.org/10.2147/LRA.S138537>
- Mukhtar, K., & Bch, M. B. (2009). TAP block Nysora. *The Journal of NYSORA*.
- Peck, T., Hill, S., & Williams, M. (2008). *Pharmacology for Anaesthesia and Intensive Care*. Cambridge University Press.
- Sutton, C. D., & Carvalho, B. (2017). Optimal Pain Management After Cesarean Delivery. In *Anesthesiology Clinics*. <https://doi.org/10.1016/j.anclin.2016.09.010>
- Ueshima, H., Otake, H., & Lin, J. A. (2017). Ultrasound-Guided Quadratus Lumborum Block: An Updated Review of Anatomy and Techniques. In *BioMed Research International*. <https://doi.org/10.1155/2017/2752876>
- EPM. (2016). *Essential Pain Management*. Available: fpm.anzca.edu.au. (2016). Essential Pain Management. [online] Available at: <http://fpm.anzca.edu.au/documents/epm-participant-manual-082016.pdf> [Accessed 6 Aug. 2019]. Last accessed 6 Aug 2019