

Daftar Pustaka

- Abe, R. et al., 2010. Gram negative bacteremia induces greater magnitude of inflammatory response than gram positive bacteremia. *Critical care*, 14(2), pp. 1-7.
- Angus, D. & Poll, T., 2013. Severe Sepsis and Septic Shock. *New England Journal of Medicine*, Volume 369, pp. 840-51.
- Anon., 2016. *Data Ruang Resusitasi*. Surabaya: Data Internal RSUD dr Soetomo.
- Barcia, M. A. & Harris, H. W., 2016. Triglyceride-rich lipoproteins as agents of innate immunity. *Oxford Journals*, 2005(41), pp. 498-503.
- Black, L. P. et al., 2020. Quantitative and Qualitative Assessments of Cholesterol Association with Bacterial Infection type in Sepsis and Septic Shock. *Journal of Intensive Care Medicine*, XX(X), pp. 1-10.
- Centers for Disease Control and Prevention, January 2017. Pneumonia (Ventilator|VAP|and non-ventilator-associated Pneumonia |PNEU|) Event. Volume pp. 6-1-16.
- Cirstea, M. et al., 2017. Decreased high-density lipoprotein cholesterol level is an early prognostic marker for organ dysfunction and death in patients with suspected sepsis. *Journal of critical care*, Volume 38, pp. 289-294.
- Colangiulo, a. et al., 2017. Serum lipid profile and survival in patients with sepsis. *Nutrition, metabolism and cardiovascular disease*, 27(1), p. 15.
- Eichbaum, E., Harris, H., Kane, J. & Rapp, J., 1991. Chylomicron can inhibit endotoxin activity in vitro. *Journal of Surgical and resuscitation*, Volume 51, pp. 413-6.
- Esteve, E., Ricart, W. & Fernandez-Real, J. M., 2005. Dyslipidemia and inflammation; an evolutionary conserved mechanism. *Clinical nutrition*, Volume 24, pp. 16-31.
- Feng, Q. P., Wei, Q. W. & Chaugai, S., 2019. Cholesterol Levels and Risk for Sepsis Among Patients Admitted to the Hospital with infectino. *JAMA Network*.
- Gouni, I., Oka, K., Etienne, J. & Chan, L., 1993. Endotoxin-induced hypertriglyceridemia is mediated by suppression of lipoprotein lipase at a post transcriptional level.. *Journal lipid Res*, Volume 34, pp. 139-46.
- Grunfeld, C. et al., 1999. Lipoproteins inhibit macrophage activation by lipotechoic acid. *J. Lipid Res*, 40(2), pp. 245-252.
- Guoying, Z. et al., 2016. The Delta high-density lipoprotein cholesterol ratio: a novel parameter for gram negative sepsis. *Springer Plus*, 5(1), pp. 1-12.
- Harvey, R. A., 2013. *Lippincott's Illustrated Reviews Microbiology*. Philadelphia: Lippincott Williams & Wilkins.
- Jawad, L., Luksic, I. & Rafnsson, S., 2012. Assessing available information on the burden of sepsis: Global estimates of incidence, prevalence and mortality.. *Journal of Global Health*, 2(1), p. 010404.

- Jung-Yien, C., Jih-Shuin, J., Chong-Jen, Y. & Pan-CHyr, Y., 2005. Low serum level of high-density lipoprotein cholesterol is a poor prognostic factor for severe sepsis. *Critical care medicine*, 33(8), pp. 1688-93.
- Khovidunkit, W. et al., 2004. Effects of infection and inflammation on lipid and lipoprotein metabolism: mechanism and consequences to the host. *Journal of Lipid Research*, Volume 45, pp. 1169-1196.
- Lee, S. H. et al., 2015. Prognostic implications of serum lipid metabolism over time during sepsis. *Biomed Research International*, pp. 1-8.
- Leeuwen, H. et al., 2003. Lipoprotein metabolism in patients with severe sepsis. *Critical care medicine*, 31(5), pp. 1359-1366.
- Levels, JH; Marquart, JA; Abraham, PR; van den Ende, AE; Malhuizen, HO; van Deventer, SJ, 2005. Lipopolysaccharide is transferred from high density to low density lipoprotein by lipopolysaccharide-binding protein and phospholipid transfer protein.. *Infection and immunity*, 73(4), pp. 2321-6.
- Martin, G. S., Mannino, D. M., Eaton, S. & Moss, M., 2003. The epidemiology of Sepsis in the United States from 1979 through 2000. *New England Journal of Medicine*, Volume 348, pp. 1546-1554.
- Memon, R., Grunfeld, C., Moser, A. & Feingold, K., 1993. Tumor necrosis factor mediates the effects of endotoxin on cholesterol and triglyceride metabolism in mice. *Endocrinology*, Volume 132, pp. 2246-53.
- Mendall, M., Patel, P. & Asante, M., 1997. Relation of serum cytokine concentrations to cardiovascular risk factors and coronary heart disease. *Heart*, Volume 78, pp. 273-7.
- Pazzini, A. et al., 2019. The impact of bacteremia on lipoprotein concentrations and patient's outcome: a retrospective analysis. *Journal of clinical microbiology & infectious disease*, pp. 1-8.
- Pirillo, A., Catapano, A. L. & Norata, G. D., 2015. HDL in Infectious disease and Sepsis. In: E. A, ed. *Handbook of experimental pharmacology*. s.l.:s.n., pp. 485-500.
- Putri, R. F. A., Setiawan, P. & Airlangga, P. S., 2018. Hubungan antara profil lipid dengan skor sofa dan mortalitas pasien sepsis di ruang resusitasi dan observasi intensif RSUD dr Soetomo Surabaya. pp. 1-121.
- Rhodes, A. & Evans, L., 2017. Surviving Sepsis Campaign: International Guidelines for Management of Septic and Septic Shock: 2016. *Critical Care Medicine*, 45(3), pp. 1-67.
- Sherwin, R., Winters, M., Vilke, G. & Wardi, G., 2017. Does early and appropriate antibiotic administration improve mortality in emergency department patients with severe sepsis or septic shock?. *Journal of emergency medicine*, 53(4), pp. 588-595.
- Singer, M. & Deutchman, C., 2016. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *Journal of American Medical Association*, 315(8).
- Struthers, K., 2017. *Clinical microbiology*. NW Washington: Taylor & Francis.

- Suwondo, V. N., Jatmiko, H. D. & Hendrianingtyas, M., 2015. Karakteristik dasar pasien sepsis yang meninggal di ICU RSUP dr Kariadi Semarang Periode 1 Januari - 31 Desember 2014. *Media Medika Muda Undip*, 4(4), pp. 1586-1596.
- Tachyla, S. A., MArochkov, A. V., Lipnitski, A. L. & Nikiforova, Y. G., 2017. The prognostic value of procalcitonin, C-reactive protein and cholesterol in patients with an infection and multiple organ dysfunction. *Korean Journal of Anesthesiology*, 70(3), pp. 305-310.
- Thaveeratitham, P., Plengpnich, W. & Naen Udom, W., 2007. Effects of human apolipoprotein A-I on endotoxin-induced leukocyte adhesion on endothelial cells in vivo and on the growth of *Escherichia coli* in vitro. *Journal of endotoxin reserach*, 13(1), pp. 58-64.
- Topchly, Elena; Cirstea, Mihai; Hye-Jin, Julia Kong; Boyd, John H; Yingjin, Wang; Russell, James A; Walley, Keith R, 2016. Lipopolysaccharide is cleared from the circulation by hepatocytes via the low density lipoprotein receptors. *PLOS One*, 11(5), pp. 1-15.
- Vincent, Jean Louis; Rello, J; Marshall, J; Silva, E; Anzueto, A; Martin, CD, 2009. International study of the prevalence and outcomes of infection in intensive care units.. *JAMA*, 302(21), pp. 2323-9.
- Wailey, K. et al., 2015. The central role of Proprotein convertase Substisillin/Kexin type 9 in septic pathogen lipid clearance. *American Journal of Respiratory and Critical Care Medicine*, 192(1), pp. 1275-1286.
- Wang, JAcob E; Dahle, Maria K; McDonald, Michelle; Foster, Simon J; Aasen, Ansgar O; Thiemermann, Christoph, 2003. Peptidoglycan and Lipothecoic acid in gram positive bacterial sepsis: receptors, signal transductoin, biological effects, and synergism. *Shock*, 20(5), pp. 402-414.
- Wang, J. et al., 2003. Peptidoglycan and lipotechoic Acid in Gram-Positive Bacterial Sepsis: Receptors, Signal Transduction, Biologiocl Effects, and Synergism. *Shock*, 20(5), pp. 402-414.
- Yamano, Shuhei; Shimizu, Kentaro; Hiroshi, Ogura; Tomoya Hirose; Hamasaki, Toshimitsu; Takeshi, Shimazu; Osamu, Tasaki, 2016. Low total cholesterol and high total bilirubiin are associated with prognostic in patients with prolonged sepsis. *Journal of critical care*, 31(1), pp. 36-40.
- Zou, G. et al., 2016. The delta high-density lipoprotein cholesterol ratio; anovel parameter for gram negative sepsis. *Springer Plus*, 1044(5), pp. 1-12.