

## DAFTAR PUSTAKA

- A., G., S., G. and R., U. (2017) ‘Study on the impact of family history of diabetes among type 2 diabetes mellitus patients in an urban area of Kancheepuram district, Tamil Nadu’, *International Journal Of Community Medicine And Public Health*, 4(11), p. 4151. doi: 10.18203/2394-6040.ijcmph20174819.
- Ahmadvaniha, R., Mansouri, M., Yunesian, M., Omidifar, K., Jeddi, M. Z., Larijani, B., Mesdaghinia A. and Rastkari N (2014) ‘Association of urinary bisphenol a concentration with type-2 diabetes mellitus’, *Journal of Environmental Health Science and Engineering*, 12(1), pp. 1–6. doi: 10.1186/2052-336X-12-64.
- Alonso-Magdalena, P., Ropero, Ana B., Soriano, S., Quesada, Ivan N. and Angel. (2010) ‘Bisphenol a exposure during pregnancy disrupts glucose homeostasis in mothers and adult male offspring’, *Environmental Health Perspectives*, 118(9), pp. 1243–1250. doi: 10.1289/ehp.1001993.
- Arnold, S. M., Clark, K. E., Staples, C. A., Klecka, G. M., Dimond, S. S., Caspers, N. and Hentges S. G. (2013) ‘Relevance of drinking water as a source of human exposure to bisphenol A’, *Journal of Exposure Science and Environmental Epidemiology*, 23(2), pp. 137–144. doi: 10.1038/jes.2012.66.
- Barboza, L. G. A., Cunha, S. C., Monteiro, C., Fernandes, J. O. and Guilhermino, L. (2020a) ‘Bisphenol A and its analogs in muscle and liver of fish from the North East Atlantic Ocean in relation to microplastic contamination. Exposure and risk to human consumers’, *Journal of Hazardous Materials*. doi: 10.1016/j.jhazmat.2020.122419.
- Barboza, L. G. A., Cunha, S. C., Monteiro, C., Fernandes, J. O. and Guilhermino, L. (2020b) ‘Bisphenol A and its analogs in muscle and liver of fish from the North East Atlantic Ocean in relation to microplastic contamination. Exposure and risk to human consumers’, *Journal of Hazardous Materials*, 393(February), p. 122419. doi: 10.1016/j.jhazmat.2020.122419.
- Bays, H. E., Chapman, R. H. and Grandy, S. (2007) ‘The relationship of body mass index to diabetes mellitus, hypertension and dyslipidaemia: Comparison of data from two national surveys’, *International Journal of Clinical Practice*, 61(5), pp. 737–747. doi: 10.1111/j.1742-1241.2007.01336.x.
- Bhatt, H., Saklani, S. and Upadhyay, K. (2016) ‘Anti-oxidant and anti-diabetic activities of ethanolic extract of Primula Denticulata Flowers’, *Indonesian Journal of Pharmacy*, 27(2), pp. 74–79. doi: 10.14499/indonesianjpharm27iss2pp74.
- Bi, Y., Wang, W., Xu, Min., Wang, T., Lu, J., Xu, Y., Dai, M., Chen, Y., Zhang, D., Sun, W., Ding, L., Chen, Y., Huang, X., Lin, L., Qi, L., Shenghan, L. and Ning, G. (2016) ‘Diabetes genetic risk score modifies effect of bisphenol a exposure on deterioration in glucose metabolism’, *Journal of Clinical Endocrinology and Metabolism*, 101(1), pp. 143–150. doi: 10.1210/jc.2015-3039.
- Biedermann, S., Tschudin, P. and Grob, K. (2010) ‘Transfer of bisphenol A from thermal printer paper to the skin’, *Analytical and Bioanalytical Chemistry*. doi: 10.1007/s00216-010-3936-9.
- Caban, M. and Stepnowski, P. (2020) ‘Determination of bisphenol A in size fractions of indoor dust from several microenvironments’, *Microchemical Journal*, 153(November 2019), p. 104392. doi: 10.1016/j.microc.2019.104392.
- Cao, X. L., Locas, C. P., Dufresne, G., Clement, G., Popovic, S., Beraldin, F., Dabeka,

- R. W. and Feeley, M. (2011) ‘Concentrations of bisphenol a in the composite food samples from the 2008 Canadian total diet study in Quebec City and dietary intake estimates’, *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment*, 28(6), pp. 791–798. doi: 10.1080/19440049.2010.513015.
- Care, D. and Suppl, S. S. (2021) ‘2. Classification and diagnosis of diabetes: Standards of medical care in diabetes-2021’, *Diabetes Care*, 44(January), pp. S15–S33. doi: 10.2337/dc21-S002.
- Corrales, J., Kristofco, L. A., Steele, W. B., Yates, B. S., Breed, C. S., Williams, E. S. and Brooks, B. W. (2015) ‘Global Assessment of Bisphenol A in the Environment : Review and Analysis of Its Occurrence and Bioaccumulation’, (September), pp. 1–29. doi: 10.1177/1559325815598308.
- Duan, Y., Yao, Y., Wang, B., Han, L., Wang, L., Sun, H. and Chen, H. (2018) ‘Association of urinary concentrations of bisphenols with type 2 diabetes mellitus: A case-control study’, *Environmental Pollution*, 243, pp. 1719–1726. doi: 10.1016/j.envpol.2018.09.093.
- European Commission (2018) ‘Commission Regulation (EU) 2018/213 of 12 February 2018 on the use of bisphenol A in varnishes and coatings intended to come into contact with food and amending Regulation (EU) No 10/2011 as regards the use of that substance in plastic food contact materi’, *Official Journal of the European Union*, 2001(May), pp. 20–30. Available at: <https://apps.fas.usda.gov/PSDOnline/app/index.html#/app/home.>
- Fahmi, U. and Wulandari, R. A. (2014) ‘Paradigma Epidemiologi Kesehatan Lingkungan’, *Kesehatan Lingkungan*, pp. 1–31.
- Flint, S., Markle, T., Thompson, S. and Wallace, E. (2012) ‘Bisphenol A exposure, effects, and policy: A wildlife perspective’, *Journal of Environmental Management*, 104, pp. 19–34. doi: 10.1016/j.jenvman.2012.03.021.
- Garnita, D. (2012) ‘Faktor Resiko Diabetes Melitus di Indonesia (Analisis Data Sakerti 2007). Skripsi Fakultas Kesehatan Masyarakat Program Studi Kesehatan Masyarakat Universitas Indonesia’, p. 189.
- Geens, T., Aerts, D., Barthot, C., Bourguignon, J. P., Goeyens, L., Lecomte, P., Rogister, G. M., Pironnet, A. M., Pussemier, L., Scippo, M. L., Loco, J. V. and Covaci, A. (2012) ‘A review of dietary and non-dietary exposure to bisphenol-A’, *Food and Chemical Toxicology*, 50(10), pp. 3725–3740. doi: 10.1016/j.fct.2012.07.059.
- Geyer, R., Jambeck, J. R. and Law, K. L. (2017) ‘Production, use, and fate of all plastics ever made’, *Science Advances*, 3(7), pp. 25–29. doi: 10.1126/sciadv.1700782.
- Hassanpour, M. and Unnisa, S. A. (2017) ‘Plastics; Applications, Materials, Processing and Techniques’, *Plastic Surgery and Modern Techniques*, 2(1), pp. 0–5. doi: 10.29011/2577-1701.100009.
- Herman, H., Dungan, M., Wolffenbuttel, R., Buse, B., Fahrbach, L., Jiang, H. and Martin, S. (2009) ‘Racial and ethnic differences in mean plasma glucose, hemoglobin A 1c, and 1,5-anhydroglucitol in over 2000 patients with type 2 diabetes’, *Journal of Clinical Endocrinology and Metabolism*, 94(5), pp. 1689–1694. doi: 10.1210/jc.2008-1940.
- Hormann, A. M., Saal, F. S. V., Nagel, S. C., Stahlhut, R. W., Moyer, C. L., Ellersiek, M. R., Welshons, W. V., Toutain, P. L. and Taylor, J. A. (2014) ‘Holding thermal receipt paper and eating food after using hand sanitizer results in high serum bioactive and urine total levels of bisphenol A (BPA)’, *PLoS ONE*,

- 9(10), pp. 1–12. doi: 10.1371/journal.pone.0110509.
- Huang, R. Ping, Liu, Z. H., Yuan, S. F., Yin, H., Dang, Zhi. and Wu, P. Xiao. (2017) ‘Worldwide human daily intakes of bisphenol A (BPA) estimated from global urinary concentration data (2000–2016) and its risk analysis’, *Environmental Pollution*, 230, pp. 143–152. doi: 10.1016/j.envpol.2017.06.026.
- Huebschmann, A. G., Huxley, R. R., Kohrt, W. M., Zeitler, P., Regensteiner, J. G. and Reusch, J. E. B. (2019) ‘Sex differences in the burden of type 2 diabetes and cardiovascular risk across the life course’, *Diabetologia*, 62(10), pp. 1761–1772. doi: 10.1007/s00125-019-4939-5.
- Hwang, S., Lim, Jung E., Choi, Y. and Jee, Sun Ha (2018) ‘Bisphenol A exposure and type 2 diabetes mellitus risk: A meta-analysis 11 Medical and Health Sciences 1117 Public Health and Health Services’, *BMC Endocrine Disorders*, 18(1). doi: 10.1186/s12902-018-0310-y.
- Igo, R. P., Kinzy, T. G. and Cooke Bailey, J. N. (2019) ‘Genetic Risk Scores’, *Current Protocols in Human Genetics*, 104(1), pp. 1–20. doi: 10.1002/cphg.95.
- Jain, S., Gupta, R. P., Gupta, D. and Jain, M. K. (2014) ‘A study on body mass index and its correlation with type 2 diabetes’, *International Journal of Research in Medical Sciences*, 2(4), p. 1638. doi: 10.5455/2320-6012.ijrms20141173.
- Jambeck, J. R., Geyer, R., Wilcox, C., Siegler, T. R., Perryman, M., Andrady, A., Narayan, R. and Law, K. L. (2015) ‘Plastic waste inputs from land into the ocean’, *Science*, 347(6223), pp. 764–768. Available at: <http://www.sciencemag.org/cgi/doi/10.1126/science.1260879>.
- Kawamura, Y., Etoh, M., Hirakawa, Y., Abe, Y. and Mutsuga, M. (2014) ‘Bisphenol A in domestic and imported canned foods in Japan’, *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment*, 31(2), pp. 330–340. doi: 10.1080/19440049.2013.874047.
- KEMENKES RI (2019) ‘Hari Diabetes Sedunia Tahun 2018’, *Pusat Data dan Informasi Kementerian Kesehatan RI*, pp. 1–8.
- Kementerian Kesehatan Republik Indonesia. ‘Peraturan Menteri Kesehatan Indonesia No 1077/Menkes/PER/2011’.
- Kim, K. and Park, H. (2013) ‘Association between urinary concentrations of bisphenol A and type 2 diabetes in Korean adults: A population-based cross-sectional study’, *International Journal of Hygiene and Environmental Health*, 216(4), pp. 467–471. doi: 10.1016/j.ijheh.2012.07.007.
- Kimber, I. (2017) ‘Bisphenol A and immunotoxic potential: A commentary’, *Regulatory Toxicology and Pharmacology*, 90, pp. 358–363. doi: 10.1016/j.yrtph.2017.08.022.
- Kolb, H. and Martin, S. (2017) ‘Environmental/lifestyle factors in the pathogenesis and prevention of type 2 diabetes’, *BMC Medicine*, 15(1), pp. 1–11. doi: 10.1186/s12916-017-0901-x.
- Konieczna, A., Rutkowska, A. and Rachon, D. (2015) ‘Health risk of exposure to Bisphenol A (BPA)’, *Roczniki Państwowego Zakładu Higieny*, 66(1), pp. 5–11.
- LaKind, J. S., Goodman, M. and Naiman, D. Q. (2012) ‘Use of NHANES Data to Link Chemical Exposures to Chronic Diseases: A Cautionary Tale’, *PLoS ONE*, 7(12). doi: 10.1371/journal.pone.0051086.
- Lee, I.. Park, Y. J., Kim, M. J., Kim, S., Choi, S., Park, J., Cho, Y. Hee., Hong, S., Yoo, J., Park, H., Cheon, G. Jeon., Choi, K. and Moon, M. Kyong. (2021) ‘Associations of urinary concentrations of phthalate metabolites, bisphenol A, and parabens with obesity and diabetes mellitus in a Korean adult population:

- Korean National Environmental Health Survey (KoNEHS) 2015–2017’, *Environmental International*, 146, p. 106227. doi: 10.1016/j.envint.2020.106227.
- Li, A. J., Xue, J., Lin, S., Maliki, A. L. A., Ghamdi, M. A. A., Kumosani, T. A. and Kannan K. (2018) ‘Urinary concentrations of environmental phenols and their association with type 2 diabetes in a population in Jeddah, Saudi Arabia’, *Environmental Research*, 166(May), pp. 544–552. doi: 10.1016/j.envres.2018.06.040.
- Liu, M., Jia, S., Dong, T., Han, Y., Xue, J., Wanjaya, E. Riani. and Fang, M. (2019) ‘The occurrence of bisphenol plasticizers in paired dust and urine samples and its association with oxidative stress’, *Chemosphere*, 216, pp. 472–478. doi: 10.1016/j.chemosphere.2018.10.090.
- Horber, M., Schecter, A., Paepke, O., Shropshire, W., Christensen, K. and Birnbaum, L. (2015) ‘Exposure assessment of adult intake of bisphenol A (BPA) with emphasis on canned food dietary exposures’, *Environment International*, 77, pp. 55–62. doi: 10.1016/j.envint.2015.01.008.
- Ma, R. C. W. and Chan, J. C. N. (2013) ‘Type 2 diabetes in East Asians: Similarities and differences with populations in Europe and the United States’, *Annals of the New York Academy of Sciences*, 1281(1), pp. 64–91. doi: 10.1111/nyas.12098.
- Ma, Y., Liu, H., Wu, J., Yuan, L., Wang, Y., Du, X., Wang, R., Marwa, P. W., Petlulu, P., Chen, X. and Zhang, H. (2019) ‘The adverse health effects of bisphenol A and related toxicity mechanisms’, *Environmental Research*, 176(June). doi: 10.1016/j.envres.2019.108575.
- Matsumoto, A., Kunugita, N., Kitagawa, K., Isse, T., Oyama, T., Foureman, G. L., Morita, M. and Kawamoto, T. (2003) ‘Bisphenol A levels in human urine.’, *Environmental health perspectives*, 111(1), pp. 101–104. doi: 10.1289/ehp.5512.
- Mikołajewska, K., Stragierowicz, J. and Gromadzińska, J. (2015) ‘Bisphenol A – Application, sources of exposure and potential risks in infants, children and pregnant women’, *International Journal of Occupational Medicine and Environmental Health*, 28(2), pp. 209–241. doi: 10.13075/ijomeh.1896.00343.
- Ning, G., Bi, Y., Wang, T., Xu, M., Xu, Y., Huang, Y., Li, M., Li, X., Wang, W., Chen, Y., Wu, Y., Hou, J., Song, A., Liu, Y. and Lai, S. (2011) ‘Relationship of urinary bisphenol A concentration to risk for prevalent type 2 diabetes in Chinese adults: a cross-sectional analysis.’, *Annals of internal medicine*, 155(6), pp. 368–374.
- Page, J., McKenzie, E., Bossuyt, M., Boutron, I., Hoffmann, T C., Mulrow, D., Shamseer, L., Tetzlaff, M., Akl, A., Brennan, E., Chou, R., Glanville, J., Grimshaw, M., Hróbjartsson, A., Lalu, M., Li, T., Loder, E W., Mayo-Wilson, E., McDonald, S., McGuinness, L A., Stewart, Lesley A., Thomas, J., Tricco, C., Welch, A., Whiting, P. and Moher, David (2021) ‘The PRISMA 2020 statement: An updated guideline for reporting systematic reviews’, *The BMJ*, 372. doi: 10.1136/bmj.n71.
- Piecha, R., Svacina, S., Maly, M., Vrbik, K., Lacinova, Z., Haluzki, M., Pavlouskova, J., Vavrouš, A., Matejkova, D., Mullerova, D., Mraz, M. and Matoulek M. (2016) ‘Urine levels of phthalate metabolites and bisphenol a in relation to main metabolic syndrome components: Dyslipidemia, hypertension and type 2 diabetes a pilot study’, *Central European Journal of Public Health*, 24(4), pp. 297–301. doi: 10.21101/cejph.a4704.

- Prager, W. and Dill, E. H. (1960) ‘An Introduction to Plasticity’, *Physics Today*, 13(3), pp. 48–48. doi: 10.1063/1.3056868.
- Provvisiero, D. P. et al. (2016) ‘Influence of bisphenol a on type 2 diabetes mellitus’, *International Journal of Environmental Research and Public Health*, 13(10). doi: 10.3390/ijerph13100989.
- Rajankar, P., Mohapatra, P. and Mathur, K. (2018) ‘Toxic Impressions-BPA in thermal paper’, *Toxics Link*.
- Rowley, J. and Slack, F. (2004) ‘Conducting a literature review’, *Management Research News*, 27(6), pp. 31–39. doi: 10.1108/01409170410784185.
- Roy, J. R., Chakraborty, S. and Chakraborty, T. R. (2009) ‘Estrogen-like endocrine disrupting chemicals affecting puberty in humans - A review’, *Medical Science Monitor*, 15(6), pp. 137–145.
- Safety, M. and Sheet, D. (2016) ‘CAS-No .: 80-05-7 MSDS MATERIAL SAFETY DATA SHEET ( MSDS )’, pp. 1–9.
- Scott M., Clark, K E., Staples, C A., Klecka, G M., Dimond, S S., Caspers, N. and Hentges, Steven G. (2013) ‘The link between family history and risk of type 2 diabetes is not explained by anthropometric, lifestyle or genetic risk factors: The EPIC-InterAct study’, *Diabetologia*, 56(1), pp. 60–69. doi: 10.1007/s00125-012-2715-x.
- Shankar, A. and Teppala, S. (2011) ‘Relationship between urinary bisphenol A levels and diabetes mellitus’, *Journal of Clinical Endocrinology and Metabolism*, 96(12), pp. 3822–3826. doi: 10.1210/jc.2011-1682.
- Silver, M. K., O'Neill, Marie S., Sowers, Mary Fran R. and Park, S K. (2011) ‘Urinary Bisphenol a and type-2 diabetes in U.S. Adults: Data from NHANES 2003–2008’, *PLoS ONE*, 6(10). doi: 10.1371/journal.pone.0026868.
- Šimková, M., Tichý, A., Dušková, M. Bradna, P. (2020) ‘Dental composites - a low-dose source of bisphenol A?’, *Physiological research*, 69, pp. S295–S304. doi: 10.33549/physiolres.934518.
- Staples, Charles A., Dorn, Philip B., Klecka, Gary M., O'Block, Sondra T., Branson, Dean R. and Harris, Lynne R. (2000) ‘Bisphenol A concentrations in receiving waters near US manufacturing and processing facilities’, *Chemosphere*, 40(5), pp. 521–525. doi: 10.1016/S0045-6535(99)00288-X.
- Suiraoaka, I. (2012) ‘Penyakit Degeneratif: Mengenal, Mencegah dan Mengurangi faktor resiko 9 Penyakit Degenaratif’, *Nuha Medica*, pp. 1–123.
- Sun, Qi, Cornelis, Marilyn C., Townsend, Mary K., Tobias, Deirdre K., Heather Eliassen, A., Franke, Adrian A., Hauser, Russ and Hu, Frank B. (2014) ‘Association of urinary concentrations of bisphenol A and phthalate metabolites with risk of type 2 diabetes: A prospective investigation in the nurses’ health study (NHS) and NHSII cohorts’, *Environmental Health Perspectives*, 122(6), pp. 616–623. doi: 10.1289/ehp.1307201.
- Sungur, Ş., Koroğlu, M. and Özkan, A. (2014) ‘Determination of bisphenol a migrating from canned food and beverages in markets’, *Food Chemistry*, 142(January), pp. 87–91. doi: 10.1016/j.foodchem.2013.07.034.
- Suryadi, H., Rasyid, A. H. and Harmita (2018) ‘Analysis of bisphenol a in indonesian canned food by gas chromatography’, *International Journal of Applied Pharmaceutics*, 10(Special Issue 1), pp. 19–23. doi: 10.22159/ijap.2018.v10s1.05.
- Takeuchi, Toru, Tsutsumi, Osamu, Ikezuki, Yumiko, Takai, Yasushi, and Taketani, Y. (2004) ‘Positive relationship between androgen and the endocrine disruptor, bisphenol A, in normal women and women with ovarian

- dysfunction', *Endocrine Journal*, 51(2), pp. 165–169. doi: 10.1507/endocrj.51.165.
- Tsai, W. T. (2006) 'Human health risk on environmental exposure to bisphenol-A: A review', *Journal of Environmental Science and Health - Part C Environmental Carcinogenesis and Ecotoxicology Reviews*. doi: 10.1080/10590500600936482.
- Vandenberg, Laura N., Hauser, Russ, Marcus, Michele, Olea, Nicolas and Welshons, Wade V. (2007) 'Human exposure to bisphenol A (BPA)', *Reproductive Toxicology*. doi: 10.1016/j.reprotox.2007.07.010.
- Wang, B., Li, Mian., Zhao, Zhiyun., Lu, Jiel., Chen, Yuhong., Xu, Yu., Xu, Min., Wang, Weiqing., Wang, Tiange., Bi, Yufang., Ning, G. (2019) 'Urinary bisphenol A concentration and glucose homeostasis in non-diabetic adults: a repeated-measures, longitudinal study', *Diabetologia*, 62(9), pp. 1591–1600. doi: 10.1007/s00125-019-4898-x.
- Wang, T., Li, Mian., Chen, Bing., Xu, Min., Xu, Yu., Huang, Yun., Lu, Jiel., Chen, Yuhong., Wang, Weiqing., Li, Xiaoying., Liu, Yu., Bi, Yufang., Lai, Shenghan and Ning, G. (2012) 'Urinary bisphenol A (BPA) concentration associates with obesity and insulin resistance', *Journal of Clinical Endocrinology and Metabolism*, 97(2), pp. 223–227. doi: 10.1210/jc.2011-1989.
- Wang, W., Abualnaja, Khalid O., Asimakopoulos, Alexandros G., Covaci, Adrian., Gevao, Bondi., Johnson-Restrepo, Boris., Kumosani, Taha A., Malarvannan, Govindan., Minh, Tu Binh., Moon, Hyo Bang., Nakata, Haruhiko., Sinha, Ravindra K., Kannan, K. (2015) 'A comparative assessment of human exposure to tetrabromobisphenol A and eight bisphenols including bisphenol A via indoor dust ingestion in twelve countries', *Environment International*, 83, pp. 183–191. doi: 10.1016/j.envint.2015.06.015.
- WHO (2014) *Updated Management Of Type 2 Diabetes In Adults At Primary Care Level*, Health Department: REPUBLIC OF SOUTH AFRICA. Available at: [https://extranet.who.int/ncdccs/Data/ZAF\\_D1\\_Management\\_of\\_type\\_2\\_Diabetes-Electronic\\_copy\\_2014.pdf](https://extranet.who.int/ncdccs/Data/ZAF_D1_Management_of_type_2_Diabetes-Electronic_copy_2014.pdf).
- WHO (2018) 'Глобальный доклад по диабету [Global report on diabetes]. Женева: Всемирная организация здравоохранения.', p. 88.
- WHO, F. and A. O. of the U. N. (2010) 'Toxicological and Health Aspects of Bisphenol A', *World Health Organization*, (November), p. 60.
- Wisudanti (2016) 'Aplikasi Terapeutik Geranin Dari Ekstrak Kulit Rambutan (*Nephelium lappaceum*) Sebagai Anti Hiperglikemik Melalui Aktivitasnya Sebagai Antioksidan Pada Diabetes Melitus Tipe 2', *Nurseline Journal*, 1(*Nephelium lappaceum*), pp. 1–19.
- Xue, J., Wan, Y. and Kannan, K. (2016) 'Occurrence of bisphenols, bisphenol A diglycidyl ethers (BADGEs), and novolac glycidyl ethers (NOGEs) in indoor air from Albany, New York, USA, and its implications for inhalation exposure', *Chemosphere*, 151, pp. 1–8. doi: 10.1016/j.chemosphere.2016.02.038.
- Zhang, Wenxin., Xia, Wei., Liu, Wenyu., Li, Xinping., Hu, Jie., Zhang, Bin., Xu., Shunqing., Zhou, Yanqiu., Li, Jiufeng., Cai, Zongwei and Li, Yuanyuan. (2019) 'Exposure to bisphenol A substitutes and gestational diabetes mellitus: A prospective cohort study in China', *Frontiers in Endocrinology*, 10(APR). doi: 10.3389/fendo.2019.00262.