

# Cytotoxic Constituents from the Stem Bark of Chisocheton pentandrus

*by Mulyadi Tanjung*

---

**Submission date:** 05-Aug-2021 01:15PM (UTC+0800)

**Submission ID:** 1627934009

**File name:** xic\_constituens\_from\_the\_stem\_bark\_of\_chisocheton\_pentandrus.pdf (697.21K)

**Word count:** 7668

**Character count:** 33123



## Cytotoxic Constituents from the Stem Bark of *Chisocheton pentandrus*

Rurini Retnowati<sup>1,\*</sup>, Hermin Sulistyarti<sup>1</sup>, Nikmatus Zahro Wahidah<sup>1</sup>, Anisa Lailatusy Syarifah<sup>2</sup>, Suprianto Salam<sup>3</sup>, Nurlelasari<sup>3</sup>, Agus Safari<sup>3</sup>, Desi Harneti<sup>3</sup>, Mulyadi Tanjung<sup>4</sup>, Ace Tatang Hidayat<sup>3,5</sup>, Rani Maharani<sup>3,5</sup>, Unang Supratman<sup>3,5</sup>, and Yoshihito Shiono<sup>6</sup>

7

<sup>1</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Brawijaya, Malang 65145, Indonesia

<sup>2</sup>Academy of Pharmacy of Putra Indonesia, Malang 65123, Indonesia

<sup>3</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences,  
Universitas Padjadjaran, Jatinangor 45363, Indonesia

<sup>4</sup>Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya 60286, Indonesia

<sup>5</sup>Central Laboratory, Universitas Padjadjaran, Jatinangor 45363, Indonesia

<sup>6</sup>Department of Food, Life, and Environmental Science, Faculty of Agriculture,  
Yamagata University, Tsuruoka, Yamagata 997-8555, Japan.

**Abstract** – Eight cytotoxic constituents, consisting of six triterpenoids, cabralealactone (**1**), cabraleadiol (**2**), prototiamin A (**3**), 23-desmethyllymocin B (**5**), melianodiol (**7**) and indicalilacol (**8**) along with one limonoid, neemfruitins A (**4**) and one protolimonoid, protoxylocarpin G (**6**), were isolated from the extract of *n*-hexane of the stem bark of *Chisocheton pentandrus*. The chemical structures were identified on the basis of spectroscopic evidence and compared to previously reported spectra. These isolated compounds appear for the first time in the plant. Compounds **1** - **8** were evaluated for their cytotoxic effect against MCF-7 breast cancer lines *in vitro*. Among the isolated compounds, melianodiol (**7**) showed the strongest cytotoxic activity with IC<sub>50</sub> values of 16.8 μM.

**Keywords** – *Chisocheton pentandrus*, MCF-7 cell lines, Meliaceae, limonoid, triterpenoids

### Introduction

The genus *Chisocheton* has approximately 150 species that are widely distributed throughout the tropical regions such as Thailand, Malaysia, and Indonesia.<sup>1,2</sup> Many *Chisocheton* plants have been traditionally used by local people to treat various diseases,<sup>3,4</sup> and previous studies on the chemical constituents of this genus have led to the isolation and determination of a number of compounds, such as triterpenoids<sup>5-7</sup> and limonoids.<sup>8-12</sup> In addition, biological effects of the constituents have been reported, such as cytotoxic, anti-inflammatory, antifungal, antimarial and antimycobacterial effects.<sup>2,8,13-15</sup> During the course of our continuing search for anticancer substances from Indonesia *Chisocheton* plants,<sup>1,7,16,17</sup> the methanol extract of the stem bark of *Chisocheton pentandrus* was found to inhibit significant cytotoxic against MCF-7 breast cancer

lines.


*C. pentandrus* is distributed in Sulawesi and Sumatera islands of Indonesia and previous investigation have led to the isolation of cytotoxic triterpenoids and limonoids.<sup>17,18</sup> In the further search for cytotoxic constituent from this plant, we investigated the *n*-hexane extract of the bark of *C. pentandrus* and obtained six triterpenoids and two limonoids (Fig. 1). The cytotoxic effects of these compounds against MCF-7 breast cancer cells were evaluated. Here, we described the structural elucidation of the isolates and their cytotoxic activity.

### Experimental

**General experimental procedures** – The IR spectra were obtained in KBr on a SHIMADZU IR Prestige-2. Mass spectra were measured with a Water QTOF-HR-TOFMS-XEV™ mass spectrometer. NMR spectra was recorded using a JEOL ECZ-500 and ECZ-600 spectrometer using tetramethyl silane (TMS) as an internal standard. Furthermore, chromatographic separations was carried out on silica gel 60 (70-230 mesh and 230-400 mesh, Merck). The PTLC glass plates were precoated

\*Author for correspondence

Rurini Retnowati, Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Brawijaya, Malang 65145, Indonesia  
Tel: +62-341-361876; E-mail: rretnowati@ub.ac.id;  
unang.supratman@unpad.ac.id

**Fig. 1.** Chemical structures of compounds 1 - 8.

with silica gel GF<sub>254</sub> (Merck, 0.25 mm). Similarly, the TLC plates were precoated with silica gel GF<sub>254</sub> (Merck, 0.25 mm), and the detection was carried out using 10% H<sub>2</sub>SO<sub>4</sub> in ethanol, followed by heating.

**Plant materials** – The stem bark of *C. pentandrus* was collected at Bogor Botanical Garden, West Java Province, Indonesia in June 2016. The plant was identified by the staff of the Herbarium Bogoriense, Bogor and an exhibition specimen (No. Bo-104) was deposited in the herbarium.

**Extraction and isolation** – The dried stem bark of *C. pentandrus* (1.8 kg) was ground to powder and extracted with methanol ( $4 \times 4$  L, 4 days each) at room temperature, and concentrated using a rotary evaporator, yielding a concentrated extract (340 g). Furthermore, about 300 g of the methanol extract was suspended in water (600 mL) and successively partitioned with *n*-hexane, ethyl acetate and *n*-butanol. The mixture was then evaporated under reduced pressure, resulting in 10.9, 25.2, and 228.6 g of crude extracts, respectively. In addition, 10.0 g of the *n*-hexane fraction was column chromatographed (CC) using silica gel (70-230 mesh), and eluted with *n*-hexane-ethyl acetate-methanol (10% stepwise), resulting to eight fraction groups (A-H).

Fraction B (439 mg) was subjected to silica gel CC (50 g, 230-400 mesh) and eluted with *n*-hexane-dichloro-

methane-ethyl acetate (5% stepwise) to give seven subfraction groups (B1-B7). Subsequently, part of the group, B3 (230 mg) was purified by crystallization in methanol to yield 1 (86.2 mg), while subfraction B7 (132.3 mg) was chromatographed using silica gel CC (10 g, 230-400 mesh) with gradient elution of *n*-hexane- dichloromethane-ethyl acetate (5% stepwise) to produce seven subfractions (B7A-B7F). Therefore, subfraction B7B (13.2 mg) was separated over silica gel CC (5 g, 230-400 mesh), and eluted with *n*-hexane: dichloromethane (1:1) to yield 2 (2.3 mg). Similarly, subfraction B7C (43.2 mg) was separated over silica gel CC (6 g, 230-400 mesh), and eluted with *n*-hexane-dichloromethane-ethyl acetate (5:4.5:0.5) to produce 3 (14.2 mg), while Fr. B7D (20.1 mg) was purified by silica gel CC (5 g, 230-400 mesh), and eluted with *n*-hexane: dichloromethane (6:4) to generate 4 (8.1 mg).

Fraction C (700 mg) was separated over silica gel CC (70 g, 70-230 mesh) with a gradient eluent of *n*-hexane-dichloromethane-ethyl acetate (5% stepwise) to produce five subfractions (C1-C5). Therefore, subfraction C3 (300 mg) was further separated over silica gel CC (30 g, 230-400 mesh), and eluted with *n*-hexane: dichloromethane (4:1) to yield four subfractions (C3A-C3D). Subsequently, subfraction C3C (110 mg) was purified by silica gel CC (15 g, 230-400 mesh), and eluted with dichloromethane :

ethyl acetate (7.5:2.5) to generate **5** (14.2 mg), similarly, subfraction C3D (14.5 mg) was separated over silica gel CC (5 g, 230-400 mesh), and eluted with chloroform: ethyl acetate (6:4) for the production of **6** (1.9 mg).

Fraction D (725 mg) was separated on CC silica gel (80 g, 70-230 mesh) using an elution gradient consisting of *n*-hexane-chloroform-ethyl acetate (5% stepwise) to generate seven sub-fractions (D1-D7). Furthermore, sub-fraction D4 (141.5 mg) was separated with CC (15 g, 230-400 mesh), and eluted with *n*-hexane: chloroform : ethyl acetate (7:1.5:1.5), to yield five subfractions (D4A-D4E), while subfraction D4C (41.2 mg) was purified on CC (5 g, 230-400 mesh), and eluted with dichloromethane-chloroform-ethyl acetate (6.0:1.5:2.5) to produce **7** (11.2 mg) and eluted with dichloromethane: ethyl acetate (7.5:2.5) to obtain **8** (2.1 mg).

**Cabralealactone (1)** – colorless crystals; m.p. 138-140 °C; IR (KBr)  $\nu_{\text{max}}$  cm<sup>-1</sup>: 2937, 2870, 1754, 1720, 1464, 1379, 1280, 1056; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz):  $\delta_{\text{H}}$  2.52 (1H, d, *J* = 10 Hz, H-23a), 2.01 (1H, m, H-22b), 1.95 (1H, m, H-5), 1.91 (1H, m, H-12b), 1.90 (1H, m, H-15a), 1.62 (1H, d, *J* = 10 Hz, H-23b), 1.56 (1H, m, H-7b), 1.53 (1H, m, H-13), 1.52 (1H, m, H-16a), 1.51 (1H, m, H-11b), 1.50 (1H, m, H-1b), 1.49 (1H, m, H-12a), 1.47 (1H, m, H-22a), 1.47 (1H, m, H-2b), 1.41 (1H, dd, *J* = 2.4, 13.2 Hz, H-9), 1.40 (1H, m, H-2a), 1.39 (1H, m, H-7a), 1.37 (1H, m, H-6b), 1.33 (3H, s, CH<sub>3</sub>-21), 1.32 (1H, m, H-6a), 1.23 (1H, m, H-17), 1.20 (1H, m, H-11a), 1.17 (1H, m, H-1a), 1.15 (1H, m, H-16b), 1.10 (1H, m, H-15b), 0.92 (3H, s, CH<sub>3</sub>-18), 0.91 (3H, s, CH<sub>3</sub>-28), 0.87 (3H, s, CH<sub>3</sub>-30), 0.82 (3H, s, CH<sub>3</sub>-19), 0.81 (3H, s, CH<sub>3</sub>-29); <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 125 MHz), Table 1; HR-TOFMS, *m/z* 415.3311 [M+H]<sup>+</sup> (calcd. C<sub>22</sub>H<sub>42</sub>O<sub>3</sub> *m/z* 414.3134).

**Cabraleadiol (2)** – colorless needle crystals; m.p. 178-180 °C;  $\nu_{\text{max}}$  cm<sup>-1</sup>: 3450, 2942, 1471, 1387, 1075; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz):  $\delta_{\text{H}}$  4.33 (1H, ddd, *J* = 2.4, 6.8, 9.6 Hz, H-3), 3.62 (1H, dd, *J* = 4.8, 10.2 Hz, H-24), 2.62 (1H, m, H-23b), 2.01 (1H, m, H-22b), 1.90 (1H, m, H-15b), 1.85 (1H, m, H-23a), 1.83 (1H, m, H-17), 1.75 (1H, m, H-12b), 1.62 (1H, m, H-13), 1.55 (1H, m, H-2b), 1.53 (1H, m, H-11b), 1.51 (1H, m, H-16b), 1.50 (1H, m, H-7b), 1.49 (1H, m, H-12a), 1.46 (1H, m, H-16a), 1.44 (1H, dd, *J* = 2.4, 13.2 Hz, H-9), 1.42 (1H, m, H-1b), 1.39 (1H, m, H-6b), 1.38 (1H, m, H-2a), 1.34 (1H, m, H-7a), 1.32 (1H, m, H-6a), 1.24 (1H, m, H-5), 1.22 (1H, m, H-22a), 1.20 (1H, m, H-11a), 1.17 (1H, m, H-1a), 1.17 (3H, s, CH<sub>3</sub>-26), 1.13 (3H, s, CH<sub>3</sub>-21), 1.09 (3H, s, CH<sub>3</sub>-27), 1.04 (1H, m, H-15a), 0.95 (3H, s, CH<sub>3</sub>-18), 0.92 (3H, s, CH<sub>3</sub>-28), 0.87 (3H, s, CH<sub>3</sub>-30), 0.84 (3H, s, CH<sub>3</sub>-19), 0.82 (3H, s, CH<sub>3</sub>-29); <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 125 MHz), Table 1; HR-

TOFMS *m/z* 461.3271 [M+H]<sup>+</sup>, (calcd. C<sub>30</sub>H<sub>52</sub>O<sub>3</sub> *m/z* 460.3916).

**Prototiamin A (3)** – White amorphous powder; IR (KBr)  $\nu_{\text{max}}$  cm<sup>-1</sup>: 3430, 2945, 1720, 1639, 1456, 1387, 1074, 847; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 600 MHz):  $\delta_{\text{H}}$  7.14 (1H, d, *J* = 10.2 Hz, H-1), 5.86 (1H, d, *J* = 10.2 Hz, H-2), 5.28 (1H, br.s, H-15a), 5.05 (1H, m, H-7), 4.47 (1H, dd, *J* = 9.6, 1.8 Hz, H-23), 3.92 (1H, d, *J* = 9.6 Hz, H-24), 2.71 (1H, m, H-20), 2.53 (1H, m, H-6b), 2.25 (1H, m, H-6a), 2.20 (1H, m, H-5), 2.19 (1H, m, H-17), 2.19 (1H, m, H-9), 2.05 (1H, m, H-22b), 1.94 (3H, s, CH<sub>3</sub>-2'), 1.82 (1H, m, H-22a), 1.82 (1H, m, H-16b), 1.76 (1H, m, H-12b), 1.63 (1H, m, H-13), 1.57 (1H, m, H-12a), 1.50 (1H, m, H-11b), 1.44 (3H, s, CH<sub>3</sub>-27), 1.35 (3H, s, CH<sub>3</sub>-26), 1.22 (1H, m, H-11a), 1.17 (1H, m, H-16a), 1.16 (3H, s, CH<sub>3</sub>-30), 1.16 (3H, s, CH<sub>3</sub>-18), 1.07 (3H, s, CH<sub>3</sub>-19), 1.06 (3H, s, CH<sub>3</sub>-28), 1.06 (3H, s, CH<sub>3</sub>-29); <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 150 MHz), Table 1. HR-TOFMS, *m/z* 543.1260 [M+H]<sup>+</sup>, (calcd. C<sub>32</sub>H<sub>46</sub>O<sub>7</sub> *m/z* 542.1244).

**Neemfruitin A (4)** – Colorless solid; IR (KBr)  $\nu_{\text{max}}$  cm<sup>-1</sup>: 3360, 2939, 1725, 1710, 1458, 1380, 1109; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 600 MHz):  $\delta_{\text{H}}$  5.34 (1H, t, *J* = 2.4 Hz, H-1), 5.33 (1H, t, *J* = 2.4 Hz, H-7), 5.28 (1H, dd, *J* = 1.8, 3.6 Hz, H-15), 4.57 (1H, dd, *J* = 9.6, 1.8 Hz, H-23), 4.04 (2H, d, *J* = 9.6 Hz, H-21), 2.70 (1H, m, H-20), 2.53 (1H, m, H-6b), 2.50 (1H, dd, *J* = 8.0, 10.4 Hz, H-5), 2.25 (1H, m, H-6a), 2.20 (1H, m, H-9), 2.17 (1H, m, H-22a), 2.16 (1H, m, H-22b), 2.04 (1H, m, H-16a), 1.97 (3H, s, CH<sub>3</sub>-2''), 1.93 (1H, m, H-2a), 1.93 (1H, m, H-11a), 1.91 (3H, s, CH<sub>3</sub>-2'), 1.73 (1H, m, H-12b), 1.70 (1H, dt, *J* = 3.0, 10.2 Hz, H-17), 1.61 (1H, m, H-2b), 1.56 (1H, m, H-11b), 1.54 (1H, m, H-12a), 1.45 (1H, m, H-16b), 1.16 (3H, s, CH<sub>3</sub>-30), 1.16 (3H, s, CH<sub>3</sub>-19), 1.09 (3H, s, CH<sub>3</sub>-29), 1.06 (3H, s, CH<sub>3</sub>-28), 1.00 (3H, s, CH<sub>3</sub>-18); <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 150 MHz), Table 1. HR-TOFMS, *m/z* 517.4498 [M+H]<sup>+</sup>, (calcd. C<sub>30</sub>H<sub>44</sub>O<sub>7</sub> *m/z* 516.4087).

**23-Desmethyllymocin B (5)** – white solid; IR (KBr)  $\nu_{\text{max}}$  cm<sup>-1</sup>: 3300, 2949, 1705, 1620, 1457, 1380, 1080; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 600 MHz):  $\delta_{\text{H}}$  7.14 (1H, d, *J* = 10.2 Hz, H-1), 5.86 (1H, d, *J* = 10.2 Hz, H-2), 5.33 (1H, t, *J* = 2.4 Hz, H-7), 5.28 (1H, dd, *J* = 1.8, 3.6 Hz, H-15), 4.57 (1H, dd, *J* = 9.6, 1.8 Hz, H-23), 4.04 (2H, d, *J* = 9.6 Hz, H-21), 2.70 (1H, m, H-20), 2.53 (1H, m, H-6b), 2.50 (1H, dd, *J* = 8.0, 12.4 Hz, H-5), 2.25 (1H, m, H-6a), 2.20 (1H, m, H-10), 2.17 (1H, m, H-22a), 2.16 (3H, s, CH<sub>3</sub>-2'), 2.16 (1H, m, H-22b), 2.04 (1H, m, H-16a), 1.93 (1H, m, H-11a), 1.92 (1H, m, H-16b), 1.73 (1H, m, H-12b), 1.70 (1H, dt, *J* = 3.0, 10.2 Hz, H-17), 1.58 (1H, m, H-11b), 1.54 (1H, m, H-12a), 1.16 (3H, s, CH<sub>3</sub>-19), 1.06 (3H, s, CH<sub>3</sub>-29), 1.06 (3H, s, CH<sub>3</sub>-30), 1.00 (3H, s, CH<sub>3</sub>-18); <sup>13</sup>C-

**Table 1.**  $^{13}\text{C}$ -NMR data for compounds **1 - 8**

Position carbon	Compounds							
	<b>1*</b> $\delta_{\text{C}}$ (mult.)	<b>2*</b> $\delta_{\text{C}}$ (mult.)	<b>3*</b> $\delta_{\text{C}}$ (mult.)	<b>4*</b> $\delta_{\text{C}}$ (mult.)	<b>5**</b> $\delta_{\text{C}}$ (mult.)	<b>6**</b> $\delta_{\text{C}}$ (mult.)	<b>7**</b> $\delta_{\text{C}}$ (mult.)	<b>8**</b> $\delta_{\text{C}}$ (mult.)
1	35.2 (t)	33.7 (t)	158.8 (d)	75.4 (d)	158.8 (d)	158.8 (d)	31.2 (t)	31.2 (t)
2	33.7 (t)	25.4 (t)	125.7 (d)	25.6 (t)	125.7 (d)	125.7 (d)	25.6 (t)	25.6 (t)
3	218.2 (s)	73.2 (d)	204.6 (s)	218.3 (s)	204.6 (s)	204.6 (s)	218.4 (s)	72.4 (d)
4	37.3 (s)	37.3 (s)	44.6 (s)	39.9 (s)	44.6 (s)	44.6 (s)	40.3 (s)	40.3 (s)
5	55.4 (d)	49.6 (d)	46.2 (d)	49.8 (d)	49.8 (d)	46.2 (d)	44.6 (d)	44.6 (d)
6	18.3 (t)	18.3 (t)	27.5 (t)	27.5 (t)	27.5 (t)	27.5 (t)	24.8 (t)	24.8 (t)
7	34.7 (t)	35.4 (t)	75.5 (d)	75.6 (d)	74.4 (d)	34.9 (t)	118.8 (d)	118.8 (d)
8	40.6 (s)	40.7 (s)	42.8 (s)	43.0 (s)	43.0 (s)	42.8 (s)	145.3 (s)	145.3 (s)
9	50.4 (d)	50.7 (d)	37.4 (d)	37.4 (d)	37.4 (d)	37.4 (d)	48.9 (d)	48.9 (d)
10	37.7 (s)	37.7 (s)	39.7 (s)	47.2 (s)	47.2 (s)	39.7 (s)	34.6 (s)	34.6 (s)
11	25.4 (t)	21.7 (t)	34.1 (t)	16.2 (t)	16.2 (t)	34.1 (t)	17.4 (t)	17.4 (t)
12	21.3 (t)	27.1 (t)	38.3 (t)	40.0 (t)	40.0 (t)	38.3 (t)	31.2 (t)	31.2 (t)
13	43.2 (d)	42.8 (d)	46.8 (s)	50.2 (s)	50.2 (s)	46.8 (s)	43.8 (s)	43.8 (s)
14	50.3 (s)	50.2 (s)	157.9 (s)	158.9 (s)	158.9 (s)	157.9 (s)	50.7 (s)	50.7 (s)
15	31.2 (t)	31.5 (t)	119.0 (d)	118.0 (d)	118.0 (d)	119.0 (d)	33.9 (t)	33.9 (t)
16	25.1 (t)	25.9 (t)	33.8 (t)	34.6 (t)	34.6 (t)	33.8 (t)	23.9 (t)	23.9 (t)
17	49.5 (d)	49.8 (d)	58.2 (d)	55.3 (d)	55.3 (d)	58.2 (d)	47.2 (d)	47.2 (d)
18	15.6 (q)	16.2 (q)	16.2 (q)	19.7 (q)	19.7 (q)	16.2 (q)	23.5 (q)	23.5 (q)
19	16.1 (q)	16.6 (q)	19.1 (q)	15.3 (q)	15.3 (q)	19.1 (q)	13.1 (q)	13.1 (q)
20	90.3 (s)	86.7 (s)	44.7 (s)	37.6 (d)	37.6 (d)	44.7 (s)	40.7 (d)	40.7 (d)
21	25.4 (q)	27.3 (q)	178.6 (s)	70.3 (d)	72.5 (d)	72.5 (d)	99.3 (d)	178.3 (s)
22	31.3 (t)	35.3 (t)	33.1 (t)	34.2 (t)	34.2 (t)	33.1 (t)	29.6 (t)	29.6 (t)
23	29.3 (t)	26.4 (t)	78.6 (d)	99.8 (d)	99.2 (d)	78.2 (d)	86.4 (d)	86.4 (d)
24	176.9 (s)	86.3 (d)	72.4 (d)	-	-	74.4 (d)	71.3 (d)	71.3 (d)
25	-	70.3 (s)	70.5 (s)	-	-	72.5 (s)	70.3 (s)	70.3 (s)
26	-	27.9 (q)	26.9 (q)	-	-	26.9 (q)	26.5 (q)	26.5 (q)
27	-	24.1 (q)	23.8 (q)	-	-	23.8 (q)	27.0 (q)	27.0 (q)
28	28.4 (q)	28.4 (q)	20.0 (q)	21.4 (q)	-	20.0 (q)	21.6 (q)	21.6 (q)
29	22.2 (q)	22.2 (q)	21.4 (q)	27.1 (q)	21.4 (q)	21.4 (q)	27.4 (q)	27.4 (q)
30	16.4 (q)	15.6 (q)	27.1 (q)	27.5 (q)	27.1 (q)	27.1 (q)	27.8 (q)	27.8 (q)
1'	-	-	170.1 (s)	170.1 (s)	27.5 (q)	170.1 (s)		
2'	-	-	21.2 (q)	21.1 (q)	170.1 (s)	21.2 (q)		
1''	-	-		170.0 (s)				
2''	-	-		21.0 (q)				

\*Measured in 500 MHz for  $^1\text{H}$  and 125 MHz for  $^{13}\text{C}$

\*\* Measured in 600 MHz for  $^1\text{H}$  and 150 MHz for  $^{13}\text{C}$

NMR ( $\text{CD}_3\text{OD}$ , 150 MHz), Table 1. HR-TOFMS,  $m/z$  455.0712 [ $\text{M}+\text{H}$ ] $^+$ , (calcd.  $\text{C}_{28}\text{H}_{40}\text{O}_5$   $m/z$  456.2876).

**Protoxylcarpin G (6)** – White solid; IR (KBr)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 3420, 2949, 1715, 1645, 1457, 1380, 1080;  $^1\text{H}$ -NMR ( $\text{CD}_3\text{OD}$ , 600 MHz):  $\delta_{\text{H}}$  7.14 (1H, d,  $J=10.2$  Hz, H-1), 5.86 (1H, d,  $J=10.2$  Hz, H-2), 5.28 (1H, br.s, H-15), 4.16 (2H, d,  $J=9.6$  Hz, H-23), 3.92 (1H, d,  $J=9.6$  Hz, H-24), 2.71 (H, m, H-20), 2.53 (1H, m, H-6b), 2.25

(1H, m, H-6a), 2.20 (IH, m, H-5), 2.19 (1H, m, H-17), 2.19 (1H, m, H-9), 1.94 (3H, s,  $\text{CH}_3$ -2'), 1.87 (1H, m, H-16a), 1.78 (1H, m, H-12a), 1.69 (1H, m, H-7b), 1.65 (m, H-7a), 1.57 (1H, m, H-12b), 1.52 (1H, m, H-11a), 1.50 (1H, m, H-11b), 1.44 (3H, s,  $\text{CH}_3$ -27), 1.35 (3H, s,  $\text{CH}_3$ -26), 1.17 (1H, m, H-16b), 1.16 (3H, s,  $\text{CH}_3$ -30), 1.16 (3H, s,  $\text{CH}_3$ -18), 1.01 (3H, s,  $\text{CH}_3$ -19), 1.06 (3H, s,  $\text{CH}_3$ -28), 1.06 (3H, s,  $\text{CH}_3$ -29);  $^{13}\text{C}$ -NMR ( $\text{CD}_3\text{OD}$ , 150 MHz)

Table 1. HR-TOFMS,  $m/z$  527.2984 [M+H]<sup>+</sup>, (calcd. C<sub>32</sub>H<sub>48</sub>O<sub>6</sub>  $m/z$  528.2951).

**Melianodiol (7)** – Colorless needle crystal; m.p. 216–218°C; IR (KBr)  $\nu_{\text{max}}$  cm<sup>-1</sup>: 3350, 2935, 1725, 1645, 1457, 1380, 1055; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 600 MHz):  $\delta_{\text{H}}$  5.26 (1H, dd,  $J$ =2.6, 5.8 Hz, H-7), 4.63 (1H, d,  $J$ =5.6 Hz, H-21), 4.05 (1H, br.s, H-24), 3.64 (1H, ddd,  $J$ =1.3, 6.2, 8.7 Hz, H-23), 2.67 (1H, ddd,  $J$ =5.7, 8.4, 11.7 Hz, H-20), 2.38 (1H, m, H-22a), 2.34 (1H, m, H-17), 2.24 (1H, m, H-9), 2.20 (1H, ddd,  $J$ =6.2, 8.4, 11.7 Hz, H-22b), 2.09 (1H, d,  $J$ =14.7, H-6a), 1.93 (1H, m, H-2a), 1.78 (1H, m, H-5), 1.76 (1H, m, H-12a), 1.73 (1H, m, H-16b), 1.63 (1H, m, H-11a), 1.61 (1H, m, H-2b), 1.61 (1H, m, H-12b), 1.57 (1H, m, H-16a), 1.53 (1H, m, H-11b), 1.52 (1H, m, H-15b), 1.49 (1H, m, H-1a), 1.41 (1H, m, H-6b), 1.39 (3H, s, CH<sub>3</sub>-27), 1.38 (1H, dt,  $J$ =3.2, 12.2 Hz, H-1b), 1.32 (3H, s, CH<sub>3</sub>-26), 1.06 (1H, m, H-15a), 1.01 (3H, s, CH<sub>3</sub>-30), 0.93 (3H, s, CH<sub>3</sub>-18), 0.93 (3H, s, CH<sub>3</sub>-28), 0.92 (3H, s, CH<sub>3</sub>-29), 0.78 (3H, s, CH<sub>3</sub>-19); <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 150 MHz) Table 1. HR-TOFMS,  $m/z$  487.4535 [M+H]<sup>+</sup>, (calcd. C<sub>32</sub>H<sub>50</sub>O<sub>7</sub>  $m/z$  488.4502).

**Indicalilacol B (8)** – White solid; IR (KBr)  $\nu_{\text{max}}$  cm<sup>-1</sup>: 3430, 2930, 1760, 1655, 1420, 1378, 1210, 1110; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 600 MHz):  $\delta_{\text{H}}$  5.26 (1H, dd,  $J$ =2.6, 5.8 Hz, H-7), 4.63 (1H, d,  $J$ =5.6 Hz, H-21), 4.32 (1H, d,  $J$ =5.4 Hz, H-3), 4.05 (1H, br.s, H-24), 3.64 (1H, ddd,  $J$ =1.3, 6.2, 8.7 Hz, H-23), 2.67 (1H, ddd,  $J$ =5.7, 8.4, 11.7 Hz, H-20), 2.38 (1H, m, H-22a), 2.34 (1H, m, H-17), 2.24 (1H, m, H-9), 2.20 (1H, ddd,  $J$ =6.2, 8.4, 11.7 Hz, H-22b), 2.09 (1H, d,  $J$ =14.7, H-6a), 1.93 (1H, m, H-2a), 1.78 (1H, m, H-5), 1.76 (1H, m, H-12a), 1.73 (1H, m, H-16b), 1.63 (1H, m, H-11a), 1.61 (1H, m, H-12b), 1.61 (1H, m, H-2b), 1.57 (1H, m, H-16a), 1.53 (1H, m, H-11b), 1.52 (1H, m, H-15b), 1.49 (1H, m, H-1a), 1.41 (1H, m, H-6b), 1.39 (3H, s, CH<sub>3</sub>-27), 1.38 (1H, dt,  $J$ =3.2, 12.2 Hz, H-1b), 1.32 (3H, s, CH<sub>3</sub>-26), 1.06 (1H, m, H-15a), 1.01 (3H, s, CH<sub>3</sub>-30), 0.93 (3H, s, CH<sub>3</sub>-28), 0.92 (3H, s, CH<sub>3</sub>-29), 0.84 (3H, s, CH<sub>3</sub>-18), 0.78 (3H, s, CH<sub>3</sub>-19); <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 160 MHz) see Table 1. HR-TOFMS,  $m/z$  487.4535 [M+H]<sup>+</sup>, (calcd. C<sub>32</sub>H<sub>50</sub>O<sub>7</sub>  $m/z$  488.4502).


**Cytotoxicity assay** – The MCF-7 Cells were plated at a density of  $5 \times 10^3$  cells/well in 96-well plate, incubated at 37°C overnight and treated in triplicate with decreasing concentrations isolated compound from *C. pentandrus* extract. Furthermore, all the extracted were dissolved in DMSO and *Cis*-platin (Sigma) was used as a positive control. After 48 hours of treatment, cell viability was evaluated using the MTT reagent 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide was added according to the manufactured protocols, followed by a 4-hour

incubation. The results were expressed as relative viable cells compared to the controls (untreated cells). The 50% inhibitory concentration (IC<sub>50</sub>) values of antiproliferative activity were calculated and converted into the IC<sub>50</sub> Dose.


## Results and Discussion

The methanol extract of the stem bark was partitioned successfully with *n*-hexane, ethyl acetate and *n*-butanol. Column chromatography was repeated using silica gel of the soluble fraction in *n*-hexane led to the isolation of eight isolated compounds (Fig. 1). The structures of the isolated compounds were identified by spectroscopic methods including 1D, 2D NMR and HR-TOFMS. To the best our knowledge, compounds **1** - **8**, were isolated from *C. pentandrus* for the first time.

Compound **1** was obtained as colorless crystals (MeOH). The molecular composition was determined was C<sub>27</sub>H<sub>42</sub>O<sub>3</sub> based on HR-TOFMS ([M+H]<sup>+</sup>  $m/z$  415.3311, calcd. for C<sub>27</sub>H<sub>42</sub>O<sub>3</sub>  $m/z$  414.3334) and NMR analysis (Table 1), and seven degrees of unsaturation is required. Furthermore, IR spectra showed absorption peaks at 2937 and 2870 cm<sup>-1</sup> (C-H sp<sup>3</sup>), 1754 cm<sup>-1</sup> (C=O), 1720 (C=O, lactone) 1464 cm<sup>-1</sup> (C=C), 1379-1280 cm<sup>-1</sup> (gem-dimethyl), and 1056 cm<sup>-1</sup> (C-O groups). The <sup>1</sup>H-NMR spectrum showed the presence of six tertiary methyl groups, resonating at  $\delta_{\text{H}}$  1.33 (CH<sub>3</sub>-21), 0.92 (CH<sub>3</sub>-18), 0.91 (CH<sub>3</sub>-28), 0.87 (CH<sub>3</sub>-30), 0.82 (CH<sub>3</sub>-19) and 0.81 (CH<sub>3</sub>-29). The aliphatic protons in the upfield region were observed also in the <sup>1</sup>H-NMR spectra. The <sup>13</sup>C-NMR, HMQC and DEPT 135° spectra of **1** showed the presence of six methyl groups, exhibiting the characteristics of dammarane-type triterpenoid, one oxygenated quaternary carbon at  $\delta_{\text{C}}$  90.3 (C-20), one carbonyl lactone at  $\delta_{\text{C}}$  176.9 (C-24) and one carbonyl ketone at  $\delta_{\text{C}}$  218.2 (C-3). These functions represented two of the total of seven degrees of unsaturation, and the remaining five degrees of unsaturation matched the dammarane-type triterpenoid framework with an addition of a five-membered lactone ring.<sup>19,20</sup> The <sup>1</sup>H-<sup>1</sup>H correlations (Fig. 2) from H-1/H-2, H-5/H-6/H-7, H-9/H-11/H-12/H-13, H-15/H-16/H-17 and H-22/H-23 were aided by the presence of the dammarane-type in compound **1**. The HMBC cross peaks (Fig. 2) from H-1 ( $\delta_{\text{H}}$  1.17 and 1.50) and H-2 ( $\delta_{\text{H}}$  1.40 and 1.47) to the carbonyl ketone at C-3 ( $\delta_{\text{C}}$  218.2) indicated the presence of a ketone group at C-3. Protons signal H-17 ( $\delta_{\text{H}}$  1.23) and CH<sub>3</sub>-21 ( $\delta_{\text{H}}$  1.33) were correlated to C-20 ( $\delta_{\text{C}}$  90.3), whereas methylene protons at C-23 ( $\delta_{\text{H}}$  2.52 and 2.62) was correlated to C-24 ( $\delta_{\text{C}}$  176.9) indicated the presence of five-membered lactone ring at C-17. The relative



**Fig. 2.** Selected  $^1\text{H}$ - $^1\text{H}$  COSY and HMBC correlations of compounds **1**, **2**, **3** and **5**.



**Fig. 3.** Selected NOESY correlations for compounds **1** and **2**.

configuration of **1** was determined by NOESY experiments, which is supported by the presence of dammarane-type triterpenoids in *Chisocheton* species.<sup>19,20</sup> The NOESY correlations (Fig. 3) from H-17/CH<sub>3</sub>-30 identified the  $\gamma$ -lactone ring at C-17 as  $\beta$ -oriented, while the cross peak observed between CH<sub>3</sub>-30/H-17/CH<sub>3</sub>-21, indicates the CH<sub>3</sub>-21 as  $\alpha$ -oriented. A comparison of the NMR data of **1** with cabralealactone obtained from the whole plant of *Cleome africana*,<sup>19</sup> revealed that the structures of the two compounds were very similar, consequently, compound **1** was identified as cabralealactone.

Compound **2** was obtained as a colorless needle crystal. Its molecular composition C<sub>30</sub>H<sub>52</sub>O<sub>3</sub>, was determined from HR-TOFMS spectra ( $m/z$  461.3271 [M+H]<sup>+</sup>, calculated, C<sub>30</sub>H<sub>52</sub>O<sub>3</sub>  $m/z$  460.3274) and NMR data (Table 1). The IR spectra showed absorption peaks at 3450 cm<sup>-1</sup> (OH), 2942 cm<sup>-1</sup> (C-H sp<sup>3</sup>), 1471 and 1387 cm<sup>-1</sup> (gem-dimethyl), and 1075 cm<sup>-1</sup> (C-O) groups. The  $^1\text{H-NMR}$  spectrum showed the presence of eight tertiary methyl groups at  $\delta_{\text{H}}$  1.17 (CH<sub>3</sub>-26), 1.13 (CH<sub>3</sub>-21), 1.09 (CH<sub>3</sub>-27), 0.95 (CH<sub>3</sub>-18), 0.92 (CH<sub>3</sub>-28), 0.87 (CH<sub>3</sub>-30), 0.84 (CH<sub>3</sub>-19) and 0.82 (CH<sub>3</sub>-29) resonances. Two oxygenated

methines resonances at  $\delta_H$  4.33 (1H, ddd,  $J=2.4, 6.8, 9.6$  Hz, H-3) and  $\delta_H$  3.62 (1H, dd,  $J=4.8, 10.2$  Hz, H-24) were also observed in the  $^1\text{H-NMR}$  spectra. The  $^{13}\text{C-NMR}$  spectra showed 30 carbons and were classified by DEPT 135° and HMQC experiments as eight methyl groups, exhibiting the characteristics of dammarane-type triterpenoid,<sup>19,20</sup> an oxymethine group at  $\delta_C$  86.3 (C-24) and 73.2 (C-3) as well as an oxygenated quaternary carbon at  $\delta_C$  70.3 (C-25). The  $^1\text{H-}^1\text{H}$  correlations (Fig. 2) from H-1/H-2/H-3, H-5/H-6/H-7, H-9/H-11/H-12/H-13, H-15/H-16/H-17 and H-22/H-23 were aided by the presence of the dammarane-type in compound **2**.<sup>19,20</sup> The HMBC cross peaks (Fig. 2) from CH<sub>3</sub>-28 ( $\delta_H$  0.92), CH<sub>3</sub>-29 ( $\delta_H$  0.82), and the methylene protons at H-2 ( $\delta_H$  1.55 and 1.38) to the oxymethine carbon at C-3 ( $\delta_C$  73.2) indicated the presence of a hydroxy group at C-3. The correlation of H-22 ( $\delta_H$  1.22 and 2.01) and H-23 ( $\delta_H$  2.52 and 2.62) to C-24 ( $\delta_C$  86.3) and C-20 ( $\delta_C$  86.7) indicates that the position of furan ring at C-20/C-24. Furthermore, methyl protons at  $\delta_H$  1.17 (CH<sub>3</sub>-26) and 1.09 (CH<sub>3</sub>-27) as well as the oxygenated proton at  $\delta_H$  3.62 correlated with oxygenated quaternary carbon at  $\delta_C$  86.3 (C-24), indicating that an isopropyl alcohol moiety was located at C-24. In addition, the key NOESY correlation (Fig. 3) between CH<sub>3</sub>-21 and CH<sub>3</sub>-26, CH<sub>3</sub>-27 specifies the isopropyl group at C-24 as a  $\alpha$ -orientation, which is supported by the presence of dammarane-type triterpenoids in *Chisocheton* species.<sup>19,20</sup> The NMR spectra of **2** were in good agreement with that reported cabraleadiol isolated from *Chisocheton penduliflorus*,<sup>20</sup> therefore, compound **2** was identified as a cabraleadiol.

Compound **3** was isolated as a white amorphous powder. Its molecular composition was assigned as C<sub>32</sub>H<sub>46</sub>O<sub>7</sub> on the basis of the HRTOFMS ( $m/z$  543.1260 [M+H]<sup>+</sup>, (calcd. C<sub>32</sub>H<sub>46</sub>O<sub>7</sub>  $m/z$  542.1244 NMR data (Table 1), therefore required ten double-bond equivalents. IR spectra showed absorption peaks at 3430 cm<sup>-1</sup> (OH), 2945 cm<sup>-1</sup> (C-H sp<sup>3</sup>), 1720 (C=O), 1639 cm<sup>-1</sup> (C=C), 1456 and 1378 cm<sup>-1</sup> (gem-dimethyl) and 1076 cm<sup>-1</sup> (C-O) groups. The  $^1\text{H-NMR}$  spectrum showed the presence of seven tertiary methyl groups, resonance at  $\delta_H$  1.44 (CH<sub>3</sub>-27), 1.35 (CH<sub>3</sub>-26), 1.16 (CH<sub>3</sub>-30), 1.16 (CH<sub>3</sub>-18), 1.07 (CH<sub>3</sub>-19), 1.06 (CH<sub>3</sub>-28) and 1.06 (CH<sub>3</sub>-29). The signal of one acetyl at  $\delta_H$  1.94 (CH<sub>3</sub>-2'), two oxygenated methines at  $\delta_H$  4.47 (1H, dd,  $J=9.6, 1.8$  Hz, H-23), 3.92 (1H, d,  $J=9.6$  Hz, H-24), three methine sp<sup>2</sup> at  $\delta_H$  7.14 (1H, d,  $J=10.2$  Hz, H-1), 5.86 (1H, d,  $J=10.2$  Hz, H-2) and 5.28 (1H, br.s, H-15) were observed also in the  $^1\text{H-NMR}$  spectrum. The  $^{13}\text{C-NMR}$  spectrum (Table 1) showed the presence 32 carbons, which were assigned to

one ketone ( $\delta_C$  204.6), one ester ( $\delta_C$  170.1), one lactone ( $\delta_C$  178.6), four sp<sup>2</sup> carbons ( $\delta_C$  158.8, 157.9, 125.7 and 119.0), eight methyls ( $\delta_C$  27.1, 26.9, 23.8, 21.4, 21.2, 20.0, 19.1 and 16.2), six methylenes and seven methines. These properties represented a total of five or ten degrees of unsaturation, and the remaining five degrees of unsaturation corresponded to the pentacyclic framework of the apotirucallan-type triterpenoid.<sup>21</sup> The proton pairing was also confirmed with the  $^1\text{H-}^1\text{H}$  COSY spectrum (Fig. 2) and supported the presence of apotirucallan-type triterpenoid in **3**. The HMBC crosspeaks (Fig. 2) from H-2 ( $\delta_H$  5.86) to C-1 ( $\delta_C$  158.8) and carbonyl at C-3 ( $\delta_C$  204.6), indicated the presence of  $\alpha,\beta$ -unsaturated ketone at C-1, C-2 and C-3, respectively. Correlation which was arising from H-7 ( $\delta_H$  5.33) to C-1' ( $\delta_C$  170.1) and C-6 ( $\delta_C$  27.5) as well as from CH<sub>3</sub>-2" ( $\delta_H$  1.94) to C-1' ( $\delta_C$  170.1), indicate that acetyl group was attached at C-7. The correlation from H-15 ( $\delta_H$  5.28) to C-16 ( $\delta_C$  33.8) and C-14 ( $\delta_C$  157.9), indicated that another double-bond was located at C-14 and C-15, respectively. An oxygenated proton at H-23 ( $\delta_H$  4.47) and methine proton at H-17 ( $\delta_H$  2.19) was correlated to carbonyl at C-21 ( $\delta_C$  178.6), indicate the lactone ring was attached at C-17. Furthermore, the correlation from CH<sub>3</sub>-26 ( $\delta_H$  1.35) to C-24 ( $\delta_C$  72.4) and C-25 ( $\delta_C$  70.5), suggesting that 1,2-diol side chain was attached at C-23. The detailed examination of the NMR data and comparison with those reported for prototiamin A, an apotirucallane-type triterpenoid isolated from the bark of *Entandrophragma congoense*,<sup>21</sup> showed that the structures of the two compounds were very similar. Therefore, compound **3** was identified as prototiamin A.

Compound **4** was isolated as colorless solid, with the molecular formula determined as C<sub>30</sub>H<sub>44</sub>O<sub>7</sub> based on HRTOFMS ( $m/z$  517.4498 [M+H]<sup>+</sup>, (calcd. C<sub>30</sub>H<sub>44</sub>O<sub>7</sub>  $m/z$  516.4087) and NMR spectra (Table 1). In addition,  $^{13}\text{C-NMR}$  analysis show that nine degrees of unsaturation is required, while, the IR absorption bands at 3560, 2939, 1725 and 1109 cm<sup>-1</sup> imply the presence of hydroxyl, aliphatic, carbonyl ester and ether functionalities. The  $^1\text{H-NMR}$  spectrum showed the presence of five methyl tertiary signals at  $\delta_H$  1.16 (CH<sub>3</sub>-19), 1.16 (CH<sub>3</sub>-30), 1.06 (CH<sub>3</sub>-28), 1.06 (CH<sub>3</sub>-29), 1.00 (CH<sub>3</sub>-18) and two acetyl signals at  $\delta_H$  1.91 and 1.97. One olefinic proton at  $\delta_H$  5.28 (1H, dd,  $J=1.8, 3.6$  Hz, H-15) and three oxygenated methane signals at  $\delta_H$  [5.39 (1H, m, H-1), 5.33 (1H, m, H-7) and 4.57 (1H, dd,  $J=9.6, 1.8$  Hz, H-23)] were also observed in the  $^1\text{H-NMR}$  spectrum. Moreover, the  $^{13}\text{C-NMR}$  spectrum (Table 1) recognized 30 nonequivalent carbon signals, which are characterized by one carbonyl

ketone ( $\delta_C$  218.2), two acetyl ( $\delta_C$  170.1 and 170.0), two  $\text{sp}^2$  carbon ( $\delta_C$  158.9 and 118.0), seven methyl signals at ( $\delta_C$  27.5, 27.1, 21.4, 21.1, 21.0, 19.7 and 15.3), seven  $\text{sp}^3$  methylenes (including one oxygenated methylene type at  $\delta_C$  70.3), seven  $\text{sp}^3$  methynes (including two oxygenated types at  $\delta_C$  75.4, 75.6 and a hemiacetal at  $\delta_C$  99.8) and four quaternary carbons. These properties represented a total of four or nine degrees of unsaturation, and the remaining five degrees of unsaturation corresponded to limonoid framework.<sup>16,17,22</sup> Comprehensive analyses of the above mentioned  $^1\text{H}$  and  $^{13}\text{C}$  NMR data exhibited that the structure of **4** was similar to that of neemfrutin,<sup>22</sup> a limonoid isolated from *Azadirachta indica*. Consequently, compound **4** was identified as neemfrutin A.

Compound **5** was obtained as a white solid with a proposed molecular formula of  $C_{28}\text{H}_{40}\text{O}_5$  based on HRTOFMS ( $m/z$  455.0712 [ $\text{M}+\text{H}$ ] $^+$ , (calcd.  $C_{28}\text{H}_{40}\text{O}_5 m/z$  456.2876) and NMR data (Table 1), therefore nine indices of hydrogen deficiency were required. Furthermore, the IR absorption bands at 3300, 1705 and 1080  $\text{cm}^{-1}$  show the presence of hydroxyl, carbonyl and ether properties, while the NMR data was highly similar to **4**. However, differences were observed in the disappearance of one of acetyl group at [ $\delta_H$  1.91 (3H, s),  $\delta_C$  21.7, 170.1], oxygenated methyne at [ $\delta_H$  5.39 (1H, m),  $\delta_C$  75.4] and the appearance of olefinic signals at [ $\delta_H$  7.14 (1H, d,  $J=10.2$  Hz), 5.86 (1H, d,  $J=10.2$  Hz);  $\delta_C$  158.8, 125.7], suggesting **5** as a 1-deacetyl derivative of **4**. The proton pairing was also confirmed with the  $^1\text{H}$ - $^1\text{H}$  COSY spectrum (Fig. 2) and supported the presence of limonoid structure in **5**.<sup>22,23</sup> The HMBC crosspeaks (Fig. 2) from H-2 ( $\delta_H$  5.86) to C-1 ( $\delta_C$  158.8) and carbonyl at C-3 ( $\delta_C$  204.6), indicated the presence of  $\alpha,\beta$ -unsaturated ketone at C-1, C-2 and C-3, respectively. Correlation which was arising from H-7 ( $\delta_H$  5.33) to C-1' ( $\delta_C$  170.1) and C-6 ( $\delta_C$  27.5) as well as from  $\text{CH}_3$ -2" ( $\delta_H$  1.94) to C-1' ( $\delta_C$  170.1), indicate that acetyl group was attached at C-7. The above-mentioned NMR data suggested that compound **5** is a limonoid, for which NMR data (Table 1) showed many similarities to those of 23-desmethyllymocin-B, a limonoid isolated from the seed of *Azadirachta indica*.<sup>23</sup> Therefore, compound **5** was identified as 23-desmethyllymocin-B.

Compound **6** was obtained as a white solid with a molecular composition of  $C_{32}\text{H}_{48}\text{O}_6$ , based on HRTOFMS ( $m/z$  527.2984 [ $\text{M}+\text{H}$ ] $^+$ , (calcd.  $C_{32}\text{H}_{48}\text{O}_6 m/z$  528.2951) and NMR analysis, therefore nine degrees of unsaturation are required. Furthermore, the IR spectrum showed the absorption bands for hydroxyl (3420  $\text{cm}^{-1}$ ), aliphatic (2949 and 2860  $\text{cm}^{-1}$ ), carbonyl (1715  $\text{cm}^{-1}$ ) and ether (1080  $\text{cm}^{-1}$ ) moieties, while the NMR data observed in

**Table 2.** IC<sub>50</sub> inhibition values of compounds **1** - **8** against MCF-7 breast cancer cell line

Compounds	IC <sub>50</sub> ( $\mu\text{M}$ )
Cabrealeolactone ( <b>1</b> )	61.18
Cabreadiol ( <b>2</b> )	33.12
Prototiamin A ( <b>3</b> )	76.08
Neemfrutins A ( <b>4</b> )	181.12
Desmethyllymocin B ( <b>5</b> )	98.18
Protoxylocarpin G ( <b>6</b> )	90.24
Melianodiol ( <b>7</b> )	16.84
Indicalilacol B ( <b>8</b> )	20.23
Cisplatin*	13.20

\*positive control

Table 1 was highly similar to **5**. However, the difference was identified in the absence of a hemiacetal group at [ $\delta_H$  4.57 (1H, dd,  $J=9.6, 1.8$  Hz),  $\delta_C$  99.2], and also the appearance of a newly oxygenated methynes at [ $\delta_H$  4.47 (1H, dd,  $J=9.6, 1.8$  Hz), 3.92 (1H, d,  $J=9.6$  Hz),  $\delta_C$  78.2, 74.4], two methyls at [ $\delta_H$  1.35 (3H, s), 1.44 (3H, s),  $\delta_C$  26.9, 23.8] and one oxygenated quaternary carbon at  $\delta_C$  72.5. These observations suggest that **6** is a 2-methylbutane-2,3-diol derivative of **5** and is attached to C-23. The aforementioned data as well as biogenetic considerations suggested that compound **6** is protolimonoid, with a structure similar to that of protoxylocarpin G, a protolimonoid isolated from *Xylocarpus granatum*.<sup>24</sup> Consequently, compound **6** was identified as protoxylocarpin G.

Compound **7** was obtained as a colorless crystal. Its molecular composition  $C_{30}\text{H}_{48}\text{O}_5$ , was determined from HRTOFMS ( $m/z$  487.4535 [ $\text{M}+\text{H}$ ] $^+$ , (calcd.  $C_{32}\text{H}_{50}\text{O}_7 m/z$  488.4502) and NMR data (Table 1), therefore seven unsaturated degrees were required. The IR spectra showed absorption peaks at 3350  $\text{cm}^{-1}$  (OH), 2935  $\text{cm}^{-1}$  (C-H  $\text{sp}^3$ ), 1725 (C=O), 1645 (C=C), 1457 and 1380  $\text{cm}^{-1}$  (*gem*-dimethyl groups), and 1055  $\text{cm}^{-1}$  (C-O) groups. The  $^1\text{H}$ -NMR spectrum showed the presence of seven tertiary methyl groups at  $\delta_H$  1.39 ( $\text{CH}_3$ -27), 1.32 ( $\text{CH}_3$ -26), 1.01 ( $\text{CH}_3$ -30), 0.93 ( $\text{CH}_3$ -28), 0.92 ( $\text{CH}_3$ -29), 0.84 ( $\text{CH}_3$ -18) and 0.78 ( $\text{CH}_3$ -19). One olefinic proton at  $\delta_H$  5.26 (1H, dd,  $J=2.6, 5.8$  Hz, H-7), two oxygenated methyne signals at  $\delta_H$  3.64 (1H, ddd,  $J=1.3, 6.2, 8.7$  Hz, H-23), 4.06 (1H, br.s, H-24) and one hemiacetal signal at  $\delta_H$  4.63 (1H, d,  $J=5.6$  Hz) were also observed in the  $^1\text{H}$ -NMR spectrum. The  $^{13}\text{C}$ -NMR spectrum of **7** showed 30 carbons and were classified as seven methyl groups at  $\delta_C$  27.8, 27.4, 27.0, 26.5, 23.5, 21.6 and 13.1, one carbonyl ketone at  $\delta_C$  218.4 (C-3), two oxymethine signals at  $\delta_C$  86.4 (C-23) and 71.3 (C-24), one oxygenated quaternary

carbon at  $\delta_{\text{C}}$  70.3 (C-24), hemiacetal signal at  $\delta_{\text{C}}$  99.3 (C-21) and two sp<sup>2</sup> carbon at  $\delta_{\text{C}}$  118.8 (C-7) and 145.3 (C-8), respectively. These properties represented two of seven total degrees of unsaturation, and the remaining five degrees of unsaturation were consistent with the limonoid structure.<sup>24,25</sup> The above data showed general features similar to those of melianodiol, a limonoid isolated from the leaves of *Aglaia andamanica*.<sup>25</sup> Therefore, compound 7 was identified as melianodiol.

Compound 8 was obtained as a white solid having a molecular composition of C<sub>30</sub>H<sub>48</sub>O<sub>5</sub>, based on HRTOFMS (*m/z* 487.4535 [M+H]<sup>+</sup>, (calcd. C<sub>30</sub>H<sub>48</sub>O<sub>7</sub> *m/z* 488.4502) along with NMR analysis, therefore seven degrees of unsaturation are required. Furthermore, the IR spectrum showed the absorption bands for hydroxyl (3430 cm<sup>-1</sup>), aliphatic (2930), carbonyl (1760 cm<sup>-1</sup>), gem-dimethyl (1420 and 1378 cm<sup>-1</sup>) and ether (1080 cm<sup>-1</sup>) moieties, while the NMR data observed in Table 1 was highly similar to 7. However, the difference was identified in the absence of a hemiacetal signal at [ $\delta_{\text{H}}$  4.63 (1H, d, *J*=5.6 Hz),  $\delta_{\text{C}}$  99.3] and ketone signal at  $\delta_{\text{C}}$  218.4, and appearance of a newly carbonyl lactone at  $\delta_{\text{C}}$  178.3 (C-21) and hydroxyl signal at [ $\delta_{\text{H}}$  4.32 (1H, d, *J*=5.4 Hz),  $\delta_{\text{C}}$  72.4]. These observations indicate that 8 is 3-hydroxy and lactone derivative of 7. The above data revealed that 8 is a trirucallane-type triterpenoid similar to indicalilacol B, isolated from *Azadiracta indica*.<sup>26</sup> Consequently, compound 8 was identified as indicalilacol B.

The isolated compounds, 1 - 8, were evaluated for their cytotoxic against MCF-7 breast cancer lines using a previously described method,<sup>17,18,27</sup> using *cis*-Platin as a positive control.<sup>28,29</sup> Among the compounds tested, melianodiol (7), exhibited the strongest cytotoxicity with an IC<sub>50</sub> value of 16.84  $\mu\text{M}$ . Interestingly, Compounds 7 and 8 share the same structure and only differed at the functional of the cyclopentane ring, the cytotoxic activities of compounds 7 and 8 differed greatly, suggested the presence of hemiacetal group increase the cytotoxic activity. These results were supported from previously studies that the presence of a hemiacetal group in triterpenoids structure can increase cytotoxic activity.<sup>17,24,25</sup> In addition, despite close structural similarities to 7, a low activity was observed in 6, which indicates the possible contribution of an acetyl group to cytotoxicity.<sup>25</sup> This results was supported from previously studies that the presence of a hemiacetal group in triterpenoids structure can increase cytotoxic activity.<sup>25</sup> Therefore, the structural comparison of compound 7 with the other compounds indicates that carbonyl, hydroxyl and acetyl groups present contribute to the cytotoxic activity.

## Acknowledgments

This investigation was financially supported by Universitas Brawijaya, Indonesia (PPKI Grant, No: 455.5/UN10.C10/PN/2020 by RR); (Universitas Padjadjaran, Indonesia (PPKI Grant, No: 1427/UN6.3.1/LT/2020 by US) and Universitas Airlangga, Indonesia (PPKI Grant No: 305/UN3.14/PT/2020 by MT).

## References

- (1) Katja, D. G; Farabi, K.; Nurlelasari.; Hameti, D.; Mayanti, T.; Supratman, U.; Awang, K.; Hayashi, H. *J. Asian Nat. Prod. Res.* **2017**, *19*, 194-200.
- (2) Shilpi, J. A.; Saha, S.; Chong, S. L.; Nahar, L.; Sarker, S. D.; Awang, K. *Chem. Biodivers.* **2016**, *13*, 483-503.
- (3) Awang, K.; Lim, C. S.; Mohamad, K.; Hiroshi, M.; Hirasawa, Y.; Takeya, K.; Thoison, O.; Hadi, A. H. A. *Bioorg. Med. Chem.* **2007**, *15*, 5997-6002.
- (4) Tan, Q. G.; Luo, X. D. *Chem. Rev.* **2011**, *111*, 7437-7522.
- (5) Yang, M. H.; Wang, J. S.; Luo, J. G.; Wang, X. B.; Kong, L. Y. *J. Nat. Prod.* **2009**, *72*, 2014-2018.
- (6) Mohamad, K.; Hirasawa, Y.; Litaudon, M.; Awang, K.; Hadi, A. H. A.; Takeya, K.; Ekasari, W.; Widyawaruyanti, A.; Zaini, N. C.; Morita, H. *Bioorg. Med. Chem.* **2009**, *17*, 727-730.
- (7) Supratman, U.; Naibaho, W.; Salam, S.; Maharani, R.; Hidayat, A. T.; Hameti, D.; Nurlelasari.; Shiono, Y. *Phytochem. Lett.* **2019**, *30*, 81-87.
- (8) Chong, S. L.; Hematpoor, A.; Hazni, H.; Sofian-Azirun, M.; Litaudon, M.; Supratman, U.; Murata, M.; Awang, K. *Phytochem. Lett.* **2019**, *30*, 69-73.
- (9) Najmuldeen, I. A.; Hadi, A. H. A.; Mohamad, K.; Awang, K.; Ketuly, K. A.; Mukhtar, M. R.; Taha, H.; Nordin, N.; Litaudon, M.; Guérinne, F.; Nugroho, A. E.; Morita, H. *Heterocycles* **2012**, *84*, 1265-1270.
- (10) Fang, X.; Di, Y. T.; Hao, X. *J. Current Organic Chemistry* **2011**, *15*, 1363-1391.
- (11) Haldar, S.; Kolet, S. P.; Thulasiram, H. V. *Green Chemistry* **2013**, *15*, 1311-1317.
- (12) Nagoor, N. H.; Mutiah, N. S. J.; Lim, C. S.; In, L. L. A.; Mohamad, K.; Awang, K. *PLoS One*. **2011**, *6*, e23661.
- (13) Wong, C. P.; Shimada, M.; Nagakura, Y.; Nugroho, A. E.; Hirasawa, Y.; Kaneda, T.; Awang, K.; Hadi, A. H. A.; Mohamad, K.; Shiro, M.; Morita, H. *Chem. Pharm. Bull.* **2011**, *59*, 407-411.
- (14) Yang, M. H.; Wang, J. S.; Luo, J. G.; Wang, X. B.; Kong, L. Y. *Bioorg. Med. Chem.* **2011**, *19*, 1409-1417.
- (15) Bordoloi, M.; Saikia, B.; Mathur, R. K.; Goswami, B. N. *Phytochemistry* **1993**, *34*, 583-584.
- (16) Nurlelasari.; Katja, D. G.; Harneti, D.; Wardayo, M. M.; Supratman, U.; Awang, K. *Chem. Nat. Compd.* **2017**, *53*, 83-87.
- (17) Supriatno.; Nurlelasari.; Herlina, T.; Hameti, D.; Maharani, R.; Hidayat, A. T.; Mayanti, T.; Supratman, U.; Azmi, M. N.; Shiono, Y. *Nat. Prod. Res.* **2018**, *25*, 1-7.
- (18) Supratman, U.; Salam, S.; Naibaho, W.; Fajar, M.; Nurlelasari.; Katja, D. G.; Harneti, D.; Maharani, R.; Hidayat, A. T.; Lesmana, R.; Nafiah, M. A.; Shiono, Y. *Phytochem. Lett.* **2020**, *35*, 63-67.
- (19) Nagaya, H.; Tobita, Y.; Nagae, T.; Itokawa, H.; Takeya, K.; Halim, A. F.; Abdel-Halim, O. B. *Phytochemistry* **1997**, *44*, 1115-1119.
- (20) Phongmaykin, J.; Kumamoto, T.; Ishikawa, T.; Sutisri, R.; Saifah, E. *Arch. Pharm. Res.* **2008**, *31*, 21-27.
- (21) Happi, G. M.; Kouam, S. F.; Talontsi, F. M.; Lamshöft, M.; Zühlke,

- S.; Bauer, J. O.; Strohmann, C.; Spitteler, M. *J. Nat. Prod.* **2015**, *78*, 604-614.
- (22) Chianese, G.; Yerbanga, S. R.; Lucantoni, L.; Habluelzel, A.; Basilico, N.; Taramelli, D.; Fattorusso, E.; Taglialatela-Scafati, O. *J. Nat. Prod.* **2010**, *73*, 1448-1452.
- (23) Kumar, C. S. S. R.; Srinivas, M.; Yakkundi, S. *Phytochemistry* **1996**, *43*, 451-455.
- (24) Pudhom, K.; Sommit, D.; Nuclear, P.; Ngamrojanavanich, N.; Petsom, A. *J. Nat. Prod.* **2009**, *72*, 2188-2192.
- (25) Puripattanavong, J.; Weber, S.; Brecht, V.; Frahm, A. W. *Planta Med.* **2000**, *66*, 740-745.
- (26) Kurimoto, S.; Takaishi, Y.; Ahmed, F. A.; Kashiwada, Y. *Fitoterapia* **2014**, *92*, 200-205.
- (27) Skehan, P.; Storeng, R.; Scudiero, D.; Monks, A.; McMahon, J.; Vistica, D.; Warren, J. T.; Bokesch, H.; Kenney, S.; Boyd, M. R. *J. Natl. Cancer Inst.* **1990**, *82*, 1107-1112.
- (28) Hadisaputri, Y. E.; Miyazaki, T.; Suzuki, S.; Yokobori, T.; Kobayashi, T.; Tanaka, N.; Inose, T.; Sohda, M.; Kuwano, H. *Ann. Surg. Oncol.* **2012**, *19*, S589-S596.
- (29) Chavoshi, H.; Vahedian, V.; Saghaei, S.; Pirouzpanah, M. B.; Raeisi, M.; Samadi, N. *Asian Pac. J. Cancer Prev.* **2017**, *18*, 2243-2247.

Received November 17, 2020

Revised February 3, 2021

Accepted February 9, 2021

# Cytotoxic Constituents from the Stem Bark of Chisocheton pentandrus

ORIGINALITY REPORT



PRIMARY SOURCES

1	<a href="http://www.tandfonline.com">www.tandfonline.com</a>	1 %
2	<a href="http://mdpi.com">mdpi.com</a>	1 %
3	<a href="http://applbiolchem.springeropen.com">applbiolchem.springeropen.com</a>	1 %
4	<a href="http://coek.info">coek.info</a>	1 %
5	<a href="http://www.e-sciencecentral.org">www.e-sciencecentral.org</a>	1 %
6	<a href="http://docplayer.net">docplayer.net</a>	1 %
7	<a href="http://doaj.org">doaj.org</a>	1 %
8	<a href="http://www.thieme-connect.de">www.thieme-connect.de</a>	1 %
9	Nenajdenko, V.G.. "The Ugi reaction with 2-substituted cyclic imines. Synthesis of	1 %

substituted proline and homoproline derivatives", Tetrahedron, 20060619

Publication

- 
- 10 [www.fedoa.unina.it](http://www.fedoa.unina.it) 1 %  
Internet Source
- 
- 11 Unang Supratman, Supriatno Salam, Wiro Naibaho, Mohamad Fajar et al. "New cytotoxic limonoids from the stem bark of Chisocheton pentandrus (Blanco) Merr", Phytochemistry Letters, 2020 1 %  
Publication
- 
- 12 [arkat-usa.org](http://arkat-usa.org) 1 %  
Internet Source
- 
- 13 [journals.plos.org](http://journals.plos.org) 1 %  
Internet Source
- 
- 14 [patents.google.com](http://patents.google.com) 1 %  
Internet Source
- 
- 15 [mafiadoc.com](http://mafiadoc.com) 1 %  
Internet Source
- 
- 16 Ace Tatang Hidayat, Kindi Farabi, Desi Harneti, Rani Maharani et al. " Cytotoxicity and Structure Activity Relationship of Dammarane-Type Triterpenoids from the Bark of against P-388 Murine Leukemia Cells ", Natural Product Sciences, 2017 <1 %  
Publication

17	edoc.ub.uni-muenchen.de Internet Source	<1 %
18	www.mdpi.com Internet Source	<1 %
19	jurnal.ugm.ac.id Internet Source	<1 %
20	link.springer.com Internet Source	<1 %
21	core.ac.uk Internet Source	<1 %
22	pubs.rsc.org Internet Source	<1 %
23	jtpc.farmasi.unmul.ac.id Internet Source	<1 %
24	Hongquan Duan, Yoshihisa Takaishi, Hiroshi Momota, Yasukazu Ohmoto et al. "Immunosuppressive terpenoids from extracts of <i>Tripterygium wilfordii</i> ", Tetrahedron, 2001 Publication	<1 %
25	patents.justia.com Internet Source	<1 %
26	ir-library.ku.ac.ke Internet Source	<1 %

27	<a href="http://www.j3.jstage.jst.go.jp">www.j3.jstage.jst.go.jp</a> Internet Source	<1 %
28	<a href="http://thieme-connect.com">thieme-connect.com</a> Internet Source	<1 %
29	<a href="http://jurnal.uns.ac.id">jurnal.uns.ac.id</a> Internet Source	<1 %
30	<a href="http://pubs.acs.org">pubs.acs.org</a> Internet Source	<1 %
31	<a href="http://eprints.usm.my">eprints.usm.my</a> Internet Source	<1 %
32	<a href="http://mdpi.org">mdpi.org</a> Internet Source	<1 %
33	<a href="http://www.yumpu.com">www.yumpu.com</a> Internet Source	<1 %
34	Jarinporn Phongmaykin. "A new sesquiterpene and other terpenoid constituents of <i>Chisocheton penduliflorus</i> ", Archives of Pharmacal Research, 01/2008 Publication	<1 %
35	Phuwapraisirisan, P.. "Feroniellides A and B, apotirucallane triterpenes with novel cyclic acetals from <i>Feroniella lucida</i> ", Tetrahedron Letters, 20070122 Publication	<1 %
36	<a href="http://edepot.wur.nl">edepot.wur.nl</a> Internet Source	

<1 %

- 
- 37 Kulcitki, V.. "Ring B functionalization of scalarane sesterterpenes by radical relay halogenation", Tetrahedron, 20070806  
Publication <1 %
- 
- 38 cmuj.cmu.ac.th Internet Source <1 %
- 
- 39 sphinxsai.com Internet Source <1 %
- 
- 40 www.biotech-asia.org Internet Source <1 %
- 
- 41 baadalsg.inflibnet.ac.in Internet Source <1 %
- 
- 42 www.freepatentsonline.com Internet Source <1 %
- 
- 43 acgpubs.org Internet Source <1 %
- 
- 44 espace.library.uq.edu.au Internet Source <1 %
- 
- 45 hoshi.academia.edu Internet Source <1 %
- 
- 46 ir.scsio.ac.cn Internet Source <1 %
-

47

rd.springer.com

Internet Source

<1 %

48

www.arkat-usa.org

Internet Source

<1 %

49

Ojika, M.. "Aplyronine A, a potent antitumor macrolide of marine origin, and the congeners aplyronines B and C: isolation, structures, and bioactivities", Tetrahedron, 20070409

Publication

<1 %

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

On

# Cytotoxic Constituents from the Stem Bark of Chisocheton pentandrus

---

## GRADEMARK REPORT

---

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---