

# Chromanone Acid Derivatives from the Stem Bark of *Calophyllum incrassatum*

*by* Mulyadi Tanjung

---

**Submission date:** 05-Aug-2021 01:12PM (UTC+0800)

**Submission ID:** 1627933149

**File name:** Hasanah\_2019\_IOP\_Conf.\_Ser.\_Earth\_Environ.\_Sci.\_217.pdf (217.74K)

**Word count:** 2425

**Character count:** 11130

PAPER · OPEN ACCESS

## Chromanone Acid Derivatives from the Stem Bark of *Calophyllum incrassatum*

To cite this article: U Hasanah *et al* 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **217** 012010



View the [article online](#) for updates and enhancements.



## Chromanone Acid Derivatives from the Stem Bark of *Calophyllum incrassatum*

U Hasanah<sup>1</sup>, T S Tjahjandarie<sup>1</sup>, M Tanjung<sup>1</sup>

<sup>1</sup>Natural Products Chemistry Research Group, Organic Chemistry Division, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia.

ulfalunks15@gmail.com; mulyadi-t@fst.unair.ac.id

**Abstract.** Two chromanone acid derivatives, calofolic acid B (**1**) and apetalic acid (**2**) were isolated from the stem bark of *Calophyllum incrassatum*. The structures of both compounds were elucidated by UV, IR, HRESIMS, 1D and 2D NMR spectroscopies and comparison of their data with literatures. Compounds **1-2** were evaluated for their cytotoxic activity against P-388 cells, compound (**1**) showing value IC<sub>50</sub> 1.14 µg/mL.

**Keywords:** *Calophyllum incrassatum*, Calofolic acid B, Apetalic acid, P-388 cells.

### 1. Introduction

The genus *Calophyllum* (Calophyllaceae) comprises approximately 150 species that grow in the restrictive area of South East Asia. In Indonesia, the plants has many advantages, used as traditional treatment for rheumatism, inflammation, eye disease, and skin disease [1]. This genus is well known to produce a variety secondary metabolites such as xanthenes [2], 4-phenylcoumarins [3], benzofuran-3-ones [4], biflavonoids [5] and chromanone acid [6-8], which have some biological activities such as anticancer, antibactery, and antitumor. Xanthone and 4-phenylcoumarin derivatives containing a isoprenyl side chain in aromatic region revealed as a key element to enhance their biological activities. In continuation of our work on phytochemical examination of chromanone acid on *Calophyllum* from West Kalimantan, we report the isolation of calofolic acid B (**1**), and apetalic acid (**2**) from the methanol extract of the stem bark of *Calophyllum incrassatum*. Compounds **1** and **2** were established by UV, IR, HRESIMS, 1D and 2D NMR, as well as by comparison with those related compounds previously reported. The cytotoxic activity against murine leukemia P-388 cells of isolated compounds from this species is also briefly described.

### 2. Experimental Methode

#### 2.1 General experimental

UV spectra was measured by Shimadzu series 1800 UV-VIS spectrophotometer with methanol (Kyoto, Japan). IR spectra was recorded in KBr on a One Perkin Elmer instrument (Waltham, MA, USA). Mass spectra was recorded on Synapt G2 mass spectrometer (Manchester, UK). NMR spectra (<sup>1</sup>H-NMR, <sup>13</sup>C-NMR, HMQC, and HMBC) was obtained with a JEOL JNM-ECA 400 MHz FTNMR spectrophotometer (Tokyo, Japan) with CDCl<sub>3</sub> as the solvent and internal standard. Column chromatography and planar radial chromatography was carried out using Si gel 60 Cat. No. 1.07734.1000 and Si gel 60 PF<sub>254</sub> No. 1.07749.1000 (Merck, Darmstadt, Germany).

#### 2.2 Plant Material

The stem bark of *C. incrassatum* was collected from Mandor sub-district, Landak regency, West Kalimantan, Indonesia. The sampel was identified by Mr. Ismail Rachman from the Herbarium Bogoriense, Bogor, Center of Biological Research and Development.



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

### 2.3 Extraction and isolation

The dried stem bark of *C. incrassatum* (3.0 kg) was macerated in methanol twice at room temperature for 24 h. The solvent was evaporated under reduced pressure to obtain a dark brown residue (280 g). The methanol extract was then partitioned with *n*-hexane and ethyl acetate. The ethyl acetate fraction (43 gr) was subjected to vacuum liquid chromatography over silica gel and eluted with *n*-hexane-ethyl acetate (from 9:1 to 7:3) to give fractions A-C. Fraction C (5.44 g) was applied to column chromatography and eluted with *n*-hexane-ethyl acetate (from 9:1 to 8:2) to produce subfractions C<sub>1</sub>-C<sub>2</sub>. Subfraction C<sub>2</sub> (1.2 g) was purified by planar radial chromatography using *n*-hexane-chloroform (from 9:1 to 3:7) yielded compound **1** (76 mg).

The *n*-hexane fraction (30 gr) was subjected to vacuum liquid chromatography over silica gel and eluted with *n*-hexane-ethyl acetate (from 9:1 to 8:2) to give fractions D and E. Fraction D (20 g) was refractionated using column chromatography and eluted with *n*-hexane-ethyl acetate (from 9:1 to 8:2) gave sub-fractions D<sub>1</sub>-D<sub>2</sub>. Subfraction D<sub>1</sub> (2.84 g) was purified by planar radial chromatography using *n*-hexane-ethyl acetate (9:1) to yielded sub-fraction D<sub>11</sub>-D<sub>13</sub>. Sub-fraction D<sub>12</sub> (487 mg) was purified by planar radial chromatography using *n*-hexane-ethyl acetate (9:1) to give D<sub>121</sub>-D<sub>125</sub>. Subfraction D<sub>125</sub> (227.2 mg) was purified by planar radial chromatography using *n*-hexane-ether (9:1) to yielded compound **2** (30 mg).

### 2.4 Cytotoxic assay

Compounds **1**-**2** were tested to *in vitro* evaluating against murine leukemia P-388 cells by using MTT colorimetric method as previously reported [9-11]. Artonin E was used as the positive control.

## 3. Result and Discussion

**Calofolic acid B (1)**, a yellow solid, m.p. 117-118 °C, showed a quasimolecular ion [M+H]<sup>+</sup> at *m/z* 359.1538 corresponding to the molecular formula C<sub>20</sub>H<sub>24</sub>O<sub>6</sub>. The UV absorption at λ<sub>max</sub> (nm) (log ε) = 225 (3.35), 274 (3.89), 289 (3.27), 305 (3.33), 312 (3.35), and 338 (2.60) typical for a chromanone acid [5]. The IR spectrum of **1** indicated absorptions for hydroxyl (3423 cm<sup>-1</sup>), conjugated carbonyl (1620 cm<sup>-1</sup>), aromatic (1577 and 1545 cm<sup>-1</sup>) and ether (1188 cm<sup>-1</sup>) groups, respectively. The <sup>1</sup>H NMR spectrum of **1** (Table 1) showed the presence of a 2,3-dimethylchromanone ring which represent by a pair of methine signals [δ<sub>H</sub> 4.51 (1H, *dq*, *J* = 6.5 and 3.2 Hz) and 2.55 (1H, *dq*, *J* = 7.2 and 3.2 Hz)] and two proton signals of methyl [δ<sub>H</sub> 1.35 (3H, *d*, *J* = 6.5 Hz) and δ<sub>H</sub> 1.13 (3H, *d*, *J* = 7.3 Hz)], a chelated hydroxyl group (δ<sub>H</sub> 12.37 (*s*, 5-OH)), a dimethylpyrano ring group [δ<sub>H</sub> 6.60 (1H, *d*, *J* = 10.0 Hz), δ<sub>H</sub> 5.46 (1H, *d*, *J* = 10.0 Hz), and *gem* dimethyl group at δ<sub>H</sub> 1.38 (3H, *s*), δ<sub>H</sub> 1.45 (3H, *s*)], and an alkyl carboxylic group consist of a methine [δ<sub>H</sub> 3.77 (1H, *m*)], a methylene [δ<sub>H</sub> 2.67 (1H, *dd*, *J* = 7.4, 15.2 Hz), δ<sub>H</sub> 2.82 (1H, *dd*, *J* = 8.3, 15.2 Hz)], and a methyl [δ<sub>H</sub> 1.26 (3H, *d*, *J* = 7.2 Hz)]. The <sup>13</sup>C NMR spectrum of **1** showed 20 carbon signals and the structure of **1** were confirmed by HMQC and HMBC spectra (Figure 1). The <sup>13</sup>C NMR spectrum of **1** corresponding to the chromanone acid. The 1D and 2D NMR spectra data are consistent with the previous published data [8].

**Apetalic acid (2)**, a yellow solid, m.p. 117-118 °C, [α]<sub>D</sub><sup>20</sup> +56° (*c* = 0.0005, MeOH). The UV spectrum λ<sub>max</sub> (nm) (log ε) = 227 (3.39), 274 (3.96), 300 (3.41), 305 (3.39), 311 (3.41) and 336 (2.48) and IR spectrum gave absorptions of 3436, 1622, 1598, 1496, 1446 and 1157 cm<sup>-1</sup> similar to those of **1**. The <sup>1</sup>H NMR spectrum of **2** very similar with **1** and the main difference is alkyl carboxylic group at C-10. The <sup>1</sup>H NMR spectrum of **2** (Table 1) showed the presence of a 2,3-dimethylchromanone ring consist of a pair of methine [δ<sub>H</sub> 4.50 (1H, *dq*, *J* = 6.5 and 3.2 Hz) and 2.53 (1H, *dq*, *J* = 7.3 and 3.2 Hz)] and two proton signals of methyl [δ<sub>H</sub> 1.35 (3H, *d*, *J* = 6.6 Hz) and δ<sub>H</sub> 1.13 (3H, *d*, *J* = 7.2 Hz)], a chelated hydroxyl group (δ<sub>H</sub> 12.39, *s*, 5-OH), a dimethylpyrano ring group [δ<sub>H</sub> 6.60 (1H, *d*, *J* = 10.0 Hz), δ<sub>H</sub> 5.45 (1H, *d*, *J* = 10.0 Hz), and *gem* dimethyl group δ<sub>H</sub> 1.37 (3H, *s*), δ<sub>H</sub> 1.44 (3H, *s*)], and an alkyl carboxylic group consist of a methine [δ<sub>H</sub> 3.68 (1H, *m*)], three methylene [δ<sub>H</sub> 2.66 (1H, *dd*, *J* = 6.9, 15.2 Hz), δ<sub>H</sub> 2.82 (1H, *dd*, *J* = 8.7, 15.2 Hz), δ<sub>H</sub> 1.54 (1H, *m*), δ<sub>H</sub> 1.82 (1H, *m*), δ<sub>H</sub> 1.19 (2H, *m*)], and a methyl [δ<sub>H</sub> 0.85 (3H, *t*, *J* = 7.3 Hz)]. The <sup>13</sup>C NMR (Table.1) of **2** showed 22 carbon signals. The assignment of <sup>13</sup>C NMR spectrum was

confirmed by HMQC and HMBC spectra (Figure 1). The 1D and 2D NMR spectra data are consistent with those of published data [12].

<sup>22</sup> Table-1. NMR spectroscopic data of compounds **1-2** in CDCl<sub>3</sub>

No	1 Calofolic acid B (1)			Apetalic acid (2)		
	$\delta$ u (mult, J in Hz)	$\delta$ c	HMBC	$\delta$ u (mult, J in Hz)	$\delta$ c	HMBC
1	-	-	-	-	-	-
2	4.51 (dq, 3.2; 6.5)	76.1	C-4; C-16	4.50 (dq, 3.2; 6.5)	76.1	C-4; C-16
3	2.55 (dq, 3.2; 7.2)	44.2	C-4; C-16	2.53 (dq, 3.2; 7.3)	44.3	C-4; C-16
4	-	20.1	-	-	20.1	-
5	-	15.7	-	-	15.7	-
6	6.60 (d, 10.0)	11.5	C-8; C-13; C-14	6.60 (d, 10.0)	11.5	C-8; C-13; C-14
7	5.46 (d, 10.0)	12.5	C-8; C-13; C-17; C-18	5.45 (d, 10.0)	12.7	C-8; C-13; C-17; C-18
8	-	78.2	-	-	78.2	-
9	-	-	-	-	-	-
10	-	11.0	-	-	10.8	-
11	-	15.9	-	-	15.7	-
12	-	10.4	-	-	10.3	-
13	-	10.2	-	-	10.6	-
14	-	15.9	-	-	15.8	-
15	1.35 (d, 6.5)	16.2	C-2; C-3	1.35 (d, 6.6)	16.3	C-2; C-3
16	1.13 (d, 7.3)	9.4	C-2; C-3; C-4	1.13 (d, 7.2)	9.4	C-2; C-3; C-4
17	1.38 (s)	28.1	C-7; C-8; C-18	1.44 (s)	28.6	C-7; C-8; C-18
18	1.45 (s)	28.5	C-7; C-8; C-17	1.37 (s)	28.1	C-7; C-8; C-17
19	3.77 (m)	25.6	C-10; C-14; C-20; C-22	3.68 (m)	30.2	C-10; C-14; C-20; C-21; C-22
20	2.67 (dd, 7.4;	39	C-10; C-19; C-	2.66 (dd, 6.9; 15.2)	38	C-10; C-19;

	15.2) 2.82 ( <i>dd</i> , 8.3; 15.2)	.5	21; C-22	2.82 ( <i>dd</i> , 8.7; 15.2)	.6	C-21; C-22
21	-	17 9. 6	-	-	17 9. 5	-
22	1.26 ( <i>d</i> , 7.2)	19 .3	C-10; C-19; C-20	1.54 ( <i>m</i> ) 1.82 ( <i>m</i> )	35 .5	C-10; C-19; C-20; C-23; C-24
23	-	-	-	1.19 ( <i>m</i> )	20 .8	C-19; C-22; C-24
24	-	-	-	0.85 ( <i>t</i> , 7.3)	14 .1	C-22; C-23
5-OH	12.37 ( <i>s</i> )	-	C-5; C-12; C-13	12.39	-	C-5; C-12; C-13




Figure 1. HMBC correlations for compounds 1 and 2

The cytotoxic activity of compounds 1-2 was evaluated for their cytotoxicity by MTT assay against murine leukemia P-388 (Table 2). Artonin E was used as the positive control. Those cytotoxic data for chromanone acid suggested that compound 1 has active and compound 2 was inactive.

Table-2. Cytotoxic activity data of compounds 1-2 by MTT assay.

N o	Compound	IC <sub>50</sub> (µg/mL)
1	Calofolic acid B (1)	1.14
2	Apetalic acid (2)	> 100
3	Artonin E	1.33

#### 4. Conclusions

Chemical study on the stem bark of *C. incrasatum* yielded two chromanone acid derivatives, calofolic acid B (1) and apetalic acid (2). Calofolic acid B (1) showed very active against murine leukemia P-388 cells.

#### References

- [1] Heyne, K., 1987, The useful Indonesian plants research and development agency. Jakarta: Ministry of forestry.

- [2] Kar, W.L., Zamakshshari, N.H., Ee, G.C.L., Mah, S.H., and Mohd, N.S.M., 2018, Isolation and structural modification of ananixanthone from *Calophyllum teysmannii* and their cytotoxic activities, *J. Nat. Prod.*, **32**(18), 2147-2151.
- [3] Daud, S.B., Ee G.C.L., Malek, E.A., Teh, S.S., See, I., 2014, A new coumarin from *Calophyllum hosei*, *Nat. Prod. Res.*, **28**(19), 1534-1538.
- [4] Tanjung, M., Rachmadiarti, F., Prameswari, A., Agyani, V.U.W., Saputri, R.D., Tjahjandarie, T.S., and Syah, Y.M. 2018 Airlangin A-B, two new isoprenylated benzofuran-3-ones from the stem bark of *Calophyllum soulattri*, *Nat. Prod. Res.*, **32**(13), 1493-1498.
- [5] Ferchichi, L., Derbre, S., Mahmood, K., Toure, K., Guilet, D., Litaudon, M., Awang, K., Hadi, A.H., A., Le Ray, A.M., Richomme, P. 2012, Bioguided fraction and isolation of natural inhibitors of advance glycation end-products (AGEs) from *Calopyllum flavoramulum* *Phytochem.*, **78**, 98-106.
- [6] Lim, C.K., Subramaniam, H., Say, Y.H., Jong, V.Y., Khaledi, H., Chee, C.F., 2015, A New chromanone acid from the stem bark of *Calophyllum teysmannii*, *Nat. Prod. Res.*, **29** (21) 1970–1977.
- [7] Ha, L.D., Hansen, P. E., Duus, F., Pham, H.D., Nguyen, L.D., 2012, A New chromanone acid from the bark of *Calophyllum dryobalanoides*, *Phytochem. Lett.*, **5**(2), 287-291.
- [8] Nugroho, A.E., Sasaki, T., Kaneda, T., Hadi, A.H.A., Morita, H., 2017 Calofolic acids A-F chromanones from the bark of *Calophyllum scriblitifolium* with vasorelaxation activity, *Bioorg. Med. Chem. Lett.*, **27**(10), 2124-2128.
- [9] Tanjung, M., Hakim, E.H., Syah, Y.M., 2017, Prenylated dihydrostilbenes from *Macaranga rubiginosa*. *Chem Nat. Compd.*, **53**(2), 215-218.
- [10] Tanjung, M., Saputri, R.D., Wahjoedi, R.A., Tjahjandarie, T.S., 2017, 4-Methoxy-3-(3-methylbut-2-en-1-yl)-7-((3-methylbut-2-en-1-yl)oxy) quinolin-2(1H)-one from *Melicope moluccana* T.G. Hartley, *Molbank*, **2**, M939.
- [11] Tanjung, M., Rachmadiarti, F., Saputri, R.D., Tjahjandarie, T.S., 2018, Mesucalophylloidin, a new isoprenylated 4-phenylcoumarin from *Mesua calophylloides* (Ridl.) Kosterm., *Nat. Prod. Res.*, **32**(9), 1062-1067.
- [12] Shen, Y., Wang, L., Khalil, A. T., Kuo, Y. 2004 Chromanones and dihydrocoumarins from *Calophyllum blancoi* *Chem. Pharm. Bull* **52** (4) 402-405.

# Chromanone Acid Derivatives from the Stem Bark of *Calophyllum incrassatum*

---

## ORIGINALITY REPORT

---

19%

SIMILARITY INDEX

10%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1 Maneerat, Wisanu, Thunwadee Ritthiwigrom, Sarot Cheenpracha, Trinop Promgool, Kulsiri Yossathera, Suwanna Deachathai, Wong Phakhodee, and Surat Laphookhieo. "Bioactive Carbazole Alkaloids from *Clausena wallichii* Roots", *Journal of Natural Products*, 2012.  
Publication 1%
  - 2 [www.lens.org](http://www.lens.org)  
Internet Source 1%
  - 3 Susilawati, Yasmiwar, Ricky Nugraha, Ahmad Muhtadi, Supriyatna Soetardjo, and Unang Supratman. "(S)-2-Methyl-2-(4-methylpent-3-enyl)-6-(propan-2-ylidene)-3,4,6,7-tetrahydropyrano[4,3-g]chromen-9(2H)-one", *Molbank*, 2015.  
Publication 1%
  - 4 [www.google.co.in](http://www.google.co.in)  
Internet Source 1%
-



5	<p>Si-Min Kim. "DIASTEREOSELECTIVE ALKYLATION AND ALDOL REACTIONS MEDIATED BY THE d-MANNITOL-DERIVED OXAZOLIDIN-2-ONE AS A CHIRAL AUXILIARY", <i>Synthetic Communications</i>, 2002</p> <p>Publication</p>	1 %
6	<p><a href="http://www.allfordrugs.com">www.allfordrugs.com</a></p> <p>Internet Source</p>	1 %
7	<p>Hai-Qiang Huang, Xi Zhang, Yun-Heng Shen, Juan Su, Xiao-Hua Liu, Jun-Min Tian, Sheng Lin, Lei Shan, Wei-Dong Zhang. " Polyacetylenes from ", <i>Journal of Natural Products</i>, 2009</p> <p>Publication</p>	1 %
8	<p>A Cezairliyan, F Righini. "Issues in high-speed pyrometry", <i>Metrologia</i>, 1996</p> <p>Publication</p>	1 %
9	<p>Anelise S. Nazari, Suziane A. Dias, Willian F. da Costa, Ciomar A. Bersani-Amado et al. " Anti-inflammatory and Antioxidant Activities of . and Major Constituents ", <i>Pharmaceutical Biology</i>, 2008</p> <p>Publication</p>	1 %
10	<p>Hirromichi Fujioka. "Concise Asymmetric Total Synthesis of Scyphostatin, a Potent Inhibitor of Neutral Sphingomyelinase", <i>Chemistry - A European Journal</i>, 12/17/2007</p> <p>Publication</p>	1 %

11 Eun-Kyoung Seo, Monroe E. Wall, Mansukh C. Wani, Hernan Navarro et al. "Cytotoxic constituents from the roots of *Tovomita brevistaminea*", *Phytochemistry*, 1999  
Publication 1 %

---

12 Shigetada Kozai, Aya Yorikane, Tokumi Maruyama. " A NEW METHOD FOR THE SYNTHESIS OF -ALKYLGUANOSINES USING MITSUNOBU REACTION AS A KEY STEP ", *Nucleosides, Nucleotides and Nucleic Acids*, 2001  
Publication 1 %

---

13 [www.acgpubs.org](http://www.acgpubs.org)  
Internet Source 1 %

---

14 M. Mostafa, Nilufar Nahar, M. Mosihuzzaman, Selestin D. Sokeng, Naheed Fatima, Atta-ur-Rahman, M. Iqbal Choudhary. " Phosphodiesterase-I inhibitor quinovic acid glycosides from ", *Natural Product Research*, 2006  
Publication 1 %

---

15 Wei Yuan†, Huixin Zhu†, Kedi Cheng, Zhentai Huang, Yong Qin, Jinling Yang, Ping Zhu. " A novel butenoate derivative from ", *Natural Product Research*, 2006  
Publication 1 %

---

16	Young Sook Yun, Haruhiko Fukaya, Takahisa Nakane, Akihito Takano, Shigeru Takahashi, Yuji Takahashi, Hideshi Inoue. " A New Bis- - abietane Diterpenoid from Pohl ex Benth ", <i>Organic Letters</i> , 2014 Publication	1 %
17	Fatmawati, , and Hengki Tasman. "An optimal control strategy to reduce the spread of malaria resistance", <i>Mathematical Biosciences</i> , 2015. Publication	1 %
18	Hong Shen, Ai-Jun Hou. " Prenylated 2- arylbenzofurans from ", <i>Natural Product Research</i> , 2008 Publication	1 %
19	Van Cuong Pham, Akino Jossang, Thierry Sévenet, Bernard Bodo. "Cytotoxic Acylphenols from <i>Myristica maingayi</i> ", <i>Tetrahedron</i> , 2000 Publication	1 %
20	<a href="http://acgpsubs.org">acgpsubs.org</a> Internet Source	1 %
21	<a href="http://repository.ugm.ac.id">repository.ugm.ac.id</a> Internet Source	1 %
22	Kim, S.. "Edulisonones A and B, two epimeric benzo[b]oxepine derivatives from the bark of <i>Aglaia edulis</i> ", <i>Tetrahedron Letters</i> , 20051226	<1 %

23

Yoshikazu Yamamoto, Yasuhiro Kinoshita, G  
Ran Thor, Masahiro Hasumi et al.

"Isofuranonaphthoquinone derivatives from  
cultures of the lichen *Arthonia cinnabarina*  
(DC.) Wallr.", *Phytochemistry*, 2002

Publication

---

24

Lan Li, Mei-Ping Xie, Hua Sun, An-Qi Lu, Bo  
Zhang, Dan Zhang, Su-Juan Wang. " Bioactive  
phenolic acid-substituted glycoses and  
glycosides from rhizomes of ", *Journal of  
Asian Natural Products Research*, 2019

Publication

---

25

Vatcharin Rukachaisirikul, Nanthaphong  
Khamthong, Yaowapa Sukpondma, Souwalak  
Phongpaichit et al. "Cyclohexene,  
diketopiperazine, lactone and phenol  
derivatives from the sea fan-derived fungi  
*Nigrospora* sp. PSU-F11 and PSU-F12",  
*Archives of Pharmacal Research*, 2010

Publication

---

26

125.235.3.98

Internet Source

---

27

R.M. Rohini, Amit Kumar Das. " Triterpenoids  
from the stem bark of ", *Natural Product  
Research*, 2010

Publication

---

<1 %

<1 %

<1 %

<1 %

<1 %

28

[dokumen.pub](http://dokumen.pub)

Internet Source

&lt;1 %

29

[www.arkat-usa.org](http://www.arkat-usa.org)

Internet Source

&lt;1 %

30

[www.beilstein-journals.org](http://www.beilstein-journals.org)

Internet Source

&lt;1 %

31

Ibrahim, Sabrin R.M., and Gamal A. Mohamed. "Ingenines C and D, new cytotoxic pyrimidine- $\beta$ -carboline alkaloids from the Indonesian sponge *Acanthostrongylophora ingens*", *Phytochemistry Letters*, 2016.

Publication

&lt;1 %

32

Sergey V. Shevyakov, Olga I. Davydova, Dmitry G. Pershin, Mikhail Krasavin et al. "Natural products as templates for bioactive compound libraries: synthesis of biaryl derivatives of ( $\pm$ )-vasicine", *Natural Product Research*, 2006

Publication

&lt;1 %

33

Li. "Dichotomains A and B: Two New Highly Oxygenated Phenolic Derivatives from *Dicranopteris dichotoma*", *Organic Letters*, 04/2006

Publication

&lt;1 %

34

[dissertations.umi.com](http://dissertations.umi.com)

Internet Source

&lt;1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# Chromanone Acid Derivatives from the Stem Bark of *Calophyllum incrassatum*

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---