

Public Awareness for Antimicrobial Resistance from Escherichia coli Isolated from Beef Sold on Several Wet Market in Surabaya, Indonesia

by Wiwiek Tyasningsih Tyasningsih

Submission date: 08-Nov-2021 03:41PM (UTC+0800)

Submission ID: 1696468916

File name: Public_Awareness_for_Antimicrobial....pdf (874.05K)

Word count: 3183

Character count: 17267

Public Awareness for Antimicrobial Resistance from *Escherichia coli* Isolated from Beef Sold on Several Wet Market in Surabaya, Indonesia

Mustofa Helmi Effendi¹, Risi Sicilia², Jola Rahmahani³, Wiwiek Tyasningsih³

¹Department of Veterinary Public Health, Faculty of Veterinary Medicine, Airlangga University, Indonesia,

²Agriculture Quarantine Center, Palu, Central Sulawesi Province, Indonesia, ³Department of Veterinary Microbiology, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

Abstract

Objective: This study aims to phenotypically identify and confirm the presence of multi-drug resistant (MDR) and Extended Spectrum Beta-Lactamase (ESBL) Producing *Escherichia coli* in the swab surface samples of beef using VITEK-2 method.

Materials and Methods: Swab samples were taken from five wet markets: Pucang market, Wonokromo market, Keputran market, Pabean market, and Manukan market; 10 swab samples of beef were collected from each market. Then, isolation and identification in terms of bacteria using selective media and biochemical test were conducted. Resistance testing using disc diffusion method was performed on 6 types of antibiotics: Ampicillin, Cefazolin, Ceftriaxone, Cefotaxime, Ceftazidime, Tetracyclin. Positive isolates resistant to ≥ 2 types of Beta-Lactam antibiotics using disc diffusion method then were tested using VITEK-2 method.

Results: Positive samples containing *Escherichia coli* are found in 29 samples out of 50 swab samples. Of the 29 *Escherichia coli* isolates, 17 isolates are found resistant to the disc diffusion method. After testing those 17 isolates using VITEK-2 method, 5% (1/17) of ESBL-producing *Escherichia coli* are obtained which is resistant to all Beta-Lactam antibiotics. Besides, this study also reveals that 35% (6/17) of the *Escherichia coli* are positive multidrug resistance. Those *E.coli* MDR are found to be resistant towards antibiotic class Beta-Lactams, Aminoglycoside, Quinolone, and Sulfonamida.

Conclusion: This study has encouraged the need for public awareness for the understanding that beef from wet market can be potential as reservoir to spread multi-drug resistant bacteria that can cause health problems in humans.

Key words: MDR, ESBL, *Escherichia coli*, Beef, Vitek-2 System, Public Health

Introduction

Inappropriate use of antibiotics is one of the main factors in the occurrence of antibiotic resistance⁽¹⁾. Antibiotic resistance is a change in the ability of bacteria

to become resistant to antibiotics. Antibiotic resistance has now become a global public health problem and has been reported by the global agenda as one of the biggest threats to health⁽²⁾. Antibiotic resistance in bacteria can cause complications, longer treatment periods, treatment failure and death due to infection with resistant bacteria⁽³⁾. Humans can be infected by bacteria that are resistant to antibiotics through direct contact, consumption of contaminated meat, and the environment⁽⁴⁾. Extended spectrum Beta-Lactamase (ESBL) is an enzyme produced by gram negative bacteria and is a cause of resistance to almost all Beta-Lactam group antibiotics

Corresponding author:

Mustofa Helmi Effendi,
Department of Veterinary Public Health, Faculty of
Veterinary Medicine, Airlangga University, Surabaya,
Indonesia. Post Code: 60115. Telp : +62315992785.
Email : mheffendi@yahoo.com

such as Penicillin, Cephalosporin and Monobactam Aztreonam⁽⁵⁾. *Escherichia coli* is a normal flora of the mammalian digestive tract which can also be a cause of diseases such as gastroenteritis, cystitis, pneumonia, septicemia in humans and animals⁽⁶⁾.

Escherichia coli can also act as a reservoir for the spread of antibiotic resistance because it can easily move resistance genes to other bacteria⁽⁷⁾. ESBL producing *Escherichia coli* has been isolated from food from animals, hospital environments, plants, and feces⁽⁸⁾. Some studies also report the high prevalence of ESBL-producing *Escherichia coli* in food-consuming animals⁽⁹⁾, food products⁽¹⁰⁾, and the environment⁽¹¹⁾. *Escherichia coli* is a contaminant bacterium commonly found in meat⁽¹²⁾. The chemical composition and moisture of the meat is ideal for the life process of bacteria, this causes the meat can not last long when stored at room temperature⁽¹³⁾. This study focuses on identifying and confirming the presence of ESBL-producing *Escherichia coli* in beef through its resistance to Beta-lactam type antibiotics using the Vitek-2 method.

Vitek-2 advance expert system (BioMerieux) is an automated system used to show the prototype of the isolates tested and this method is able to determine the sensitivity or resistance of an isolate to antibiotics. The Vitek-2 test method has proven to be more reliable in detecting bacterial resistance to antibiotics because there is no subjective interpretation of the results⁽¹⁴⁾.

Materials and Methods

Ethical approval

Fresh beef samples were used in this study; hence, ethical approval was not necessary. Fresh beef samples were collected from Surabaya wet market.

Sampling

Sampling uses purposive sampling method. Samples were taken from five wet markets with criteria such as the market environment and the condition of dirty beef stalls. The number of samples is 10 samples of beef swabs from each market. The total samples examined were 50 samples of beef swabs. Swab results are labeled ¹⁵ sample swabs must be carried out aseptically. Using sterile swab sticks (Oxoid, Bangistoke, UK) which are placed in tubes containing media transport. After the

sample is taken, the sample is stored in the cooler box and taken to the laboratory.

Isolation and Identification of *Escherichia coli* Bacteria

Each sample produced by beef swabs was planted on Brilliant Green Bile Broth (BGBB) media (E. Merck, Darmstadt, Germany) then incubated at 37°C for 24 hours. Positive results are indicated by the presence of gas bubbles in the Durham tube and the change in color of the media to cloudy green. After being positive, it was then planted in Eosin Methylene Blue Agar (EMBA) media (E. Merck, Darmstadt, Germany) by streaking and incubated at 37°C for 18-24 hours. Typical colonies of *Escherichia coli* on metallic EMBA media.

Typical colonies of *Escherichia coli* grown in EMBA are planted again in the Triple Sugar Iron Agar (TSIA) media (E. Merck, Darmstadt, Germany) and Pepton Water Buffers and then incubated at 37°C for 24 hours. The Pepton Water 1 buffer, which has been incubated, is then dropped with a Kovach reagent of two or three drops. A positive test for *Escherichia coli* is characterized by the formation of a red ring on the surface of 1% Pepton Water^(15; 16).

Antibiotic Sensitivity Test

The *Escherichia coli* colonies found on the EMBA media were planted in a test tube containing 8 ml of physiological NaCl, homogenized using vortex until the same turbidity was obtained with standard McFarland 0.5. Then 0.2 ml was taken and gently rubbed on the entire surface of Mueller Hinton Agar (MHA) media (E. Merck, Darmstadt, Germany) using sterile cotton swabs.

³⁰ Sensitivity tests using the disc diffusion method were performed on 6 types of antibiotic disks namely Ampicillin 10 µg (Oxoid CT0003, UK), cefotaxime 30 µg (Oxoid CT0166, UK), ceftazidime 30 µg (Oxoid CT0412, UK), ceftriaxone 30 µg (Oxoid CT0417, UK), tetracycline 30 µg (Oxoid CT0054, UK), cefazolin 30 µg (Oxoid CT0011, UK) (CLSI, 2016). Culture of bacteria was incubated at 37°C for 24 hours. The results of the tests are interpreted based on the provisions of the Standard Laboratory Clinical Institute (CLSI, 2016). Positive isolates resistant to ≥ 2 types of Beta-Lactam antibiotics using disc diffusion method then were tested

using VITEK-2 method ⁽¹⁴⁾.

ESBL confirmation with Vitek-2 system

All isolates tested with the disk diffusion ¹⁸ were then identified and confirmed phenotypically using the Vitek-2 system (BioMerieux, Marcy L'Etoile, France) at the Microbiology Laboratory at Hospital of Airlangga University, Surabaya. Tests with Vitek-2 were carried out based on the factory protocol (BioMerieux, Marcy L'Etoile, France) that had been printed on the device.

Findings

The results of isolation and identification of bacteria from a total of 50 beef swab samples taken from 5 wet markets found 29 (58%) samples that were positive for *Escherichia coli* (Fig. 1). The high level of *Escherichia coli* contamination in beef found in this study is in accordance with the study conducted by Chuku et al ⁸ (17) which reported that *Escherichia coli* contamination levels in beef sold in traditional markets in Nigeria reached 90%.




Figure 1. E coli seen metallic green on EMBA media

Factors that cause high levels of *Escherichia coli* contamination in beef sold on the wet market are table surfaces that are in contact with meat, cleanliness of stalls is ¹⁰ not maintained ⁽¹⁸⁾. The sensitivity test (Fig. 2.) results using the disk diffusion test method of a total of 29 positive *Escherichia coli* samples showed 17 (58%) *Escherichia coli* isolates suspected of producing ESBL due to en 2 types of Beta-Lactam antibiotics with a resistance pattern shown in table 1.




Figure 2. Antibiotic sensitivity test results using the method for the disk diffusion test.

Table 1. Pattern of *Escherichia coli* Resistance to Antibiotics using Disk Diffusion Test

Antibiotics	Isolate number	Pattern of Antibiotic Resistance		
		S % (n)	I % (n)	R % (n)
Ampicillin	29	0	0	100 (29)
Cefazolin	29	10 (3)	31 (9)	58 (17)
Ceftriaxone	29	89 (26)	6 (2)	3 (1)
Cefotaxime	29	86 (25)	10 (3)	3 (1)
Ceftazidime	29	93 (27)	0 (0)	6 (2)
¹⁰ Tetracyclin	29	6 (2)	3 (1)	89 (26)

Description: R=Resistant, I= Intermediate, S= Susceptible.

A total of 17 isolates were then confirmed by phenotype using the Vitek-2 system. Confirmation test results using the Vitek-2 system showed that from 17 *Escherichia coli* isolates suspected of producing ESBL in the disc diffusion test, only 1 (5%) ESBL-producing positive *Escherichia coli* isolates were found in the

pattern of resistance shown in Table 3. Although in the last three years there has been a dynamic increase in research on antibiotic resistance caused by ESBL^(19; 20), there have been relatively few publications that report the presence of ESBL-producing Gram Negative Bacteria⁽²¹⁾.

26
Table 2. Identification of ESBL producing *E. coli* using Vitek-2 System

No	Isolate	Location	Antibiotics Resistance
1	A4	Pabean Market	AMX, AMP, SAM, CZ, CAZ, CRO, CTX, FEP, ATM, GM, CIP, SXT

The available data mostly discusses the existence of these microorganisms in livestock animals^(22; 23), only a few data about ESBL-producing *Escherichia coli* contamination in meat and processed meat products⁽⁹⁾. Research on meat is mostly done on chicken meat compared to meat from other animal species such as cattle⁽²⁴⁾. From the results found in this study, it can be confirmed phenotypically the presence of ESBL producing *Escherichia coli* in beef sold in wet markets using the Vitek-2 method shown in table 2.

Table 3. Identification of MDR *E. coli* using Vitek-2 System

No	Isolate	Location	Antibiotics Resistance
1	C4	Pucang Market	AMX, AMP, GM, CIP, SXT
2	C5	Pucang Market	AMX, AMP, CIP, SXT
3	C7	Pucang Market	AMX, AMP, CIP, SXT
4	C9	Pucang Market	AMX, AMP, CIP, SXT, SAM
5	C10	Pucang Market	AMX, AMP, CIP, SXT
6	E3	Keputran Market	AMX, AMP, CIP, SXT

4
Description: AMX = Amoxicillin, AMP = Ampicillin, SAM = Ampicillin-sulbactam, CZ = Cefazolin, CAZ = Cefazidime, CRO = Ceftriaxone, CTX = Cefotaxime, FEP = Cefepime, ATM = Aztreonam, GM = Gentamycin, CIP = Ciprofloxacin, SXT = Trimethoprim-Sulfamethoxazole.

This study also succeeded in obtaining 6 (35%) positive multidrug resistance *E. coli* isolates, MDR *E. coli* isolates were found to be resistant to Amoxicillin antibiotics by 35% (6/17), Gentamycin

5% (1/17), Ciprofloxacin 35% (6/17), Trimethoprim-Sulfamethoxazole 35% (6 /17), and Ampicillin-sulbactam 5% (1/17) shown in Table 3. These results are consistent with the study by Adinepekun et al⁽²⁵⁾

which states that resistance to a number of antibiotics such as aminoglycoside, Beta-Lactam, cephalosporin, fluoroquinolones, sulfonamide, tetracyclin, and trimethoprim have been found in *Escherichia coli* isolated from food from animal origin. Multidrug Resistance (MDR) is a condition where bacteria are resistant to ≥ 3 types of antibiotics. MDR has recently spread widely, especially in gram negative bacteria such as *Escherichia coli* (7).

Food contamination by antibiotic-resistant bacteria can be a serious threat to public health, the presence of resistant genes in plasmids, transposons and integrons facilitates the rapid spread of resistance genes between bacteria. Many resistant genes in *Escherichia coli* MDR are located on plasmids, which increase the likelihood of spreading these genes in the community (26).

Conclusion

In this study testing using the Vitek-2 method has succeeded in confirming the presence of ESBL-producing *Escherichia coli* with the discovery of one isolate positive ESBL-producing *Escherichia coli* that is resistant to all Beta-Lactam group antibiotics, Penicillin, Cephalosporin and Aztreonam. This study also succeeded in confirming the presence of *Escherichia coli* as MDR bacterial isolated from fresh beef swab samples which amounted to 35% of 17 samples examined using the Vitek-2 method. This results can be concluded that the need for public awareness for the understanding that beef from wet market can be potential as reservoir to spread multi-drug resistant bacteria that can cause health problems in humans.

Conflicts of Interest : The authors declare no conflict of interest.

Acknowledgments : The authors would like to thank the Rector of Airlangga University for providing research funds grant numbers; 892/UN3/2017 on the research title: PENANGGULANGAN ANTIBIOTICS RESISTANT BERSUMBER PANGAN ASAL HEWAN (PAH) MELALUI DETEKSI DINI. This article is part of the research.

References

1. Gross M. Antibiotics in crisis. *Curr Biol*, 2013; 23(24): R1063–R1065.

2. Centers for Disease Control and Prevention, Office of Infectious Disease. Antibiotic resistance threats in the United States, 2013; Available at: <http://www.cdc.gov/drugresistance/threat-report-2013>. Accessed January 28, 2015.
3. O'Brien G., Mary C., Altmann D. M., Dombrowski O., Morpeth S., Bhutta Z.A. Surveillance for antimicrobial drug resistance in under-resourced countries. *Emerg. Infect. Dis.* 2014; 20 (3): 20434–441. 10.3201/EID2003.121157.
4. Food and Agriculture Organization. Antibiotics in farm animal production. Publichealth and animal welfare. *FAO*, 2011; [Internet]. [leadmiuser_upload/animal_welfare_antibiotics_in_animal_farming.pdf](#).
5. Clinical and Laboratory Standards Institute. Performances Standards for Antimicrobial Susceptibility Testing. 2016; M100S 26th ed.
6. Dutta A, Jalal MS, Nath SK, Dhar PK, Das A, Uddin MM. Multi-drug resistance pattern of *Escherichia coli* isolated from hospital effluent and determination of tetracycline resistance gene. *J. Inf. Mol. Biol.* 2017;(3): 49-53.
7. Smolzka, A and Nagy, B. Multidrug resistant commensal *Escherichia coli* in animals and its impact for public health. 2013; doi: 10.3389/fmicb.2013.00258.
8. Shaheen BW, Nayak R, Foley SL, Kweon O, Deck J, Park M. Molecular characterization resistance to extended-spectrum cephalosporins in clinical *Escherichia coli* isolates from companion animals in the United States. *Antimicrob Agents.* 2011; 55:5666–75.
9. Ojer-Usoz E., Gonzalez D., Vitas A., Leiva J., Garcia-Jalon I., Febles Casquero A., de la Soledad Escolano M. Prevalence of extended spectrum beta-Lactamase producing enterobacteriaceae in meat products sold in Navarra, Spain. *Meat Sci*, 2013; 93, 316-321.
10. Schmid A., Hormansdorfer S., Messelhauser U., Kasbohrer A., Santer-Louise C., Mansfeld R. Prevalence of extended spectrum Beta-Lactamase producing *Escherichia coli* on Bavarian dairy and beef cattle farms. *Appl Environment Microbiol.* 2013; 79(9):3027-3032
11. Reinthaler FF, Feierl G, Galler H, Haas D, Leitner E, Mascher F, Melkes A, Posch J, Winter I, Zarfel G, Marth E. ESBL-producing

- Escherichia coli* in Austrian sewage sludge. Water Res. 2010; 44: 1981-1985. Doi: 10.1016/j.watres.2009.11.052
12. Sabarinath A , Tiwari K P, Deallie C , Belot G , Vanpee G , Matthew V , Sharma R , Hariharan H. Antimicrobial Resistance and Phylogenetic Groups of Commensal *Escherichia Coli* Isolates from Healthy Pigs in Grenada . WebmedCentral VETERINARY MEDICINE, 2011; 2(5):WMC001942
 13. Prihharsanti, A.H.T. Population of Bacteria and Fungi in Beef with Low Temperature Storage. Sains Peternakan, 2009; 7 (2): 66-72 ISSN 1693-8828
 14. Munoz-Dávila MJ, Roig M, Yagüe G, Blázquez A, Salvador C, Segovia M. Comparative evaluation of Vitek 2 identification and susceptibility testing of urinary tract pathogens directly and isolated from chromogenic media. Eur J Clin Microbiol Infect Dis 2013;32:773–780. 10.1007/s10096-012-1806-4.
 15. Effendi, MH, Harijani, N, Yanestria, SM and Hastutiek, P. Identification Of Shiga Toxin-Producing *Escherichia Coli* In Raw Milk Samples From Dairy Cows In Surabaya, Indonesia. Philipp. J. Vet. Med. 2018; 55(SI): 109-114
 16. Yanestria, YM, Rahmaniar, RP, Wibisono, FJ and Effendi, MH. Detection of *invA* gene of *Salmonella* from milkfish (*Chanos chanos*) at Sidoarjo wet fish market, Indonesia, using polymerase chain reaction technique. *Veterinary World*, 2019;12(1): 170-175.
 17. Chuku, A. , Lawrence B. Etim, LB, Obande, GA, Asikong, BE, Bashir Eya Sani, BE. Bacteriological Quality of Fresh Raw Beef and Chevron Retailed in Lafia Metropolis, Nigeria. Journal of Microbiology Research, 2016; 6(2): 29-34.
 18. Ismaïl, R., Aviat, F., Michel, V., Le Bayon, I., Gay-Perret, P., Kutnik, M., and Fédérighi, M. Methods for recovering microorganisms from solid surfaces used in the food industry: a review of the literature. *Int J Environ Res Public Health*. 2013;10(11):6169–6183. doi: 10.3390/ijerph10116169.
 19. Raponi G, Ghezzi MC, Gherardi G, Lorino G, Dicuonzo G. Analysis of methods commonly used for glycopeptide and oxazolidinone susceptibility testing in *Enterococcus faecium* isolates. *J. Med. Microbiol.* 2010; 59:672–678. 10.1099/jmm.0.016444-0.
 20. European Food Safety Authority (EFSA). Scientific opinion on the public health risks of bacterial strains producing extended– spectrum Beta lactamases and/ or Amp C Beta lactamases in food and food producing animals. 2011; EFSA V: 9, 2322
 21. Effendi, MH, Bintari, IG, Aksono, EB, and Hermawan, IP. Detection of *bla_{TEM}* Gene of *Klebsiella pneumoniae* Isolated from Swab of Food-Producing Animals in East Java. *Tropical Animal Science Journal*, 2018; 41(3):174-178
 22. Bortolaia, V. Guardabassi, L. Trevisani, M. Bisgaard, M. Venturi, L. And Bojesen, A. High diversity of Extended Spectrum beta Lactamase in *Escherichia coli* fecal sample isolates from healthy chickens. *Antimicrob Agents Chemother*, 2010; 54, 1624-1626.
 23. Kallau NHG, Wibawan IWT, Lukman DW, Sudarwanto MB. Detection of multi-drug resistant (MDR) *Escherichia coli* and tet gene prevalence at a pig farm in Kupang, Indonesia. *J Adv Vet Anim Res* 2018; 5(4):388–96.
 24. Sheikh A., Checkley S., Avery B., Chalmers G., Bophaychuk V., Boerlin P., Reid Smith R., Aslam M. Antimicrobial resistance and resistance genes in *Escherichia coli* isolated from retail meat purchased in Alberta, Canada. *Foodborne Pathol Dis*, 2012; 9, 625–631.
 25. Adenipekun, E.O, Jackson, C.R, Oluwadun, A, Iwalokun, B.A, Frye, J.G, Barrett, J.B, Hiott, L.M, and Woodley, T.A. Prevalence and antimicrobial resistance in *Escherichia coli* from food animals in Lagos, Nigeria. *Microb Drug Resist*. 2015; 21: 358-365.
 26. Chen X., He L., Li Y., Zeng Z., Deng Y., Liu Y., and Liu J.H. Complete sequence of a F2:A-B-plasmid pHN3A11 carrying *rmtB* and *qepA*, and its dissemination in China. *Vet Microbiol*, 2014; 174: 267-271.

Public Awareness for Antimicrobial Resistance from Escherichia coli Isolated from Beef Sold on Several Wet Market in Surabaya, Indonesia

ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

18%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	docplayer.net Internet Source	1%
2	www.hindawi.com Internet Source	1%
3	Bernard Wasiński, Hanna Róžańska, Jacek Osek. "Occurrence of extended spectrum β -lactamase and AmpC-producing Escherichia coli in meat samples", Bulletin of the Veterinary Institute in Pulawy, 2013 Publication	1%
4	Melano, R.G.. "Cephalosporin resistance in Klebsiella pneumoniae from Nova Scotia, Canada", Diagnostic Microbiology & Infectious Disease, 200610 Publication	1%
5	res.mdpi.com Internet Source	1%
6	www.ncbi.nlm.nih.gov Internet Source	1%

7	Qurat ul Ain Zahra, Zhaofeng Luo, Rizwan Ali, Muhammad Imran Khan, Fenfen Li, Bensheng Qiu. "Advances in Gold Nanoparticles-Based Colorimetric Aptasensors for the Detection of Antibiotics: An Overview of the Past Decade", <i>Nanomaterials</i> , 2021 Publication	1 %
8	www.researchsquare.com Internet Source	1 %
9	"Technical program", <i>Journal of the American Oil Chemists Society</i> , 02/1976 Publication	1 %
10	Olanrewaju E. Fayemi, Gabriel B. Akanni, James A. Elegbeleye, Olusimbo O. Aboaba, Patrick M. Njage. "Prevalence, characterization and antibiotic resistance of Shiga toxigenic <i>Escherichia coli</i> serogroups isolated from fresh beef and locally processed ready-to-eat meat products in Lagos, Nigeria", <i>International Journal of Food Microbiology</i> , 2021 Publication	1 %
11	www.virox.com Internet Source	1 %
12	bmcmicrobiol.biomedcentral.com Internet Source	1 %

13	jyoungpharm.org Internet Source	1 %
14	bmcvetres.biomedcentral.com Internet Source	1 %
15	K. Chishimba, B. M. Hang'ombe, K. Muzandu, S. E. Mshana, M. I. Matee, C. Nakajima, Y. Suzuki. " Detection of Extended-Spectrum Beta-Lactamase-Producing in Market-Ready Chickens in Zambia ", International Journal of Microbiology, 2016 Publication	<1 %
16	es.scribd.com Internet Source	<1 %
17	jidc.org Internet Source	<1 %
18	pure.uva.nl Internet Source	<1 %
19	Magdalena Nüesch-Inderbinnen, Roger Stephan. "Epidemiology of Extended-Spectrum β -Lactamase-Producing Escherichia coli in the Human-Livestock Environment", Current Clinical Microbiology Reports, 2016 Publication	<1 %
20	Murat Duman, Hakan Abacioglu, Meral Karaman, Nuray Duman, Hasan Ozkan. "beta-lactam antibiotic resistance in aerobic	<1 %

commensal fecal flora of newborns",
Pediatrics International, 2005

Publication

21

oatao.univ-toulouse.fr

Internet Source

<1 %

22

A.M. Aristizábal-Hoyos, E.A. Rodríguez, L. Arias, J.N. Jiménez. "High clonal diversity of multidrug-resistant and extended spectrum beta-lactamase-producing *Escherichia coli* in a wastewater treatment plant", *Journal of Environmental Management*, 2019

Publication

<1 %

23

Altaf Hussain, Wang, Sun, Gu, Liu, Yu, Ahmad, Jiang, Hou. "Molecular Characterization Of Pathogenic *Salmonella* Spp From Raw Beef In Karachi, Pakistan", *Antibiotics*, 2020

Publication

<1 %

24

Ben Sallem, Rym, Karim Ben Slama, Yolanda Sáenz, Beatriz Rojo-Bezares, Vanesa Estepa, Ahlem Jouini, Haythem Gharsa, Naouel Klibi, Abdellatif Boudabous, and Carmen Torres. "Prevalence and Characterization of Extended-Spectrum Beta-Lactamase (ESBL)- and CMY-2-Producing *Escherichia coli* Isolates from Healthy Food-Producing Animals in Tunisia", *Foodborne Pathogens and Disease*, 2012.

Publication

<1 %

25 Isadora de Alcântara Rodrigues, Rafaela Gomes Ferrari, Pedro Henrique Nunes Panzenhagen, Sergio Borges Mano et al. "Antimicrobial resistance genes in bacteria from animal-based foods", Elsevier BV, 2020
Publication <1 %

26 Lim, Jong-Soo, Da-Som Choi, Young-Jo Kim, Jung-Whan Chon, Hong-Seok Kim, Hyun-Jung Park, Jin-San Moon, Sung-Hwan Wee, and Kun-Ho Seo. "Characterization of Escherichia coli-Producing Extended-Spectrum β -Lactamase (ESBL) Isolated from Chicken Slaughterhouses in South Korea", Foodborne Pathogens and Disease, 2015.
Publication <1 %

27 Muhammad Mustapha, Yusuf Audu, Kingsley Uwakwe Ezema, Jafar Umar Abdulkadir et al. "Antimicrobial Susceptibility Profiles of Isolates from Diarrheic Dogs in Maiduguri, Borno State, Nigeria ", Macedonian Veterinary Review, 2021
Publication <1 %

28 Rita Endriani, Elita Rafni, Ligat Pribadi Sembiring, Fajri Marindra Siregar, Budi Haryadi, Arkhan Kurniawan. "Antibiotics multi-resistant bacteria in patient with dental abscess at arifin achmad general hospital of Riau province", AIP Publishing, 2019 <1 %

29

A Umo, O John, E Ekpuk. "Detection of Plasmid-Mediated AmpC Beta-lactamase Enzyme among Escherichia coli Isolates in Livestock, South Nigeria", Asian Journal of Biology, 2018

Publication

<1 %

30

F.Y. Zhang, S.Y. Huo, Y.R. Li, R. Xie, X.J. Wu, L.G. Chen, Y.H. Gao. " A survey of the frequency of aminoglycoside antibiotic-resistant genotypes and phenotypes in in broilers with septicaemia in Hebei, China ", British Poultry Science, 2014

Publication

<1 %

31

Ioannis Galdadas, Shen Qu, Ana Sofia F Oliveria, Edgar Olehnovics et al. "Allosteric communication in Class A β -lactamases occurs via Cooperative Coupling of Loop Dynamics", Cold Spring Harbor Laboratory, 2021

Publication

<1 %

32

Sifuna Anthony Wawire, Oleg N. Reva, Thomas J. O'Brien, Wendy Figueroa, Victor Dinda, William A. Shivoga, Martin Welch. "Virulence and antimicrobial resistance genes are enriched in the plasmidome of clinical Escherichia coli isolates compared with

<1 %

wastewater isolates from western Kenya",
Infection, Genetics and Evolution, 2021

Publication

33

www.fedoa.unina.it

Internet Source

<1 %

34

"Guide to Antimicrobial Use in Animals",
Wiley, 2008

Publication

<1 %

35

Fedik A. Rantam, Purwati, Budi Setiawan,
Sony Wibisono et al. "Induced Monocytes-
Derived HSCs (CD34+) with LPS Accelerated
Homing Rat Bone Marrow-Mesenchymal Stem
Cell (BM-MSCs, CD105) in Injured Pancreas",
Journal of Biomedical Science and
Engineering, 2015

Publication

<1 %

36

Teixeira, Juliana V., Pedro Cecílio, Daniela
Gonçalves, Vítor J. P. Vilar, Eugénia Pinto, and
Helena N. Ferreira. "Multidrug-resistant
Enterobacteriaceae from indoor air of an
urban wastewater treatment plant",
Environmental Monitoring and Assessment,
2016.

Publication

<1 %

37

id.wikipedia.org

Internet Source

<1 %

38

"13th European Congress of Clinical Microbiology and Infectious Diseases", Clinical Microbiology and Infection, 2003

Publication

<1 %

39

"Abstracts cont.", Clinical Microbiology and Infection, 4/2005

Publication

<1 %

40

our-illumination.com

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Public Awareness for Antimicrobial Resistance from Escherichia coli Isolated from Beef Sold on Several Wet Market in Surabaya, Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
