


Get vaccinated and follow health protocols during this Covid-19 pandemic

Critical Care and Shock

Acute necrotizing pancreatitis associated with Covid-19: A case report

CATEGORIES

- > Abstracts
- > Case Report



Critical
Care & Shock

Editorial Board

Founder

Iqbal Mustafa (Indonesia)

Editor-in-Chief

Joseph Varon (USA)

Associate Editors Americas

Robert E. Fromm (USA)

Paul Marik (USA)

Associate Editors Asia and Pacific

Anthony McLean (Australia)

Dessmon YH Tai (Singapore)

Associate Editor Europe

Santiago Herrero Fernandez
(Spain)

Associate Editor Middle East

Abdullah Al-Shimemeri (Saudi
Arabia)

Senior Editors

E. Benjamin (USA)

GJ Dobb (Australia)

Chen Dechang (China)

Y. Koh (Korea)

Dessmon YH Tai (Singapore)

Patrick SK Tan (Malaysia)

Editorial Board Members

Pravin R. Amin (India)

DC Angus (USA)

Youzhong An (China)

D. Bihari (Australia)

R. Bellomo (Australia)

IH Chaudry (Alabama)

Susan E. Dantoni (USA)

Ahmed I. Ghali (Egypt)

KM Iqbal (Bangladesh)

GM Joynt (Hong Kong)

CM Khoa (Vietnam)

Haekyu Kim (Korea)

YL Kim (Korea)

Shin Ok Koh (Korea)

V. Kvetan (USA)

Chae-Man Lim (Korea)

J. Lipman (Australia)

T. Maekawa (Japan)

KN Maranetra (Thailand)

Xian Jun Meng (China)

S. McKinley (Australia)

K. Okada (Korea)

PJ Papadacos (USA)

MR Pinsky (USA)

Fang Qiang (China)
VG Reddy (Sultanate of Oman)
John H. Reeves (Australia)
E. Santos (Philippines)
MM Sayeed (USA)
Mervyn Singer (UK)
J. Takezawa (Japan)
J. Takala (Switzerland)
GJ Tang (Taipei-Taiwan)
IKS Tan (Hong Kong)
J-L Vincent (Belgium)

NR Webster (UK)

Managing Editor

Simon Nanlohy (Indonesia)

Assistant Editors

Ana L. Huerta-Alardin (USA)

Luciana B. Sucanco (Indonesia)

Get vaccinated and follow health protocols during this Covid-19 pandemic

Month: October 2020

Treatment of COVID-19 is critically phase specific

Low-dose heparin for sepsis-associated disseminated

CATEGORIES

- > Abstracts
- > Case Report
- > Clinical Pearls

A number of beneficial (corticosteroids and remdesivir), and potentially beneficial therapies, exist for the treatment of COVID-19. What is not widely appreciated is that patients with COVID-19 progress through several distinct phases, with each stage having a different treatment strategy (Figure 1). This fact is of critical importance in achieving the best possible outcomes of this disease. The symptomatic phase is marked by enormous viral replication reaching viral loads in the upper respiratory tract in excess of a billion virions per milliliter, with a very limited host inflammatory response. While patients may remain SARS-CoV-2 PCR positive for weeks (and in...

Authors: Paul E. Marik, Joseph Varon, Pierre Kory

intravascular coagulation and septic shock

Abstract

Background: Sepsis and septic shock are severe clinical problems with high mortality rates, characterized by a systemic inflammation with substantial procoagulant elements and activating some clotting factors. Those are possible to develop the disseminated intravascular coagulation (DIC). Heparin is a well-known anticoagulant, which also provides anti-inflammatory properties. Nevertheless, the efficacy of heparin was limited by the potential risk of bleeding in critically ill patients. In this case report, we would like to explain the therapeutic effects of low-dose heparin on sepsis-associated DIC and septic shock.

- > Commentary for Critical Care and Shock
- > EBM
- > Editorial
- > Images in Clinical Medicine
- > Obituary
- > Original Article
- > Other
- > Perspective
- > Review Article

ARCHIVES

- > December 2021
- > September 2021
- > July 2021
- > June 2021
- > April 2021
- > February 2021
- > December 2020
- > October 2020
- > August 2020
- > May 2020
- > April 2020
- > February 2020

A case of severe respiratory distress in a patient with chronic myeloid leukemia receiving dasatinib

Abstract

Introduction: Dasatinib is a multi-kinase inhibitor used primarily in the treatment of chronic myeloid leukemia (CML). The major reported side effects are pleural effusion, pulmonary hypertension, and severe infection. The most common infection among these patients is pneumonia. Here, we present a case of severe respiratory failure in a patient with CML who was taking dasatinib.

Case presentation: A 75-year-old male with CML, hypertension, hyperlipidemia presented to the emergency department

Case report: We report a case of a 60-year-old male with sepsis-associated DIC and septic shock caused by pneumonia. The patient had decreased consciousness for 10 hours before hospitalization. Standard intensive care and intravenous low-dose heparin (250 IU/h) were performed in the Intensive Care Unit (ICU). On day 12 of ICU admission, the patient's condition was fully conscious and clinically stable. Therefore, the patient discharged from ICU with no bleeding manifestation and no sequelae.

Conclusion: Low-dose heparin successful treats patient with sepsis-associated DIC and septic shock.

Authors: Arie Zainul Fatoni, Suwarman, Hori Hariyanto, Oloan E. Tampubolon, Wiwi Jaya

- > December 2019
- > October 2019
- > August 2019
- > June 2019
- > April 2019
- > February 2019
- > November 2018
- > August 2018
- > May 2018
- > February 2018
- > November 2017
- > July 2017
- > May 2017
- > February 2017
- > October 2016
- > August 2016
- > May 2016
- > February 2016
- > November 2015
- > August 2015
- > May 2015
- > March 2015
- > October 2014
- > August 2014
- > May 2014
- > February 2014

with progressively worsening shortness of breath and hemoptysis for one week. The patient's CML had been well controlled with dasatinib since his diagnosis two years ago, and the most recent BCR-ABL1 assay was undetectable. Computed tomography (CT) of the chest revealed diffuse ground glass opacity with superimposed interlobular septal thickening and intralobular lines ("crazy-paving pattern") and a moderate-sized right pleural effusion. Therapeutic thoracentesis yielded 1.8 l of lymphocyte predominant, exudative pleural effusion. Pneumocystis jirovecii polymerase chain reaction (PCR) of induced sputum was positive, which was consistent with the CT finding of "crazy-paving pattern." Dasatinib was held for the possibility of drug induced pneumonitis and pleural effusion, and the patient was successfully treated with trimethoprim-sulfamethoxazole for his pneumocystis jirovecii pneumonia (PCP).

Resuscitation incoherence after cardiac arrest due to acute intraoperative pulmonary emboli related acute right heart failure: A case report

Abstract

Resuscitation incoherence is a mismatch between macrocirculation, microcirculation, and cellular parameters after resuscitation. We report a 34-year-old obese male patient, suffering from end-stage renal disease or chronic kidney disease (CKD), underwent routine hemodialysis three times a week and hypercoagulability state with rivaroxaban therapy. He had a cardiac

- > November 2013
- > August 2013
- > May 2013
- > February 2013
- > November 2012
- > September 2012
- > July 2012
- > April 2012
- > October 2011
- > September 2011
- > July 2011
- > June 2011
- > February 2011
- > December 2010
- > November 2010
- > September 2010
- > May 2010
- > April 2010
- > March 2010
- > December 2009
- > July 2009
- > June 2009
- > March 2009
- > January 2009
- > August 2008
- > May 2008

Conclusion: Our case suggests that a common tyrosine kinase inhibitor, dasatinib, cannot only act as an effective antileukemic agent, but also can cause several adverse effects including pleural effusion and immunosuppression. Physicians should consider opportunistic infections in their differential when patients on dasatinib present with respiratory insufficiency.

Authors: Natsumi T. Hamahata, Sophie Rodrigues Pereira, Ehab G. Daoud

Are preinjury anticoagulant and antiplatelet medications a pitfall in the bleeding tendencies of elderly

arrest during kidney transplant surgery. Acute right heart failure causing cardiac arrest was presumably caused by acute intraoperative pulmonary embolism. Hemodynamic and resuscitation incoherence occurred and proper treatment was needed. At the time of cardiac arrest, hemodynamic coherence was lost and resuscitation was performed to restore this loss by correcting the possible causes of cardiac arrest. Although the return of spontaneous circulation (ROSC) was successfully achieved, a type 1 resuscitation incoherence occurred where the macrocirculation was optimal but cellular parameters were disturbed by cell hypoxia, characterized by high levels of lactate. Type 2 resuscitation incoherence was also found in this patient until the end of treatment in the intensive care unit (ICU). Cellular parameters such as lactate levels and the venoarterial carbon dioxide tension difference to arteriovenous oxygen

- > March 2008
- > December 2007
- > September 2007
- > August 2007
- > July 2007



TRENDS OF ELDERLY trauma patients in intensive care?

Abstract

Purpose: The elderly are more likely to be on anticoagulant or antiplatelet medications, which increase bleeding. We aimed to determine the effect of preinjury anticoagulant or antiplatelet medications on required blood transfusions and the trauma outcomes of elderly patients.

Methods: We retrospectively reviewed the medical records of all elderly trauma patients admitted to Chungbuk National University Hospital from January 2016 to June 2019. We compared the required number of blood transfusion units, complications, and mortality rate between those on anticoagulant or antiplatelet medications and those that were not, using the chi-squared test,

content difference ratio ($P[v-a]CO_2 / (C[a-v]O_2)$) continued to improve during ICU treatment, but cell hypoxia might occur since the central venous pressure (CVP) value as a macrocirculation parameter was likely to increase, presumed to be caused by incomplete resolved acute pulmonary embolism related chronic thromboembolic pulmonary hypertension.

Authors: Ade Susanti, Mayang Indah Lestari, Dita Aditiansih

Does preinjury anticoagulant or antiplatelet medication increase the need for blood

independent t-test, linear regression analysis, and logistic regression analysis.

Results: Out of 466 patients, 142 were on anticoagulant or antiplatelet medications while 324 were not. There was a significant statistical difference in the unit amount of red blood cells transfused within 4 hours of arriving at the hospital between the medicated and non-medicated groups (0.89 vs 1.43 units, respectively, $p=0.02$); however, multivariate analysis showed no statistical difference ($p=0.28$). The medication group showed a higher rate of complications compared to the non-medicated group (47.9% vs 29.6%, respectively, $p=0.001$), bleeding (17.6% vs 2.8%, respectively, $p=0.001$), and pneumonia (24.4% vs 14.2%, respectively, $p=0.01$). There was no statistical difference in the mortality rate (16.9% vs 22.2%, respectively, $p=0.21$).

transfusions in patients aged older

than 65 years with traumatic brain injury?

Abstract

Background: Anticoagulant or antiplatelet medications are commonly prescribed in older adults, increasing bleeding tendency and affecting traumatic brain injury (TBI)-related morbidity and mortality.

Objectives: This study aimed to determine the effects of preinjury anticoagulant or antiplatelet medication on blood transfusions and outcomes in patients aged >65 years with TBI.

Methods: We retrospectively reviewed records of patients with TBI without other injuries admitted to our hospital

Conclusion: Preinjury anticoagulant or antiplatelet medications in elderly trauma patients increased bleeding and complications such as pneumonia but did not affect transfusion requirement, or mortality rate.

Authors: Se Heon Kim, Young Hoon Sul, Jin Young Lee, Joong Suck Kim

**Future views on
nanonutrition for
critically ill patients:
The role of extra
virgin olive oil
nanoemulsion in
sepsis enteral
nutrition**

Other injuries admitted to our hospital between January 2016 and June 2019. We compared the number of blood transfusions administered and outcomes

between patients who were receiving anticoagulant/antiplatelet medication and those who were not.

Results: Overall, 82 patients (66% male) with an average \pm standard deviation age of 76.6 ± 7.29 years were enrolled. Thirty-one patients were receiving anticoagulants or antiplatelets and 51 were not. There were no differences in age, medical history, Injury Severity Score, and Glasgow Coma Scale score between the groups. International normalized ratios of patients who were on warfarin were significantly higher than those of patients who were not ($p < 0.05$). Analysis of covariance demonstrated that patients who were receiving medications needed more plasma transfusions than did those who were not ($p < 0.05$). The incidence of complications was 64.5% and 37.3% in

Abstract

Enteral nutrition (EN) can maintain the structure and function of the gastrointestinal mucosa better than parenteral nutrition. Early intervention by enteral nutrition in critically ill patient may help the patient from fatality of multiple organ failure. Oral nutrition is an attempt to provide a physiological nutrition that is expected to trigger the immune system, prevent blood stream infection from the intravenous route, and reduce cost of therapy. One of the problems that inhibits supply of enteral nutrition in critically ill patients is absorption disorders that cause the body's nutritional needs to be hampered. Administration of extra virgin olive oil (EVOO) in the form of nanoemulsion is expected to improve pharmacokinetics and pharmacodynamics in those patients. EVOO is one functional food that has a lot of health benefits. Nanoemulsion-

complications was 0.15% and 0.15% in patients who were and were not receiving medication, respectively ($p < 0.05$). Multivariate regression

analysis showed that patients who were receiving medications bled 5.62 times more than did those who were not (95% confidence interval: 1.52~20.70).

Conclusions: Bleeding incidence and plasma transfusion requirements are increased by preinjury anticoagulant or antiplatelet medication in patients aged >65 years with TBI.

Authors: Se Heon Kim, Young Hoon Sul, Jin Young Lee, Jin Bong Ye, Jin Suk Lee, Hong Rye Kim, Soo Young Yoon, Joong Suck Kim, Moon Sang Ahn

Giant extrathoracic hematoma in a

based delivery systems are proven to increase utilization of lipophilic bioactive components in food, personal care, cosmetic, and pharmaceutical applications. So far, there is no report describes the use of enteral nononutrition in critically ill patients. This review discusses the perspective view of using EVOO nanoemulsion to care the critically ill patients.

Authors: Anna Surgean Veterini, Subijanto Marto Soedarmo, Hasanul, Annis Catur Adi, Heni Rachmawati, Nancy Margarita Rehatta

Acute necrotizing encephalopathy secondary to COVID-19

COVID-19 patient

Case presentation

A 69-year-old lady presented to our emergency department with a two week history of shortness of breath, fever, and dry cough. Chest computed tomography revealed patchy ground glass opacities throughout both lungs, most pronounced in the mid-to-lower lung zones bilaterally, with peripheral distribution. She was admitted to the hospital with the presumptive diagnosis of coronavirus disease 2019 (COVID-19) pneumonia. She received our standard MATH+ protocol (hydroxychloroquine, intravenous ascorbic acid, zinc, thiamine, melatonin, azithromycin, vitamin D3, and enoxaparin). This was followed by oral anticoagulation with warfarin. On day 18, she complained of severe right breast pain. Upon examination she...

Authors: Elizabeth Gamboa, Joseph C

A 54-year-old gentleman without significant past medical history presented to our emergency department with complaints of shortness of breath and

fever. On arrival he was very disoriented, and unable to communicate well. He was found to be positive for SARS-CoV-2 and had severe pneumonia. On hospital day 8, he abruptly began to decline, initially presenting with elevation in blood pressure, and soon thereafter with severely decreased level of consciousness. This was followed by a cardiac arrest, that was treated, and he had eventual return of spontaneous circulation within 15 minutes. Despite this, the patient never regained consciousness. A head computed...

Authors: Swethen Dushianthan, Elizabeth Gamboa, Joseph Varon

Authors: Elizabeth Gamboa, Joseph C. Gathe Jr, Joseph Varon

Future views on nanonutrition for critically ill patients: The role of extra virgin olive oil nanoemulsion in sepsis enteral nutrition

Anna Surgean Veterini, Subijanto Marto Soedarmo, Hasanul, Annis Catur Adi, Heni Rachmawati, Nancy Margarita Rehatta

Abstract

Enteral nutrition (EN) can maintain the structure and function of the gastrointestinal mucosa better than parenteral nutrition. Early intervention by enteral nutrition in critically ill patient may help the patient from fatality of multiple organ failure. Oral nutrition is an attempt to provide a physiological nutrition that is expected to trigger the immune system, prevent blood stream infection from the intravenous route, and reduce cost of therapy. One of the problems that inhibits supply of enteral nutrition in critically ill patients is absorption disorders that cause the body's nutritional needs to

be hampered. Administration of extra virgin olive oil (EVOO) in the form of nanoemulsion is expected to improve pharmacokinetics and pharmacodynamics in those patients. EVOO is one functional food that has a lot of health benefits. Nanoemulsion-based delivery systems are proven to increase utilization of lipophilic bioactive components in food, personal care, cosmetic, and pharmaceutical applications. So far, there is no report describes the use of enteral nanonutrition in critically ill patients. This review discusses the perspective view of using EVOO nanoemulsion to care the critically ill patients.

Key words: Nanonutrition, critically ill patient, extra virgin olive oil nanoemulsion, oral formulation, emulsion.

Introduction

Enteral nutrition (EN) support can be determined from the supply of calories, protein, electrolytes, vitamins, minerals, trace elements, and fluids through the intestinal route. The early initiation of EN including fiber may prevent atrophy of intestinal

mucosa and attenuation of gastrointestinal peristalsis, because the energy substrates for intestinal mucosa are partially supplied via intraluminal. In addition, it is believed that bacterial translocation (BT) can be prevented by early initiation of EN. (1) Prevention of bacterial translocation is one of efforts to prevent sepsis that has high mortality rate.

Sepsis is a serious clinical condition that can result in death, as a result of systemic inflammatory response syndrome including fever leukopenia, hypotension, and failure of many organs. (2) An unbalanced pro- and anti-inflammatory response, inflammatory resolution disorder, and persistent inflammation have an important roles in the acute and/or chronic phases of sepsis. (3)

Oils are very important substances in human nutrition. However, they are sensitive to oxygen, heat, moisture, and light. In recent years, there has been a growing interest in the modification technology of oils. Methods that modify oil characteristics and make oils suitable applications have been increasingly studied. Nanotechnology has become one of the technologies that can revolutionize conventional food science and the food industry. (4,5)

From Department of Anesthesiology and Intensive Care, Medical Faculty of Universitas Airlangga - Dr. Soetomo General Academic Hospital (Anna Surgean Veterini and Nancy Margarita Rehatta), Department of Pediatric, Medical Faculty of Universitas Airlangga - Dr. Soetomo General Academic Hospital (Subijanto Marto Soedarmo), Medical Faculty of Universitas Sumatera Utara (Hasanul), Public Health Faculty of Universitas Airlangga (Annis Catur Adi), School of Pharmacy, and Bandung Institute of Technology (Heni Rachmawati).

Address for correspondence:

Nancy Margarita Rehatta
Department of Anesthesiology and Intensive Therapy, Medical Faculty of Universitas Airlangga - Dr. Soetomo General Academic Hospital, Surabaya
Mayjend Prof. Dr. Moestopo No. 6-8. Airlangga, Gubeng, Surabaya 60285, Indonesia
Tel: (+62) 811307034
Email: margaritarehatta@gmail.com

Nanotechnology, especially nanoemulsions (NEs), has gained increasing interest for researchers. NEs are important in many industries because they have small droplets with a high surface area. The formation of nano-sized emulsions can be divided into two types: high and low-energy methods. High-energy methods are high pressure homogenization, microfluidization, and ultrasonic emulsification. Spontaneous emulsification, phase inversion temperature (PIT), phase inversion composition (PIC), and the less-known phase-D emulsification (DPE) method are emphasized in the low-energy method. The NEs application is explained in three main areas namely food, cosmetics, and drug delivery. (6,7)

NEs are sub-micron sized colloidal particle systems, which act as carriers of drug molecules. As a drug delivery system they enhance the therapeutic efficacy of the drug and minimize adverse effect as well as the toxic reactions. Major application includes treatment of infection of the reticuloendothelial system (RES), enzyme replacement therapy in the liver, treatment of cancer, and vaccination. (8,9)

Early nutrition therapy is recommended for individuals in critical conditions. (10) The parenteral and enteral routes are currently prescribed for critically ill patients. (11) There is increasing evidence that significant benefits are achieved if nutrients are delivered within the gut compared with the parenteral route. However, there is increasing assistance for combined enteral-parenteral therapy in the case of sustained hypocaloric enteral nutrition. (11) Despite limited investigation, a strategic temporary minimal enteral nutrition with hypocaloric content has been recommended, with the aim of avoiding the consequences of overfeeding syndrome and protecting the gut from the threat of hypoperfusion. (12,13) Enteral nutrition may be used with caution during a period of hypotension. Evidences of poor gastrointestinal function such as increased nasogastric tube output, unexplained abdominal pain, and abdominal distention, or the development of dilated loops of bowel or intramural gas (pneumatosis intestinalis) on radiographic studies should be interpreted as potential indicators of gut ischemia. (14)

There is consistent evidence that nutrients delivered by the gut promote significant benefits when compared with the intravenous route. Enteral feeding is associated not only with lower costs but also with shortening of hospital stay and infectious complications. (15) This review article interest giving extra virgin olive oil (EVOO) nanoemulsion in sepsis condition.

The active absorption of materials, particularly sugars and amino acids, causes a major increase in oxygen consumption and is dependent on an adequate supply of oxygen and blood flow. Passive absorption appears to be primarily a function only of blood flow. It is conceivable that a portion of local vascular control during active absorption could be related to a decrease in tissue PO₂ during absorption. Although Svanvik, et al have shown that arterial PO₂ must be reduced by approximately one-half to increase intestinal blood flow to about 200% of control, this does not necessarily negate a possible role for oxygen in absorptive hyperemia as Svanvik, et al have proposed. Duling has recently demonstrated that a decrease in general tissue PO₂ causes substantial vasodilation, even though a comparable change in PO₂ at the arteriolar wall when tissue PO₂ is normal has a minor dilatory effect. The basic question is whether or not intestinal tissue PO₂ is sufficiently decreased by active absorption to cause a dilatory response. (16)

Extra virgin olive oil

The major components of olive oil are fatty acids that represent more than 98% of its composition. Fatty acids contained in olive oil consist of mono-unsaturated fatty acids (MUFAs), such as oleic acid (55-85% of the total fatty acids). Beside that, it also contains polyunsaturated fatty acids (PUFAs), such as linoleic acid and saturated fatty acids (SFAs) like stearic or palmitic acids. (17)

EVOO contains around 36 phenolic compounds including tyrosol, hydroxytyrosols, oleocanthal, and oleuropein and carotenes. These phenolic compounds enter brain and produce neuroprotective effects through their antioxidant, anti-apoptotic, and anti-inflammatory activities. (18) In addition to polyphenols, virgin olive oil contains vitamins such as α - and γ -tocopherols (around 200 ppm) and β -carotene, phytosterols, pigments, terpenic acids, flavonoids, and squalene. (19)

The consumption of olive oil is associated with a low incidence of cardiovascular diseases, neurological disorders, and breast cancer. (20) It is well known that polyphenols concentration of extra virgin olive oils may differ, depending on both cultivars, degree of ripeness of the olives as well as the extraction procedure but only tyrosol, oleuropein, gallic acid, caffeic acid, and vanillic acid are the most important agents responsible for the antioxidant activity in olive oils. (21) It is clear that the antioxidant effects in vivo of EVOO and olives are strictly linked to the bioavailability of these molecules from the diet. Nutritional trials in human and

animal models reported that olive phenols are rapidly absorbed in a dose-dependently manner from the gastrointestinal tract and distributed through all organs as conjugated forms. (22) Olive oil, tyrosol and hydroxytyrosol, are dose-dependently absorbed in humans after ingestion and that they are excreted in the urine as glucuronide conjugates. (23)

The impact of an in vitro procedure that mimics the physicochemical changes occurring in gastric and small intestinal digestion on the bioaccessibility and antioxidant activity of phenols from 10 extra virgin olive oil samples was assessed. EVOO phenols were totally extracted in the aqueous phase, which reproduces gastric fluids during the digestion procedure. EVOO with different phenolic compositions would show a different capacity to act as a potential antioxidant after ingestion. (24) Polyphenols are considered to be the main contributors to the antioxidant activity of olive oils. (25) Depending on the manufacturing process, we can find different types of oil: (17)

- Virgin olive oil: obtained directly from the ripe fruit by mechanical procedures. It is the only consumed raw without the use of any solvents, so it keeps all its properties intact. Olive oil has excellent antioxidants such as polyphenols and vitamin E, which are lost if the oil is subjected to refining processes. "Extra virgin" olive oils are those that have no taste defects and have a very low acidity rate (0.8%). They are the most expensive ones. Those labeled as "Virgin" olive oils have modest taste defects and a slightly higher acidity level (2%).
- Pure olive oil: which is a blend of refined olive oil and virgin olive oil. In the market, there are varieties of intense flavor and mild taste depending on whether more or less virgin olive oil is added, respectively.
- Olive pomace oil: which is obtained from the pulp and seeds of olives after extraction of virgin olive oil using chemical solvents. It must pass a refining process and virgin olive oil is added after to make it suitable for consumption.

Advantages and disadvantages of EVOO are shown in **Table 1**. Although EVOO is prone to oxidation during production and storage, phenolic compounds have been demonstrated to remain stable under appropriate storage conditions. (26) EVOO quality can be safeguarded by using proper packaging, ideal storage conditions (cool and dark), and having an accurate best before date. (27) Various studies (in vivo and in vitro) have demon-

strated that olive oil phenolic compounds beneficially alter microbial activity, oxidative processes, and inflammation. (26)

The antimicrobial activities of olive oil phenolics were tested against three foodborne pathogenic bacteria: *Escherichia coli* O157:H7, *Listeria monocytogenes*, and *Salmonella enteritidis* in vitro. A synergistic interaction was noted amongst various olive oil phenolic compounds and this synergism appeared to increase antimicrobial capacity compared to that of individual compounds. The study's authors concluded that the use of EVOO in foods may help to prevent foodborne disease. (28)

Emulsions and nanoemulsions

Nanoemulsions are biphasic dispersion of two immiscible liquids: either water in oil (W/O) or oil in water (O/W) droplets stabilized by an amphiphilic surfactant. (29) Nanoemulsions are considered excellent delivery systems for sensitive bioactive compounds improving the solubility and bioavailability of the bioactive compounds. (30) Nanoemulsion has droplets covering the size range of 100-600 nm. In the present work, nanoemulsions were prepared using the spontaneous emulsification mechanism, which occurs when an organic phase and an aqueous phase are mixed. (31,32)

An interesting in vitro and in vivo studies about the influence of the droplet size on the anti-allergic and anti-inflammatory activities of a curcumin NE was conducted by Onodera, et al (2015). The tested curcumin NE was prepared by a thin-film hydration method. Soybean oil and hydrogenated egg yolk phosphatidylcholine (HEPC) were dissolved in the organic solvent chloroform, as curcumin and Tween-80 separately. All three solutions were mixed and dried afterwards by rotary evaporation and vacuum desiccation. The so obtained dry thin film was hydrated with distilled water and sonicated at 25-55 °C till the desired oil droplet size of 50 nm, 100 nm, and 200 nm were obtained. Remarkable, both in vivo tests in orally fed mice and in vitro cell culture tests revealed the advantageous properties of 100 nm curcumin NE despite curcumin NE with 50 nm or 200 nm. (33)

In context to this, nanoemulsions have gained great attention and have been used as a delivery system for not only enhancing the stability, solubility as well as bioavailability of bioactive compounds in body, but also protecting them from adverse environmental conditions (pH, light, moisture, temperature, etc.). Nanoemulsions are used to reduce the separation of particles via sedimentation or creaming and improve the appearance (clear suspension) as well as other physico-chemical proper-

ties of food systems in which bioactive compounds are incorporated. (34)

The result from Rao and McClements (2011) indicated that relatively stable nanoemulsions could be formed at pH 6 and 7, and stable microemulsions at pH 5 and 6, but extensive particle growth/aggregation occurred at lower and higher pH values. Microemulsions were relatively stable to salt addition (0-200 mM NaCl), but nanoemulsions exhibited droplet aggregation/growth at ≥ 50 mM NaCl after 1 month storage at pH 7. Microemulsions formed gels at low temperatures (5 °C), were stable at ambient temperatures (23 °C), and exhibited particle growth at elevated temperatures (40 °C). Nanoemulsions were stable at refrigerator (5 °C) and ambient (23 °C) temperatures, but exhibited coalescence at elevated temperatures (40 °C). (35)

Tan and Nakajima (2005) had successfully prepared b-carotene in nanoemulsions using high-pressure homogenization. They studied the influence of phase ratio and homogenization conditions on droplet size and b-carotene content. (36) Silvia, et al (2011) prepared b-carotene in nanoemulsions using high-energy emulsification with evaporation technique. Their work presented a proof-of-concept of the use of a high-energy emulsification-evaporation technique to produce oil in-water nanoemulsions of b-carotene, without the need of using high-pressure homogenization. The results showed that time and shear rate of homogenization were the most significant processing parameters influencing nanoemulsion size distribution. (37)

Preparation of nanoemulsions

Formulation aspects and method of preparation of nanoemulsion

NEs can be prepared in three ways: (1) oil-in-water (O/W) NEs where oil is disseminated in a continuous aqueous phase; (2) water-in-oil (W/O) NEs where water is dispersed in a continuous oil phase; and (3) the bicontinuous phase. NEs have a variety of shapes as they can be seen as spherical swollen micelles or bicontinuous structures. (38)

Nanoemulsion systems can be formulated by appropriate proportions of water, surfactant, and oil. In order to determine the optimum formula of w/o nanoemulsions, various combinations of oil-surfactant-water were tested. Emulsifiers are surface-active molecules that are capable of adsorbing to droplet surfaces, facilitating droplet disruption, and protecting droplets against aggregation; thus their selection is one of the most crucial factors considering the proper design of a nanoemulsion. (39,40)

Nanoemulsions are prepared using low-energy, high-energy, and combined methods. The high-energy methods include high-shear stirring, ultrasonic emulsification, high-pressure homogenization, in particular, microfluidics and membrane emulsification. The most widely used low-energy methods include the PIT method, the emulsion inversion point (EIP) method, and the spontaneous emulsification in nonequilibrium systems. The combined methods include high-shear stirring to obtain macroemulsions containing special substances (so-called nano-emulsifiers) as the first step. (41)

In the high-energy mechanism, the average droplet size on a nanoscale is difficult to obtain. To overcome this drawback, a multipass regime must be adopted as the maximum degree of dispersion of the system is not reached by the single-pass regime and the efficacy decreases when high viscosity systems are used. (41,42)

Formulation of nanoemulsion includes active drug, additive, and emulsifier. The preparation of nanoemulsion include two methods: (a) high-energy emulsification and (b) low-energy emulsification. (8) The high-energy emulsification method includes high-energy stirring, ultrasonic emulsification, high-pressure homogenization, microfluidization, and membrane emulsification. The low-energy emulsification method includes phase inversion temperature, emulsion inversion point, and spontaneous emulsification. Using a combined method, which includes the high-energy and low-energy emulsification, it is possible to prepare reverse nanoemulsion in a highly viscous system. (8)

High-energy dan low-energy

Three methods are described for nanoemulsion preparation, namely high-energy emulsification (using homogenisers), low-energy emulsification whereby water is added to an oil solution of the surfactant, and the principle of the phase inversion temperature (PIT). (43)

In general, the high-energy process is followed by two steps: first, the deformation and disruption of macrometric droplets into the smaller droplets; second, the surfactant adsorption at their interface (to ensure the steric stabilization). (44) Low-energy process depends on the intrinsic physicochemical properties of the surfactants, cosurfactants, and excipients composing the formulation. (45) Recently, the interest in the low-energy methods for nanoemulsion generation has grown considerably being a mild process for the sensitive molecules and energy-saving process for large-scale production. (41)

Ultrasonic emulsification

Ultrasonic emulsification is very efficient in reducing droplet size. In ultrasonic emulsification, the energy is provided through sonotrodes called as sonicator probe. It contains piezoelectric quartz crystal which can expand and contract in response to alternating electric voltage. As the tip of sonicator contacts the liquid, it produces mechanical vibration and cavitation occurs. Cavitation is the formation and collapse of vapour cavities in liquid. Thus, ultrasound can be directly used to produce emulsion; it is mainly used in laboratories where emulsion droplet size as low as 0.2 micrometer can be obtained. (8) Process mechanism, advantages, and disadvantages of nanoemulsion are shown in **Table 2.** (46)

Nanoemulsion verification

Cryogenic-temperature scanning electron microscopy (cryo-SEM) is an excellent technique for imaging liquid and semi-liquid materials of high vapour pressure, which are highly viscous or contain large (>0.5 µm) aggregates. (47)

On the other side, cryogenic transmission electron microscopy (cryo-TEM) methods have recently begun to be applied to engineered nanoparticles (48). The advantages and disadvantages of cryo-TEM are shown in **Table 3.**

Nutrition in critically ill patients

Enteral nutrition is the preferred method to provide nutritional support in critically ill patients. Despite the advantages of enteral feeding, enteral nutrition may not be tolerated by all patients and may cause complications such as diarrhea, constipation, aspiration, and bowel ischemia. (49) All of these complications make the patient condition becomes worst. The idea of giving nanoemulsion as enteral nutrition to patients with sepsis based on sized and encapsulation is expected to benefit the patient.

Splanchnic ischemia of septic origin and the cascade of biochemical and biologic events leading to multiple-organ dysfunction has reached such proportions in most intensive care units (ICUs) that it is fast becoming the most common cause of death in critically ill surgical patients. Over the last three decades, nutritional support of the critically ill patient has progressed from an afterthought to a vital part of the intensivist's therapeutic armamentarium. The initial purpose of parenteral or enteral nutrition administration was the caloric cover of the energy demands of the patient. Later it became apparent that enterally administered nutritional support prevents mucosal weight loss, increases epithelial proliferation, and improves

maintenance of gut mucosal integrity. Several investigators have suggested that these beneficial effects of intraluminally supplied elements on the gut mucosa are derived from the improved supply of absorbed nutrients to the mucosal epithelium or from the release of trophic hormones. (2) In contrast, this trophic benefit could reflect improved mucosal blood flow due to food absorption-induced hyperemia. (3) Based on that explanation, nanoemulsion may be a good idea for patient in ICU.

The previous article by Sadikot, et al (2017) was one of the example reports of nanotechnology application in critically ill patient. They reported about the nanomedicine for acute respiratory distress syndrome (ARDS) caused by sepsis. ARDS represents an unmet medical need with an urgency to develop effective pharmacotherapies. Multiple promising targets have been identified that could lead to the development of potential therapies for ARDS; however, they have been limited because of difficulty with the mode of delivery, especially in critically ill patients. Nanobiotechnology is the basis of innovative techniques to deliver drugs targeted to the site of inflamed organs, such as the lungs. Nanoscale drug delivery systems have the ability to improve the pharmacokinetics and pharmacodynamics of agents, allowing an increase in the biodistribution of therapeutic agents to target organs and resulting in improved efficacy with reduction in drug toxicity. Although attractive, delivering nanomedicine to lungs can be challenging as it requires sophisticated systems. Here we review the potential of novel nanomedicine approaches that may prove to be therapeutically beneficial for the treatment of this devastating condition. (50)

The illustration we mentioned above gives us some view that nutrition still could be given closely to the physiologic condition. Enteral nutrition with nano scale of delivery system make it possible because the particle size is the primary precondition that can make interaction with immune cell. Nowadays, nanotechnology-based modulation of the immune system is presented as a cutting-edge strategy, which may lead to significant improvements in the treatment of severe diseases. The different immune responses can be elicited by tuning nanosystems properties. (51,52)

Enteral nutrition absorption in sepsis

Possible applications of nanoemulsions are strongly limited by the stability of nanoemulsions, except for nanoparticle preparation when the process of physical or chemical solidification takes place

within the period of stability of nanoemulsions. Nanoemulsions are proposed for numerous applications in pharmacy as drug delivery systems because of their capacity of solubilizing non polar active compounds. Due to the stability problems, most of proposed formulations are self-emulsifying systems and the nanoemulsions are produced just before their application. Although there have not been reported too many applications in other fields, there is a great potential for nanoemulsion applications if Oswald ripening destabilization mechanism is limited by using very insoluble oils. (53) Nanoemulsions can be used for the encapsulation and oral delivery of bioactive lipophilic components, such as nutraceuticals and pharmaceuticals. (54,55)

The previous study by Gosche, et al (1990) mentioned that the hyperemia induced by direct application of glucose to the mucosal surface of the small intestine is able to overcome the flow-restrictive effect of live E coli bacteremia. Restoration of blood flow appears to have been the result of a reversal in arteriolar vasoconstriction that occurs at multiple levels in the intestinal microcirculation during hyperdynamic sepsis. We suggest that this direct hyperemic effect of transmucosal absorption of nutrients on the intestinal microvasculature is one factor that accounts for the maintenance of mucosal integrity associated with enteral alimentation. (56)

The gastrointestinal tract may not absorb with normal efficiency (due to edema, etc.) and insufficient energy is delivered to cope with the metabolic stress. (57) Improved understanding and recent innovations in technology and informatics that are applicable to the bedside promise to advance the precision and appropriateness of our future nutritional care. The idea is to utilize modern advances in technological, molecular science, and informatics to better monitor and deliver nutritional requirements for optimized and nuanced care for the individual through all stages of critical illness. (57) When mucosal immunity is successfully induced, mucosal antibodies (IgA) are produced, and the T-cell-mediated immune response and a systemic antibody response occurs. Oral delivery is considered the gold standard at present, with better patient compliance and a lower social healthcare burden. (58) Here, enteral nanonutrition may have some good impact to the critically ill patient who has poor gastrointestinal tract function based on basic theory in the previous study about mucosal immunity. (59)

Fat metabolism in sepsis

Septicaemia is a major cause of morbidity and mortality among critically ill patients. In the United States alone, the incidence of septic shock is 200,000 per annum and 40 percent of these cases Gram-negative organisms are the cause. Lipid metabolism is altered in septic shock; Gram-negative septicaemia can induce profound change (Table 4). (60,61) Some abnormalities result from failure of specific organs, while others are part of a host response. Increasingly, artificial nutrition is used to meet the raised metabolic fuel demands in affected individuals, and lipids have become an important source of this fuel. The regulation of lipid metabolism in the patient with sepsis is complex and still unclear. In this review an attempt is made to elucidate the mechanisms influencing this regulation, the consequences for the host and the relevance in terms of nutritional support. (62)

The elevation in serum triglycerides in patients with infection caused by Gram-negative bacilli cannot be explained by an increase in absorption of dietary fat, for there were no increase in chylomicrons. However, these elevations were associated with increases in pre-beta-lipoproteins. Pre-beta lipoproteins, which contain 50 to 80 percent triglycerides, are synthesized mainly in the liver. Therefore, it is likely that the elevations in triglyceride levels were associated with decreased peripheral utilization or increase production by the liver that could have been secondary to increased mobilization of fatty acids from peripheral lipid store. (61)

Nanoemulsion with lipid based delivery system is the important thing when we try to modulate cells activity. The factors of nanoparticle that could influenced the cells are particle size, surface charge, surface hydrophilicity/lipophilicity, and antigen-adjuvant binding strength. (63) The scientific report by Bielinska, et al (2014) informed that giving nanoemulsion oil-in-water through nasal administration has a unique mechanism of action. Nanoemulsion activates innate immunity, induces specific gene transcription, and modulates NF-kB activity. (64) This article gives an idea that if we give nanoemulsion as adjunct nutrition through enteral pathway may has the same effect as the reported article by Bielinska, et al (2014). Lipid is needed to manipulate the cell surface and the size of lipid give the opportunity to be absorbed in optimum result. (65,66) The previous study by Serra, et al (2017) reported that extra virgin olive oil nanoemulsion has good effect in modulating immune response induced by dietary and endogenous cholesterol oxidation products in human immune cells and may hold benefit in controlling chronic

immune and/or inflammatory processes. (67)

Particle size and immunology response

The previous research by Bandi, et al (2020) informed that particle size was the most important factor that influence the absorption. (68) This article support that nanonutrition may give the optimal impact to the critically ill patient. Many natural chemicals in food are in the nanometer size range, and the selective uptake of nutrients with nanoscale dimensions by the gastrointestinal tract (GIT) is a normal physiological process. How if the patient is in the abnormal condition, such as sepsis that has GIT failure? We need a mode of delivery that can against the GIT barrier that in the abnormal condition caused by sepsis. The answer may come from nano technology.

The complex pathophysiology of sepsis actually provides many opportunities for treatment at a cellular level, and even at a molecular level. But since these therapies aiming a target at the molecular or cellular level are applied at a systemic level, they generally cannot be translated to safe and effective therapies. Applications of nanotechnology may serve to design better therapies that can apply treatment and can monitor results at the level of the proposed effect, and may instantly modify the amplitude and direction of therapy accordingly at the molecular or cellular level. (69)

Material sizes between 20-40 nm have been found to be an optimal measure of absorption of nanoparticles by dendritic cells. (70,71) The size of the nanoparticles (20-40 nm) penetrates the tissue barrier and circulates through the lymphatic flow faster than the larger nanoparticles (>100 nm). Large nanoparticles are retained by cells and require dendritic cells to facilitate transport to lymph nodes. So that smaller nanoparticles will begin to activate the adaptive immune response more effectively. (72)

Future obsession

Nanoemulsions are of great interest in food indus-

try finding various food applications. Many studies on nanoemulsions have been presented recently, however there are only few references found in food systems, because of the limited number of food grade surfactants able to facilitate the nanoemulsion formulation. Some surfactants are not permitted in foods whereas others may be added only at low concentrations. (73) The distribution of surfactant between the interfacial region and the oil phase depends upon a number of factors, such as: hydrophilic-lipophilic balance (HLB) value of surfactant, nature of oil, interfacial forces, thermal conditions, and interaction between the surfactant and the system composition. According to relevant studies nonionic food grade emulsifiers with HLB values ranging from 6 to 12 are able to form w/o nanoemulsion.

Very rarely patients in ICU without using nasogastric tube. Its condition gives the chance for using nanoemulsion in nutrition composition. The obstacles in the form of various pH condition in human GIT could be avoid. The hope is the absorption will works optimal without any influence of the various pH. But in fact the obstacle comes from the stomach which has pH 1-3, small intestine has pH 6-7.5, and colon pH is 5-7. (38) The challenge of nanoemulsion formulation to be well absorbed is still an ongoing research material. The application of nanonutrition in sepsis patient may be supported by transcellular mecahnisme, typically absorption occurs by an endocytosis mechanism. We hope that nanoemulsion could be expected to do immune regulation by communication between cell and nanoparticle that contain of advantageous bioactive.

Summary

Extra virgin olive oil nanoemulsion is a promising food ingredient to be given to septic patients. Nanoemulsion sized is expected to improve absorption of extra virgin olive oil, while encapsulation form will protect the bioactive content in extra virgin olive oil.

Table 1. Advantages and disadvantages of EVOO

Advantages	Disadvantages	References
<ul style="list-style-type: none"> • More economical and beneficial in the long run • Potential cancer preventative agent • Antidiabetic, cardioprotective, nephroprotective, and neuroprotective effects in the body 	<ul style="list-style-type: none"> • The price of extra virgin olive oil is much more expensive than the other types of oil • Prone to oxidation during production and storage 	17,27,74,75

Legend: EVOO=extra virgin olive oil.

Table 2. Nanoemulsion process mechanism, advantages, and disadvantages

Nanoemulsion process		Advantages	Disadvantages	References
High-energy	Ultrasound generators	<ul style="list-style-type: none"> • Less expensive than the other high-energy equipment • More flexible on surfactant • Internal structure selection than low-energy process 	Limited to small batches	41,44,45,53
	High-pressure homogenization	<ul style="list-style-type: none"> • More flexible on surfactant • Low process time • Internal structure selection than low-energy process 	<ul style="list-style-type: none"> • High cost • Not recommended for thermo- or shear- sensitive compounds 	
	Microfluidization	<ul style="list-style-type: none"> • Controlled size droplets • Allows multiple emulsion preparation 	<ul style="list-style-type: none"> • High cost • Not recommended for large scale production 	
Low-energy	Phase inversion composition	<ul style="list-style-type: none"> • Low cost • Easy to scale up • No need to heat up 	<ul style="list-style-type: none"> • Requires gradual addition of one phase to another • Requires to the presence of LC or mid-range ME phases 	41,44,45,76
	Phase inversion temperature	<ul style="list-style-type: none"> • Low cost • Easy to scale up 	<ul style="list-style-type: none"> • Limited to the nonionic surfactant • Requires to the presence of LC or mid-range ME phases • Heat energy is required 	
	Spontaneous emulsification	<ul style="list-style-type: none"> • Low cost • Easy to scale up 	<ul style="list-style-type: none"> • Limited amount of oil phase • Presence of solvent 	

Legend: LC=liquid crystal; ME=microemulsion.

Table 3. Advantages and disadvantages of cryo-TEM

Advantages	Disadvantages	References
<ul style="list-style-type: none"> • A powerful way to visualize nanostructures in 3D, which can resolve ambiguity present in 2D projection images • Can be applied to nanostructures that have a uniform and homogeneous structure, such as icosahedral viruses, the vault particle, and DNA-origami frames • Provide detailed information about cells and viruses 	<ul style="list-style-type: none"> • Image contrast in 2D cryo-micrographs can be quite low • Frozen specimens are highly beam sensitive and they can only tolerate a low electron dose before significant damage occurs to the sample 	48,77

Legend: cryo-TEM=cryogenic transmission electron microscopy; DNA=deoxyribonucleic acid.

Table 4. Effect of sepsis on lipid metabolism as shown by the concentration of glycerol, non-esterified fatty acids and triacylglycerol, and the rate of appearance of glycerol and non-esterified fatty acids

	Plasma glycerol concentration ($\mu\text{mol/l}$) (7)	Rate of appearance of glycerol ($\mu\text{mol/kg/min}$) (8)	Plasma NEFA concentration ($\mu\text{mol/l}$) (5)	Rate of appearance of NEFA ($\mu\text{mol/kg/min}$) (8)	Plasma TAG concentration ($\mu\text{mol/l}$) (5)
Normal subjects	70 (4) (n=16)	2.4 (0.2) (n=5)	970 (270) (n=5)	6.5 (0.8) (n=5)	1250 (100) (n=5)
Patients with sepsis	130 (22) (n=7)	6.3 (1.1) (n=12)	6450 (1600) (n=13)	13.1 (3.0) (n=12)	5020 (970) (n=13)

Legend: NEFA=non-esterified fatty acids; TAG=triacylglycerol. Values are mean (standard error of the mean).

References

1. Tatsumi H. Enteral tolerance in critically ill patients. *J Intensive Care* 2019;7:1-10.
2. Poulin LF, Lasseaux C, Chamaillard M. Understanding the cellular origin of the mononuclear phagocyte system sheds light on the myeloid postulate of immune paralysis in sepsis. *Front Immunol* 2018;9:1-14.
3. Ding R, Meng Y, Ma X. The Central Role of the Inflammatory Response in Understanding the Heterogeneity of Sepsis-3. *Biomed Res Int* 2018;2018:5086516.
4. Ferreira CD, Nunes IL. Oil nanoencapsulation: development, application, and incorporation into the food market. *Nanoscale Res Lett* 2019;14:9.
5. Liu Q, Huang H, Chen H, Lin J, Wang Q. Food-grade nanoemulsions: Preparation, stability and application in encapsulation of bioactive compounds. *Molecules* 2019;24:1-37.
6. Azmi NAN, Elgharbawy AAM, Motlagh SR, Samsudin N, Salleh HM. Nanoemulsions: Factory for food, pharmaceutical and cosmetics. *Processes* 2019;7:1-34.
7. Sutradhar KB, Amin L. Nanoemulsions: Increasing possibilities in drug delivery. *Eur J Nanomed* 2013;5:97-110.
8. Jaiswal M, Dudhe R, Sharma PK. Nanoemulsion: an advanced mode of drug delivery system. *3 Biotech* 2015;5:123-7.
9. Noguera-Ortega E, Blanco-Cabra N, Rabanal RM, Sánchez-Chardi A, Roldán M, Guallar-Garrido S, et al. Mycobacteria emulsified in olive oil-in-water trigger a robust immune response in bladder cancer treatment. *Sci Rep* 2016;6:1-12.
10. Heyland DK, Dhaliwal R, Drover JW, Gramlich L, Dodek P, Canadian critical care clinical practice guidelines committee. Canadian clinical practice guidelines for nutrition support in mechanically ventilated, critically ill adult patients. *JPEN J Parenter Enteral Nutr* 2003;27:355-73.
11. de Aguilar-Nascimento JE, Kudsk KA. Early nutritional therapy: The role of enteral and parenteral routes. *Curr Opin Clin Nutr Metab Care* 2008;11:255-60.
12. Malone AM. Permissive underfeeding: Its appropriateness in patients with obesity, patients on parenteral nutrition, and non-obese patients receiving enteral nutrition. *Curr Gastroenterol Rep* 2007;9:317-22.
13. Boitano M. Hypocaloric feeding of the critically ill. *Nutr Clin Pract* 2006;21:617-22.
14. McClave SA, Chang W-K. Feeding the Hypotensive Patient: Does Enteral Feeding Precipitate or Protect Against Ischemic Bowel? *Nutr Clin Pract* 2003;18:279-84.
15. de Aguilar-Nascimento JE, Dock-Nascimento DB, Bragagnolo R. Role of enteral nutrition and pharmaconutrients in conditions of splanchnic hypoperfusion. *Nutrition* 2010;26:354-8.
16. Bohlen HG. Intestinal tissue PO₂ and microvascular responses during glucose exposure. *Am J Physiol* 1980;238:H164-71.
17. Sánchez-Villegas A, Sánchez-Tainta A. The Prevention of Cardiovascular Disease Through the Mediterranean Diet [internet]. Amsterdam: Academic Press; 2018. Chapter 4, Virgin Olive Oil: A Mediterranean Diet Essential. [cited 2020 Jan 19]. Available from: https://www.researchgate.net/publication/322172994_Virgin_Olive_Oil
18. Farooqui T, Farooqui AA. Role of the Mediterranean Diet in the Brain and Neurodegenerative Diseases [internet]. Amsterdam: Academic Press; 2018. Chapter 1, Effects of Mediterranean Diet Components on Neurodegenerative Diseases. [cited 2020 Jan 26]. Available from: <http://www.sciencedirect.com/science/article/pii/B9780128119594000018>
19. Boskou D. Olive oil. *World Rev Nutr Diet* 2000;87:56-77.
20. Preedy VR, Watson RR. The Mediterranean Diet: An Evidence-Based Approach [internet]. Amsterdam: Academic Press; 2015. Chapter 13, Olive Oil [Internet]. [cited 2020 Jan 27]. Available from: <http://www.sciencedirect.com/science/article/pii/B9780124078499000130>
21. Benedetti S, Cosio MS, Scampicchio M, Mannino S. Nanoemulsions for the Determination of the Antioxidant Capacity of Oils by an Electrochemical Method. *Electroanalysis* 2012;24:1356-61.
22. Kountouri AM, Mylona A, Kaliora AC, Andrikopoulos NK. Bioavailability of the phenolic compounds of the fruits (drupes) of *Olea europaea* (olives): impact on plasma antioxidant status in humans. *Phytomedicine* 2007;14:659-67.
23. Visioli F, Galli C, Bornet F, Mattei A, Patelli

- R, Galli G, et al. Olive oil phenolics are dose-dependently adsorbed in humans. *FEBS Lett* 2000;468:159-60.
24. Dinnella C, Minichino P, D'Andrea AM, Monteleone E. Bioaccessibility and antioxidant activity stability of phenolic compounds from extra-virgin olive oils during in vitro digestion. *J Agric Food Chem* 2007;55:8423-9.
 25. Servili M, Selvaggini R, Esposto S, Taticchi A, Montedoro GF, Morozzi G. Health and sensory properties of virgin olive oil hydrophilic phenols: Agronomic and technological aspects of production that affect their occurrence in the oil. *J Chromatogr A* 2004;1054:113-27.
 26. Cicerale S, Lucas LJ, Keast RSJ. Antimicrobial, antioxidant and anti-inflammatory phenolic activities in extra virgin olive oil. *Curr Opin Biotechnol* [Internet]. 2012 [cited 2020 Feb 1];23:129-35. Available from: <http://dx.doi.org/10.1016/j.copbio.2011.09.006>
 27. Li X, Wang SC. Shelf Life of Extra Virgin Olive Oil and Its Prediction Models. *J Food Qual* 2018;2018:1-15.
 28. Karaosmanoglu H, Soyer F, Ozen B, Tokatli F. Antimicrobial and antioxidant activities of Turkish extra virgin olive oils. *J Agric Food Chem* 2010;58:8238-45.
 29. Singh Y, Meher JG, Raval K, Khan FA, Chaurasia M, Jain NK, et al. Nanoemulsion: Concepts, development and applications in drug delivery. *J Control Release* [Internet]. 2017 [cited 2020 Jan 22];252:28-49. Available from: <http://dx.doi.org/10.1016/j.jconrel.2017.03.008>
 30. Aqil M, Kamran M, Ahad A, Imam SS. Development of clove oil based nanoemulsion of olmesartan for transdermal delivery: Box-Behnken design optimization and pharmacokinetic evaluation. *J Mol Liq* [Internet]. 2016 [cited 2020 Jan 22];214:238-48. Available from: <http://dx.doi.org/10.1016/j.molliq.2015.12.077>
 31. Bouchemal K, Briçon S, Perrier E, Fessi H. Nano-emulsion formulation using spontaneous emulsification: Solvent, oil and surfactant optimisation. *Int J Pharm* 2004;280:241-51.
 32. Debnath S, Satayanarayana, Kumar GV. Nanoemulsion-A Method to Improve The Solubility of Lipophilic Drugs. *Pharmanest* 2011;2:72-83.
 33. Onodera T, Kuriyama I, Andoh T, Ichikawa H, Sakamoto Y, Lee-Hiraiwa E, et al. Influence of particle size on the in vitro and in vivo anti-inflammatory and anti-Allergic activities of a curcumin lipid nanoemulsion. *Int J Mol Med* 2015;35:1720-8.
 34. Saini A, Panesar PS, Bera MB. Valorization of fruits and vegetables waste through green extraction of bioactive compounds and their nanoemulsions-based delivery system. *Bioresour Bioprocess* [Internet]. 2019 [cited 2020 Mar 1];6:26. Available from: <https://doi.org/10.1186/s40643-019-0261-9>
 35. Rao J, McClements DJ. Food-grade microemulsions, nanoemulsions and emulsions: Fabrication from sucrose monopalmitate & lemon oil. *Food Hydrocoll* [Internet]. 2011 [cited 2020 Jan 21];25(6):1413-23. Available from: <http://dx.doi.org/10.1016/j.foodhyd.2011.02.004>
 36. Tan CP, Nakajima M. Beta-Carotene nano-dispersions: Preparation, characterization, and stability evaluation. *Food Chem* 2005;92:661-71.
 37. Silva HD, Cerqueira MA, Souza BWS, Ribeiro C, Avides MC, Quintas MAC, et al. Nanoemulsions of β -carotene using a high-energy emulsification- evaporation technique. *J Food Eng* 2011;102:130-5.
 38. McClements DJ, Rao J. Food-Grade nano-emulsions: Formulation, fabrication, properties, performance, Biological fate, and Potential Toxicity. *Crit Rev Food Sci Nutr* 2011;51:285-330.
 39. Polychniatou V, Tzia C. Study of formulation and stability of co-surfactant free water-in-olive oil nano- and submicron emulsions with food grade non-ionic surfactants. *J Am Oil Chem Soc* 2014;91:79-88.
 40. Norn V, editor. *Emulsifiers in Food Technology*. 2nd ed. USA: John Wiley & Sons Inc.; 2015.
 41. Koroleva MY, Yurtov EV. Nanoemulsions: the properties, methods of preparation and promising applications. *Russ Chem Rev* [Internet]. 2012 [cited 2020 Feb 7];81(1):21-43. Available from: <https://iopscience.iop.org/article/10.1070/RC2012v081n01ABEH004219>
 42. Wang L, Mutch KJ, Eastoe J, Heenan RK, Dong J. Nanoemulsions prepared by a two-step low-energy process. *Langmuir* 2008;24:6092-9.
 43. Tadros T, Izquierdo P, Esquena J, Solans C. Formation and stability of nano-emulsions. *Adv Colloid Interface Sci* 2004;108-109:303-18.

44. Maali A, Mosavian M. Preparation and Application of Nano-Emulsion in the Last Decade (2000-2010). *J Disper Sci Technol* 2012;34.
45. Anton N, Benoit J-P, Saulnier P. Design and production of nanoparticles formulated from nano-emulsion templates-a review. *J Control Release* 2008;128:185-99.
46. Yukuyama MN, Ghisleni DDM, Pinto TJA, Bou-Chacra NA. Nanoemulsion: Process selection and application in cosmetics - A review. *Int J Cosmet Sci* 2016;38:13-24.
47. Issman L, Talmon Y. Cryo-SEM specimen preparation under controlled temperature and concentration conditions. *J Microsc* 2012;246:60-9.
48. Burrows ND, Penn RL. Cryogenic transmission electron microscopy: aqueous suspensions of nanoscale objects. *Microsc Microanal* 2013;19:1542-53.
49. Gwon J-G, Lee Y-J, Kyoung K-H, Kim Y-H, Hong S-K. Enteral nutrition associated non-occlusive bowel ischemia. *J Korean Surg Soc* 2012;83:171-4.
50. Sadikot RT, Kolanjiyil AV, Kleinstreuer C, Rubinstein I. Nanomedicine for Treatment of Acute Lung Injury and Acute Respiratory Distress Syndrome. *Biomed Hub* 2017;2:1-12.
51. Dacoba TG, Olivera A, Torres D, Crecente-Campo J, Alonso MJ. Modulating the immune system through nanotechnology. *Semin Immunol* 2017;34:78-102.
52. O'Konek JJ, Landers JJ, Janczak KW, Goel RR, Mondrusov AM, Wong PT, et al. Nanoemulsion adjuvant-driven redirection of TH2 immunity inhibits allergic reactions in murine models of peanut allergy. *J Allergy Clin Immunol* 2018;141:2121-31.
53. Gutiérrez JM, González C, Maestro A, Solè I, Pey CM, Nolla J. Nano-emulsions: New applications and optimization of their preparation. *Curr Opin Colloid Interface Sci* 2008;13:245-51.
54. Ostertag F, Weiss J, McClements DJ. Low-energy formation of edible nanoemulsions: Factors influencing droplet size produced by emulsion phase inversion. *J Colloid Interface Sci* [Internet]. 2012 [cited 2020 Feb 7];388(1):95-102. Available from: <http://dx.doi.org/10.1016/j.jcis.2012.07.089>
55. Liu X, Zhang R, McClements DJ, Li F, Liu H, Cao Y, et al. Nanoemulsion-Based Delivery Systems for Nutraceuticals: Influence of Long-Chain Triglyceride (LCT) Type on In Vitro Digestion and Astaxanthin Bioaccessibility. *Food Biophys* 2018;13:412-21.
56. Gosche JR, Garrison RN, Harris PD, Cryer HG. Absorptive Hyperemia Restores Intestinal Blood Flow During Escherichia coli Sepsis in the Rat. *Arch Surg* 1990;125:1573-6.
57. Marini JJ, Debacker D, Gattinoni L, Ince C, Martin-Loeches I, Singer P, et al. Thinking forward: Promising but unproven ideas for future intensive care. *Crit Care* 2019;23(Suppl 1):1-7.
58. Long P, Zhang Q, Xue M, Cao G, Li C, Chen W, et al. Tomato lectin-modified nanoemulsion-encapsulated MAGE1-HSP70/SEA complex protein vaccine: Targeting intestinal M cells following peroral administration. *Biomed Pharmacother* 2019;115:108886.
59. Farazuddin M, Goel RR, Kline NJ, Landers JJ, O'Konek JJ, Baker JR Jr. Nanoemulsion adjuvant augments retinaldehyde dehydrogenase activity in dendritic cells via MyD88 pathway. *Front Immunol* 2019;10:1-11.
60. Banerjee S, Bhaduri JN. Serum Protein-Bound Carbohydrates and Lipids in Cholera. *Proc Soc Exp Biol Med* 1959;101:340-1.
61. Gallin JI, Kaye D, O'Leary WM. Serum lipids in infection. *N Engl J Med* 1969;281:1081-6.
62. Samra JS, Summers LKM, Frayn KN. Sepsis and fat metabolism. *Br J Surg* 1996;83:1186-96.
63. Yan S, Gu W, Xu ZP. Re-considering how particle size and other properties of antigen-adjuvant complexes impact on the immune responses. *J Colloid Interface Sci* 2013;395:1-10.
64. Bielinska AU, Makidon PE, Janczak KW, Blanco LP, Swanson B, Smith DM, et al. Distinct Pathways of Humoral and Cellular Immunity Induced with the Mucosal Administration of a Nanoemulsion Adjuvant. *J Immunol* 2014;192:2722-33.
65. Sravanthi V, Pallavi MCP, Bonam SR, Sathyabama S, Sampath Kumar HM. Oleic acid nanoemulsion for nasal vaccination: Impact on adjuvanticity based immune response. *J Drug Deliv Sci Technol* 2015;28:56-63.
66. Yang Y, Chen L, Sun HW, Guo H, Song Z, You Y, et al. Epitope-loaded nanoemulsion delivery system with ability of extending antigen release elicits potent Th1 response for intranasal vaccine against Helicobacter pylori. *J Nanobiotechnology* 2019;17:1-15.
67. Serra G, Deiana M, Spencer JPE, Corona G. Olive Oil Phenolics Prevent Oxysterol-Induced Proinflammatory Cytokine Secretion and Re-

- active Oxygen Species Production in Human Peripheral Blood Mononuclear Cells, Through Modulation of p38 and JNK Pathways. *Mol Nutr Food Res* 2017;61:1-7.
68. Bandi SP, Kumbhar YS, Venuganti VVK. Effect of particle size and surface charge of nanoparticles in penetration through intestinal mucus barrier. *J Nanopart Res* 2020;22:11051.
 69. Kilic YA, Kilic I, Tez M. Sepsis and multiple organ failure represent a chaotic adaptation to severe stress which must be controlled at nanoscale. *Crit Care* 2009;13:424.
 70. Boraschi D, Italiani P, Palomba R, Decuzzi P, Duschl A, Fadeel B, et al. Nanoparticles and innate immunity: new perspectives on host defence. *Semin Immunol* 2017;34:33-51.
 71. Miquel-Clopés A, Bentley EG, Stewart JP, Carding SR. Mucosal vaccines and technology. *Clin Exp Immunol* 2019;196:205-14.
 72. Smith DM, Simon JK, Baker JR Jr. Applications of nanotechnology for immunology. *Nat Rev Immunol* 2013;13:592-605.
 73. Bera A, Ojha K, Kumar T, Mandal A. Water solubilization capacity, interfacial compositions and thermodynamic parameters of anionic and cationic microemulsions. *Colloids Surfaces A* 2012;404:70-7.
 74. İnanç AL. Chlorophyll: Structural Properties, Health Benefits and Its Occurrence in Virgin Olive Oils. *Akad Gıdatr (AL İnanç)* [Internet]. 2011 [cited 2020 Jan 20];9(2):26-32. Available from: https://www.researchgate.net/publication/267786661_Chlorophyll_Structural_Properties_Health_Benefits_and_Its_Occurrence_in_Virgin_Olive_Oils
 75. Farooqui T, Farooqui AA. Role of the Mediterranean Diet in the Brain and Neurodegenerative Diseases. [Internet]. Amsterdam: Elsevier Inc; 2018. Chapter 13, Effect of Mediterranean Diet Components on Type 2 Diabetes and Metabolic Syndrome. [cited 2020 Jan 20]. Available from: <http://dx.doi.org/10.1016/B978-0-12-811959-4.00013-4>
 76. Solans C, Solé I. Nano-emulsions: Formation by low-energy methods. *Curr Opin Colloid Interface Sci* [Internet]. 2012 [cited 2020 Jan 22];17(5):246-54. Available from: <http://dx.doi.org/10.1016/j.cocis.2012.07.003>
 77. Stewart PL. Cryo-electron microscopy and cryo-electron tomography of nanoparticles. *Wiley Interdiscip Rev Nanomed Nanobiotechnol* 2017;9.