

Novel CHA2DS2-VASc-HSF is Superior to CHADS and CHA2DS-VASc Score to Predict the Risk of Severe Coronary Artery Disease

by Andrianto Andrianto

Submission date: 27-May-2021 01:00PM (UTC+0800)

Submission ID: 1595056500

File name: Score_to_Predict_the_Risk_of_Severe_Coronary_Artery_Disease.pdf (422.84K)

Word count: 5198

Character count: 27492



Novel CHA₂DS₂-VASc-HSF is Superior to CHADS₂ and CHA₂DS₂-VASc Score to Predict the Risk of Severe Coronary Artery Disease

Andrianto Andrianto^{1*}, Benny Jovie², Makhyan Jibril Al-Farabi^{1,3}, Parama Gandi^{1,2}, Khubay Alvia Shonafi¹, Rofda Lathifah⁴

¹Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga – Dr Soetomo Academic General Hospital, Surabaya, Indonesia; ²Department of Cardiology and Vascular Medicine, Dr Ramelan Navy Hospital, Surabaya, Indonesia; ³School of Healthcare Management, University College London, Gower St, Bloomsbury, London, United Kingdom; ⁴Department of Health Policy and Administration, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

Abstract

Edited by: Ksenija Bogojeva-Kostovska
Citation: Andrianto A, Jovie B, Al-Farabi MJ, Gandi P, Shonafi KA, Lathifah R. Novel CHA₂DS₂-VASc-HSF is Superior to CHADS₂ and CHA₂DS₂-VASc Score to Predict the Risk of Severe Coronary Artery Disease. Open Access Maced J Med Sci. 2020 Apr 12; 8(B):451-456. https://doi.org/10.3889/oamjms.2020.3393

Keywords: Atherosclerosis; Coronary occlusion; Gensini score; Risk stratification
***Correspondence:** Andrianto Andrianto, Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo Academic General Hospital, Surabaya, Indonesia. Tel.: +62-31-502-0251 (A). E-mail: andrianto@fk.unair.ac.id

Received: 20-Jul-2019
Revised: 07-Jan-2020
Accepted: 27-Feb-2020

Copyright: © 2020 Andrianto Andrianto, Benny Jovie, Makhyan Jibril Al-Farabi, Parama Gandi, Khubay Alvia Shonafi, Rofda Lathifah

Funding: This research did not receive any financial support

Competing Interests: The authors have declared that no competing interests exist

Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

BACKGROUND: Various risk scoring methods are available to predict the severity of coronary artery disease (CAD). However, the majority of them are complex and require advanced technologies, thus limiting its usage in primary care settings. CHA₂DS₂-VASc-HSF is a novel risk scoring which we develop from CHA₂DS₂-VASc score.

AIM: We hypothesize that CHA₂DS₂-VASc-HSF is predictive for the risk of severe CAD, and we compare its validity with previously established CHADS₂ and CHA₂DS₂-VASc score.

MATERIALS AND METHODS: A total of 210 patients who underwent elective coronary angiography were enrolled in our study. Anthropometric, laboratory, angiographic findings, and patient history were obtained from medical records and used to calculate CHA₂DS₂-VASc-HSF score. Severe CAD defined as coronary artery occlusion with the Gensini score of ≥ 20 . Statistical analyses were done using SPSS 25.0 and MedCalc 18.2.1.

RESULTS: This research showed that the patient with severe CAD has significantly higher CHADS₂, CHA₂DS₂-VASc, and CHA₂DS₂-VASc-HSF score compared to normal and mild CAD ($p < 0.001$). CHADS₂, CHA₂DS₂-VASc, and CHA₂DS₂-VASc-HSF correlated significantly with the CAD severity ($r = 0.315$, $p \leq 0.001$; $r = 0.395$, $p \leq 0.001$; $r = 0.612$, $p \leq 0.001$, respectively). CHA₂DS₂-VASc-HSF may predict the risk of severe CAD independent from other variables (odds ratio = 2.540; 95% confidence interval = 1.794–3.595; $p = 0.002$) with the cutoff value of ≥ 2.5 (sensitivity = 81.4% and specificity = 68.1%). Pairwise comparison of receiver operating characteristic curves showed that CHA₂DS₂-VASc-HSF was superior to predict severe CAD.

CONCLUSIONS: CHA₂DS₂-VASc-HSF scores may predict the risk of severe CAD better than CHADS₂ and CHA₂DS₂-VASc score. This score may easily be used in primary care physicians to predict the risk of severe CAD and provide an early referral to the cardiologist.

Introduction

Coronary artery disease (CAD) remained the top cause of mortality and morbidity for a person aged 35 and over worldwide [1], [2]. Failure to detect CAD and provide early treatment may cause CAD treatment to become a more expensive and higher mortality rate [2]. It is estimated that around one-third of the middle age's population in the USA will suffer from CAD manifestation [3]. In Indonesia, the Indonesian Ministry of Health also showed that CAD is the leading cause of morbidity, which is responsible for 12.9% of death [4].

Determining the best risk factor assessment for CAD is extremely important for early prevention and treatment. The screening for CAD using angiography is easily available in developed countries with short waiting lists. However, in developing countries, the awareness and accessibility of cardiovascular disease screening are still low [5]. To obtain cost-effective prevention and treatment of CAD at the patient level, stratification of the

cardiovascular risk using a simple method is extremely important. Cardiovascular risk screening will have a relevant implication for decision making in early referral and healthcare resource allocation [6].

At present, CHADS₂ and CHA₂DS₂-VASc scores have been established as clinical predictors for cardiac thromboembolism and indication of antithrombotic therapy [7]. Both CHADS₂ and CHA₂DS₂-VASc component has similarities with the risk factors of CAD development [8]. The components within the CHADS₂ score also had been proven in large cohort studies to be associated with CAD in with ischemic stroke patients [9], while CHA₂DS₂-VASc is the refinement of CHADS₂ score, which has been proven to outperform its predecessor in the various patient group, including AF patient who received elective electrical cardioversion [10]. This suggests CHA₂DS₂-VASc score may predict the risk for both cerebrovascular and cardiovascular diseases. However, these scores did not include the major risk factors of CAD such as smoking, hyperlipidemia, and family histories. Hence, this

research aims to improve the validity of the CHA₂DS₂-VASc score by including new major risk factor of CAD which are hyperlipidemia (H), smoking (S), and family history of CAD (F) and compare it with the previous CHADS₂ and CHA₂DS₂-VASc score to predict severe CAD in the patients.

Materials and Methods

Study design

50 This cross-sectional study involves 210 consecutive patients who underwent coronary angiography in the Cardiology Department of Ramelan Navy Hospital Surabaya during 1 year period between January-December 2018. Coronary artery occlusion was assessed from angiograms using the Gensini score. Patients with infectious processes within 2 weeks before catheterization, hepatic dysfunction, thyroid dysfunction, cancer, and chronic kidney disease were excluded from the study. This study had received ethical clearance (No.06/EC/KERS/2019) from the local ethics committee. Informed consents were obtained and details which disclose patients' identity were omitted.

Risk factor data collection

Clinical findings, 12-lead electrocardiogram, and echocardiographic examination were performed based on the American Society of Echocardiography guidelines [11]. The standard laboratory was performed to measure fasting blood glucose (FBG), total cholesterol, and renal function tests from the blood samples [12].

CHA₂DS₂-VASc-HSF score consists of congestive cardiac failure (C), hypertension (H), age >75 years (A), diabetes mellitus (D), stroke (S), vascular diseases (V), age 65–74 years (A), sex category (Sc), hyperlipidemia (H), smoking (S), and family history of cardiovascular disease (F) were obtained by medical record thorough examination. Congestive cardiac failure (C) score was given if left ventricular ejection fraction was reduced (<45%) from echocardiography examination. Hypertension (H) was defined as systolic blood pressure >140 mmHg or diastolic <90 mmHg for repeated measurement, or when the patient was taking antihypertensive medications. Diabetes mellitus (D) Type 2 was defined FBG >126 mg/dl, previous diabetes diagnosis, or when the patient was taking anti-diabetic medications. Stroke (S) was defined as the history or current diagnosis of stroke or TIA which was given by the patients. Vascular disease (V) was defined from the existence of a pathologic condition which causes stenosis of at least 50% in the

non-coronary artery. Hyperlipidemia (H) defined as a cholesterol level of more than 200 mg/dL based on the National Cholesterol Education Program or when the patient is consuming of lipid-lowering medications. Cigarette smoking (S) was defined as the habit of smoking of more than five cigarettes per-day without a quit attempt for a minimum of 1 year. Family history of cardiovascular disease (F) was defined as the presence of cardiovascular disease or sudden cardiac-related death of the first degree-relative.

CHA₂DS₂-VASc-HSF scoring

CHA₂DS₂ score was calculated by adding 1 point for the presence of chronic heart failure, age >75 years, DM and hypertension by assigning 2 points for the history of stroke or TIA. In the CHA₂DS₂-VASc score, age 65–74 was assigned for 1 point (A) and age >75 years (A₂) was assigned for 2 points. The CHA₂DS₂-VASc-HSF score put 1 point for the finding of hyperlipidemia (h), smoking (S), and family history of the cardiac disease (F).

Coronary angiography and Gensini scoring

Judkins technique 4 with 5-F catheters was used to perform cannulation of coronary arteries. Kodak 35-mm cinefilm was used to record the images at 30 frames/s. Computer-assisted coronary angiography analysis system was used to detect coronary stenosis (Mipron 1; Kontron, Tokyo, Japan). One minute after the injection of ISDN (2.5 mg/5 mL for 20 s) through the Judkins catheter, several projections were taken to observe the coronary angiography. Coronary atherosclerosis severity was measured using the Gensini scoring method, as described previously [12].

Calculation of the Gensini score was done for each patient through the severity score assignment based on coronary occlusion. Narrowing between 1 and 25% will be scored 1, 26–50% will be scored 2, 51–75% will be scored 3, 75–90% will be scored 8, 91–99% will be scored 16, and 100% will be scored 32. The score is then multiplied based on the location and importance of the artery. We multiply by factor 5 for left main coronary artery occlusion, 2.5 for both proximal circumflex artery and proximal left anterior descending artery, 1.5 for a mid-left anterior descending artery, and 1 mid or distal circumflex artery, for distal left anterior descending artery and the right coronary artery. The multiplication factor for any other branch is 0.5 [3].

7 Statistical analyses

Data analyses were performed using SPSS Statistics 25.0 and MedCalc 18.2.1. Continuous variables, presented as mean ± standard deviation (SD), were compared using ANOVA test. Correlation between

parametric variables was obtained using Spearman's Rho followed by logistic regression. Specificity and sensitivity were obtained from the receiver operating characteristic (ROC) curve and cutoff point analysis. Area under the curve (AUC) comparison was done using the pairwise comparison method as described previously [13].

Results

Clinical characteristics of the patients

The total of 210 patients was involved in this study. Table/Figure 1 shows the characteristics of the participant, which grouped based on the CAD severity. Of the 210 patients, 70 patients had normal angiogram (Gensini score = 0, 33.3%), 48 patient had mild CAD (Gensini score = 1–19, 22.9%), and 92 patients had obstructive/severe CAD (Gensini score >20, 43.8%). The comparison of the baseline demographics and characteristics of the three groups (normal coronary arteries, mild CAD, and severe CAD) is presented in Table 1.

From Table 1, significant differences between severe CAD and normal angiography groups were

observed on the age, FBG, ureum, and creatinine which are the CAD risk factors. CHADS₂, CHA₂DS₂-VASc, and CHA₂DS₂-VASc-HSF score also significantly higher on the patient with severe CAD compared to the patient with normal angiography.

Correlations between multiple variables with CAD severity

The correlation test was used to identify the factors associated with the severity of CAD. Table 2 shows the results of Spearman's correlations between Gensini score and multiple independent variables in the subjects. Spearman's correlation analysis showed that the highest correlation was identified on the CHA₂DS₂-VASc-HSF score with Gensini score, which showed a moderate to strong correlation ($r = 0.612$, $p = \leq 0.001$).

Logistic linear regression analysis of the variables to predict severe CAD

Univariate and multivariate logistic linear regression analysis was done on various variables in predicting the outcome (severe CAD) as presented in Tables 3 and 4. The analysis from Tables 3 and 4 showed that CHADS₂, CHA₂DS₂-VASc, and CHA₂DS₂-VASc-HSF were significant predictors for severe CAD.

Specificity and sensitivity test using ROC curves

From the ROC curves in Figure 1, it is suggested that CHA₂DS₂-VASc-HSF score has higher AUC compared to CHADS₂ and CHA₂DS₂-VASc score. Optimum cutoff point analysis showed that the CHA₂DS₂-VASc-HSF score ≥ 2.5 provided the highest predictive value for severe CAD (sensitivity = 81.4% and specificity = 68.1%). Pairwise comparison from Table 5 showed that the CHA₂DS₂-VASc-HSF score was found to be the best scoring scheme to predict severe CAD compared to CHADS₂ and CHA₂DS₂-VASc score.




Figure 1: Receiver operating characteristic curve of CHADS₂, CHA₂DS₂-VASc, and CHA₂DS₂-VASc-HSF score to predict severe CAD. AUC: Area under the curve; CI: Confidence interval

Table 1: Characteristics of the patients based on CAD severity

Variables*	CAD severity			p-value
	Normal angiogram (n=70)	Mild CAD (n=48)	Severe CAD (n=92)	
Age (years)	50.97 ± 8.95 ^{ab}	53.87 ± 10.90 ^c	54.65 ± 9.13 ^c	≤ 0.001
SBP (mmHg)	131.81 ± 26.17	123.00 ± 16.67	130.90 ± 23.05	0.691
DBP (mmHg)	72.81 ± 10.86	76.53 ± 9.12	77.58 ± 9.23	0.463
Weight (kg)	69.48 ± 12.43	65.27 ± 7.70	70.03 ± 9.88	0.369
Height (m)	164.06 ± 8.59	164.33 ± 3.90	165.45 ± 5.79	0.208
BMI (kg/m ²)	25.67 ± 3.26	24.22 ± 3.28	25.55 ± 3.15	0.341
Hb (g/dL)	14.01 ± 1.11	13.63 ± 1.73	14.60 ± 3.75	0.474
WBC (cells/ μ L)	7434.2 ± 1725.5	6746.0 ± 1624.1	7350.1 ± 2108.5	0.292
HCT (%)	41.63 ± 3.64	41.20 ± 4.89	42.46 ± 4.36	0.897
Platelet ($\times 10^3$ cells/ μ L)	273.00 ± 78.55	271.66 ± 47.73	258.12 ± 52.97	0.268
PT (s)	13.31 ± 1.28	13.56 ± 2.67	13.56 ± 3.46	0.301
APTT (s)	32.56 ± 3.23	31.33 ± 6.19	33.39 ± 5.27	0.097
FBG (mg/dL)	108.06 ± 37.97 ^a	114.13 ± 32.82	112.45 ± 34.39 ^a	0.032
Ureum (mg/dL)	12.45 ± 3.15 ^{bc}	16.81 ± 5.20 ^c	16.45 ± 7.70 ^c	0.005
Creatinine (mg/dL)	1.00 ± 0.31 ^{bc}	1.27 ± 0.25 ^c	1.16 ± 0.26 ^c	0.005
Ejection fraction (%)	57.65 ± 18.31	64.73 ± 5.96	52.15 ± 16.85	0.474
CHADS ₂ Score	0.68 ± 0.65 ^{bc}	0.60 ± 0.99 ^a	1.08 ± 0.94 ^a	≤ 0.001
CHA ₂ DS ₂ -VASc Score	1.32 ± 0.83 ^{bc}	1.47 ± 1.06 ^a	1.95 ± 1.04 ^a	≤ 0.001
CHA ₂ DS ₂ -VASc-HSF Score	2.06 ± 0.77 ^{bc}	2.33 ± 1.23 ^a	3.43 ± 1.20 ^{ab}	≤ 0.001

SBP: Systolic blood pressure; DBP: Diastolic blood pressure; BMI: Body mass index, Hb: Hemoglobin; WBC: White blood cells; HCT: Hematocrit; PT: Prothrombin time; APTT: Activated partial thromboplastin time; FBG: Fasting blood glucose; SD: Standard deviation. Values are presented as a mean \pm SD; different annotation showed a significant difference ($p < 0.05$) for the post hoc LSD test to normal angiogram (a), mild CAD (b) and severe CAD (c).

Discussion

The major findings of this research were: (1) CHA₂DS₂-VASc-HSF score was significantly increased both mild and severe CAD patients, (2) the CHADS₂, **Table 2: Spearman's correlations between various independent variables with CAD Gensini score**

Variables*	Correlation coefficient (r)
Age (years)	0.276**
FBG (mg/dL)	0.180*
Urem (mg/dL)	0.232**
Creatinine (mg/dL)	0.204**
WBC	0.236**
Ejection fraction (%)	-0.215**
CHA ₂ DS ₂	0.315**
CHA ₂ DS ₂ -VASc	0.395**
CHA ₂ DS ₂ -VASc-HSF	0.612**

*Significant correlation at p<0.05, **Significant correlation at p<0.01. FBG: Fasting blood glucose; WBC: White blood cells

CHA₂DS₂-VASc, and CHA₂DS₂-VASc-HSF scores have positive and significant correlation with CAD severity measured by Gensini score, (3) CHADS₂, CHA₂DS₂-VASc, **Table 3: Univariate regression analysis for the predictors of severe CAD**

Variables	Odds ratio	95% CI	p-value
Age (years)	1.036	1.007-1.066	0.013
SBP (mmHg)	1.003	0.992-1.015	0.585
DBP (mmHg)	1.015	0.990-1.042	0.242
BMI (kg/m ²)	1.016	0.933-1.106	0.721
Hb (g/dL)	1.072	0.861-1.335	0.551
WBC (cells/μL)	1.000	1.000-1.000	0.676
HCT (%)	0.995	0.899-1.102	0.929
Platelet (x 10 ³ cells/μL)	1.000	1.000-1.000	0.615
FBG (mg/dL)	1.009	1.001-1.017	0.024
Urem (mg/dL)	1.103	1.045-1.164	≤0.001
Creatinine (mg/dL)	5.000	1.756-14.233	0.005
Ejection fraction (%)	0.390	0.059-2.563	0.474
CHA ₂ DS ₂ score	1.834	1.304-2.580	≤0.001
CHA ₂ DS ₂ -VASc score	1.962	1.455-2.644	≤0.001
CHA ₂ DS ₂ -VASc-HSF score	2.716	1.996-3.696	≤0.001

SBP: Systolic blood pressure; DBP: Diastolic blood pressure; BMI: Body mass index; Hb: Hemoglobin; WBC: White blood cells; HCT: Hematocrit; PT: Prothrombin time; APTT: Activated partial thromboplastin time; FBG: Fasting blood glucose.

and CHA₂DS₂-VASc-HSF are significant predictors for severe CAD, the highest odds ratio was found on the CHA₂DS₂-VASc-HSF score, and (4) CHA₂DS₂-VASc-HSF was the best score to predict severe CAD with the cutoff point of ≥2.5.

Table 4: Multivariate regression analysis for the predictors of severe CAD

Variables	Odds ratio	95% CI	p-value
With CHADS ₂ score			
Age (years)	1.017	0.985-1.049	0.298
FBG (mg/dL)	1.002	0.994-1.011	0.588
Urem (mg/dL)	1.067	1.008-1.130	0.026
Creatinine (mg/dL)	2.828	0.893-8.960	0.077
CHADS ₂ Score	1.572	1.009-2.345	0.046
With CHA ₂ DS ₂ -VASc score			
Age (years)	1.008	0.976-1.042	0.613
FBG (mg/dL)	1.002	0.994-1.010	0.601
Urem (mg/dL)	1.065	1.006-1.129	0.032
Creatinine (mg/dL)	2.323	0.731-7.380	0.153
CHA ₂ DS ₂ -VASc score	1.569	1.098-2.240	0.013
With CHA ₂ DS ₂ -VASc-HSF score			
Age (years)	0.997	0.962-1.033	0.871
FBG (mg/dL)	0.997	0.989-1.006	0.516
Urem (mg/dL)	1.065	0.998-1.137	0.056
Creatinine (mg/dL)	1.511	0.421-5.419	0.527
CHA ₂ DS ₂ -VASc-HSF Score	2.540	1.794-3.595	0.002

FBG: Fasting blood glucose; CI: Confidence interval.

Table 5: Pairwise comparison between receiver operating characteristic curves

Variables	Differences between areas	SE	95% CI	Z-Statistic	p-value
CHADS ₂ and CHA ₂ DS ₂ -VASc	0.0496	0.0227	0.0052-0.941	2.190	0.028
CHADS ₂ and CHA ₂ DS ₂ -VASc-HSF	0.154	0.0320	0.0914-0.217	4.819	≤0.001
CHA ₂ DS ₂ -VASc and CHA ₂ DS ₂ -VASc-HSF	0.105	0.0255	0.0546-0.154	4.105	≤0.001

SE: Standard error; CI: Confidence interval.

The severe CAD may be fatal if remained undiagnosed and developed further into coronary total occlusion, which caused myocardial infarction. Hence, early detection of severe CAD is extremely important to prevent the mortality and morbidity of the patients [3]. Coronary angiography is the gold standard to diagnose the severity of stable CAD. However, early coronary angiography screening is lacking in developing countries [14]. Hence, clinicians need reliable, simple, objective, and quantitative tools to identify these risk stratifications to refer the patient for early screening, modify the risk factor, and provide early treatment [15]. Several scoring systems which involve major risk factors such as European SCORE and Framingham risk score (FRS) have been developed to assess the risk of CAD [16]. FRS is the most widely used score, which estimates the 10-year risk of developing CAD risk. However, this score cannot assess the severity of CAD. Furthermore, FRS also overestimates cardiovascular mortality rates in a low-risk population and underestimates it at the high-risk populations [17], [18]. Due to its multiplicity and complexity, FRS, SCORE, and other scoring systems are considered to be unpractical for daily use for primary care physician [19], [20]. Hence, alternatives scoring such as that CHADS₂, CHA₂DS₂-VASc, and CHA₂DS₂-VASc-HSF may offer a better alternative, which is easily be applied by the physician without any additional cost.

This study showed that CHADS₂, CHA₂DS₂-VASc, and CHA₂DS₂-VASc-HSF are having a positive and significant correlation with CAD severity measured by Gensini score. The highest correlation was found on the CHA₂DS₂-VASc-HSF score. This was in accordance with the previous research that showed both CHADS₂ and CHA₂DS₂-VASc score are significantly correlated with the Gensini score with almost similar r-value (r = 0.383, p < 0.001; r = 0.300, p = 0.001) [15]. When CHA₂DS₂-VASc score is modified by adding hyperlipidemia (H) and smoking (S), a stronger and significant correlation was found between CHA₂DS₂-VASc-HS score and Gensini score (r = 0.813, p < 0.001) [21]. Similar to this research, CHA₂DS₂-VASc-HSF also has a stronger and significant correlation with the severity of CAD measured by syntax score in NSTEMI patients compared to CHADS₂ and CHA₂DS₂-VASc [22]. This suggested that CHA₂DS₂-VASc-HSF score has a superior association with CAD severity compared to CHADS₂ and CHA₂DS₂-VASc score.

We investigated whether the CHADS₂-VASc-HSF scores could aid clinicians to predict the patient who has higher odds of severe CAD, which need immediate diagnosis and treatments. This research showed that CHA₂DS₂-VASc-HSF has the highest odd

ratio to predict the severe CAD compared to CHADS₂ and CHA₂DS₂-VASc. Previously, the CHADS₂ score, as one of the rapid and very practical scores for risk stratification for thromboembolism, has also shown able to predict CAD in ischemic stroke patients [23]. The development of CHADS₂ into CHA₂DS₂-VASc score also has been shown to have better predictive power for long-term mortality for patients with severe CAD [24]. This suggested that the improvement of CHA₂DS₂-VASc into CHA₂DS₂-VASc-HSF may provide a better prediction for the odds of having severe CAD.

The CHADS₂ score is considered as one of the rapid, wide range, and very practical for risk stratification for thromboembolism, which was also developed to predict CAD in ischemic stroke patients [23]. CHA₂DS₂-VASc is the development of the CHADS₂ score, which showed better predictive power for long-term mortality for patients with CAD [24]. When compared with TIMI, GRACE score, CHA₂DS₂-VASc showed that the capability to predict severe CAD measured by syntax score [25]. Modification of CHA₂DS₂-VASc into CHA₂DS₂-VASc-HS score also has been shown to improve its predictive value for severe CAD compared to both CHADS₂ into CHA₂DS₂-VASc with the sensitivity of 85.2% and specificity of 57.5% at the cutoff value of >2 [13]. In this research, modification of CHA₂DS₂-VASc into CHA₂DS₂-VASc-HSF score also showed a higher AUC area compared to both CHADS₂ into CHA₂DS₂-VASc score with the sensitivity of 81.4% and specificity of 68.1% at the cutoff value of >2.5. CHA₂DS₂-VASc-HSF also shown to be the best scoring scheme for severe CAD prediction compared to CHADS₂ and CHA₂DS₂-VASc score. This suggests that CHA₂DS₂-VASc-HSF score is a better scoring method which easily used by the physician to screen the patient with angina, which may require referral for coronary angiography and early treatment.

However, this study may yet to be generalized since it only involved a single-center as the source of data. This study also used consecutive samplings from all patients who were admitted for diagnostic coronary angiography. Hence, selection bias might occur. In the future, it is suggested to involve more cardiac center and stratify the sample based on several factors such as race and social status to ensure the validity of the score among various demographic characteristics.

Conclusions

The CHA₂DS₂-VASc-HSF score can predict the severe CAD with superior validity compared to CHADS₂ and CHA₂DS₂-VASc score. Suggesting that CHA₂DS₂-VASc-HSF score may be recommended for primary care physicians to easily predict severe CAD and refer them earlier without additional costs.

Authors' Contributions

1. Conceptualization, A. and M.J.A.; Methodology, R.L.; Software, M.J.A. and R.L.; Validation, A. and B.J.; Formal analysis, M.J.A. and R.L.; Investigation, I.G.P.G.S., M.J.A., and K.A.S.; Resources, A. and B.J.; Data curation, I.G.P.G.S. and K.A.S.; writing—original draft preparation, I.G.P.G.S. and K.A.S.; writing—review and editing, A. and M.J.A.; visualization, R.L.; supervision, A. and B.J.; project administration, R.L.

References

- Rosamond W, Flegal K, Furie K, Go A, Greenlund K, Haase N, et al. Heart disease and stroke statistics-2008 update: A report from the American heart association statistics committee and stroke statistics subcommittee. *Circulation*. 2008;117(4):e25-146. <https://doi.org/10.1161/circulationaha.107.187998> PMID:18086926
- Benjamin EJ, Virani SS, Callaway CW, Chamberlain AM, Chang AR, Cheng S, et al. Heart disease and stroke statistics-2018 update: A report from the American heart association. *Circulation*. 2018;137(12):e67-492. <https://doi.org/10.1161/cir.0000000000000573> PMID:29386200
- Lloyd-Jones DM, Larson MG, Beiser A, Levy D. Lifetime risk of developing coronary heart disease. *Lancet*. 1999;353(9147):89-92. [https://doi.org/10.1016/s0140-6736\(98\)10279-9](https://doi.org/10.1016/s0140-6736(98)10279-9) PMID:10023892
- Ministry of Health Indonesia. Sample Registration System 2014. Indonesia: NIHRD Library Cataloguing in Publication Data; 2015. Available from: <https://www.litbang.kemkes.go.id/wp-content/uploads/2017/10/final-report-srs-2014.pdf>. [Last accessed on 2019 Mar20].
- Kočka V. The coronary angiography-an old-timer in great shape. *Cor et Vasa*. 2015;57(6):e419-24. <https://doi.org/10.1016/j.crvasa.2015.09.007>
- Modesti PA, Agostoni P, Agyemang C, Basu S, Benetos A, Cappuccio FP, et al. Cardiovascular risk assessment in low-resource settings: A consensus document of the European society of hypertension working group on hypertension and cardiovascular risk in low resource settings. *J Hypertens*. 2014;32(5):951-60. <https://doi.org/10.1097/hjh.000000000000125> PMID:24577410
- January CT, Wann LS, Calkins H, Chen LY, Cigarroa JE, Cleveland JC Jr., et al. 2019 focused update on atrial fibrillation AHA/ACC/HRS focused update of the 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: A report of the American college of cardiology/American Heart Association task force on clinical practice guidelines and the heart rhythm society. *J Am Coll Cardiol* 2019;140:e125-e51. <https://doi.org/10.1161/cir.0000000000000665>
- Henriksson KM, Farahmand B, Johansson S, Asberg S, Terént A, Edvardsson N. Survival after stroke-the impact of CHADS₂ score and atrial fibrillation. *Int J Cardiol*. 2010;141(1):18-23. <https://doi.org/10.1016/j.ijcard.2008.11.122> PMID:19144430
- Loh E, Sutton MS, Wun CC, Rouleau JL, Flaker GC, Gottlieb SS, et al. Ventricular dysfunction and the risk of stroke after myocardial infarction. *N Engl J Med*. 1997;336(4):251-7.

- <https://doi.org/10.1056/nejm199701233360403>
PMid:8995087
10. Yarmohammadi H, Varr BC, Puwanant S, Lieber E, Williams SJ, Klostermann T, *et al.* Role of CHADS2 score in evaluation of thromboembolic risk and mortality in patients with atrial fibrillation undergoing direct current cardioversion (from the ACUTE trial substudy). *Am J Cardiol.* 2012;110(2):222-26. <https://doi.org/10.1016/j.amjcard.2012.03.017>
PMid:22503581
 11. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, *et al.* Recommendations for chamber quantification: A report from the American society of echocardiography's guidelines and standards committee and the chamber quantification writing group, developed in conjunction with the European association of echocardiography, a branch of the European Society of cardiology. *J Am Soc Echocardiogr.* 2005;18(12):1440-63. <https://doi.org/10.1016/j.echo.2005.10.005>
PMid:16376782
 12. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem.* 1972;18(6):499-502. <https://doi.org/10.1093/clinchem/18.6.499>
PMid:4337382
 13. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: A nonparametric approach. *Biometrics.* 1988;44(3):837-45. <https://doi.org/10.2307/2531595>
PMid:3203132
 14. Zhao Y, Shaista M, Nathan D. Evidence for coronary artery calcification screening in the early detection of coronary artery disease and implications of screening in developing countries. *Glob Heart.* 2014;9(4):399-407. <https://doi.org/10.1016/j.ghart.2014.10.005>
PMid:25592793
 15. Cetin M, Cakici M, Zencir C, Tasolar H, Baysal E, Balli M, *et al.* Prediction of coronary artery disease severity using CHADS2 and CHA2DS2-VASc scores and a newly defined CHA2DS2-VASc-HS score. *Am J Cardiol.* 2014;113(6):950-6. <https://doi.org/10.1016/j.amjcard.2013.11.056>
PMid:24444782
 16. Anderson KM, Odell PM, Wilson PW, Kannel WB. Cardiovascular disease risk profiles. *Am Heart J.* 1991;121(1):293-8.
PMid:1985385
 17. Menotti A, Puddu PE, Lanti M. Comparison of the framingham risk function-based coronary chart with risk function from an Italian population study. *Eur Heart J.* 2000;21(5):365-70. <https://doi.org/10.1053/euhj.1999.1864>
PMid:10666350
 18. Onat A, Can G, Hergenç G, Küçükduymaz Z, Ugur M, Yüksel H. High absolute coronary disease risk among Turks: Involvement of risk factors additional to conventional ones. *Cardiology.* 2010;115(4):297-306. <https://doi.org/10.1159/000312009>
PMid:20395681
 19. Siontis GC, Tzoulaki I, Siontis KC, Ioannidis JP. Comparisons of established risk prediction models for cardiovascular disease: Systematic review. *BMJ.* 2012;344:e3318. <https://doi.org/10.1136/bmj.e3318>
PMid:22628003
 20. D'Agostino RB Sr., Vasan RS, Pencina MJ, Wolf PA, Cobain M, Massaro JM, *et al.* General cardiovascular risk profile for use in primary care: The framingham heart study. *Circulation.* 2008;117(6):743-53. <https://doi.org/10.1161/circulationaha.107.699579>
PMid:18212285
 21. Naik G, Sekhar S, Prabhu M, Singh G, Vijay A, Subrahmanian M, *et al.* Comparison of CHA2DS2-VASc-HS score and gensini score to predict severity of coronary artery disease. *J Clin Diagn Res.* 2018;12(7):IC01-4. <https://doi.org/10.7860/jcdr/2018/28509.11802>
 22. Al-shorbagy AN, Al-Cekelly MM, Dwedar AA, Soliman MM. The predictive value of newly defined CHA2DS2-VASc-HSF score for severity of coronary artery disease in non ST segment elevation myocardial infarction. *Zagazig Univ Med J.* 2018;24(4):289-96. <https://doi.org/10.21608/zumj.2018.13209>
 23. Poci D, Hartford M, Karlsson T, Herlitz J, Edvardsson N, Caidahl K. Role of the CHADS2 score in acute coronary syndromes: Risk of subsequent death or stroke in patients with and without atrial fibrillation. *Chest.* 2012;141(6):1431-40. <https://doi.org/10.1378/chest.11-0435>
PMid:22016485
 24. Kim KH, Kim W, Hwang SH, Kang WY, Cho SC, Kim W, *et al.* The CHA2DS2VASc score can be used to stratify the prognosis of acute myocardial infarction patients irrespective of presence of atrial fibrillation. *J Cardiol.* 2015;65(2):121-7. <https://doi.org/10.1016/j.jjcc.2014.04.011>
PMid:24972564
 25. Bozbay M, Uyarel H, Cicek G, Oz A, Keskin M, Murat A, *et al.* CHA2DS2-VASc score predicts in-hospital and long-term clinical outcomes in patients with st-segment elevation myocardial infarction who were undergoing primary percutaneous coronary intervention. *Clin Appl Thromb Hemost.* 2017;23(2):132-8. <https://doi.org/10.1177/1076029616646874>
PMid:27170782

Novel CHA2DS2-VASc-HSF is Superior to CHADS and CHA2DS - VASc Score to Predict the Risk of Severe Coronary Artery Disease

ORIGINALITY REPORT

16%

SIMILARITY INDEX

10%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	en.wikipedia.org Internet Source	1%
2	www.researchsquare.com Internet Source	1%
3	www.e-dmj.org Internet Source	<1%
4	Mikail Yarlioglues, Fatih Oksuz, Damla Yalcinkaya, Mustafa Duran, Sani Namik Murat. "CHA2DS2-Vasc score and saphenous vein graft disease in patients with coronary artery bypass graft surgery", Coronary Artery Disease, 2020 Publication	<1%
5	www.middleeastmedicalportal.com Internet Source	<1%
6	J. Waller, L. A. V. Marlow, J. Wardle. "The association between knowledge of HPV and	<1%

feelings of stigma, shame and anxiety",
Sexually Transmitted Infections, 2006

Publication

7

Mohsen Kerkeni, Izabella Santos Weiss,
Stephane Jaisson, Azza Dandana, Faouzi
Addad, Philippe Gillery, Mohamed Hammami.

"Increased serum concentrations of
pentosidine are related to presence and
severity of coronary artery disease",
Thrombosis Research, 2014

Publication

8

ehnheart.org

Internet Source

9

www.banglajol.info

Internet Source

10

Eser Durmaz, Baris Ikitimur, Burcak Kilickiran
Avci, Adem Atıcı et al. "The clinical significance
of premature atrial contractions: How
frequent should they become predictive of
new - onset atrial fibrillation", Annals of
Noninvasive Electrocardiology, 2019

Publication

11

Paul L. Hess, Melissa A. Greiner, Gregg C.
Fonarow, Winslow Klaskala et al. "Outcomes
Associated With Warfarin Use in Older
Patients With Heart Failure and Atrial
Fibrillation and a Cardiovascular Implantable
Electronic Device: Findings From the ADHERE

<1 %

<1 %

<1 %

<1 %

<1 %

Registry Linked to Medicare Claims", Clinical Cardiology, 2012

Publication

12

Stéphane Jaisson, Mohsen Kerkeni, Izabella C.R. Santos-Weiss, Faouzi Addad, Mohammed Hammami, Philippe Gillery. "Increased serum homocitrulline concentrations are associated with the severity of coronary artery disease", Clinical Chemistry and Laboratory Medicine (CCLM), 2015

Publication

<1 %

13

www.spandidos-publications.com

Internet Source

<1 %

14

Ghattas, Maivel H., Eman T. Mehanna, Noha M. Mesbah, and Dina M. Abo-Elmatty. "Relaxin-3 is associated with metabolic syndrome and its component traits in women", Clinical Biochemistry, 2013.

Publication

<1 %

15

T Takezako. "Angiotensin I converting enzyme gene polymorphism and insulin resistance in patients with angina pectoris", American Journal of Hypertension, 03/1999

Publication

<1 %

16

www.thieme-connect.com

Internet Source

<1 %

17 Mengyao Ji, Tengyu Zhu, Meichen Xing, Ning Luan et al. "An Antiviral Peptide from Alopecosa nagpag Spider Targets NS2B-NS3 Protease of Flaviviruses", Toxins, 2019
Publication <1 %

18 mafiadoc.com
Internet Source <1 %

19 Iwona Gołaś, Jacek Potorski, Małgorzata Woźniak, Piotr Niewiadomski et al. "Amaranth Meal and Environmental Carnobacterium maltaromaticum Probiotic Bacteria as Novel Stabilizers of the Microbiological Quality of Compound Fish Feeds for Aquaculture", Applied Sciences, 2020
Publication <1 %

20 Kuai Yu, Binyao Yang, Haijing Jiang, Jun Li et al. "A multi-stage association study of plasma cytokines identifies osteopontin as a biomarker for acute coronary syndrome risk and severity", Scientific Reports, 2019
Publication <1 %

21 Tveit, Arnljot, Ingebjørg Seljeflot, Pal Smith, Harald Arnesen, and Jesper Hastrup Svendsen. "Angiotensin II blockade, YKL-40 and maintenance of sinus rhythm after electrical cardioversion for atrial fibrillation", Immunobiology, 2013.
Publication <1 %

22	Yao Chen, Shu-ying Fang. "Potential genetic polymorphisms predicting polycystic ovary syndrome", Endocrine Connections, 2018 Publication	<1 %
23	hqlo.biomedcentral.com Internet Source	<1 %
24	www.biomedsearch.com Internet Source	<1 %
25	www.scribd.com Internet Source	<1 %
26	d-nb.info Internet Source	<1 %
27	bmcgastroenterol.biomedcentral.com Internet Source	<1 %
28	pesquisa.bvsalud.org Internet Source	<1 %
29	www.plosone.org Internet Source	<1 %
30	www.science.gov Internet Source	<1 %
31	Cameron Griffin, Leila Eter, Nico Lanzetta, Simin Abrishami et al. " TLR4, TRIF, and MyD88 are essential for myelopoiesis and CD11c adipose tissue macrophage production	<1 %

in obese mice ", Journal of Biological
Chemistry, 2018

Publication

32

Ming-Liang Zuo, Chun-Mei Li, Yan Deng, Sanjib
Bhattacharyya, Ping Shuai, Hung-Fat Tse,
Chung-Wah Siu, Li-Xue Yin. "The impact of
cigarette smoking in predicting stroke using
CHADS2 and CHA2DS2-VASc schemas",
Neurological Sciences, 2020

Publication

<1 %

33

my.unil.ch

Internet Source

<1 %

34

preview-jneuroengrehab.biomedcentral.com

Internet Source

<1 %

35

En-Zhi Jia. "Association of the mutation for the
human carboxypeptidase E gene exon 4 with
the severity of coronary artery
atherosclerosis", Molecular Biology Reports,
02/2009

Publication

<1 %

36

Laura E. Mantella, Kiera Liblik, Amer M. Johri.
"Vascular imaging of atherosclerosis:
Strengths and weaknesses", Atherosclerosis,
2021

Publication

<1 %

37

Miroslav Stojadinovic M., Tomislav Pejovic.
"Regression tree for choledocholithiasis

<1 %

prediction", European Journal of
Gastroenterology & Hepatology, 2015

Publication

38

Mohammad Rezaeimoghaddam, Gokce Nur
Oguz, Sanser Ates, Tijen Alkan Bozkaya et al.
"Patient-specific hemodynamics of new
coronary artery bypass configurations", Cold
Spring Harbor Laboratory, 2020

Publication

39

Suryapranata, H.. "Coronary vasodilatory
action after a single dose of nicorandil", The
American Journal of Cardiology, 19880201

Publication

40

Takao Hoshino, Kentaro Ishizuka, Satoru
Shimizu, Shinichiro Uchiyama. "CHADS₂,
CHA₂DS₂-VASc, and R₂CHADS₂ Scores Are
Associated With 3-Month Functional Outcome
of Stroke in Patients With Prior Coronary
Artery Disease", Circulation Journal, 2014

Publication

41

bjcvs.org
Internet Source

Internet Source

42

bmccardiovascdisord.biomedcentral.com
Internet Source

Internet Source

43

www.cureus.com
Internet Source

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

44

www.pulsus.com

Internet Source

<1 %

45

www.remedypublications.com

Internet Source

<1 %

46

K. Hanai, T. Babazono, N. Yoshida, I. Nyumura, K. Toya, T. Hayashi, R. Bouchi, N. Tanaka, A. Ishii, Y. Iwamoto. "Gender differences in the association between HDL cholesterol and the progression of diabetic kidney disease in type 2 diabetic patients", *Nephrology Dialysis Transplantation*, 2011

Publication

<1 %

47

Minyahil Woldu, Omary Minzi, Workineh Shibeshi, Aster Shewaamare, Ephrem Engidawork. "Estimation of the 10-Year Atherosclerotic Cardiovascular Disease and The Framingham Risk Score Among Adults Living with HIV/AIDS in Addis Ababa, Ethiopia: A Hospital-Based, Observational Study.", *Research Square Platform LLC*, 2021

Publication

<1 %

48

Yap-Hang Chan, Kai-Hang Yiu, Kui-Kai Lau, Yuen-Fung Yiu, Sheung-Wai Li, Tai-Hing Lam, Chu-Pak Lau, Chung-Wah Siu, Hung-Fat Tse. "The CHADS2 and CHA2DS2-VASc scores predict adverse vascular function, ischemic stroke and cardiovascular death in high-risk

<1 %

patients without atrial fibrillation: Role of incorporating PR prolongation",
Atherosclerosis, 2014

Publication

49

peerj.com
Internet Source

<1 %

50

repository-tnmgrmu.ac.in
Internet Source

<1 %

51

www.id-press.eu
Internet Source

<1 %

52

www.nature.com
Internet Source

<1 %

53

www.termedia.pl
Internet Source

<1 %

54

Abhay Kumar Chaudhary, Vijay Pathak, Shekhar Kunal, Shubhra Shukla, Pooja Pathak. "CHA2DS2-VASc score as a novel predictor for contrast-induced nephropathy after percutaneous coronary intervention in acute coronary syndrome", Indian Heart Journal, 2019
Publication

<1 %

55

Fotios Barkas, Moses Elisaf, Panagiotis Korantzopoulos, Stavroula Tsiara, Evangelos Liberopoulos. "The CHADS 2 and CHA 2 DS 2 - VASc scores predict atrial fibrillation in

<1 %

dyslipidemic individuals: Role of incorporating low high-density lipoprotein cholesterol levels", International Journal of Cardiology, 2017

Publication

56

Muhsin Kalyoncuoglu, Semi Ozturk, Mazlum Sahin. "Does CHA2DS2-VASc Score Predict MACE in Patients Undergoing Isolated Coronary Artery Bypass Grafting Surgery?", Brazilian Journal of Cardiovascular Surgery, 2019

Publication

<1 %

57

En-Zhi Jia. "Helicobacter pylori infection is associated with decreased serum levels of high density lipoprotein, but not with the severity of coronary atherosclerosis", Lipids in Health and Disease, 2009

Publication

<1 %

58

Gokhan Cicek, Ersin Yildirim. "CHA2DS2-VASc score predicts contrast-induced nephropathy in patients with ST-segment elevation myocardial infarction, who have undergone primary percutaneous coronary intervention", Kardiologia Polska, 2017

Publication

<1 %

59

Rugheed Ghadban, Greg Flaker, Natraj Katta, Martin A. Alpert. "Anti-thrombotic therapy for atrial fibrillation in patients with chronic

<1 %

kidney disease: Current views", Hemodialysis International, 2017

Publication

60

Yue Zhang, Xiaosong Ding, Bing Hua, Qingbo Liu, Hui Gao, Hui Chen, Xue-Qiao Zhao, Weiping Li, Hongwei Li. "Predictive Effect of Triglyceride-Glucose Index on Clinical Events In Patients With Type 2 Diabetes Mellitus and Acute Myocardial Infarction: Results From an Observational Cohort Study In China", Research Square, 2020

<1 %

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Novel CHA2DS2-VASc-HSF is Superior to CHADS and CHA2DS - VASc Score to Predict the Risk of Severe Coronary Artery Disease

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
