



[Home](#) / [Editorial Team](#)

## Editor-In-Chief's

Dr. Yousef Saleh Khader Al-Gaud - Jordan  
(BDS, MSc, MSPH, MHPE, FFFPH, ScD)  
Professor (Full) - Department of Community  
Medicine, Jordan University of Science and  
Technology, Jordan

Dr. Swapnil S. Bumb - India  
(BDS, MDS, MPH, MSc, PGDHA, PDCR)  
Assistant Professor, Department of Public Health  
Dentistry, ACPM Dental College, Dhule, Maharashtra,  
India

ResearchGate Link:

[https://www.researchgate.net/profile/Swapnil\\_Bumb4](https://www.researchgate.net/profile/Swapnil_Bumb4)

ResearchGate Link:

[https://www.researchgate.net/profile/Yousef\\_Khader](https://www.researchgate.net/profile/Yousef_Khader)

PubMed Link: <http://www.ncbi.nlm.nih.gov/pubmed/?term=Swapnil+Bumb>

PubMed Link:

<http://www.ncbi.nlm.nih.gov/pubmed/?term=Yousef+Saleh+Khader>

**1. Editorial board:** Journal of Clinical Studies and  
Medical Case Reports

**2. Academic Editor:** Case Reports in Dentistry

**1. Director:** International Research Organization for  
Life & Health Sciences (IROLHS)

**1. Founder & Editor In Chief:** International Journal of  
Scientific Study

**2. Founder & Editor in Chief:** IJSS Case Reports &  
Reviews

**3. Founder & Managing Editor:** IJSS Journal of  
Surgery

**3. Technical Editor:** World Journal of Diabetes

**4. Technical Editor:** International Journal of Osteoporosis and Metabolic Disorders

.Recipient of several research Grants from international and national universities/organizations

### **Associate Editors Specialty - Anthropology**

**Dr. Kaushik Sankar Bose - India (PhD, MSc, BSc - University of Cambridge)**

Professor, Biological Anthropology, Department of Anthropology, Vidyasagar University, Midnapore

- Editorial Board Member of Several reputed International and National journals indexed in PubMed & Medline.
- Peer Reviewer in several reputed International Journals. To mention few are Biomedical Central (BMC) Public Health, Biomedical Central (BMC) Nutrition, Homo, The Lancet, Journal of Public Health, Journal of Public Health and Epidemiology, and many more.
- Recipient of several research Grants from international and national universities/organizations.
- Authors of Several Textbooks and 200+ articles published in several national and international journals.

### **Specialty - Public Health**

**Dr. Yong Zhao - China (B.Sc.Med, M.Sc.Med)**

Associate professor, School of Public Health and Management, Chongqing Medical University. China.  
# ResearchGate Link: [https://www.researchgate.net/profile/Yong\\_Zhao12](https://www.researchgate.net/profile/Yong_Zhao12)

Recipient of 50+ research Grants as Principal and Co Investigator from international and national universities/organizations.

- Authored 50+ international and national publications in reputed journals.
- Authored 20 Books and Book Chapters for international and national publishers.
- Recipient of 20+ Awards for academics related activities.
- Editorial Board Members and Reviewers for several high impact factor journals.

### **Senior Editorial Board - Public Health (Medical Specialty)**

**Dr. Rita Caroline Isaac ( MD, MPH )**

Professor & Head, RUHSA Department, Christian Medical College & Adjunct Associate Professor, Public Health and

Community Medicine, Tufts University, Vellore

#ResearchGate Link: [https://www.researchgate.net/profile/Rita\\_Isaac](https://www.researchgate.net/profile/Rita_Isaac)

**Dr. Sadhu Charan Panda ( M.D )**

Associate Professor, Department of Community Medicine, V. S. S. Medical College, Burla

**Senior Editorial Board - Public Health (Dental Specialty)****Dr. Shivalingaswamy Hiremath (M.D.S)**

Professor & Head, Department of Public Health Dentistry, Oxford Dental College, Bangalore.

**Dr. D J Bhaskar (M.D.S, M.P.H)**

Professor & Head, Department of Public Health Dentistry, Teerthanker Mahaveer Dental College & Research Centre, Moradabad

**Dr. Manjunath P. Puranik (M.D.S)**

Professor & Head, Department of Public Health Dentistry, Government Dental College & Research Centre, Bangalore.

**Dr. Sahana Hegde (M.D.S)**

Professor & Head, Department of Public Health Dentistry, D Y Patil Dental College, Pune

**Dr. Arun Doddamani (M.D.S)**

Principal & Professor & Head, Department of Public Health Dentistry, ACPM Dental College, Dhule

**Dr. Pradeep Tangde (M.D.S)**

Professor & Head, Department of Public Health Dentistry, Kothiwal Dental College, Moradabad

**Dr. Ajith Krishnan C G (M.D.S)**

Principal & Professor, Department of Public Health Dentistry, K M Shah Dental College & Hospital, Vadodara

**Dr. Ramesh N (M.D.S)**

Professor & Head, Department of Public Health Dentistry, Institute of Dental Sciences, Bhubaneswar.

**Dr. Aswini Y B (M.D.S)**

Assistant Professor, Department of Public Health Dentistry, Maulana Azad Institute of Dental Sciences, New Delhi

**Dr. Chandan R Agali (M.D.S)**

Reader, Department of Public Health Dentistry, Teerthanker Mahaveer Dental College & Research Centre,  
Moradabad

**Dr. Nilesh Arjun Torwane (M.D.S)**

Assistant Professor, Department of Public Health Dentistry People's Dental Academy, Bhopal

**Assistant Editors**

**Dr. Shilpkumar Arora (MPH, M.D)**

Icahn School of Medicine at Mount Sinai, New York, St Luke's – Roosevelt Hospital Center Program

**Dr. Khan Amir Maroof (M.D)**

Assistant Professor cum Medical Officer of Urban Health Centre, Department of Community Medicine,  
University College of Medical Sciences, Delhi

**Dr. Safalya Kadtane (M.D.S)**

Assistant Professor, Department of Public Health Dentistry, Maharashtra Institute of Dental Science &  
Research Dental College, Latur, Maharashtra

**Dr. Subha Soumya Dany (M.D.S)**

Public Health Dentistry.

[Make a Submission](#)



**grammarly**



**turnitin®**

## Information



[For Readers](#)

[For Authors](#)

[For Librarians](#)

## Browse

## Current Issue



Open  
Access



INTERNATIONAL COMMITTEE of  
MEDICAL JOURNAL EDITORS



**Publisher:** [International Research Organization for Life & Health Sciences \(IROLHS\)](#), L 214, Mega Center, Solapur Road, Magarpatta, Pune, Maharashtra 411028

**International Journal of Preventive and Public Health Sciences (IJPPHS)** is licensed under a [Creative Commons Attribution 4.0 International License](#).

**Website** © 2021. Article copyright remains as specified within the article.

Platform &  
workflow by  
**OJS / PKP**



[Home](#) [About](#) [Editorial Team](#) [Indexing](#) [Current](#) [Archives](#) [For Authors](#)  
[For Reviewers](#) [Contact](#)

[Home](#) / [Archives](#) / Vol 2 No 4 (2016): VOLUME: 2 | ISSUE: 4 2016

**Published:** 2016-07-30

## Table of Contents

### **Differences in the Effect of Various Intervention Cream Sprouts (*Phaseolus radiatus*) Extract to Wound Healing Process on Postpartum Mother**

Wiulin Setiowati, Hadi, Supriyana, Bahiyatun

1-4



### **Clinical Spectrum and Biochemical Profile of Chronic Kidney Disease Patients Attending a Tertiary Hospital Setting in India**

Dr. Richa Mishra, Chandra Pati Mishra, Rana Gopal Singh, Shivendra Singh, Prem Nath Tiwari

5-11



### **Utilization of Lindur Fruit Flour and Tempeh Flour as a Healthy Snack Alternative for Diabetic**

Made Mita Dwi Saraswati, Annis Catur Adi, Ikeu Ekayanti

12-17



### **Knowledge, Attitude, and Practice of Budding Doctors in Prevention of Thalassemia**

Dr. Mausumi Basu, Sita Chatterjee, Tushar Kanti Monda, Afifa Ahamed, Ishita Sarkar, Kaushik Sarkar, Bhaskar Shahbabu

18-24



### **Sexual Education as Knowledge of Improvement Efforts on Adolescent Sexuality in X Junior High School North Raman District of East Lampung**

Ristu Wiyani, Hendy Muagiri Margono, Hari Basuki Notobroto

25-27



### **Effect Red Guava Juice (*Psidium guajava* L) on Blood Glucose Levels Fasting Wistar Rats Induced Pre-diabetes Dexamethasone**

Jasmani, Bambang Wirjatdmadi, Merryana Adriani

28-31



### **Galactagogue Instant Powder Combination of Papaya Leaves and Red Ginger for Breastfeeding Mother**

Filia Startia Setyono, Annis Catur Adi, Rita Ismawati

32-36



[Make a Submission](#)



# Various Formulations of Lepa as a Zinc-rich Food for Primary School Children

Oskar Karyantono<sup>1</sup>, Annis Catur Adi<sup>2</sup>, Merryana Adriani<sup>2</sup>

<sup>1</sup>Master Program Of Public Health Science, Faculty Of Public Health Airlangga University, Surabaya, <sup>2</sup>Department of Nutrition, Airlangga University, Surabaya, Indonesia

## ABSTRACT

**Introduction:** Nutrient that is responsible in the incidence of stunting is zinc. Lower intake of zinc can be attributed to the amount and quality of major food as well as low nutritional value in supplementary food that includes zinc. To improve the quality of food, it is needed to create the new formulations by adding certain nutritional content to enrich the food. Lepa is a traditional cake belongs to the people of rote and is preferred by children. The basic ingredients consisted of palm sugar water, cornstarch, and grated coconut. Lepa considered as poor in nutritional value. Formulations by adding Moringa leaf powder and anchovy flour, sesame and peanuts enrich the content of zinc in the cake.

**Purpose:** The purpose of this study was to analyze the level of preference and zinc content in Lepa.

**Methods:** This study applied experimental study design using completely randomized design with six repetitions. Ingredients used were palm sugar water, grated coconut, cornstarch, flour of Moringa leaves, anchovy flour, sesame, and peanuts. There were four formulations made: Control formula ( $F_0$ ) = 42%: 37%: 21%.  $F_1$  = 42%: 13%: 9%: 4%: 13%: 12%: 7%.  $F_2$  = 42%: 10%: 10%: 6%: 13%: 12%: 7%.  $F_3$  = 42%: 8%: 9%: 9%: 13%: 12%: 7%. Data concerning to organoleptic was processed using Kruskal–Wallis test to determine the differences among formulas and was described descriptively.

**Results:** An organoleptic test that consisted of limited panelist demonstrating the preference level of formulation for Lepa was between “somewhat liked” and “highly liked” (3.63-4.18). There was no difference in the level of preference between  $F_0$ ,  $F_1$ ,  $F_2$ , and  $F_3$ . The content of zinc indicated per 60 g were  $F_1$  = 2.81 g,  $F_2$  = 4.25 g, and  $F_3$  = 3.65 g.

**Conclusion:** Lepa can be used as supplementary food which is rich in zinc.

**Key words:** Lepa, Rote, Supplementary food

## INTRODUCTION

One of the unfinished nutritional problems in school children is stunted.<sup>1</sup> The prevalence of stunting in primary school children (6-12 years) in 2010 was 35% and was classified as “high.”<sup>2</sup> One of the direct causes of stunting is due to a lack of energy and protein in long term.<sup>3</sup> Besides the lack of energy and protein, another nutrient that is responsible in the incidence of stunting is zinc.<sup>4</sup> The same report that the cause of malnutrition in Africa was due to micronutrient deficiencies, zinc included.<sup>5</sup> In Thailand reported, the incidence of stunting is associated with a lack of energy intake, protein and several micronutrients such as calcium, phosphorus, iron, Vitamin A, and zinc. Macro and micronutrients deficiencies cause a decrease in linear growth.<sup>6</sup> In

Lombok Island, Indonesia found that child stunting experienced deficit intake of zinc (<70%).<sup>7</sup> The cause of the lack of nutrients in food is due to inadequate quantity and quality of the major food and supplementary food as well as less varied and low nutritional value foods consumed.<sup>8</sup> To meet the nutritional needs of school children, the government implemented supplementary food program (Indonesian: Pemberian Makanan tambahan). Supplementary food must be nutritious, balanced and varied, and preferably taken from local food or local ingredients.<sup>9</sup> The provision of supplementary food to primary school children should meet nutritional requirements which are minimum of 10-15% of recommended dietary allowance (RDA).<sup>10</sup>

Lepa is a traditional cake belongs to the people of Rote and is widely preferred by children and found in many. Lepa

### CORRESPONDING AUTHOR:

S G Z Oskar Karyantono,

Jl. Mojoklanggru II No. 20 Kel. Mojo Kec. Gubeng Surabaya, East Java Indonesia. Phone: +081241929317.

E-mail: karyantono@yahoo.com

Submission: 05-2016; Peer Review: 06-2016; Acceptance: 07-2016; Publication: 08-2016

cake contains low of zinc since made only of palm sugar water (*Borassus flabellifer*), coconut (*Cocos nucifera*), and cornstarch (*Zea mays*) which are poor sources of zinc. The content of zinc in the cake is only reached 2.14% of the nutritional value required for being a supplementary food and fulfills only 4.8-6.11% of nutritional adequacy for primary school children.<sup>11</sup> Formulation with Moringa leaf powder (*Moringa oleifera*), anchovy flour (*Stolephorus* sp.), sesame (*Sesamum indicum*), and peanut (*Arachis hypogaea*) increases the content of zinc on the cake. The purpose of this study was to determine the level of preference and zinc content of Lepa that has been formulated.

## METHODS

### Study Design

This study applied experimental design of study using completely randomized design with six repetitions. This study carried out formulations for Lepa with the addition of Moringa leaf powder, anchovy flour, sesame, and peanuts.

### Location and Time

All process that includes making Moringa leaf powder and anchovy flour, making Lepa, and analyzing the zinc levels were conducted at Nutrition Laboratory in Public Health Faculty Airlangga University, Surabaya. Organoleptic tests with limited panelist were conducted to four lecturers from the Department of Health Nutrition in Public Health Faculty Airlangga University, Surabaya, who are the expertise in organoleptic, food and nutrition. The study was conducted from February to June 2016.

### Analysis of Nutritional Content

Analysis of zinc content was done using zinc method of analysis, SNI 6989.7.2009.

### Tools

Tools used for making Moringa leaf powder and anchovy flour were: 100 mesh sieve, oven, blender, griddle and digital scales. Tools used for making Lepa were: Frying pan, spatula, cake molder, and gas stoves.

### Ingredients

Ingredients selected for making Lepa contains a high level of zinc, excluding the basic ingredients; cornstarch, coconut, and palm sugar water.

1. Flour leaves of Moringa (*M. oleifera*): 1.3 mg of zinc
2. Anchovy flour (*Stolephorus* sp.): 5.8 mg of zinc
3. Cornstarch (*Z. mays*): 1.8 mg of zinc
4. Grated coconut (*C. nucifera*) providing aroma and savory taste
5. Sesame (*S. indicum*): 7.8 mg of zinc
6. Peanut (*A. hypogaea*) 3.3 mg of zinc
7. Palm sugar water (*B. flabellifer*) as binder.

### Preparation

Moringa leaf powder: Sort Moringa leaves, wash in flowing water, blanch at a temperature of 90-100°C for 10-15 min, remove and drain for several hours, then dry using the oven with a temperature of 60°C for 6-8 h to make sure the powder was properly dried. Finally, sift the powder using a 100 mesh sieve. To make anchovy flour: Soak the anchovy for about 3 h, wash and boil in 80-90°C temperature for about 30-60 min, drain and dry at 100°C for 6 h till properly dried then sift using a 100 mesh sieve. To make cornstarch: Sort the corn seeds,

roast until golden-or brown-colored, mill into powder and sift with a 100 mesh sieve. Roast grated coconut, sesame, and peanuts separately until browned-colored and fragrant. Finally, blend the peanut.

### Method for Processing Lepa

Put in toasted coconut, cornstarch, Moringa leaves powder, anchovy flour and sesame in one bowl and mixed thoroughly. Cook the palm sugar water until it thickens (1.5-2.5 min) with a temperature of 95°C-105°C and can be higher when cooking Lepa cake for up to 120°C. Reduce the heat, put mixed ingredients, stir well. Stir constantly, until cooked and thickened (and mixed) for 3-6 min. Mold using cake molder and pack when ready.

### Processing and Data Analysis

Organoleptic data were analyzed using Kruskal-Wallis test to determine the differences in treatment between groups, while if differences existed then being tested using the Mann-Whitney U-test. The zinc content was processed using Microsoft Excel 2007. Data were being presented through tables and were being analyzed descriptively.

## RESULTS

### Formulations for Lepa

The formulation was done by selecting zinc-rich ingredients, the results of the formulation presented in Table 1.

The addition of Moringa leaf powder in  $F_1$ ,  $F_2$ , and  $F_3$  was 4%, 6%, and 9%, respectively. While the addition of anchovy flour, sesame and peanut were 13%, 12% and 7%, respectively, in each formulation.

### Hedonic Test Results

#### Color

The results of the organoleptic test with limited panelists regarding the color of Lepa were presented in Figure 1.

Figure 1 presented that the highest ratings were  $F_3$  with an average value of 4.25 (liked), while the lowest ones were  $F_1$  with an average value of 3.5 (somewhat liked).

#### Aroma

The results of the organoleptic test with limited panelists regarding to the aroma of Lepa were presented in Figure 2.

Figure 2 presented that most panelists preferred  $F_1$  and  $F_3$  with average values of 4.25 (liked), the lowest ratings was  $F_2$  with an average value of 3.75 (somewhat liked).

#### Texture

The results of the organoleptic test with limited panelists regarding the texture Lepa were presented in Figure 3.

**Table 1:** Formulations for making Lepa

Ingredients	$F_0$ %	$F_1$ %	$F_2$ %	$F_3$ %
Palm sugar water	42	42	42	42
Coconut	37	13	10	8
Cornstarch	21	9	10	9
Moringa leaf powder	0	4	6	9
Anchovy powder	0	13	13	13
Sesame	0	12	12	12
Peanut	0	7	7	7
Sum	100	100	100	100

Figure 3 presented that the panelists provided the highest ratings to F<sub>3</sub> with an average value of 4.25 (liked), and the lowest ones were F<sub>1</sub> with an average value of 3 (somewhat liked).

**Taste**

The results of the organoleptic test with limited panelists regarding to the taste of Lepa were presented in Figure 4.

Figure 4 presented that the panelists provided the highest rating to F<sub>2</sub> with an average value of 4.5 (liked), while the lowest ratings were F<sub>1</sub> with an average value of 3.75 (somewhat liked).

**Acceptability**

The average ratings for acceptability of Lepa were presented in Table 2.

Table 2 presented that the panelists provided the highest ratings to F<sub>5</sub> with an average value of 4.1 (liked) and the lowest ones were F<sub>2</sub> with an average value of 3.6 (somewhat liked).

Figure 5 presented that in terms of color, aroma, texture and taste, panelists preferred F<sub>3</sub> with ingredients of palm sugar water 42%, grated coconut 10%, cornstarch 10%, flour Moringa leaves 6%, anchovy flour 13%, sesame 12% and 7% peanuts.

**The zinc content**

The zinc content of Lepa was presented in Table 3.

Table 3 presented that the highest content of zinc was in F<sub>2</sub>; as much as 4.25 mg, followed by F<sub>3</sub> as the second; as much as 3.65 mg followed on the third by F<sub>1</sub> as much as 2.81 mg while F<sub>0</sub> was the lowest with only 1.1 mg.

**DISCUSSION**

The formulation of Lepa was intended to increase the content of zinc by adding zinc-rich food ingredients. Moringa leaf powder is a nutrient-rich food source. Moringa leaf powder has a high content of zinc (1.3 mg).<sup>12</sup> Besides, according to some studies Moringa leaves are often being used to address the incidence of malnutrition because of its richness in nutrients and active compounds. Moringa leaves have been used in several African countries to tackle malnutrition in children, pregnant and lactating mothers.<sup>5</sup> By adding 7 g of Moringa leaf powder into toddler foods every day for 2 months, can increase the child's height. Further explanation, that the average height of children increased by 3.79 cm than the previous ones.<sup>13</sup> Anchovy also has a high content of zinc (5.8 mg).<sup>11</sup> Children who are provided with anchovy based-biscuit had average height growth higher than the control group.<sup>14</sup> Likewise, sesame and peanuts have fairly high content of zinc with 7.8 mg and 3.3 mg, respectively.<sup>11</sup>




Figure 1: Preference level of Lepa regarding to the color




Figure 3: Preference levels of Lepa regarding to the texture




Figure 2: Preference level of Lepa regarding to the aroma




Figure 4: Preference level of Lepa regarding to the taste of Lepa




Figure 5: Preference level of the panelists to Lepa

Table 2: Average level of acceptability of Lepa

Formulation	Organoleptic scoring of Lepa				Total	Average
	Color	Aroma	Texture	Taste		
F <sub>0</sub>	5.00	4.75	3.50	4.00	17.25	4.30
F <sub>1</sub>	3.50	4.25	3.00	3.75	14.50	3.60
F <sub>2</sub>	4.00	3.75	2.25	4.50	14.50	3.60
F <sub>3</sub>	4.25	4.25	4.25	4.00	16.75	4.10

Table 3: The zinc content of Lepa per 60 g

Formulation	Zinc (mg)
F <sub>0</sub>	1.12
F <sub>1</sub>	2.81
F <sub>2</sub>	4.25
F <sub>3</sub>	3.65

## Characteristics of Lepa

### Color

Formula used in F<sub>1</sub>, F<sub>2</sub>, and F<sub>3</sub> produce darker green appearance. F<sub>3</sub> used 9% Moringa powder and had a darker green color than F<sub>2</sub> (6%). F<sub>1</sub> formula used 4% Moringa powder and had brighter colors than both F<sub>2</sub> and F<sub>3</sub>. The more Moringa leaf powder used, the darker the color of the cake. The green color in Moringa leaves comes from the chlorophyll. Blanching process causes the color changes to dark green or yellowish-green due to chlorophyll content in leaves degraded by heat.<sup>15</sup> In the control formula (F<sub>0</sub>), the cake appeared brownish to yellowish because of no Moringa leaf powder used. The brown color comes from heating palm sugar water which produces sticky and brown-colored caramel hence the effect on the cake's appearance.<sup>16</sup>

### Aroma

F<sub>1</sub>, F<sub>2</sub>, and F<sub>3</sub> were dominated by the aroma of sesame which is produced from pyrazine compounds contained in sesame while roasting triggers evaporation of pyrazine compounds.<sup>17</sup> F<sub>0</sub> was dominated by the aroma of toasted coconut, derived from bounded fatty acids in the form of ester which triggers distinctive coconut aroma.<sup>18</sup>

### Texture

In F<sub>1</sub>, F<sub>2</sub> and F<sub>3</sub> produced more solid texture. The proportion of Moringa leaf powder as much as 4%, 6%, and 9% provides an effect on the texture of the cake. The more Moringa leaf powder is used; the more solid texture obtained.<sup>19</sup>

### Taste

The flavors derived from palm sugar water which contains sucrose. Sucrose provides the sweet taste.<sup>20</sup> In F<sub>1</sub>, F<sub>2</sub> and F<sub>3</sub>, the addition of Moringa leaf powder and anchovy flour does not provide fishy nor unpleasant taste/smell. Blanching process lowers the levels of phytate in Moringa leaf that causes unpleasant odor. While fishy smell was not noticeable because of being covered with sesame and tea-like flavor from Moringa leaves. For F<sub>0</sub>, the sweetness was mixed with savory flavors derived from coconuts.

### Acceptability

The results of the organoleptic test with limited panelists toward 4 formulations indicated that the color, aroma, texture and taste of the cake were acceptable. For the texture and flavor, F<sub>2</sub> and F<sub>3</sub> were being more preferred by the panelists compared with the control. This was confirmed by Kruskal tests indicating that there was no difference in the level of preference among the treatments in terms of color, aroma, texture and taste of Lepa. This also means that the formulation of Lepa using F<sub>1</sub>, F<sub>2</sub>, and F<sub>3</sub> formulations do not provide a significant influence on the level of preference of color, aroma, texture and taste of the cake.

### Nutritional value

Table 3 shows that the F<sub>1</sub>, F<sub>2</sub>, and F<sub>3</sub> formulations have a higher content of zinc and fulfill nutritional value required as supplementary food. The need for zinc for elementary school children aged 7-9 years can be fulfilled at 15.33-23.18%. Male children aged 10-12 years can be fulfilled with 12.04-18.21%, and female children aged 9-10 years can be fulfilled with 12.97-19.62%. Therefore, F<sub>1</sub>, F<sub>2</sub>, and F<sub>3</sub> as Lepa formulations can be used as zinc-rich supplementary food for elementary children.

The role of zinc in the growth process is that it is involved in the bioactivity of insulin-like growth factor 1 (IGF-1) on bone cells. Some studies have found that zinc plays a role in the work of IGF-1 and promotes the synthesis of endogenous IGF-1.<sup>21</sup> Growth occurs through cell division and requires the synthesis of DNA, RNA, and protein. Growth is regulated by hormones especially by growth hormone and IGF-1.<sup>22</sup>

## CONCLUSION

Organoleptic of Lepa was being accepted by the panelists. Statistical analysis showed that Lepa formulations did not influence the preference level of the panelists.

The zinc content in all formula meets nutritional value required for being supplementary foods and fulfills the needs of the zinc as much as 12.04-23.18% for primary school children aged 7-12 years, both males and females.

Generally speaking, the formulations were being accepted by the panelists and were in compliance to nutritional value for being supplementary foods, and RDA therefore could be used as zinc-rich supplementary foods for elementary school children.

## REFERENCES

- Minister of Health of Indonesia, Gizi Kurang Penyebab Stunting; 2013. Available from: [http://www.dinkes.sumselprov.go.id/download/unggah/stunting\\_anak-2016-01-04.pdf](http://www.dinkes.sumselprov.go.id/download/unggah/stunting_anak-2016-01-04.pdf).
- Salimar S, Kartono D, Fuada NF, Setyawati B. Stunting among school-age children in Indonesia by characteristics of family. *J Nutr Food Res* 2013;36:121-6.
- Blössner M, De Onis M. Malnutrition: Quantifying the Health Impact at National and Local Levels. Environmental Burden of Disease Series, No. 12. Geneva: WHO, Nutrition for Health and Development, Protection of the Human Environment; 2005.
- Grider A. Zinc, copper and manganese. In: Stipanuk MH, editor. *Biochemical, Physiological & Molecular Aspects of Human Nutrition*. 2<sup>nd</sup> ed. St. Louis, MO: Elsevier Inc.; 2006.
- Fuglie LJ. The Moringa Tree: A Local Solution to Malnutrition. Senegal: Church World Service; 2005.
- Gibson RS, Manger MS, Krittaphol W, Pongcharoen T, Gowachirapant S, Bailey KB, et al. Does zinc deficiency play a role in stunting among primary school children in NE Thailand? *Br J Nutr* 2007;97:167-75.
- Taufiqurrahman T, Hadi H, Julia M, Herman S. Vitamin A and zinc deficiency as risk factor in children in event of stunting at West Nusa Tenggara. *Media Penelitian dan Pengembangan Kesehatan* 2009;10:84-94.
- WHO, World Health Organisation. *Childhood Stunting: Context, Causes and Consequences WHO Conceptual Framework*; 2013. Available from: <http://www.who.int>. [Last accessed on 2016 Jun 08].
- Ministry of Home Affairs of Indonesia, Peraturan Menteri Dalam Negeri Nomor 18 Tahun 2011. Tentang Pedoman Penyediaan Makanan Tambahan Anak Sekolah. Available from: <http://www.binapemdes.kemendagri.go.id>. [Last accessed on 2016 Jun 08].
- Minister of Health of Indonesia. Regulation of Minister of Health of Indonesia No. 899/Menkes/SK/X/2009 on Technical Specification Requirements of Quality Nutritional Supplementary Food 2-5 Years Childhood, School Age Children Primary and Pregnancy. 2009.
- Earhard, J. *Nutrisurvey for Windows*. Jakarta: SEAMEO-TROPMED. University of Indonesia; 2007.
- Witt KA. The Nutrient Content of *Moringa oleifera* Leaves. Messiah College Department of Nutrition and Dietetics. ECHO Research Note No. 1. 2013. p. 1-6. Available from: <http://www.miracletrees.org/moringa-doc/nutrient-content-of-moringa-oleifera-leaves.pdf>. [Last accessed on 2016 Jun 08].
- Juhartini. The Effect of Biscuits of Government Programs and Slurry Mixture Moringa Oleifera to Increased Weight, Height and Hemoglobin in Under Nutrition Children Under Five. [Tesis]. Universitas Airlangga Surabaya; 2015.
- Affandy A. dkk. Effect of biscuits anchovy on growth toward under two years children under nutrition in the district Tanete Rilau. *J MKMI* 2008;4: 175-82.
- Madalena M, Heriyanto H, Hastuti SP, Limantara L. The effect of heating time to the content of pigments and Vitamin A in Cassava (*Manihot Esculenta* Crantz) and Ceara-Rubber (*Manihot glaziovii* Muell. Arg) leaves. *Indonesian J Chem* 2007;7:105-10.
- McGee H. *On Food and Cooking: The Science and Lore of The Kitchen*. New York: Scribner; 2004.
- Shah NC. *Sesamum indicum* (Sesame or Til): Seeds and Oil-A Historical and Scientific Evaluation from Indian Perspective. *Asian Agri History* 2016;20:151-74.
- Saittagaroon S, Kawakishi S, Namiki M. Aroma constituents of roasted coconut. *Agric Biol Chem* 1984;48:2301-7.
- Dachana KB, Rajiv J, Indrani D, Prakash J. Effect of dried Moringa (*Moringa Oleifera* Lam) leaves on rheological, microstructural, nutritional, textural and organoleptic characteristics of cookies. *J Food Qual* 2010;33:660-77.
- Winarno FG. *Kimia Pangan dan Gizi*. Jakarta: Gramedia; 1992.
- Yamaguchi M, Hashizume M. Effect of beta-alanyl-L-histidinato zinc on protein components in osteoblastic MC3T3-E1 cells: Increase in osteocalcin, insulin-like growth factor-I and transforming growth factor-beta. *Mol Cell Biochem* 1994;136:163-9.
- MacDonald RS. The role of zinc in growth and cell proliferation. *J Nutr* 2000;130 5S Suppl:1500S-8.

### HOW TO CITE THIS ARTICLE:

Karyantono O, Adi AC, Adriani M. Various Formulations of Lepa as a Zinc-rich Food for Primary School Children. *Int J Prevent Public Health Sci* 2016;2(2):18-22.