

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies](#) policy.

[Close this notification](#)

[IOP Science home](#)

- [Skip to content](#)
- [Accessibility Help](#)

Search

[Journals](#)

[Books](#)

[Publishing Support](#)

[Login](#)

[IOP Conference Series: Earth and Environmental Science](#)

Table of contents

Volume 441

2020

[Previous issue](#)[Next issue](#)

2nd International Conference on Fisheries and Marine Science 26 September 2019, Surabaya, Indonesia

Accepted papers received: 16 January 2020
Published online: 24 February 2020

[Open all abstracts, in this issue](#)

Preface

011001

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Preface](#)

[Open abstract](#), [Preface](#) [View article](#), [Preface PDF](#), [Preface](#)

011002

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Peer review statement](#)

[Open abstract](#), [Peer review statement](#) [View article](#), [Peer review statement PDF](#), [Peer review statement](#)

Papers

Earth and Environmental Science

012001

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The grow-out of abalone \(*Haliotis squamata*\) at different shelter shape on growth and survival and its marine environmental influences at Lembongan Bay coastal waters](#)

I Ardi, E Setiadi, Rasidi and W S Pranowo

[Open abstract](#), The grow-out of abalone (*Haliotis squamata*) at different shelter shape on growth and survival and its marine environmental influences at Lembongan Bay coastal waters [View article](#), The grow-out of abalone (*Haliotis squamata*) at different shelter shape on growth and survival and its marine environmental influences at Lembongan Bay coastal waters [PDF](#), The grow-out of abalone (*Haliotis squamata*) at different shelter shape on growth and survival and its marine environmental influences at Lembongan Bay coastal waters

012002

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Stocking density of Rotifera, *Hexarthra mira* on water quality and production](#)

I Ardi, S Cahyaningsih and E Setiadi

[Open abstract](#), Stocking density of Rotifera, *Hexarthra mira* on water quality and production [View article](#), Stocking density of Rotifera, *Hexarthra mira* on water quality and production [PDF](#), Stocking density of Rotifera, *Hexarthra mira* on water quality and production

012003

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The effect addition of kappa carrageenan flour to the level of gel strength and acceptability of dumpling from threadfin bream fish \(*Nemipterus nematophorus*\) surimi](#)

D M Astutik, L Sulmartiwi, E Saputra and D Y Pujiastuti

[Open abstract](#), The effect addition of kappa carrageenan flour to the level of gel strength and acceptability of dumpling from threadfin bream fish (*Nemipterus nematophorus*) surimi [View article](#), The effect addition of kappa carrageenan flour to the level of gel strength and acceptability of dumpling from threadfin bream fish (*Nemipterus nematophorus*) surimi [PDF](#), The effect addition of kappa carrageenan flour to the level of gel strength and acceptability of dumpling from threadfin bream fish (*Nemipterus nematophorus*) surimi

012004

THE FOLLOWING ARTICLE ISOPEN ACCESS

Maximizing production of a male offspring in *Moina macrocopa* culture through manipulation of rice bran suspension concentration

A S Mubarak, D Jusadi, M Zairin Jr and M A Suprayudi

Open abstract, Maximizing production of a male offspring in *Moina macrocopa* culture through manipulation of rice bran suspension concentration View article, Maximizing production of a male offspring in *Moina macrocopa* culture through manipulation of rice bran suspension concentration PDF, Maximizing production of a male offspring in *Moina macrocopa* culture through manipulation of rice bran suspension concentration

012005

THE FOLLOWING ARTICLE ISOPEN ACCESS

The Concept of the Right to Management of Coastal Communities in the Regional Autonomy Era: Experience from Community Assistance to Obtain the Right to Manage Sea Cucumbers in Sunsak Bay, East Lombok.

A Wahyono and M Illiyani

Open abstract, The Concept of the Right to Management of Coastal Communities in the Regional Autonomy Era: Experience from Community Assistance to Obtain the Right to Manage Sea Cucumbers in Sunsak Bay, East Lombok. View article, The Concept of the Right to Management of Coastal Communities in the Regional Autonomy Era: Experience from Community Assistance to Obtain the Right to Manage Sea Cucumbers in Sunsak Bay, East Lombok. PDF, The Concept of the Right to Management of Coastal Communities in the Regional Autonomy Era: Experience from Community Assistance to Obtain the Right to Manage Sea Cucumbers in Sunsak Bay, East Lombok.

012006

THE FOLLOWING ARTICLE ISOPEN ACCESS

Selective breeding technique: Pandu and Kunti tilapia (*Oreochromis niloticus*) broodstock candidates at PBIAT Janti, Klaten-Central Java

S H Samara, A W Fathurrozi and Sutarno

Open abstract, Selective breeding technique: Pandu and Kunti tilapia (*Oreochromis niloticus*) broodstock candidates at PBIAT Janti, Klaten-Central Java View article, Selective breeding technique: Pandu and Kunti tilapia (*Oreochromis niloticus*) broodstock candidates at PBIAT Janti, Klaten-Central Java PDF, Selective breeding technique: Pandu and Kunti tilapia (*Oreochromis niloticus*) broodstock candidates at PBIAT Janti, Klaten-Central Java

012007

THE FOLLOWING ARTICLE ISOPEN ACCESS

Potential of *Caulerpa racemosa* extracts as sunscreen creams

E B Ersalina, A A Abdillah and L. Sulmartiwi

THE FOLLOWING ARTICLE IS OPEN ACCESS

Immature Platelet Level in Stable Coronary Heart Disease (CHD) Patients with Diabetes Mellitus compared to Stable CHD Patients without Diabetes Mellitus

D R Balti, **A Andrianto**, B B Dharmadjati and D N Asmarani

[Open abstract](#), Immature Platelet Level in Stable Coronary Heart Disease (CHD) Patients with Diabetes Mellitus compared to Stable CHD Patients without Diabetes Mellitus [View article](#), Immature Platelet Level in Stable Coronary Heart Disease (CHD) Patients with Diabetes Mellitus compared to Stable CHD Patients without Diabetes Mellitus [PDF](#), Immature Platelet Level in Stable Coronary Heart Disease (CHD) Patients with Diabetes Mellitus compared to Stable CHD Patients without Diabetes Mellitus

012166

THE FOLLOWING ARTICLE IS OPEN ACCESS

Correlation Between Arterial and Left Ventricle Diastolic Stiffness in Hypertensive Patients

D Sumantha, E R Utami and B S Pikir

[Open abstract](#), Correlation Between Arterial and Left Ventricle Diastolic Stiffness in Hypertensive Patients [View article](#), Correlation Between Arterial and Left Ventricle Diastolic Stiffness in Hypertensive Patients [PDF](#), Correlation Between Arterial and Left Ventricle Diastolic Stiffness in Hypertensive Patients

012167

THE FOLLOWING ARTICLE IS OPEN ACCESS

Neglected Cases of Hypertension in Rural Indonesia: A Cross-Sectional Study of Prevalence and Risk Factors on Adult Population

E P B Mulia and S Prajitno

[Open abstract](#), Neglected Cases of Hypertension in Rural Indonesia: A Cross-Sectional Study of Prevalence and Risk Factors on Adult Population [View article](#), Neglected Cases of Hypertension in Rural Indonesia: A Cross-Sectional Study of Prevalence and Risk Factors on Adult Population [PDF](#), Neglected Cases of Hypertension in Rural Indonesia: A Cross-Sectional Study of Prevalence and Risk Factors on Adult Population

012168

THE FOLLOWING ARTICLE IS OPEN ACCESS

The Challenges of Managing Deep Vein Thrombosis in the Elderly: A Narrative Review

E P B Mulia and H Firdausi

[Open abstract](#), The Challenges of Managing Deep Vein Thrombosis in the Elderly: A Narrative Review [View article](#), The Challenges of Managing Deep Vein Thrombosis in the Elderly: A Narrative Review [PDF](#), The Challenges of Managing Deep Vein Thrombosis in the Elderly: A Narrative Review

012169

THE FOLLOWING ARTICLE IS OPEN ACCESS

Effect of Black Tea (*camellia sinensis*) on Serum Adiponectin Level in Atherogenic Diet Rats

M Muqsith, B S Pikir and M Rifqi

[Open abstract](#), [The Correlation Between Strain and Strain Rate Left Ventricle with Functional Capacity in Post Myocardial Infarction Patients with Heart Failure](#) [View article](#), [The Correlation Between Strain and Strain Rate Left Ventricle with Functional Capacity in Post Myocardial Infarction Patients with Heart Failure](#) [PDF](#), [The Correlation Between Strain and Strain Rate Left Ventricle with Functional Capacity in Post Myocardial Infarction Patients with Heart Failure](#)

012195

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Predicting the Likelihood for Severe CAD and CABG Indication on Elective Patients: Comparison of Novel CHA2DS2-VASc-HSF with CHA2DS2 and CHA2DS2-VASc Score](#)

M J Al-Farabi, I G P G Semita, K A Shonafi, R Ramadhiani, B Jovie and A Andrianto

[Open abstract](#), [Predicting the Likelihood for Severe CAD and CABG Indication on Elective Patients: Comparison of Novel CHA2DS2-VASc-HSF with CHA2DS2 and CHA2DS2-VASc Score](#) [View article](#), [Predicting the Likelihood for Severe CAD and CABG Indication on Elective Patients: Comparison of Novel CHA2DS2-VASc-HSF with CHA2DS2 and CHA2DS2-VASc Score](#) [PDF](#), [Predicting the Likelihood for Severe CAD and CABG Indication on Elective Patients: Comparison of Novel CHA2DS2-VASc-HSF with CHA2DS2 and CHA2DS2-VASc Score](#)

012196

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Early and Late Initiation Time of Statin Administration Effects on Early and Long-Term Outcome in Acute Coronary Syndrome Patients: A Literature Review](#)

M Q A'yun

[Open abstract](#), [Early and Late Initiation Time of Statin Administration Effects on Early and Long-Term Outcome in Acute Coronary Syndrome Patients: A Literature Review](#) [View article](#), [Early and Late Initiation Time of Statin Administration Effects on Early and Long-Term Outcome in Acute Coronary Syndrome Patients: A Literature Review](#) [PDF](#), [Early and Late Initiation Time of Statin Administration Effects on Early and Long-Term Outcome in Acute Coronary Syndrome Patients: A Literature Review](#)

012197

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Predictors of Arteriovenous Fistula Early Failure in End-Stage Renal Disease Patients: Real-World Data in Surabaya](#)

M R Amadis, J N E Putranto, I Maghfirah and N Mardiana

[Open abstract](#), [Predictors of Arteriovenous Fistula Early Failure in End-Stage Renal Disease Patients: Real-World Data in Surabaya](#) [View article](#), [Predictors of Arteriovenous Fistula Early Failure in End-Stage Renal Disease Patients: Real-World Data in Surabaya](#) [PDF](#), [Predictors of Arteriovenous Fistula Early Failure in End-Stage Renal Disease Patients: Real-World Data in Surabaya](#)

012198

THE FOLLOWING ARTICLE IS OPEN ACCESS

Different Grades of Body Mass Index are Correlated with Left Atrium and Ventricle Structures in Patients with Hypertensive Heart Disease

M S Bhisma, M J Al-Farabi and B S Pikiir

[Open abstract](#), [Different Grades of Body Mass Index are Correlated with Left Atrium and Ventricle Structures in Patients with Hypertensive Heart Disease](#) [View article](#), [Different Grades of Body Mass Index are Correlated with Left Atrium and Ventricle Structures in Patients with Hypertensive Heart Disease](#) [PDF](#), [Different Grades of Body Mass Index are Correlated with Left Atrium and Ventricle Structures in Patients with Hypertensive Heart Disease](#)

JOURNAL LINKS

- [Journal home](#)
- [Information for organizers](#)
- [Information for authors](#)
- [Search for published proceedings](#)
- [Contact us](#)
- [Reprint services from Curran Associates](#)
- [IOP Science home](#)

- [Journals](#)
- [Books](#)
- [About IOPscience](#)
- [Contact us](#)
- [Developing countries access](#)
- [IOP Publishing open access policy](#)
- [IOP Publishing home](#)

- [© Copyright 2020 IOP Publishing](#)
- [Terms & conditions](#)
- [Disclaimer](#)
- [Privacy & cookie policy](#) [This link opens in a new tab.](#)
- [This site uses cookies. By continuing to use this site you agree to our use of cookies.](#)

PAPER • OPEN ACCESS

Immature Platelet Level in Stable Coronary Heart Disease (CHD) Patients with Diabetes Mellitus compared to Stable CHD Patients without Diabetes Mellitus

To cite this article: D R Balti *et al* 2020 *IOP Conf. Ser.: Earth Environ. Sci.* 441 012165

View the [article online](#) for updates and enhancements.

Immature Platelet Level in Stable Coronary Heart Disease (CHD) Patients with Diabetes Mellitus compared to Stable CHD Patients without Diabetes Mellitus

D R Balti¹, A Andrianto^{1,2*}, B B Dharmadjati^{1,2} and D N Asmarani¹

¹Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

²Department of Cardiology and Vascular Medicine, Soetomo General Hospital, Surabaya, Indonesia

*Corresponding author: andricor9@yahoo.com

Abstract. Coronary heart disease (CHD) causes high mortality and morbidity in many countries. Immature platelet reactivity is one of the major causes of CHD thrombus formation due to its larger size and stronger aggregation properties than normal platelets. Though still controversial, diabetes mellitus (DM), which is one of the CHD risk factors, is mentioned to have a strong association with a high immature platelet level. Since studies on immature platelet in stable CHD are limited, the authors intended to search for the difference between immature platelet level in DM and non-DM patients with stable CHD. This research was an observational analytic study with a cross-sectional design; 31 samples of CHD patients who underwent coronary catheterization were obtained. The characteristics of the data was collected by completing a pre-operation form and the immature platelets were counted using an automated cytometry tool, SYSMEX XE 2100. They were analyzed using Pearson's statistical analysis. There was no significant correlation between the immature platelet levels of DM and non-DM patients with stable CHD ($p > 0.05$). There was a significant positive correlation between HbA1C and the immature platelet level (Immature Platelet Fraction, $p < 0.05$; Immature Platelet Count, $p < 0.01$) in stable CHD patients.

1. Introduction

Coronary heart disease (CHD) is one of the chronic diseases that is a major contributor to the morbidity rate, mortality rate and health care spending in both developing and developed countries. The National Heart, Lung and Blood Institute (NHLBI) explained that the enhancement risk for CHD occurs in those aged over 40 years in men and women by 49% and 32% respectively. From 2005 to 2008, NHLBI estimated that 16,300,000 Americans ≥ 20 years old had CHD and that they had a mortality rate as big as 126 over 100,000 populations [1]. Platelet reactivity related to immature platelets is the main driver of thrombus formation in CHD [2]. An immature platelet is a granule rich young platelet which contains the messenger RNA (mRNA) residue of megakaryocyte and Ribonucleic Acid (RNA). This makes protein biosynthesis possible. Immature platelets are also able to express cyclooxygenase-2 (COX-2) and they are of a bigger size than normal platelets [2-4].

Various research has proven that there is a relationship between peripheral immature platelets with coronary heart disease incidence. Immature platelets increase in the comorbid factors of CHD, such as



diabetes mellitus and smoking. Another research study stated that immature platelets are an independent predictor of mortality incidence in the Acute Coronary Syndrome (ACS) patients who underwent PCI for 12 months [2].

Immature platelets were counted using the Immature Platelet Fraction (IPF) and Immature Platelet Count (IPC) parameters. The IPC value, which was obtained by multiplying IPF with total platelet count, is considered to be the new marker of IPF for the stratification risk of a major cardiovascular event in CHD [5]. IPC is also related to platelet reactivity and it is an excellent predictor to use to evaluate the patient antiplatelet response to thienopyridine therapy, which is one of the primary therapies to prevent CHD restenosis [4,6]. Antiplatelets in CHD patients are one of the main therapies but according to some research, high immature platelet levels will decrease the effectiveness of antiplatelet therapy [5,7].

Up until today, a study comparing the immature platelet level of diabetes mellitus (DM) patients to non-diabetic mellitus patients with stable CHD has not been conducted yet. Therefore, researchers are interested in knowing more about the immature platelet level of diabetes mellitus patients and how it affects CHD incidence.

2. Methods

This research was an observational analytical study using a cross-sectional design. This study analyzed the difference in immature platelet level between the CHD patients with DM compared to those with CHD who were non-DM patients.

The samples were CHD patients in Surabaya who underwent heart catheterization at Instalasi Diagnostik Intervensi Kardiovaskular (IDIK) Dr. Soetomo General Hospital from September 2017 until December 2017. We included stable CHD patients who agreed to sign the informed consent form. We excluded the CHD patients who had hematologic abnormalities causing thrombocytopenia and those who refused to sign the informed consent.

The calculation of the minimum sample size was done using the hypothesis test formula for the analysis of the comparative numerical data with 2 non-pair group, with a confidence level of 95%. The power of the test was 95%, the standard deviation was 1.82 and the significant mean difference between immature platelets considered by the author was 1.5. The minimum sample size was 31 samples and the sampling method was total sampling.

The sample's characteristic data was collected when the samples showed up for catheterization preparation through anamneses and a physical and laboratory examination. A blood sample was collected at IDIK Dr. Soetomo General Hospital. The sample characteristics were observed using the pre-cath form and the immature platelet levels were observed using laboratory tools and automated flow cytometry, SYSMEX XE 2100 at Gedung Diagnostic Center (GDC) of Dr. Soetomo General Hospital.

The data was analyzed using SPSS 22.0. The sample characteristic data included age, sex, CHD family history, history of diabetes mellitus, hypertension, dyslipidemia and smoking habits. These characteristics were processed as descriptive statistic data. They were shown descriptively as the mean and standard deviation or as the median and frequency in percentage form. The data distribution was tested using the Saphiro Wilk test. The T2 independent sample comparison test was conducted only if both sample group were homogenous. The α value was 0.05, which had mean that the IPC and IPF difference between two sample groups were significant only if the p value were < 0.05 . The Pearson method was conducted to analyze the correlation between immature platelet level and diabetes mellitus.

3. Results

This study was conducted in the IDIK of Dr. Soetomo General Hospital from October 2017 until December 2017. This study enrolled 31 CHD patients who underwent catheterization that met the research criteria.

3.1 Samples Characteristics

There were 32 subjects that consisted of 23 males (71.9%) and 9 females (28.1%). The youngest age was 34 years old while the oldest was 78 years old. The mean age was 57 years old. The risk factors for CHD obtained from the subjects were hypertension (53.1%), hypercholesterolemia (62.5%), smoking (56.3%) and DM (46.9%). The subject's baseline characteristic data has been shown in Table 1.

Table 1. Baseline characteristics of the samples.

Variables	Non-diabetes mellitus (N=17)	Diabetes mellitus (N=15)	<i>p</i>
Unmodifiable Factors			
Males, n (%)	14 (60.9%)	9 (39.1%)	0.453 ^a
Females, n (%)	4 (44.4%)	5 (55.6 %)	
Age, mean ± SD	57.44 ± 11.036	8 ± 6.109	0.809 ^b
Modifiable Factors			
Hypertension, n (%)	7 (41.2%)	10 (58.8%)	0.067 ^a
Hypercholesterolemia, n (%)	8 (40%)	12 (60%)	0.017 ^a
Smoking, n (%)	12 (66.7%)	6 (33.3 %)	0.178 ^a
Aspilet Consumption, n (%)	13 (52.0%)	12 (48.0 %)	0.426 ^b
Clopidogrel Consumption, n (%)	11 (52.4%)	10 (47.6 %)	0.691 ^b

^a Pearson's chi square test

^b Fisher's test

The table implies that there is a significant relationship between hypercholesterolemia and DM. Furthermore, we analyzed the correlation between hypercholesterolemia and IPF/IPC. The result was a non-significant correlation between hypercholesterolemia with IPF and IPC ($P > 0.05$) as shown in Table 2. Therefore, the significant correlation between hypercholesterolemia and DM could be ignored in this research.

Table 2. The relationship between hypercholesterolemia and IPF/IPC.

Variable	Immature platelet fraction (<i>p</i>)	Immature platelet count (<i>p</i>)
Hypercholesterolemia	0.537	0.537

3.2. Analysis Result

The researchers divided the immature platelets into two values, namely IPF and IPC. The results showed there to be a statistically significant correlation between IPF and IPC level, implying that the higher IPF level, the higher the IPC level would be in turn (Table 3).

The mean IPF and IPC levels in the CHD non-DM group were 3.89 ± 1.689 and 994 ± 409.86 , respectively. The mean IPF and IPC on CHD in the DM group was 4.050 ± 2.002 and 1075 ± 564.76 respectively. Further analysis of the correlation between DM and the IPF/IPC level showed that DM influences the IPF and IPC level, although it was not significant (Table 4).

Table 3. Correlation between IPF and IPC.

Variable	Immature platelet count (N=32)	
	<i>r</i>	<i>p</i>
Immature Platelet Fraction (N=32)	0.781	<0.001

Table 4. IPF and IPC level on CHD with DM and CHD without DM comorbid.

Variable	Non DM (Mean \pm SD)	DM (Mean \pm SD)	p
Immature Platelet Fraction	3.89 \pm 1.689	4.050 \pm 2.002	0.667
Immature Platelet Count	994 \pm 409.860	1075 \pm 564.760	0.896

We also analyzed the correlation between HbA1C and IPF/IPC using a cross-tabulated test. There was a significant correlation between HbA1C and the IPF level, in that the higher the HbA1C level, the higher the IPF level would be in turn ($P < 0.05$) (Table 5).

Table 5. Correlation between HbA1C and IPF/IPC

Variable	HbA1C	
	r	p
Immature Platelet Fraction	0.781	<0.001
Immature Platelet Count	0.413	0.067

Furthermore, we tried to analyze the independent contribution of each of these variables to the immature platelet level using the multivariate regression test. The correlation between IPF and some of the risk factors were analyzed, including smoking, hypercholesterolemia, blood glucose level, HbA1C, creatinine serum, clopidogrel and aspirin therapy. Clopidogrel was found to be the most influence on the IPF level. The clopidogrel value was negative ($\beta = -0.576$) and significant ($p < 0.05$). Subjects who received clopidogrel had the lowest immature platelet level.

Table 6. Multivariate analysis table.

Variable	Immature platelet fraction	
	β	p
Clopidogrel	-0.576	0.01

4. Discussion

4.1 Characteristics of subjects

This study enrolled 32 subjects with stable CHD who met the research criteria. The gender proportion of the subjects was 71.9% male and 28.1% female. We can conclude that the male gender is one of the risk factors for CHD. The average age for all of the subjects was 57 years old. The data implied that the manifestation of CHD starts at the age of 40 [8].

Various modifiable risk factors were found in the subjects, including hypertension (53.1%), hypercholesterolemia (62.5%), smoking (56.3%) and diabetes (46.9%). The hypertensive condition affects the vascular endothelial layer directly, causing endothelial dysfunction and leading to the formation of atherosclerotic plaque. Hypercholesterolemia is known as the main component of atherosclerosis development. This research showed that hypercholesterolemia had the highest proportion. The high proportion of smoking in this study also supports the theory that smoking can increase the risk of CHD incidence [9].

Almost 50% of subjects had DM as a CHD risk factor. Diabetes mellitus is related to abnormalities in the lipid metabolism, obesity, systemic hypertension and the raised risk of thrombogenesis that contributes to the development of CHD [10].

4.2 Immature platelet

Two values for the immature platelet parameters were obtained; IPF (3.959 ± 1.8) and IPC (1029.88 ± 477.0). The Pearson correlation test found there to be a highly significant positive

correlation (coefficient 0.781) between the IPF level and the IPC level. This result means that the higher the IPF level, the higher the IPC level would be as well. This result is in line with the previous research that said that the IPC value is obtained by multiplying IPF with the platelet count [6,11]. It could be concluded that given the strong relationship between IPF and IPC, an immature platelet level can be interpreted just by using one of the parameters.

From this study, we found that DM had a positive effect on IPF and IPC but that this effect was not statistically significant. In other words, DM is associated with an increase in immature platelet level but it cannot be generalized to all populations aside from the research sample. This insignificant correlation could result from the small sample size, the controlled glycemic index of some of the DM subjects, the presence of other factors which would affect immature platelets (i.e. hypercholesterolemia, hypertension, and smoking history) or the possibility of non-valid measurement tools [12]. Compared to the patients with DM status, the correlation analysis between HbA1c and IPF/IPC level had a different result. HbA1c, a parameter used to evaluate and control the glycemic index for a 3-month period, gave positive and statistically significant correlation results. This means that if the HbA1c level is high, then the IPF level would also be high.

Some of the mechanisms that are involved in the hyperglycemic condition related to an immature platelet level are thromboxane synthesis and calcium mobilization. This will cause adhesion at the cell's surface, which has a direct effect on the C protein given the hyper-reactivity of the platelet. The osmotic effect generates activity in the platelets and it suppresses the effect of NO and PGI₂, which are low in the DM condition. The raised osmosis also causes a bigger platelet size [13].

4.3 Clopidogrel

Clopidogrel had a stronger correlation to IPF than the other clopidogrels. It had a negative ($\beta = -0.576$) and significant correlation to the IPF level. Subjects undergoing clopidogrel therapy had a lower IPF level. On the contrary, aspirin didn't have a better correlation with a lower IPF level compared to clopidogrel.

Aspirin has a lower relationship associated with high platelet levels in circulation [14] and it only has a short half-life (2–6 hours). Aspirin binds irreversibly to platelets in the first 2-6 hours and it will last during the age of the platelet. If the platelet age becomes shorter, as in the immature platelet condition (<1 day), then there will be many more new platelets produced by the spinal cord. This would result in a lot of platelets that will not bind to the aspirin.

5. Conclusion

There is a non-significant difference in the immature platelet levels between stable CHD in DM patients and the stable CHD in non-DM patients. However, there is a significant positive correlation between glycemic control and the immature platelet fraction level.

References

- [1] Roger V, Go A, Lloyd-Jones D and Al E 2011 Heartdisease stroke statistics – 2011 update: Areport from the American Heart Association. *Circulation* **123** e18–e209
- [2] Cesari F, Marcucci R, Gori A M, Caporale R, Fanelli A, Casola G, Balzi D, Barchielli A, Valente S, Giglioli C, Gensini G F and Abbate R 2013 Reticulated platelets predict cardiovascular death in acute coronary syndrome patients: Insights from the AMI-florence 2 study *Thromb. Haemost.* **109** 846–53
- [3] Kieffer N, Guichard J, Farcet JP, Vainchenker W and BretonGJ 1987 Biosynthesis of major platelet proteins in human blood platelets *Eur. J. Biochem.* **164** 189–95
- [4] Würtz M, Grove E L, Wulff L N, Kalsoft A K, Tilsted H H, Jensen L O, Hvas A M and Kristensen S D 2010 Patients with previous definite stent thrombosis have a reduced antiplatelet effect of aspirin and a larger fraction of immature platelets *JACC Cardiovasc. Interv.* **3** 828–35

- [5] Ibrahim H, Schutt R C, Hannawi B, DeLao T, Barker C M and Kleiman N S 2014 Association of immature platelets with adverse cardiovascular outcomes *J. Am. Coll. Cardiol.* **64** 2122–9
- [6] Stratz C, Amann M, Valina C, Nuehrenberg T, Trenk D, Neumann F J and Hochholzer W 2016 Immature platelet count or immature platelet fraction as optimal predictor of antiplatelet response to thienopyridine therapy *J. Am. Coll. Cardiol.* **67** 206
- [7] Guthikonda S, Alviar C L, Vaduganathan M, Arikian M, Tellez A, DeLao T, Granada J F, Dong J F, Kleiman N S and Lev E I 2008 Role of reticulated platelets and platelet size heterogeneity on platelet activity after dual antiplatelet therapy with aspirin and clopidogrel in patients with stable coronary artery disease *J. Am. Coll. Cardiol.* **52** 743–9
- [8] Braunwald E and Bonow R O 2012 *Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine* (Philadelphia: Elsevier Saunders).
- [9] Grundy S M, Pasternak R, Greenland P, Smith Jr S and Fuster V 1999 Assessment of cardiovascular risk by use of multiple-risk-factor assessment equations a statement for healthcare professionals from the American Heart Association and the American College of Cardiology *Circulation* **100** 1481–92
- [10] Fuster V, Moreno P R, Fayad Z A, Corti R and Badimon J J 2005 Atherothrombosis and high-risk plaque: Part I: Evolving concepts *J. Am. Coll. Cardiol.* **46** 937–54
- [11] Eikelboom J W and Warkentin T E 2014 Immature platelet count: Part of the cardiologist's complete blood count? *J. Am. Coll. Cardiol.* **64** 2130–2
- [12] Widhiarso W 2010 *Model Analisis dengan Menggunakan Variabel Moderator* (Yogyakarta: UGM) pp 1–2
- [13] Keating F K, Sobel B E and Schneider D J 2003 Effects of increased concentrations of glucose on platelet reactivity in healthy subjects and in patients with and without diabetes mellitus *Am. J. Cardiol.* **92** 1362–5
- [14] Vaduganathan M and Lev E I 2014 Aspirin resistance antiplatelet *Ther. Cardiovasc. Dis.* **367** 277–84