

Identification of Legionella Pneumophila Serogroups as Zoonotic Disease Agent Distributed in Water Sources of East Java

by Eduardus Bimo Aksono

Submission date: 30-Jun-2021 10:59AM (UTC+0800)

Submission ID: 1613988250

File name: Identification_of_Legionella-Q3.pdf (200.25K)

Word count: 1491

Character count: 7817

Identification of *Legionella Pneumophila* Serogroups as Zoonotic Disease Agent Distributed in Water Sources of East Java

Eduardus Bimo Aksono¹, Kadek Rachmawati, Retno Bijanti and Herinda Pertiwi

Institute of Tropical Disease, Faculty of Veterinary Medicine, Faculty of Vocational Studies Universitas Airlangga - Kampus C Mulyorejo, Surabaya, Indonesia. 60115.

(Received : June, 2019 207/19 Accepted : August, 2019)

Abstract

The aims of this study was to understand the differences of *L.pneumophila* serogroups distribution in well water, tap water, ice cubes, hospital water and hotel water in East Java-Indonesia. A total of 60 water samples were tested by polymerase chain reaction and then it was analyzed by phylogenetic tree. Out of the 60 water samples collected, 12% of the samples were contaminated with *L. pneumophila*. The phylogenetic tree revealed *L.pneumophila* contamination in well water from Surabaya and tap water from Sidoarjo and the ice cubes from Sidoarjo, while the bacterial contamination in 2 well water isolate from Surabaya classified into their own group.

Key words: *L.pneumophila*, serogroup contamination distribution.

Legionella spp. are gram-negative bacteria, found in nature, and spreads by air upto 300 m radius. These bacteria are zoonotic and, infect the lungs leading to pneumonia which can be

life-threatening, by inhalation of contaminated droplets of water (Ghotaslou *et al*, 2013). These bacteria thrives well in any environments; 56% in domestic hot water system, 44% in water cooling system and their ideal conditions are at the pH 6.9 and temperature 35-37°C with availability of ferric oxide as nutrients in the water. The bacteria are prevalent in any buildings with cooling and hot water system such as in office building and hotels. That is why the illness caused by the bacteria is also called “sick building syndrome”.

Materials and Methods

60 samples of water (200ml) were collected from different environments in Surabaya-Sidoarjo-Kediri East Java such as well water (25), tap water (5), ice cubes (5), hospital water (16), and hotel water (9). Samples were put in a sterile vials then filtered using a Millipore membrane 0.22 µm, and they were put in a 50 ml conical flask and rinsed with 1 ml Phosphate Buffer solution (PBS) vortex for 10 minutes. 1 ml of this was transferred into into

¹Corresponding author : Email : eduardus-b-a-h@fkh.unair.ac.id

Table I. Presence of *Legionella* bacteria in water samples from East Java by PCR

Origin of Samples	Results		Species
	Negative	Positive	
Well water	23/25	2/25	<i>L. pneumophila</i>
Tap water	4/5	1/5	<i>L. pneumophila</i>
Ice cubes	4/5	1/5	<i>L. pneumophila</i>
Hospital water	16/16	0/16	<i>L. pneumophila</i>
Hotel water	6/9	3/9	<i>L. pneumophila</i>
TOTAL	53/60	7/60	<i>L. pneumophila</i>

ependorf tubes, and centrifuged at 13000 rpm for 3 minutes. The supernatant was discarded and the sediment was used for DNA extraction using DNA extraction kit (QIAamp®DNA mini kit Qiagen) as per the manufacturer's instructions. Phylogenetic analysis was done by using software Genetix Mac Ver. 10.0

Results and Discussion

Table I showed that using PCR method for the 60 water samples collected in East Java, 12% of the samples (7/60) were contaminated by *L. pneumophila*. In details, 8% of the well water samples (2/25), 2% of the tap water samples (1/5), 2% of the ice cubes samples(1/5), 0% of the hospital water samples (0/16) and 33.33% of the hotel water samples (3/9) were positive for *L. pneumophila*.

Results of Phylogenetic analysis showed that *Legionella pneumophila* contamination from well water was 1 from Surabaya and in tap water from Sidoarjo which were closer to *L.pneumophila* serogroup 2, 3, 4, 6, 9, 10, 12, isolates reported from Brazil, China, Spain and Australia. *L.pneumophila* contamination in the ice cubes from Sidoarjo were closer to serogroup 1, 7, 8, 11, 13, 14, while the bacteria contaminating well water were 2 and water from hotels of Surabaya (hotel water isolate 1, 2 and 3).

The *Legionella* bacteria contamination was found in little quantity (>1,000 CFU/liter) (Qin *et al*, 2012) in tap water and it is not dangerous, (Aksono and Hermadi, 2017). However, they can multiply fast in flood waters.

The infection is transmitted through water spray or aerosol contaminated with

Legionella microbes. The infected person will fall ill after five or six days from the exposure. *Legionella* infection risk is less in small houses than the big building. In family housing, the infection was found only 6-30%. Prevention of *L.pneumophila* transmission could be contained by added chlorine >2 mg/liter of water (Moran-Gilad *et al*, 2014; Sánchez-Busó *et al*, 2015).

In this research, *L.pneumophila* distribution of serogroups from the natural water (such as well water) was different from different water source (such as water tank and drinking water system in hotels). These results are in line with the previous research conducted in Japan and Canada (Amemura-Maekawa *et al*, 2012; Reimer *et al*, 2009), they may correlated with water source characteristic differences (such as,




Fig 1. Results of PCR *L.pneumophila* in water samples from East Java by electrophoresis gel in 2% (403 bp). M (marker); K+ (positive control); K-(negative control); 1 (positive from tap water); 3 (positive from ice cubes).

temperature and pH level). There are various types of water sources, while cooling tower and drinking water system tend to have similar characteristics due to same water treatments. *Legionella* has high prevalence, due to their faster intracellular growth, and in diverse hot water sources (Qin *et al.*, 2013).

L. pneumophila isolated from hot water tank in hotels had higher genetic diversity, compared with natural water and drinking water. This result may be related to growth in amoeba which adapts in different environment. It has been reported that the growth of *L. Pneumophila* in amoebae host depends on the bacterial genetics (Buse *et al.*, 2012; Dey *et al.*, 2009).

Summary

L.pneumophila serogroups isolated from environment water in East Java had a closer correlation with serogroups from Brazil, China, Spain and Australia and some samples were comparable to the local serogroups.

Acknowledgement

We would like to express our gratitude to Rector of Universitas Airlangga and the Ministry of Research, Technology and Higher Education for PDUPTresearch grants given in 2018.

References

Aksono, E.B., and Hermadi, H.A. (2017) Rapid detection of *Legionella pneumophila* in the water environment in Surabaya-Indonesia. *International J. of Environ. Biology.*7(1) : 5-10.

Amemura-Maekawa, J., Kikukawa, K., Helbig, J.H., Kaneko, S., Suzuki-Hashimoto, A., Furuhata, K., Chang, B., Murai, M., Ichinose, M., Ohnishi, M., and Kura, F.(2012) Distribution of

monoclonal antibody subgroups and sequence-based types among *Legionella pneumophila* serogroup 1 isolates derived from cooling tower water, bathwater, and soil in Japan. *Appl. Environ. Microbiol.* 78:4263–4270.

Buse, H.Y., and Ashbolt, N.J. (2011) Differential growth of *Legionella pneumophila* strains within a range of amoebae at various temperatures associated with in-premise plumbing. *Lett. Appl. Microbiol.* 53:217–224.

Dey, R., Bodennec J., Mameri, M.O., and Pernin, P (2009) Free-living freshwater amoebae differ in their susceptibility to the pathogenic bacterium *Legionella pneumophila*. *FEMS Microbiol. Lett.*290:10–17.

Ghotaslou, R., Sefidan, F.Y., Akhi, M.T., Soroush, M.H., and Hejazi, M.S. (2013) Detection of *Legionella* Contamination in Tabriz Hospitals by PCR Assay. *Adv Pharm Bull.* 3(1): 131–134.

Moran-Gilad, J., Mentasti, M., Lazarovitch, T., Huberman, Z., Stocki, T., Sadik, C., Shahar, T., Anis, E., Valinsky, L., Harrison, TG., and Grotto I., (2014) ESCMID Study Group for *Legionella* Infections (ESGLI). Molecular epidemiology of Legionnaires' disease in Israel. *Clin Microbiol Infect* 20:690–696.

Reimer, A.R., Au, S., Schindle, S., and Bernard, K.A. (2010) *Legionella pneumophila* monoclonal antibody subgroups and DNA sequence types isolated in Canada between 1981 and 2009: laboratory component of national surveillance. *Eur. J. Clin. Microbiol. Infect. Dis.*29:191–205.

Sánchez-Busó, L., Coscollà, M., Palero, F., Camaró, M.L., Gimeno, A., Moreno, P., Escribano, I., López Perezagua, M.M., Colomina, J., Vanaclocha, H., and González-Candelas, F. (2015) Geographical and temporal structures of *Legionella pneumophila* sequence types in Comunitat Valenciana (Spain), 1998 to 2013. *Appl Environ Microbiol* 81:7106–7113.

Qin, T., Tian, Z., Ren, H., Hu, G., Zhou, H., Lu, J., Luo, C., Liu, Z., and Shao, Z. (2012) Application of EMA-qPCR as a complementary tool for the detection and monitoring of *Legionella* in different water systems. *World J. Microbiol. Biotechnol.*28:1881–1890.

Qin, T., Yan, G., Ren, H., Zhou, H., Wang, H., Y., Zhao, M., Guan, H., Li, M., and Shao, Z. (2013) High prevalence, genetic diversity and intracellular growth ability of *Legionella* in hot spring environments. *PLoS One* 8(3): e59018.

Identification of Legionella Pneumophila Serogroups as Zoonotic Disease Agent Distributed in Water Sources of East Java

ORIGINALITY REPORT

14%

SIMILARITY INDEX

8%

INTERNET SOURCES

7%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Tian Qin, Haijian Zhou, Hongyu Ren, Hong Guan, Machao Li, Bingqing Zhu, Zhujun Shao. "Distribution of Sequence-Based Types of Legionella pneumophila Serogroup 1 Strains Isolated from Cooling Towers, Hot Springs, and Potable Water Systems in China", Applied and Environmental Microbiology, 2014
Publication 3%
- 2 ivj.org.in
Internet Source 2%
- 3 eprints.nottingham.ac.uk
Internet Source 1%
- 4 "Detection of Legionella pneumophila from domestic water and their antibiotic resistance profiles", Journal of Coastal Life Medicine, 2015
Publication 1%
- 5 Submitted to Universitas Airlangga
Student Paper 1%

6

[bioone.org](https://www.bioone.org)

Internet Source

1 %

7

Boyang Cao, Xiangqian Liu, Xiang Yu, Min Chen, Lu Feng, Lei Wang. "A New Oligonucleotide Microarray for Detection of Pathogenic and Non-Pathogenic Legionella spp.", PLoS ONE, 2014

Publication

1 %

8

Jingyu Guo, Ting Liang, Chaohui Hu, Ruichen Lv, Xianwei Yang, Yujun Cui, Youtao Song, Ruifu Yang, Qingyi Zhu, Yajun Song. "Sequence types diversity of Legionella pneumophila isolates from environmental water sources in Guangzhou and Jiangmen, China", Infection, Genetics and Evolution, 2015

Publication

1 %

9

K. Sreenath, Rama Chaudhry, E. V. Vinayaraj, A. B. Dey, S. K. Kabra, Bhaskar Thakur, Randeep Guleria. "Distribution of Virulence Genes and Sequence-Based Types Among Legionella pneumophila Isolated From the Water Systems of a Tertiary Care Hospital in India", Frontiers in Public Health, 2020

Publication

1 %

10

[nutriweb.org.my](https://www.nutriweb.org.my)

Internet Source

1 %

11 www.neliti.com 1 %
Internet Source

12 David K. Boamah, Guangqi Zhou, Alexander W. Ensminger, Tamara J. O'Connor. "From Many Hosts, One Accidental Pathogen: The Diverse Protozoan Hosts of Legionella", *Frontiers in Cellular and Infection Microbiology*, 2017 1 %
Publication

13 dspace.alquds.edu 1 %
Internet Source

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Identification of Legionella Pneumophila Serogroups as Zoonotic Disease Agent Distributed in Water Sources of East Java

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
