

Detection of the extended
spectrum β -lactamase
produced by *Escherichia coli*
from dairy cows by using the
Vitek-2 method in Tulungagung
regency, Indonesia

by Wiwiek Tyasningsih

Submission date: 05-Nov-2021 04:40PM (UTC+0800)

Submission ID: 1693749298

File name: Detection_of_the_extended_spectrum....pdf (690.75K)

Word count: 3287

Character count: 17788

Detection of the extended spectrum β -lactamase produced by *Escherichia coli* from dairy cows by using the Vitek-2 method in Tulungagung regency, Indonesia

A.R. Putra¹, M.H. Effendi², S. Koesdarto³, S. Suwarno⁴,
W. Tyasningsih⁵ and A.T. Estoe pangestie⁶

¹ Postgraduate Program on Veterinary Public Health, ^{2,6} Department of Veterinary Public Health, ³ Department of Veterinary Parasitology, ^{4,5} Department of Veterinary Microbiology, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia, Email: ¹ akyunrozaqis@gmail.com, ² mheffendi@yahoo.com, ³ skoedarto@yahoo.com, ⁴ snow_arno@yahoo.co.id, ⁵ witya_kh@yahoo.com, ⁶ soelih.estoe pangestie@gmail.com

(Received May 15, 2019; Accepted July 10, 2019)

Abstract

The use of antibiotics in inappropriate on food producing animals can lead to resistance many of the pathogenic bacteria to the various types of antibiotics, one of which is the *Escherichia coli* (*E. coli*) which produces extended spectrum β -lactamase (ESBL). Antibiotic resistance in animals and humans has become a global problem that needs attention and immediate management by using specific antibiotics that used for therapeutic the infected animals. The aim of this study was to isolate and detect *E. coli* producing ESBL. All *E. coli* from the surface of dairy cow rectal swabs in Sendang District, Tulungagung Regency, Indonesia using the Vitek-2 method. The number of rectal swab samples used in the present study was 50. The results of this study showed that all the samples were suspected of being *E. coli*, based on the morphological growth of colonies on EMBA media. The isolates were identified by using the biochemical tests. All the samples were positive. In this study the double disc synergy test (DDST) method was using to confirm the ESBL. The antibiotics were used ampicillin-clavulanate, ceftazidime and cefotaxime for DDST. In additional ESBL confirmation test was used the Vitek-2 method. The presence of ESBL producing by *E. coli* isolated from rectal dairy swabs in tulungagung was 6% (3/50).

Keywords: *E. coli*, ESBL, Vitek-2, Dairy Cows

Available online at <http://www.vetmedmosul.com>, © 2020, College of Veterinary Medicine, University of Mosul. This is an open access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

الكشف عن إنزيم بيتا-اكتام ممتد الطيف المنتج من الايشيريكيا القولونية من أبقار حلب بوساطة طريقة الـVitek-2 في تولونججونج، اندونيسيا

أكيون روزاقي بوترا¹، مصطفى حلمي إيفندي¹، سيتاوان كويسدارتو²، سوارنو سوارنو³، ويويك تياسنينكسيه³
و أكنس ايستويبانكيسي¹

¹ فرع الصحة العامة البيطرية، ² فرع الطفيليات البيطرية، ³ فرع الاحياء المجهرية البيطرية،
كلية الطب البيطري، جامعة آيرلانكا، سورابايا، إندونيسيا

الخلاصة

يمكن أن يؤدي الاستخدام غير الملائم للمضادات الحيوية في الحيوانات المنتجة للغذاء إلى مقاومة العديد من الجراثيم الممرضة لأنواع مختلفة من المضادات الحيوية، منها الايشيريكيا القولونية المنتجة لانزيم بيتا-اكتام ممتد الطيف (ESBL). لقد أصبحت مقاومة الجراثيم للمضادات الحيوية في كل من الإنسان والحيوانات مشكلة عالمية تتطلب الانتباه لها وإجراء تدابير أنية لتفاديها من خلال استخدام مضادات

حياتية محددة لعلاج الحيوانات المصابة. هدفت هذه الدراسة إلى الكشف عن الإيشيريكيا القولونية المنتجة لإنزيم ESBL وعزلها. عزلت الإيشيريكيا القولونية من مسحات مستقيم أبقار حلب في مقاطعة سيندنج، تولونججونج، اندونيسيا باستخدام طريقة Vitek-2. بلغ عدد المسحات في هذه الدراسة ٥٠ مسحة. بيّنت النتائج أن العينات جميعها مشتبه بأنها إيشيريكيا قولونية بالاعتماد على المظهر الشكلي للمستعمرات النامية على وسط EMBA. مُيزت العزلات بوساطة الاختبارات الكيموحيوية، وكانت موجبة جميعها. استخدمت طريقة DDST لتأكيد إنزيم ESBL. شملت المضادات الحيوية التي استخدمت في طريقة DDST كلٌّ من: الأموكسيسيلين-كلافولانيت، السيفتازيديم، والسيفوتاكسيم. شكّل وجود إنزيم ESBL المنتج من الإيشيريكيا القولونية المعزولة من مسحات مستقيم الأبقار الحلوب في تولونججونج نسبة ٦% (٥٠/٣).

Introduction

The occurrence of antibiotic resistance is caused by the excessive and inaccurate use of antibiotics in the treatment. The Bacteria show the resistance to the antibiotics when their growth cannot be inhibited by using the antibiotics at the maximum doses. The development³² of the resistance of the microorganisms to antibiotics is a consequence of the wrong use of antibiotics. The bacteria resistance to antibiotics is due to occur the mutation or resistance of the gene to specific antibiotics. The antibiotic resistance is currently a public health problem and has been reported by the WHO that the antibiotic resistance poses a threat to human health (1). The bacteria resistance to antibiotics have a different mechanism, one of them is through a gene transfer via a plasmid. The gene responsible for the production of β -lactamase is located in a plasmid or chromosome of the most important enzymes carried by plasmids. The bacteria that produce the betalactamase enzyme can be called¹⁵ the bacterium ESBL. The ESBL is an enzyme which has the ability to hydrolyze first, second, and third generation cephalo⁹prin antibiotics, and aztreonam groups. Carbapenem is not inhibited by beta-lactamase inhibitors such as clavular¹³, sulbactam and tazobactam (2). The resistance caused by ESBL bacteria is often associated with resistance to other antibiotics used in humans (1). The infected animals play an important role to spread bacteria resistant to antibiotics by contaminating the farms through the feces, slaughterhouses, and workers which are direct contact with the infected animals. In addition, The animals may be exposed to the contamination by drinking the water contains the bacteria resistant to antibiotic (3). The pets at home play an important role to infect the human with bacteria resistant to antibiotics, especially the animals which are closely related to human⁷ (4). Many of the studies showed There is similar of the ESBL producing *E. coli* which isolated in humans, livestock, wildlife (5).

The⁶ aims of the study are to isolate and detect the presence of ESBL producing *E. coli* through its resistance to β -lactam antibiotics using the Vitek-2 method. This method is very superior for detecting betalactam enzymes. The method of Vitek-2 is regarded as an automatic identification system for microorganism²⁴ show the phenotype of the isolates tested and able to determine the sensitivity or resistance of an isolate to antibiotics (6).

Materials and methods

Sampling method

The rectal swab samples were taken from November 2018 to March 2019 from dairy farms in Sendang district, Tulungagung regency. The number of samples taken from rectal swab samples in dairy cows was 50. The swab results are labeled and all the samples must be aseptic by using a sterile stick swab then put it in an eppendorf tube containing 1% pepton water buffer media. All the²⁸ samples are stored in the ice box and immediately transport to the laboratory for inspection (7).

Isolation and identification of *E. coli*

The swab rectal samp²⁵ were planted on brilliant green bile broth (BGBB) media (Merck, Darmstadt, Germany). All the samples were incubated at 37 °C for 18-24 hours (8). The positive results are indicated by the presence of gas bubbles in the durham tube and green changes to cloudy green. ⁸ after being positive, the *E. coli* isolates were planted in eosin methylene blue (EMBA) media (Merck, Darmstadt, Germany) by streaking and incubating at 37 °C for 24 hours. The suspect colonies of *E. coli* on EMBA media are metallic green. The colonies of *E. coli* grown in EMBA were planted again in 1% pepton water buffer media, then incubated at 37 °C for 24 hours. The incubated pepton water media is dripped with Kovach reagents in two or three drops. A positive result for *E. coli* is indicated by the formation of a red ring on the surface of the media at 1% pepton water buffer. Furthermore, the positive isolates of *E. coli* were planted on EMBA and incubated at 37 °C for 24 hours (9).

ESBL confirmation: Double Disk Synergy Test (DDST)²⁷

The *E. coli* ESBL confirmation test was carried out by using DDST method which produced a qualitative category with sensitive, inter²mediate and resistant assessment using an antibiotic disk (OXOID, Basinstoke, United Kingdom) amoxicillin-clavulanate 30 μ g (CT0223B), cefotaxime 30 μ g (CT0166), ceftazidime 30 μ g (CT0412). Planting on the plate to be carried out by taking the *E. coli* isolates in the standard turbidity test Mc Farland 0.5 was then gently rubbed with a sterile swab on the entire surface of the mueller hinton agar (MHA) media. Interpretation of² results by measuring the inhibition of the diameter of the inhibitory

zone formed in accordance with the Clinical and Laboratory Standards Institutions (CLSI) (10).

Vitek-2 compact

All the resistance category of the *E. coli* isolates were tested for the phenotype identification and confirmation using the Vitek-2 compact system method and worked according to the protocol (BioMerieux, Marcy L'Eyoily, France). The results automatically came out in the form of a pint out.

Results

The result of this study showed that all the *E. coli* isolated from the rectal dairy were positive 100%. All the *E. coli* showed a change in bright green to cloudy green and gas in the durham tube on BGBB media. In the EMBA media all the isolates showed a change in color to metallic green. There is a red ring on the pepton water buffer media when added with the kovach reagent in the pepton water buffer media as many as 50 positive samples of *E. coli*. The positive isolates of *E. coli* were purified by eosin methylene blue agar (EMBA) media which appeared metallic green (Figure 1).

The ESBL confirmation test of the *E. coli* isolates using DDST and Vitek-2 methods. In this method there were three positive samples of ESBL 6% (Table 1). In the DDST method there is an enlargement zone with the synergy pattern of the three antibiotics with amoxycylin-clavulanate as its inhibitor (Figure 2).




Figure 1: Presumptive *E. coli* in EMBA media with the metallic green color.



Figure 2: The double disc synergy test for ESBL confirmation.

Table 1: Data of the *E. coli* isolates in this study from Tulungagung farms

Location	Tulungagung regency
No. samples	50
Positive <i>E. coli</i>	50
ESBL Confirmation by DDST	3
ESBL Confirmation by Vitek-2	3

Discussion

The infected animals that have closely related to the human that becomes the main reservoir to the spread of bacteria resistant to an antibiotic in the environment and infected the human (11). O'brien (12) said that *E. coli* plays a role in the spread of genes resistant to bacterial populations between animal and humans through foodborne.

Screening of ESBL producing bacteria is usually done in humans only and in the hospital environment, but there are many studies show that the emergence of pathogenic bacteria had found in the livestock and animal food (13). The presence of ESBL producing Enterobacteriaceae bacteria in the feces samples which taken of livestock may raise the risk of carcass contamination at the time of cutting in the slaughterhouse, thus lead to potentially contaminating meat products (14). ESBL producing Enterobacteriaceae, including *E. coli*, which contaminate products of animal origin have the potential to cause health risks even though the level of risk is difficult to quantify (15). The ESBL penngssil infection through food consumption from animal

origin can cause limited choices in handling patients. This situation can extend the period of treatment, increase the cost of treatment, increase the incidence of disease, and the death (16).

Many of research reported on cases of ESBL producing *E. coli* in animals have been widely carried out in the various countries. Babypedmini and Appalaraju (17) showed that ESBL producing *E. coli* was 41% and *K. pneumonia* was 40%. The prevalence of ESBL producing *E. coli* in Malaysia was 5.6%, Japan 8.1%, Philippines 13.14, Taiwan 16.7% and Indonesia 23% (18). The prevalence of ESBL producing *E. coli* from beef feces in Bogor, Indonesia was 15.8% (19). In addition, Haenni *et al.* (20) showed that the prevalence of ESBL producing *E. coli* from 491 calf stools isolated from 12 slaughterhouses 29.4%.

Schmid *et al.* (21) reported that the ESBL producing *E. coli* contamination originating from samples of dairy and beef cattle farms taken from 2011-2012 in Bavaria, Germany was 32,8 %. In another study S. Bina *et al.* (22) obtained a higher yield of the incidence of ESBL producing *E. coli* contamination originating from dairy farms in Germany was 75,6 %. Many of research conducted in Indonesia regarding the level of contamination of ESBL producing bacteria is still focused on contamination which occurs in humans. The incidence of ESBL producing bacterial infections in the hospitals in Surabaya from 2006-2012, namely 34.84% (23), 6.7% (24), and 94.5% (25).

According to CLSI (26) explained that the Vitek-2 compact system tool is a system for identifying and testing semi-automatic resistance for bacteria making it possible to determine MIC rapidly by analyzing the kinetics of bacterial growth with antimicrobials using a card to test. This Vitek-2 method has proven to be very good at detection bacterial resistance to an antibiotic because the results are very accurate (27). In a study conducted by Spanu *et al.* (28) showed that the Vitek-2 method has a sensitivity of 98.1% and a specificity of 99.5%. The compact serining vitek-2 method is used as a reference in detecting ESBL producing *Enterobacteriaceae* and this method from several references and studies is highly recommended to identify ESBL producing bacteria using this method.

Conclusion

The detection of ESBL producing *E. coli* from dairy farms in Tulungagung Regency was 3 (6%) for confirmation using DDST and Vitek-2. These results indicate that ESBL producing *E. coli* can threaten animal and human health which can spread quickly and widely. The results of this study indicate that the high prevalence of ESBL producing *E. coli* on Sendang, Tulungagung Regency, Indonesia.

34

Acknowledgements

This study was supported in part with Hibah Mandat, Fund Airlangga University, Indonesia.

References

1. WHO. Critically important antimicrobials for human medicine. 5th ed. Geneva; WHO Press; 2016. 1-2 p.
2. Paterson DL, Bonomo RA. Extended-spectrum beta-lactamases: A clinical update. *Clin Microbiol Rev.* 2005;18(4):657-686. doi: 10.1128/CMR.18.4.657-686.2005
3. Price LB, Graham JP, Lackey LG, Roess A, Vailes R, Silbergeld E. Elevated risk of carrying gentamicin-resistant *Escherichia coli* among US Poultry workers. *Environ Health Perspect.* 2007;115(12):1738-1742. doi: 10.1289/ehp.10191
4. Guerra B, Fischer J, Helmuth R. An emerging public health problem: acquired aarbaenemase-producing microorganisms are present in food-producing animals, their environment, companion animals and wild birds. *Vet Microbiol.* 2014;171(3-4):290-297. doi: 10.1016/j.vetmic.2014.02.001
5. Hordijk J, Schoormans A, Kwakernaak M, Duim B, Broens E, Dierikx C, Dik M, Jaap AW. High prevalence of fecal carriage of extended spectrum beta-lactamase/AmpC-producing *Enterobacteriaceae* in cats and dogs. *Front Microbiol.* 2013;4 :274. Doi: 10.3389/fmicb.2013.00242
6. Sanders A, Amanda M. Vitek 2 compact-identification and susceptibility testing. USA: Standard Operating Procedure; 2018. 4-24 p.
7. Safitri RD, Cicilia R, Bintari IG, Hermawan IP, Effendi MH, Ernawati R, Rahmahani J. Detection of encoding gene extended spectrum beta lactamase on *Escherichia coli* isolated from broiler chicken meat in traditional market Surabaya. *Inter J Develop Res.* 2017;07(11):17254-17357. ISSN : 2230-9926.
8. Effendi MH, Harjani N, Yanestria SM, Hastutie P. Identification of shiga toxin-producing *Escherichia coli* in raw milk samples from dairy cows in Surabaya, Indonesia. *Philippine J Ved Med.* 2018;55(SI):109-114.
9. Effendi MH, Harjani N, Budiarto, Triningtya NP, Tyasningsih W, Plumeriastusi H. Prevalensi of pathogenic *Escherichia coli* isolated from subclinical mastitis in east Java province, Indonesia. *Indian Vet J.* 2019;96(03):22-25.
10. CSLI. Performances standards for antimicrobial susceptibility testing. 26th ed. New York: Clinical and Laboratory Standards Institute; 2016. 1-34 p.
11. Santos LL, Moura RA, Agilar-Ramires P, Castro AP, Lincopan N. Current status of extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriaceae* in animals. 1st ed. USA: Formatex; 2013. 1600-1607 p.
12. O'Brien TF. Emergence, spread, and environmental effect of antimicrobial resistance: how use of an antimicrobial anywhere can increase resistance to any antimicrobial anywhere else. *Clin Infect Dis.* 2002;3:78-84. doi: 10.1086/340244
13. Ghatak S, Singha A, Sen A, Guha C, Ahuja A, Bhattacharjee U, Das S, Pradhan NR, Puro K, Jana C. Detection of New Delhi metallobeta-lactamase and extended spectrum beta-lactamase genes in *Escherichia coli* isolated from mastitic milk samples. *Transbound Emerg Dis.* 2013;60(5):385-9. doi: 10.1111/tbed.12119
14. Geser N, Stephan R, Kuhnert P, Zbinden R, Kaeppli U, Cernela N, Haechler H. Fecal carriage of extended-spectrum beta-lactamase-producing *Enterobacteriaceae* in swine and cattle at slaughter in Switzerland. *J Food Protect.* 2011;74(3):446-449. doi: 10.4315/0362-028X.JFP-10-372
15. EFSA. Scientific opinion on the public health risks of bacterial strains producing extended-spectrum beta-lactamases and/or AmpC beta-lactamase

- in food and food-producing animals. EFSA J. 2011;9(8):1-95. doi: 10.2903/j.efsa.2011.2322
16. Khosbayan T, Munguntsetseg B, Ochbadrakh B, Udval U, Batbaatar G, Wu J, Yong D. Plasmid analysis of ESBL producing gram negative bacilli in Mongolia. Mongolian J Hea Sci. 2013;10(1):90-100.
 17. Babypedmini S, Appalaraju B. Extended spectrum β -lactamases in urinary isolates of *Escherichia coli* and *Klebsiella pneumoniae* prevalence and susceptibility pattern in a tertiary care hospital. Indian J Med Microbiol. 2004;22(3):172-4.
 18. Parasakthi N, Arifin H, Kamarulzaman A, Ibrahim HSM, Adnan A, Choeng I. Consensus guidelines for the management of infections by ESBL-producing bacteria. Malaysian Soc Infect Dis Chemother. 2001;7:3.
 19. Sukmawinata E. Tingkat kejadian *Escherichia coli* penghasil extended spectrum β -lactamase di feses sapi di rumah potong hewan ruminansia kota Bogor [MSc thesis]. Bogor: Institut Pertanian Bogor; 2015. 1-185 p.
 20. Haenni M, Châtre P, Métayer V, Bour M, Signol E, Madec JY, Gay E. Comparative prevalence and characterization of ESBL producing Enterobacteriaceae in dominant versus subdominant enteric flora in veal calves at slaughterhouse, France. Vet Microbiol. 2014;171:321-327. doi: 10.1016/j.vetmic.2014.02.023
 21. Schmid A, Hörmansdorfer S, Messelhäusser U, Käsbohrer A, Sauter-Louis C, Mansfeld R. Prevalence of extended-spectrum β -lactamase-producing *Escherichia coli* on Bavarian dairy and beef cattle farms. Appl Environ Microbiol. 2013;79(9):3027-3032. doi: 10.1128/AEM.00204-13
 22. Odenthal S, Akinenden O, Usleber E. Extended-Spectrum β -Lactamase producing Enterobacteriaceae in bulk tank milk from German dairy Fam. Inter J Food Microbiol. 2016;238:72-78. doi: 10.1016/j.ijfoodmicro.2016.08.036
 23. Kuntaman, Mertiasih NM, Hadi U. Multiresistance pattern of extended spectrum β -lactamase (ESBL)-*Escherichia coli* and *Klebsiella pneumoniae* strains. Folia Med Indones. 2006;42(1):40-46.
 24. Bramantono PH. The prevalence of extended spectrum betalactamase (ESBL) in third generation cephalosporin usage among sepsis patients in the department of internal medicine RSUD Dr Soetomo Surabaya. Folia Med Indones. 2012;49(4):244-251.
 25. Severin JA, Mertaniasih NM, Kuntaman K, Lestari ES, Purwanto M, Toom NL, Duerink DO, Hadi U, Belkum A, Verburg HA, Goessens WH. Molecular characterization of extended-spectrum β -lactamases in clinical *Escherichia coli* and *Klebsiella pneumoniae* isolates from Surabaya, Indonesia. J Antimicrob Chemother. 2010;65:465-469. doi:10.1093/jac/dkp471
 26. CLSI. Performance standards for antimicrobial susceptibility testing. 24th ed. Pennsylvania: Clinical Laboratory and Standard Institute; 2014. 1-25.
 27. Munoz-Davila MJ, Roig M, Yague G, Blazquez A, Salvador C, Segovia M. Comparative evaluation of Vitek-2 identification and susceptibility testing of urinary tract pathogens directly and isolated from chromogenic media. Eur J Clin Microbiol Infect Dis. 2013;32:773-780. doi: 10.1007/s10096-012-1806-4
 28. Spanu T, Sanguinetti M, Tumbarello M, D'Inzeo T, Fiori B, Posteraro B, Santangelo R, Cauda R, Fadda G. Evaluation of the new VITEK-2 Extended-Spectrum Beta-Lactamase (ESBL) test for rapid detection of ESBL production in Enterobacteriaceae isolates J Clin Microbiol. 2006;44(9):3257-3262. doi: 10.1128/JCM.00433-06

Detection of the extended spectrum β -lactamase produced by Escherichia coli from dairy cows by using the Vitek-2 method in Tulungagung regency, Indonesia

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

16%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1	Seyed Mahdi Hosseini, Ye Tingzhu, Majid Pasandideh, Aixin Liang et al. "Genetic Association of PPARGC1A Gene Single Nucleotide Polymorphism with Milk Production Traits in Italian Mediterranean Buffalo", BioMed Research International, 2021 Publication	1%
2	Submitted to Universitas Airlangga Student Paper	1%
3	www.envirobiotechjournals.com Internet Source	1%
4	discovery.ucl.ac.uk Internet Source	1%
5	www.veterinaryworld.org Internet Source	1%
6	worldwidescience.org Internet Source	1%

geb.uni-giessen.de

7	Internet Source	1 %
8	www.ijpmonline.org Internet Source	1 %
9	ajms.alameenmedical.org Internet Source	<1 %
10	Daiga Gāliņa, Andris Balins, Anda Valdovska. "The Prevalence and Characterization of Fecal Extended-Spectrum-Beta-Lactamase- Producing Escherichia coli Isolated from Pigs on Farms of Different Sizes in Latvia", Antibiotics, 2021 Publication	<1 %
11	Ebru Şebnem Yılmaz, Ayşe Dolar. "Detection of Extended-Spectrum β -Lactamases in Escherichia coli From Cage Birds", Journal of Exotic Pet Medicine, 2017 Publication	<1 %
12	Laube, H., A. Friese, C. von Salviati, B. Guerra, A. Kasbohrer, L. Kreienbrock, and U. Roesler. "LONGITUDINAL MONITORING OF ESBL/AMPC-PRODUCING ESCHERICHIA COLI IN GERMAN BROILER CHICKEN FATTENING FARMS", Applied and Environmental Microbiology, 2013. Publication	<1 %

13

Rocha-Gracia, R.C., G. Cortés-Cortés, P. Lozano-Zarain, F. Bello, Y. Martínez-Laguna, and C. Torres. "Faecal Escherichia coli isolates from healthy dogs harbour CTX-M-15 and CMY-2 β -lactamases", *The Veterinary Journal*, 2015.

Publication

14

text-id.123dok.com

Internet Source

15

"Escherichia coli, a Versatile Pathogen", Springer Science and Business Media LLC, 2018

Publication

16

Geser, N., R. Stephan, P. Kuhnert, R. Zbinden, U. Kaeppli, N. Cernela, and H. Haechler. "Fecal Carriage of Extended-Spectrum β -Lactamase-Producing *Enterobacteriaceae* in Swine and Cattle at Slaughter in Switzerland", *Journal of Food Protection*, 2011.

Publication

17

Neelima Angaali, Laxmi Vemu, Chavali Padmasri, Neeraja Mamidi, Vijay Dharma Teja. "Direct identification and susceptibility testing of Gram-negative bacilli from turbid urine samples using VITEK2", *Journal of Laboratory Physicians*, 2020

Publication

<1 %

<1 %

<1 %

<1 %

<1 %

18	hal.inrae.fr Internet Source	<1 %
19	media.neliti.com Internet Source	<1 %
20	vetdergi.kafkas.edu.tr Internet Source	<1 %
21	www.benthamopen.com Internet Source	<1 %
22	www.tandfonline.com Internet Source	<1 %
23	123dok.com Internet Source	<1 %
24	Marilyn Martinez, Peter Silley. "Chapter 10 Antimicrobial Drug Resistance", Springer Science and Business Media LLC, 2010 Publication	<1 %
25	Maryam Rajaei, Mir-Hassan Moosavy, Sahar Nouri Gharajalar, Seyed Amin Khatibi. "Antibiotic resistance in the pathogenic foodborne bacteria isolated from raw kebab and hamburger: phenotypic and genotypic study", BMC Microbiology, 2021 Publication	<1 %
26	doc.rero.ch Internet Source	<1 %

27	docplayer.net Internet Source	<1 %
28	hdl.handle.net Internet Source	<1 %
29	journals.plos.org Internet Source	<1 %
30	repositorio.utad.pt Internet Source	<1 %
31	www.frontiersin.org Internet Source	<1 %
32	Ágnes Nagy, Rita Székelyhidi, Erika Hanczné Lakatos, Viktória Kapcsándi. "Review on the occurrence of the mcr-1 gene causing colistin resistance in cow's milk and dairy products", <i>Heliyon</i> , 2021 Publication	<1 %
33	Asinamai Athliamai Bitrus, Peter Anjili Mshelia, Iliya Dauda Kwoji, Mohammed Dauda Goni, Saleh Mohammed Jajere. "Extended-spectrum beta-lactamase and ampicillin Class C beta lactamase-producing <i>Escherichia coli</i> from food animals: A review", <i>International Journal of One Health</i> , 2019 Publication	<1 %
34	L. Romero. "Long-term study of the frequency of <i>Escherichia coli</i> and <i>Klebsiella pneumoniae</i>	<1 %

isolates producing extended-spectrum beta-lactamases", *Clinical Microbiology and Infection*, 8/2005

Publication

35

Shinyoung Lee, Raies A. Mir, Si Hong Park, Donghyuk Kim, Hae-Yeong Kim, Raoul K. Boughton, J. Glenn Morris, Kwangcheol C. Jeong. "Prevalence of extended-spectrum β -lactamases in the local farm environment and livestock: challenges to mitigate antimicrobial resistance", *Critical Reviews in Microbiology*, 2020

Publication

36

T. Winstanley, P. Courvalin. "Expert Systems in Clinical Microbiology", *Clinical Microbiology Reviews*, 2011

Publication

37

Ursula Kaspar, Alexa von Lützu, Andreas Schlattmann, Uwe Roesler, Robin Köck, Karsten Becker. "Zoonotic multidrug-resistant microorganisms among small companion animals in Germany", *PLOS ONE*, 2018

Publication

38

Thorsten Schauss, Stefanie P. Glaeser, Alexandra Gütschow, Wolfgang Dott, Peter Kämpfer. "Improved Detection of Extended Spectrum Beta-Lactamase (ESBL)-Producing *Escherichia coli* in Input and Output Samples

<1 %

<1 %

<1 %

<1 %

of German Biogas Plants by a Selective Pre-Enrichment Procedure", PLOS ONE, 2015

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Detection of the extended spectrum β -lactamase produced by Escherichia coli from dairy cows by using the Vitek-2 method in Tulungagung regency, Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
