




Heliyon





Heliyon

Open access

[Latest issue](#)

[All issues](#)

[Submit your article](#) ↗

Search in this journal



Journal info

[Aims and scope](#)

[Editorial board](#)

Lead Editor

Christian Schulz

Assistant Editors

Elizabeth Hancock

Marion Thibaudeau

Senior Editorial Assistant

Chris Russell

Editorial Assistants

Charles Booty

Poppy Craig

Ailish Cronin

Zoe Gooden

On Ching Lo

Harry Mcgee

James Morgan

Dimitri Pantelli

Laura Salisbury

Elizabeth Wetherell

Section Editors

Professor Jinrong Min

University of Toronto, Toronto, Ontario, Canada

Professor Nicola Zambrano

University of Naples Federico II Department of Molecular Medicine and Medical Biotechnology, Naples, Italy

Associate Editors

Dr. Anna-Lena Ackfeldt

Aston University, Birmingham, United Kingdom

Assoc. Professor Gholamreza Amin

University of New Brunswick, Fredericton, New Brunswick, Canada

Dr. Alessio Emanuele Biondo

University of Catania, Catania, Italy

Dr. Joanne Connell

University of Exeter Department of Management Studies, Exeter, United Kingdom

Assoc. Professor Pavlos Delias

Eastern Macedonia and Trace Institute of Technology, Kavala, Greece

Dr. Eugene Ezebilo

Papua New Guinea National Research Institute, Boroko, Papua New Guinea

Dr. Bernard Njindan Iyke

Deakin Graduate School of Business, Burwood, Australia

Dr. Eleonora Pantano

University of Bristol, Bristol, United Kingdom

Dr. Rudra Pradhan

Indian Institute of Technology Kharagpur, Kharagpur, India

Professor Ernesto Santibanez Gonzalez

University of Talca, Department of Industrial Engineering, Maule, Chile

Dr. Serje Schmidt

FEEVALE University, NOVO HAMBURGO, Brazil

Professor Alfonso Sousa-Poza

University of Hohenheim, Stuttgart, Germany

Associate Editors

Assoc. Professor Carlos Javier Alméciga Díaz

Pontifical Javeriana University Institute of Inborn Errors of Metabolism, Bogotá, Colombia

Dr. Timo Kuosmanen

Aalto University, Aalto, Finland

Advisory Board Members- Business & Economic

Ben Marshall

Vesa Puhakka

Martin Falk

S. Mostafa Rasoolimanesh

Gianpaolo Iazzolino

Margaret Raviro Mabugu

Ming-Long Lee

Lincoln C. Wood

Giray Gozgor

Fabio Nonino

Encarnación Cerdón Lagares

Paolo Pin

Ulrich Berger

Section Editors

Dr. Francesco Epifano

Associate Editors

Dr. Anthony Curtis

Keele University, Newcastle, United Kingdom

Dr. Dario Donno

University of Turin, Torino, Italy

Dr. Serena Fiorito

Gabriele d'Annunzio University of Chieti and Pescara, Chieti, Italy

Dr. Ramesh L. Gardas

Indian Institute of Technology Madras Department of Chemistry, Chennai, Tamil Nadu, India

Dr. Salvatore Genovese

Gabriele d'Annunzio University of Chieti and Pescara, Chieti, Italy

Dr. Boris-Marko Kukovec

University of Split, Split, Croatia

Dr. Károly Lázár

MTA Centre for Energy Research

Dr. Gianfranco Pasut

University of Padua, Padova, Italy

Prof. Dr. Yolanda Picó

University of Valencia Faculty of Pharmacy, Burjassot, Spain

Professor Ming Hua Xu

Shanghai Institute of Materia Medica Chinese Academy of Sciences, Shanghai, China

Advisory Board Member - Chemistry

Jussara Miranda

Federal University of Rio de Janeiro, Rio de Janeiro, Brazil

Nitin Chattopadhyay

Jadavpur University, Kolkata, India

Makoto Tsunoda

The University of Tokyo, Bunkyo-Ku, Tokyo, Japan

Anjanapura Venkataramanaiah Raghu

Jain University, Bangalore, India

Praveen Nekkar Rao

University of Waterloo, Waterloo, Ontario, Canada

Jorge Masini

University of Sao Paulo, São Paulo, Brazil

Pellegrino Musto

National Research Council of Italy, Bari, Italy

Silvia Antonia Brandán

National University of Tucuman, San Miguel De Tucuman, Argentina

Wojciech Domagala

Silesian University of Technology, Gliwice, Poland

Agostinho Almeida

University of Porto, Porto, Portugal

K. Miraz Rahman

King's College London, London, United Kingdom

Zhaoyan Sun

Chinese Academy of Sciences, Beijing, China

Hakan Arslan

Mersin University, Mersin, Turkey

Peter Kusch

University of Applied Sciences Bonn-Rhein-Sieg Department of Natural Sciences, Rheinbach, Germany

Maria Grazia Francesconi

University of Hull, Hull, United Kingdom

Luiz Fernando Cappa De Oliveira

Federal University of Juiz de Fora, JUIZ DE FORA, Brazil

Andrei Kovalevsky

University of Aveiro, Aveiro, Portugal

Section Editors

Dr. Gary Adams

University of Nottingham, Nottingham, United Kingdom

Dr. Carolyn Mackintosh-Franklin

The University of Manchester, Manchester, United Kingdom

Associate Editors

Dr. Seraphim Alvanides

GESIS Institute of Social Sciences Cologne Branch, Koln, Germany

Professor Barbara Canlon

Karolinska Institute, Stockholm, Sweden

Dr. Manoj Das

The INCLEN Trust International, New Delhi, India

Dr. Richard Gillis

University of Nottingham, Nottingham, United Kingdom

Dr. Philemon Gyasi-Antwi

University of Nottingham, Nottingham, United Kingdom

Professor Kenji Hashimoto

Chiba University, Chiba, Japan

Dr. Marina Ivanisevic

ProNatal Polyclinic, Zagreb, Croatia

Dr. Shahwar Imran Jiwani

University of Nottingham, Nottingham, United Kingdom

Dr. Devika Kapuria

University of New Mexico, Albuquerque, New Mexico, United States

Dr. Avril Mansfield

University Health Network, Toronto, Ontario, Canada

Dr. Giuseppe Musumeci

University of Catania, Catania, Italy

Professor Graham Pawelec

Eberhard Karls University Tübingen, Tübingen, Germany

Dr. Salvatore Pepe

Murdoch Childrens Research Institute, Parkville, Australia

Dr. Ferdinand Schweser

University at Buffalo - The State University of New York, Buffalo, New York, United States

Dr. Ekaterina Silina

I M Sechenov First Moscow State Medical University, Moskva, Russian Federation

Dr. Sushma Sonale

Thai Moogambigai Dental College and Hospital, Chennai, India

Assist. Prof. Georgios Tsoulfas

Aristotle University of Thessaloniki, Thessaloniki, Greece

Dr. Corinna Walsh

University of the Free State, Bloemfontein, South Africa

Section Editors

Dr. Jonathan Chan

King Mongkut's University of Technology Thonburi, Bangkok, Thailand

Associate Editors

Dr. Gong Cheng

Northwestern Polytechnical University, Xian, China

Dr. Joana Dias

University of Coimbra, Coimbra, Portugal

Assist. Prof. Shang Gao

Swedish Business School at Orebro University, Örebro, Sweden

Dr. Byung-Gyu Kim

Sookmyung Women's University, Yongsan-gu, Korea, Republic of

Dr. Daniel Lemire

University of Quebec in Montreal, Montreal, Quebec, Canada

Dr. Eugenia Mee Wah Ng

Lingnan University, New Territories, Hong Kong

Dr. Tuan Pham

Linköping University, Linköping, Sweden

Section Editors

Dr. Socrates Kaplanis

Associate Editors

Dr. Stefano Cordiner

University of Rome Tor Vergata, Roma, Italy

Professor Michele Dassisti

Polytechnic University of Bari, Bari, Italy

Dr. Eleni Kaplani

University of East Anglia, Norwich, United Kingdom

Dr. António Cardoso Marques

University of Beira Interior, Covilha, Portugal

Dr. Tingzhen Ming

Wuhan University of Technology, Wuhan, China

Assoc. Professor Yongheng Yang

Aalborg University, Aalborg, Denmark

Engineering

Dr. Andrea F. Morabito

University of Reggio Calabria, Reggio Calabria, Italy

Dr. Mohammad Mehdi Rashidi

Bu Ali Sina University, Hamedan, Iran, Islamic Republic of

Associate Editors

Professor Serge Abrate

Southern Illinois University Carbondale, Carbondale, Illinois, United States

Dr. Aires Camões

University of Minho Department of Civil Engineering, Guimaraes, Portugal

Dr. Pablo Gomez

Western Michigan University, Kalamazoo, Michigan, United States

Dr. Mark Jermy

University of Canterbury, Christchurch, New Zealand

Dr. Xian Fang Li

Central South University School of Civil Engineering, Changsha, China

Susana Lopez-Querol

University College London, London, United Kingdom

Dr. Oronzio Manca

University of Campania Luigi Vanvitelli Department of Industrial Engineering and Information Technology, Aversa, Italy

Professor Yogendra K. Mishra

University of Southern Denmark, Mads Clausen Institute, NanoSYD, Sønderborg, Denmark

Dr. Togay Ozbakkaloglu

Texas State University, San Marcos, Texas, United States

Prof. Dr. Fatih Sen

Kütahya Dumlupınar University Department of Biochemistry, Kütahya, Turkey

Professor Bart Van der Bruggen

KU Leuven Science and Technology Group Department of Chemical Engineering, Heverlee, Belgium

Section Editors

Professor Frederic Coulon

Cranfield University, Cranfield, Bedford, United Kingdom

Professor Christian Sonne

Aarhus University Department of Environmental Science, Roskilde, Denmark

Associate Editors

Dr. Julian Blasco

Institute of Marine Science of Andalusia, Puerto Real, Spain

Dr. Jianmin Chen

Fudan University Department of Environmental Science and Engineering, Shanghai, China

Professor Hefa Cheng

Peking University, Beijing, China

Assoc. Professor Pavlos Delias

Eastern Macedonia and Trace Institute of Technology, Kavala, Greece

Professor Xinbin Feng

Institute of Geochemistry Chinese Academy of Sciences, Guiyang, China

Professor Ashantha Goonetilleke

Queensland University of Technology, Brisbane, Queensland, Australia

Dr. Jun Huang

Tsinghua University, Beijing, China

Dr. Olga-Ioanna Kalantzi

University of the Aegean Department of Environment, Mytilini, Greece

Professor Tsair-Fuh Lin

National Cheng Kung University, Tainan, Taiwan

Dr. Keith Maruya

Southern California Coastal Water Research Project, Costa Mesa, California, United States

Professor Huu Hao Ngo

University of Technology Sydney School of Civil and Environmental Engineering, Broadway, Australia

Professor Yong Sik Ok

Korea University Division of Environmental Science and Ecological Engineering, Seoul, Korea, Republic of

Dr. Patryk Oleszczuk

Maria Curie-Sklodowska University, Lublin, Poland

Professor Wei Ouyang

Beijing Normal University, Beijing, China

Thomas Panagopoulos

University of the Algarve Faculty of Science and Technology, Faro, Portugal

Prof. Dr. Paulo Alexandre da Silva Pereira

Mykolas Romeris University, Institute of Public Administration, Vilnius, Lithuania

Prof. Dr. Yolanda Picó

University of Valencia Faculty of Pharmacy, Burjassot, Spain

Prof. Dr. Jörg Rinklebe

University of Wuppertal, Wuppertal, Germany

Dr. Admir Créso Targino

Federal Technological University of Parana Department of Physics, Londrina, Brazil

Assoc. Professor Daniel Tsang

The Hong Kong Polytechnic University Department of Civil and Environmental Engineering, Hong Kong, Hong Kong

Dr. Daniel A. Wunderlin

National University of Cordoba, Cordoba, Argentina

Assoc. Professor Yeomin Yoon

University of South Carolina Department of Civil & Environmental Engineering, Columbia, South Carolina, United States

Professor Shuzhen Zhang

Chinese Academy of Sciences, Beijing, China

Section Editors

Dr. Lilian Mariutti

State University of Campinas Faculty of Food Engineering, Campinas, São Paulo, Brazil

Associate Editors

Dr. Cinthia Betim

State University of Campinas, Campinas, São Paulo, Brazil

Professor Nour-Eddine Es-Safi

Mohammed V University of Rabat, Rabat, Morocco

Dr. Ana Luisa Fernando

New University of Lisbon Department of Engineering and Environmental Sciences, Caparica, Portugal

Assist. Prof. Efstathios (Stathis) Giaouris

University of the Aegean, Food Science and Nutrition, Lemnos, Greece

Dr. Angela Meireles

State University of Campinas Faculty of Food Engineering, Campinas, São Paulo, Brazil

Dr. Donatella Restuccia

University of Calabria, Cosenza, Italy

Dr. Apollinaire Tsopmo

Carleton University, Ottawa, Ontario, Canada

Section Editors

Professor Luis M. Gandía

Associate Editors

Professor Saibal Basu

Homi Bhabha National Institute, Mumbai, India

Professor Yingchao Dong

Dalian University of Technology School of Environmental Science and Technology, Dalian, China

Professor Pierre Dumont

Computer Science Laboratory for Image Processing and Information Systems, Villeurbanne, France

Dr. Fabien Grasset

Université de Rennes I, UMR CNRS 6226, Rennes cedex, France

Assoc. Professor Changdong Gu

Zhejiang University School of Materials Science and Engineering, Hangzhou, China

Dr. Aloysius F. Hepp

NASA John H Glenn Research Center, Cleveland, Ohio, United States

Assoc. Professor Rui Miranda Guedes

University of Porto, Porto, Portugal

Dr. Alessandro Polini

Institute of Nanotechnology, Monteroni di Lecce, Italy

Professor Rafael José Rodríguez Trías

University of Navarra, Navarra, Spain

Dr. Oleksandr Stroyuk

Research Centre Jülich, Jülich, Germany

Section Editors

Dr. Dragana (Dana) Stanley

Central Queensland University, Rockhampton, Australia

Associate Editors

Professor Cristobal Aguilar

Autonomous University of Coahuila, Saltillo, Mexico

Dr. Uffe Christian Braae

State Serum Institute, København, Denmark

Professor Marina Eremeeva

Georgia Southern University Jiann-Ping Hsu College of Public Health, Statesboro, Georgia, United States

Dr. Volker Gurtler

RMIT University School of Applied Sciences- Bundoora Campus, Bundoora, Victoria, Australia

Assoc. Professor Florian Krammer

Icahn School of Medicine at Mount Sinai, New York, New York, United States

Dr. Xiaobo Liu

University of Hong Kong, Pokfulam, Hong Kong

Dr. Alexander Pasternak

University of Amsterdam, Amsterdam, Netherlands

Section Editors

Professor Jacob Raber

Oregon Health & Science University, Portland, Oregon, United States

Associate Editors

Dr. Ilan Dar-Nimrod

The University of Sydney, Sydney, New South Wales, Australia

Dr. Charlotte D'Mello

University of Calgary, Calgary, Alberta, Canada

Dr. Kioko Guzmán-Ramos

Metropolitan Autonomous University Lerma Campus, Lerma de Villada, Mexico State, Mexico

Dr. Andreas Löw

Helmut-Schmidt-University / University of the Armed Forces Hamburg, Hamburg, Germany

Dr. Julian Macoveanu

Copenhagen University Hospital, København, Denmark

Dr. Michael Noll-Hussong

Saarland University Hospital Department of Psychiatry and Psychotherapy, Homburg, Germany

Dr. Janice Pluth

University of Nevada Las Vegas, Las Vegas, Nevada, United States

Dr. Ladislao Salmerón

University of Valencia, Valencia, Spain

Dr. Paul Sauseng

Ludwig Maximilians University Munich, Munich, Germany

Dr. Stefan Sütterlin

Oslo University Hospital, Oslo, Norway

Dr. Mario Tiberi

Ottawa Hospital Research Institute, Ottawa, Ontario, Canada

Dr. Mitchell Turker

Oregon Health & Science University, Portland, Oregon, United States

Section Editors

Professor Emilio Clementi

University of Milan, Milano, Italy

Dr. Martin Leonard

Toxicology Department, Centre for Radiation, Chemical and Environmental Hazards, Public Health England, Oxfordshire, United Kingdom

Associate Editors

Dr. Jorge Bernardino de la Serna

Imperial College London Faculty of Medicine, London, United Kingdom

Dr. Nicolas Clere

University of Angers, Angers, France

Dr. Maxime Culot

Artois University Blood Brain Barrier Laboratory, Lens, France

Dr. Dimitrios Lamprou

Queen's University Belfast, Belfast, United Kingdom

Section Editors

Dr. Gerald Cleaver

Baylor University, Waco, Texas, United States

Associate Editors

Dr. Sebastian Brauchi

University of Southern Chile, Valdivia, Chile

Dr. Saurya Das

University of Lethbridge, Lethbridge, Alberta, Canada

Dr. Jordi Faraudo

Institute of Materials Science of Barcelona, Barcelona, Spain

Dr. Suhairul Hashim

Universiti Teknologi Malaysia, Johor, Malaysia

Dr. Gunar Schnell

University of the Basque Country, Bilbao, Spain

Dr. Jun-Jun Xiao

Harbin University of Science and Technology, Haerbin, China

Associate Editor

Dr. Mercedes Regadío

The University of Sheffield, Sheffield, United Kingdom

Professor Sergio Llana Fúnez

University of Oviedo, Oviedo, Spain

Section Editor

Dr. Andrea De Martino

University of Rome La Sapienza, Roma, Italy

ISSN: 2405-8440

Copyright © 2020 Elsevier Ltd.



[About ScienceDirect](#)

[Remote access](#)

[Shopping cart](#)

[Advertise](#)

[Contact and support](#)

[Terms and conditions](#)

[Privacy policy](#)

We use cookies to help provide and enhance our service and tailor content and ads. By continuing you agree to the **use of cookies**.

Copyright © 2020 Elsevier B.V. or its licensors or contributors. ScienceDirect® is a registered trademark of Elsevier B.V.

ScienceDirect® is a registered trademark of Elsevier B.V.





Open access

[Latest issue](#) [All issues](#) [Submit your article ↗](#)

[Search in this journal](#)

Volume 5, Issue 9

September 2019

[← Previous vol/issue](#)

[Next vol/issue >](#)

Receive an update when the latest issues in this journal are published

[Sign in to set up alerts](#)

Research article *Open access*

S(-) and R(+) species derived from antihistaminic promethazine agent: structural and vibrational studies

María Eugenia Manzur, Silvia Antonia Brandán

Article e02322

[↓ Download PDF](#) [Article preview](#) 

Research article *Open access*

Minimally-Invasive midline posterior interbody fusion with cortical bone trajectory screws compares favorably to traditional open transforaminal interbody fusion

Charles H. Crawford, Roger K. Owens, Mladen Djurasovic, Jeffrey L. Gum, ... Leah Y. Carreon

Article e02423

[Download PDF](#) Article preview 

Research article *Open access*

Screening of bacteriocin associated genes of *Streptococcus uberis* strains

Mirta Lasagno, María de los Angeles Navarro, Melina Moliva, Elina Reinoso

Article e02393

[Download PDF](#) Article preview 

Research article *Open access*

Concerns about the pathology job market dominate a popular online pathology forum and likely deterred American medical students from pursuing pathology residency

Ryan Philip Jajosky, Mara Banks, Audrey N. Jajosky

Article e02400

[Download PDF](#) Article preview 

Research article *Open access*

Silver/quartz nanocomposite as an adsorbent for removal of mercury (II) ions from aqueous solutions

Rasha S. El-Tawil, Shaimaa T. El-Wakeel, Ashraf E. Abdel-Ghany, Hanaa A.M. Abuzeid, ... Ahmed M. Hashem

Article e02415

[Download PDF](#) Article preview 

Research article *Open access*

Motivation and E-Learning English as a foreign language: A qualitative study

Fredy Geovanni Escobar Fandiño, Luz Dary Muñoz, Angela Juliette Silva Velandia

Article e02394

[Download PDF](#) Article preview 

Research article *Open access*

The transition state conformational effect on the activation energy of ethyl acetate neutral hydrolysis

Febdian Rusydi, Nufida D. Aisyah, Rizka N. Fadilla, Hermawan K. Dipojono, ... Andriwo Rusydi
Article e02409


[Download PDF](#) Article preview 

Research article *Open access*

The impact of culture and product on the subjective importance of user experience aspects

Harry B. Santoso, Martin Schrepp

Article e02434


[Download PDF](#) Article preview 

Research article *Open access*

Sensor-based occupancy detection using neutrosophic features fusion

N.S. Fayed, Mervat Abu-Elkheir, E.M. El-Daydamony, A. Atwan

Article e02450


[Download PDF](#) Article preview 

Research article *Open access*

Contrast enhancement of surface layers with fast middle-infrared scanning

Tim Kümmel, Tobias Teumer, Patrick Dörnhofer, Frank-Jürgen Methner, ... Matthias Rädle

Article e02442

[Download PDF](#) Article preview 

Research article *Open access*

Structural, electronic and magnetic properties of M_xPt_{1-x} , (M= Co, Ni and V) binary alloys

A.M. Alsaad, A.A. Ahmad, Hamzah A. Qattous

Article e02433

[Download PDF](#) Article preview 

Research article *Open access*

Peptidylglycine monooxygenase activity of monomeric species of growth hormone

John Donlon, Patrick Ryan

Article e02436

[Download PDF](#) Article preview 

Research article *Open access*

Study of novel triazolo-benzodiazepine analogues as antidepressants targeting by molecular docking and ADMET properties prediction

Assia Belhassan, Hanane Zaki, Mohamed Benlyas, Tahar Lakhli, Mohammed Bouachrine

Article e02446

[Download PDF](#) [Article preview](#) 

Research article *Open access*

Pyroprocessing and the optimum mix ratio of rice husks, broken bricks and spent bleaching earth to make pozzolanic cement

Protus Nalobile, Jackson Muthengia Wachira, Joseph Karanja Thiong'o, Joseph Mwiti Marangu

Article e02443

[Download PDF](#) [Article preview](#) 

Research article *Open access*

Supersymmetric approach to coherent states for nonlinear oscillator with spatially dependent effective mass

M. Tchoffo, F.B. Migueu, M. Vubangsi, L.C. Fai

Article e02395

[Download PDF](#) [Article preview](#) 

Research article *Open access*

Investigation effects of extracted compounds from shell and cluster of pistachio nut on the inactivation of free radicals

Morteza Mohammadi, Mohammad Ghorbani, Adel Beigbabaei, Samira Yeganehzad, Alireza Sadeghi-Mahoonak

Article e02438

[Download PDF](#) [Article preview](#) 

Research article *Open access*

Unsteady computational fluid dynamics analysis of the hydrodynamic instabilities in a reversible Francis turbine used in a storage plant

S. Mauro, R. Lanzafame, S. Brusca, M. Messina

Article e02441

[Download PDF](#) [Article preview](#) 

Research article *Open access*

Preparation and evaluation of ZnO nanoparticles by thermal decomposition of MOF-5

Shirin Hajiashrafi, Negar Motakef Kazemi

Article e02152

[Download PDF](#) [Article preview](#) 

Research article *Open access*

Polypropylene and tire powder composite for use in automotive industry

Kelly C. de Lira Lixandrão, Fabio F. Ferreira

Article e02405

[Download PDF](#) [Article preview](#) 

Research article *Open access*

Effect of Mo on the catalytic activity of Ni-based self-organizing catalysts for processing of dichloroethane into segmented carbon nanomaterials

Yury I. Bauman, Yulia V. Rudneva, Ilya V. Mishakov, Pavel E. Plyusnin, ... Roman A. Buyanov

Article e02428


[Download PDF](#) [Article preview](#) 

Research article *Open access*

Coverage, quality, and correlates of childhood immunization in slums under national immunization program of India: A cross-sectional study

Sanjeev Singh, Damodar Sahu, Ashish Agrawal, Lakshmanan Jeyaseelan, ... Meeta Dhaval Vashi

Article e02403

[Download PDF](#) [Article preview](#) 

Review article *Open access*

A systematic review of behavioral health interventions for suicidal and self-harming individuals in prisons and jails

Natalie Winicov

Article e02379

[Download PDF](#) [Article preview](#) 

Research article *Open access*

Numerical integration strategies of PFR dynamic models with axial dispersion and variable superficial velocity: the case of CO₂ capture by a solid sorbent

A. Di Giuliano, E. Pellegrino

Article e02040


[Download PDF](#) Article preview 

Review article *Open access*

A meta-analysis of suprapatellar versus infrapatellar intramedullary nailing for the treatment of tibial shaft fractures

Heng Xu, Fushun Gu, Jindang Xin, Chengguang Tian, Fei Chen

Article e02199

[Download PDF](#) Article preview 

Research article *Open access*

Semi-synthesis of β -keto-1,2,3-triazole derivatives from ethinylestradiol and evaluation of the cytotoxic activity

Thayane M. Queiroz, Erika V.M. Orozco, Valdenizia R. Silva, Luciano S. Santos, ... André L.M. Porto

Article e02408

[Download PDF](#) Article preview 

Erratum *Open access*

Corrigendum to “CFC delta-beta is related with mixed features and response to treatment in bipolar II depression” [Heliyon 5 (6) (June 2019) e01898]

Sermin Kesebir, Rüştü Murat Demirer, Nevzat Tarhan

Article e02286

[Download PDF](#)

Research article *Open access*

Predicting the degradation potential of Acid blue 113 by different oxidants using quantum chemical analysis

Anam Asghar, Mustapha Mohammed Bello, Abdul Aziz Abdul Raman, Wan Mohd Ashri Wan Daud, ... Sharifuddin Bin Md Zain

Article e02396

[Download PDF](#) Article preview 

Research article *Open access*

Opsonic monoclonal antibodies enhance phagocytic killing activity and clearance of *Mycobacterium tuberculosis* from blood in a quantitative qPCR mouse model

Clara J. Sei, Bong-Akee Shey, Richard F. Schuman, Nimisha Rikhi, ... Gerald W. Fischer
Article e02260


[Download PDF](#) Article preview 

Research article Open access

Biodiversity, *Leishmania* genetic typing and host identification of phlebotomine species in endemic foci of southeastern Iran

Ismail Amiri Ghannat Saman, Mohammad Saaid Dayer, Majid Pirestani

Article e02369


[Download PDF](#) Article preview 

Research article Open access

Interlayer structure and magnetic field-induced orientation of modified nanoclays in polymer aqueous solution

Min Kwan Kang, Eun Jung Cha, Hyun Hoon Song, Yang Ho Na

Article e02421


[Download PDF](#) Article preview 

Research article Open access

Paclitaxel-loaded sodium deoxycholate-stabilized zein nanoparticles: characterization and *in vitro* cytotoxicity

Agnese Gagliardi, Sonia Bonacci, Donatella Paolino, Christian Celia, ... Donato Cosco

Article e02422

[Download PDF](#) Article preview 

Research article Open access

Disruptions in the female reproductive system on consumption of calcium carbide ripened fruit in mouse models

Enitome E. Bafor, Emmanuella Greg-Egor, Osemelomen Omoruyi, Ejiroghene Ochoyama, Glory U. Omogiade

Article e02397

[Download PDF](#) Article preview 

Review article Open access

The phytochemical and pharmacological actions of *Entada africana* Guill. & Perr.

A.J. Yusuf, M.I. Abdullahi

Article e02332

[Download PDF](#) Article preview 

Review article *Open access*

Optimization of quercetin extraction method in *Dendrobium officinale* by response surface methodology

Yingpeng Zhu, Jiangliu Yu, Chunyan Jiao, Jinfeng Tong, ... Yongping Cai

Article e02374

[Download PDF](#) Article preview 

Research article *Open access*

Cyclic β -(1, 2)-glucan blended poly DL lactic co glycolic acid (PLGA 10:90) nanoparticles for drug delivery

Geetha Venkatachalam, Nandakumar Venkatesan, Ganesan Suresh, Mukesh Doble

Article e02289

[Download PDF](#) Article preview 

Research article *Open access*

Assessment of fluoride intake from groundwater and intake reduction from delivering bottled water in Chiang Mai Province, Thailand

Benyapa Sawangjang, Takashi Hashimoto, Aunnop Wongrueng, Suraphong Wattanachira, Satoshi Takizawa

Article e02391

[Download PDF](#) Article preview 

Research article *Open access*

Effect of gabapentin on fetal rat brain and its amelioration by ginger

Gamal M. Badawy, Marwa N. Atallah, Saber A. Sakr

Article e02387

[Download PDF](#) Article preview 

Research article *Open access*

Low weight, socioeconomics and behavioral issues: examining a population in the Northeast of Brazil

Maria de Jesus Torres Pacheco, Fernanda T. Orsati, Paulo Guirro Laurence, Héron Máximo da Cunha Gonçalves, ... Elizeu Coutinho Macedo

Article e02399


[Download PDF](#) Article preview 

Research article *Open access*

Comparison between gelatin/carboxymethyl cellulose and gelatin/carboxymethyl nanocellulose in tramadol drug loaded capsule

Ghada Kadry

Article e02404


[Download PDF](#) Article preview 

Research article *Open access*

Microglia are both a source and target of extracellular cyclophilin A

Gurkiran Kaur Flora, Ryan S. Anderton, Bruno P. Meloni, Gilles J. Guillemin, ... Sherif Boulos

Article e02390


[Download PDF](#) Article preview 

Research article *Open access*

Adsorption of heavy metal ions (Cu^{2+} , Ni^{2+} , Co^{2+} and Fe^{2+}) from aqueous solutions by natural zeolite

T.P. Belova

Article e02320

[Download PDF](#) Article preview 

Research article *Open access*

Oedometer based estimation of vertical shrinkage of expansive soil in a large instrumented soil column

Rajitha Shehan Udukumburage, Chaminda Gallage, Les Dawes

Article e02380

[Download PDF](#) Article preview 

Research article *Open access*

Hemp to limit diffusion of difenoconazole in vegetable garden soils

Clothilde Léchenault-Bergerot, Nadia Morin-Crini, Steffi Rocchi, Eric Lichtfouse, ... Grégorio Crini

Article e02392

[Download PDF](#) Article preview 

Research article *Open access*

Dynamic ecological system analysis: A holistic analysis of compartmental systems

Huseyin Coskun

Article e02347


[Download PDF](#) [Article preview](#) 

Research article *Open access*

Multi-objective future rule curves using conditional tabu search algorithm and conditional genetic algorithm for reservoir operation

Teerawat Thongwan, Anongrit Kangrang, Haris Prasanchum

Article e02401

[Download PDF](#) [Article preview](#) 

[< Previous vol/issue](#)

[Next vol/issue >](#)

ISSN: 2405-8440

Copyright © 2020 Elsevier Ltd.



[About ScienceDirect](#)

[Remote access](#)

[Shopping cart](#)

[Advertise](#)

[Contact and support](#)

[Terms and conditions](#)

[Privacy policy](#)





The transition state conformational effect on the activation energy of ethyl acetate neutral hydrolysis

Febdian Rusydi ^{a,b,c,*}, Nufida D. Aisyah ^{b,d}, Rizka N. Fadilla ^{b,d}, Hermawan K. Dipojono ^d, Faozan Ahmad ^e, Mudasir ^f, Ira Puspitasari ^{b,g}, Andriwo Rusydi ^h

^a Department of Physics, Faculty of Science and Technology, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, Indonesia

^b Research Center for Quantum Engineering Design, Faculty of Science and Technology, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, Indonesia

^c Visiting Researcher at Precision Sciences & Technology and Applied Physics, Graduate School of Engineering, Osaka University, Suita 565-0871, Japan

^d Department of Engineering Physics, Faculty of Industrial Engineering, Institut Teknologi Bandung, Bandung 40132, Indonesia

^e Department of Physics, Faculty of Mathematics and Science, Institut Pertanian Bogor, Bogor 16680, Indonesia

^f Department of Chemistry, Faculty of Mathematics and Science, Universitas Gadjah Mada, Yogyakarta 55281, Indonesia

^g Information System Study Program, Faculty of Science and Technology, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, Indonesia

^h Department of Physics, Faculty of Science, National University of Singapore, Singapore 117542, Singapore

ARTICLE INFO

Keywords:

Organic chemistry
Physical chemistry
Theoretical chemistry
Long-range correction
First-principles calculations
Neutral hydrolysis
Ester
Density functional theory
Ethyl acetate
Activation energy
Conformational effect

ABSTRACT

We report a first-principles study on ethyl acetate neutral hydrolysis in which we focus on the activation energy variation resulting from the conformational effect in the transition state. We use the conformers of ethyl formate, ethyl acetate, ethyl fluoroacetate, and ethyl chloroacetate as the ester models and one water molecule with a one-step reaction mechanism. We also consider the long-range interaction and the surrounding water in the form of PCM. Our results show that the various conformers yield a significant range of activation energy. Moreover, the gauche conformer has lower activation energy than the trans conformer. The activation energy in its own right is lowered by the halogen atoms. Finally, we remark that the long-range correction and PCM stabilize the transition state geometry but raise the activation energy.

1. Introduction

It is commonly known that neutral hydrolysis of an ester is generally slow. For instance, the rate constants (in 1/s) of methyl acetate and ethyl acetate are an order of 10^{-9} [1] and 10^{-10} [2] respectively. An ester ($R-CO_2-R'$) with a long chain also has low rate constant such as achetylcholine [$R=CH_3$, $R'=C_2H_4N^+(CH_3)_3$] with a rate constant of an order of 10^{-9} [3]. The rate constant is higher when halogenated substitution is introduced to the ester. For instance, the alkyl- and aryl-substituted trifluoroacetates (with $R'=C_nH_{2n+1}$, $n=3, 4$, and 5) has an order of 10^{-6} [4], chloromethyl chloroacetate has an order of 10^{-5} [5], while ethyl trifluorothiolacetate ($CF_3COSC_2H_5$) has a rate constant with an order of 10^{-3} [6].

Even though it is slow, the neutral hydrolysis is part of the measured rate constant. The measured rate constant, k_m , is


$$k_m = k_0 + k_a + k_b$$

where the first term is the rate constant of neutral hydrolysis and the last two terms are the rate constant of the acid- and base-induced hydrolysis, respectively. Since k_0 is experimentally difficult to obtain, it is common to apply some extrapolation technique [7]. Following an argument presented by Wolfenden and Yuan [3], neutral hydrolysis of an ester might emerge as the predominant factor at an elevated temperature. If it is the case, neutral hydrolysis cannot be ignored and becomes important, especially when one is interested to study biochemistry based on the achetylcholine hydrolysis.

However, the mechanism of neutral hydrolysis of an ester is not fully understood [8] and its theoretical study is very limited [9]. One way to respond to this challenge is by using a computational approach where a reaction is modeled and studied under first-principles calculations. A common modeling approach is to involve water molecules that act as

* Corresponding author at: Department of Physics, Faculty of Science and Technology, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, Indonesia.

E-mail address: rusydi@fst.unair.ac.id (F. Rusydi).



Note:

- (i) Isolated reactant is state **1a**; Transition State (TS) is state **1b**; isolated product is state **1c**.
 (ii) Abbreviation Et, W, and Ca are used as molecule names throughout the manuscript.

Scheme 1. The one-step reaction model in this study.

Table 1

Selected ester models based on Scheme 1. The R'-group is always C₂H₅.

R	Ester (Label)	Reaction product
H	Ethyl formate (Et1)	Ca1 + Ethanol
CH ₃	Ethyl acetate (Et2)	Ca2 + Ethanol
CH ₂ F	Ethyl fluoroacetate (Et3)	Ca3 + Ethanol
CH ₂ Cl	Ethyl chloroacetate (Et4)	Ca4 + Ethanol

a bifunctional acid-base catalyst [10, 11, 12, 13], to model the aqueous as a reaction field [14, 15], or to combine both methods [8, 16].

The study of neutral hydrolysis of an ester becomes interesting. It can be a model for a computational study of the cholinergic hypothesis. The hypothesis involves the depletion of acetylcholine molecules that may lead to Alzheimer's disease [17, 18]. Acetylcholine belongs to the ester family and has conformers [19, 20, 21, 22, 23]. The conformation affects its stability and activity [24, 25, 26, 27].

In this article, we report our finding on the conformational effect of the ester neutral hydrolysis. All aforementioned theoretical studies showed the importance of a transition state, but none has explored the different arrangement of atoms in the transition states. It is because most theoretical studies so far concern on the methyl acetate family as the ester models, hence no conformer in the transition state. A conformer is likely to exist in an ester with an R or R' group that has two or more C atoms, such as ethyl acetate. There are two previous works on ethyl acetate (the trans conformer) neutral hydrolysis, which are reported by Schmeer and Sturm [12] and Yamabe et al. [14]. The former used two H₂O molecules, one of which served as the nucleophilic reagent and the other as the acid-base catalyst, and was calculated with Hartree-Fock method. The later used one H₂O molecule and considered Onsager's model as the water surrounding and was calculated with density functional theory method. However, no conformational effect has been reported. To see the variation of conformational effect, we study conformers of four molecule from the ethyl acetate family: ethyl formate, ethyl acetate, ethyl fluoroacetate, and ethyl chloroacetate as the ester models. We also consider the long-range interaction and surrounding water as solvent effect.

2. Methods

Reaction model. We carry out the one-step mechanism to model the neutral hydrolysis as shown in Scheme 1. We use four ester models as listed in Table 1 and notations in Table 2.

The one-step mechanism is argued by Shi et al. [9] as favored if tetrahedral intermediates have a short lifetime and do not interconvert prior to breakdown. The one-step mechanism ensures one saddle point between an initial state (**1a**) and a final state (**1c**) since our concern is on the conformational effect at the saddle point (transition state, **1b**). As da Silva et al. [8] reported, the usual method for searching state **1b** may not work for this kind of study; therefore, we take a great care in ensuring that the saddle point is computationally correct.

DFT calculations. We perform the ground state calculation routines on the basis of density functional theory (DFT) [28, 29] and 6-311++G(d,p) basis set that are integrated into the Gaussian 09 software

Table 2

List of notations used throughout the manuscript.

Notation	Meaning
XC1g	B3LYP (gas phase)
XC2g	CAM-B3LYP (gas phase)
XC1s	B3LYP coupled with PCM (solvent)
XC2s	CAM-B3LYP coupled with PCM (solvent)
R(X,Y)	Bond length between X and Y atom (in Å)
A(X, Y, Z)	∠XYZ (in degree)
Expr.	Experimental value in gas phase
err. _{n0}	(XC _n g - Expr.)/Expr. (in %)
err. _{msg}	(XC _n s - XC _n g)/XC _n g (in %), n = 1, 2
err. ₂₁	(XC ₂ g - XC ₁ g)/XC ₁ g (in %)
Δ ₂₁	XC ₂ g - XC ₁ g
Δ _{sg}	XC _n s - XC _n g, n = 1, 2
range	maximum - minimum value
rat. ₂₁	XC ₂ g/XC ₁ g
rat. _{msg}	XC _n s/XC _n g, n = 1, 2
rat. _k	ratio of maximum and minimum k(T)

[30]. For the exchange-correlation functional, we employ B3LYP [31] and CAM-B3LYP [32] which have been integrated into the software as well. The former has provided good prediction in our previous studies on molecular orbital interaction problems [33, 34, 35, 36], excitation in solvent problems [37, 38], and reaction path coordinate to calculate the tunneling probability [39, 40, 41]. Meanwhile, the latter is needed to consider the correction of the long-range interaction in states **1a**, **1b**, and **1c**. We consider the surrounding water as a perturbation to the systems of interest such that we can couple the DFT calculation with the polarized continuum model (PCM) calculation [42], which is also integrated into the software. We use notations in Table 2 to simplify the writing of the exchange-correlation functionals and PCM coupling calculations.

The Transition State Searching. We set up four steps, with the first three aiming to determine the correct geometry of [Et-W] complex in the transition state (state **1b** in Scheme 1). Step one is to determine the electrophilic site where the water molecule may attack. We analyze the charge population by using NBO program [43] which is integrated into the Gaussian 09 software. Step two is to calculate the potential energy surface (PES) in the plane where the electrophilic site lies (see Fig. 1a). This is done by using partial optimization calculations where the distance of oxygen atom of water and the electrophilic site is fixed. The PES result provides a guide to construct the geometry of TS which becomes the input for step three: to perform TS optimization to predict the geometry of **1b**, which is then followed by the intrinsic reaction coordinate (IRC) calculations. The correct TS geometries require one imaginary frequency to indicate that they are in the saddle point. Step four is to optimize the geometries at two minima predicted by IRC calculations to confirm the molecular geometry in state **1a** and **1c**.

We perform the first three steps with XC1g. Once the geometry of transition state is determined (see Fig. 1b), we repeat step 3 and 4 with XC2g, XC1s, and XC2s.

Possible Transition State Conformers. Fig. 1–3 show the nomenclature we use throughout the manuscript. There are two configurations to form a conformer. First is the R' part of the ester: C3 position with respect




Fig. 1. Molecular model of [Et-W] complex: (a) Initial position of H₂O for the potential energy surface calculation to form the complex, C1-O3 distance is 3.00 Å. (b) The typical optimized geometry of the complex in the transition state. Atom C3 may be positioned at (1) trans or (2) gauche.




Fig. 2. Illustration of (a) ester, (b) carboxylic acid, and (c) alcohol for nomenclature used throughout the manuscript. Label (1) is for trans and (2) is for gauche conformation with respect to the position of C3. One H atom bound to C2 and three H atoms bound to C4 are shown to give the illustration the correct perspective.




Fig. 3. (a) Newman projection along C1-C4 bond. The "X" is the position of H atom in Et₂ and of halogen atom in Et₃ and Et₄. (b) The eclipse conformation is defined when "X" is at "a", "b", or "c"; while in staggered ones are at "d", "e", or "f".

to C2-O1-C1 plane, which are (1) trans and (2) gauche (Fig. 1 and 2). Second is the R part of the ester: Orientation of "X" with respect to O (Fig. 3) which gives eclipsed ("a", "b", or "c") and staggered ("d", "e", or "f"). Based on these two configurations, we construct conformers of [Et-W] complex at state 1b, along with their corresponding molecules at state 1a and 1c.

For simplicity, we use the following notation "molecule label(R' - R configuration)". For examples: Et2(1-a) is ethyl acetate with a trans R' and a staggered R conformation; Ct2(2) is acetic acid with a gauche conformation, Et3(2-d) is ethyl fluoroacetate with a gauche R' and an eclipsed R conformation where F atom is at "d".

Chemical Kinetic Calculations. Our interest quantity is the activation energy. The activation energy is expressed in term of the standard Gibbs activation energy, $\Delta^\ddagger G^\circ$ that is the total energy difference between state 1b and 1a. The total energy of a state is the total electronic energy with a correction from Gibbs free energy. The former is the results of DFT calculations while the latter is the results of force constant calculations which is integrated into the software. We extend the usefulness of $\Delta^\ddagger G^\circ$ to calculate the rate constant $k(T)$ based on the transition-state theory [44],

$$k(T) = \frac{k_B T}{h c^\circ} \exp \left[\frac{\Delta^\ddagger G^\circ}{RT} \right] \quad (1)$$

with k_B , h , R are the constant of Boltzmann, Planck, and molar gas, c° is the molecule's concentration from the reactant to the transition state (which we assume to be 1), and T is the temperature (which we assume to be the room temperature, 298.15 K).

3. Results and discussion

3.1. On the ground state structures

The effect of long-range correction. We use Et2(1-a), W, Ca2(1), and ethanol to investigate the effect of the long-range correction on their ground state structures. The reason is that these molecules are experimentally well studied.

Table 3 resumes geometrical parameters from the experiment and our DFT calculations (gas phase). The important value here is the percentage error with respect to the experimental value (err.). The error in general is about less than 1% which corresponds to an order of 10^{-3} Å or 0.1° . This implies both exchange-correlation functionals are excellent to study the ground state structures of the molecules of interest.

Table 4 lists the charge population at some selected atoms. The number of charges does not change significantly after the long-range correction is applied to the calculation. Since the charge population is directly related to the electronic structures of the molecule, the long-range correction does not significantly change the geometrical and electronic structures of the molecule of interest. Therefore, we can conclude that the electrophilic site clearly is at C1 for all exchange correlation functional cases. A water molecule, which has a nucleophilic site at its O3 atom, favorably attacks C1 atom. This makes the starting point of the hydrolysis mechanism shares similarity with that of base-induced ester hydrolysis (Fig. 1a). This analysis becomes our bases on selecting the PES calculation, which we carry in yz -plane where O1, C1, and O3 atoms lie. Fig. 4 shows the PES calculations where the expected entrance of water molecule is at $(z, y) = (1.0, 0.0)$ and it has the highest barrier.

We also present the charge population at C1 and C4 for all esters in Table 5. These two atoms are close to halogen atom in Et3 and Et4. The long-range correction does not change significantly for the charge population at C4, but it does at C1 in Et3 molecules. The $\Delta_{21}(C1)$ average is 0.033 e hence it makes the trend linear: the heavier halogen atom at C4, the more negative the C1 and the more positive the C4. This trend is understandable by considering that (1) the electronegativity of F and Cl (Et3 and Et4, respectively) is higher than C (Et2) and (2) the valence electrons of F is 2p while those of Cl is 3p.

The significant difference between XC1g and XC2g is the calculated total electronic energy (E). Fig. 5 shows the energy level diagram of four molecules in Table 3. The diagrams clearly show that the long-range correction consistently increases E . Except water (Fig. 5b), E increases more than 2 eV; it is almost 4 eV in the case of Et2 (Fig. 5a).

Table 3
Selected geometrical parameters of ethyl acetate, water, acetic acid, and ethanol in the ground state.

Parameter	Expr.	DFT				DFT + PCM			
		XC1g	err. ₁₀	XC2g	err. ₂₀	XC1s	err. _{sg}	XC2s	err. _{sg}
(a) Ethyl acetate, Et2(1-a) (Expr. from [46])									
R(C1,C4)	1.508	1.508	0.0	1.501	-0.5	1.505	-0.2	1.498	-0.2
R(C1,O1)	1.203	1.207	0.3	1.203	0.0	1.214	0.6	1.210	0.6
R(C1,O2)	1.345	1.351	0.5	1.343	0.2	1.343	-0.6	1.335	-0.6
R(C2,C3)	1.515	1.515	0.0	1.509	0.4	1.514	-0.1	1.508	-0.1
R(O2,C2)	1.448	1.450	0.1	1.440	-0.6	1.456	0.4	1.445	0.4
A(O1,C1,C4)	125.4	125.5	0.1	125.5	0.1	125.3	-0.2	125.3	-0.2
A(C4,C1,O2)	110.8	111.0	0.2	111.3	0.4	111.5	0.4	111.7	0.4
A(C3,C2,O2)	123.8	123.4	-0.3	123.2	-0.5	123.3	-0.1	123.0	-0.1
A(O2,C2,C3)	108.2	107.6	-0.5	107.6	-0.6	107.6	0.0	107.6	0.0
A(C1,O2,C2)	117.3	116.6	-0.6	116.5	-0.7	117.4	0.7	117.3	0.7
(b) Water (Expr. from [45])									
R(O3,H)	0.958	0.962	0.5	0.960	0.3	0.964	0.2	0.962	0.2
A(H1,O3,H2)	104.5	105.0	0.5	105.5	0.9	104.5	-0.5	104.9	-0.6
(c) Acetic acid, Ca2(1) (Expr. from [45])									
R(C1,C4)	1.520	1.504	-1.1	1.498	-1.5	1.501	-0.2	1.495	-0.2
R(C1,O3)	1.214	1.205	-0.7	1.200	-1.1	1.212	0.5	1.207	0.6
R(C1,O1)	1.364	1.359	-0.4	1.350	-1.0	1.351	-0.6	1.343	-0.5
R(C4,H)	1.100	1.088	-1.1	1.087	-1.2	1.087	0.0	1.086	0.0
A(C4,C1,O3)	126.6	126.2	-0.3	126.2	-0.4	125.9	-0.2	125.9	-0.2
A(C4,C1,O1)	110.6	111.5	0.8	111.7	1.0	111.9	0.4	112.1	0.4
(d) Ethanol (Expr. from [45])									
R(C2,C3)	1.512	1.517	0.3	1.511	-0.1	1.517	0.0	1.511	0.0
R(C3,H)	1.090	1.093	0.2	1.091	0.1	1.093	0.0	1.092	0.1
R(C2,H)	1.100	1.099	-0.1	1.097	-0.3	1.097	-0.2	1.095	-0.2
R(C2,O2)	1.431	1.431	0.0	1.422	-0.6	1.438	0.5	1.430	0.5
R(O2,H1)	0.971	0.962	-1.0	0.960	-1.2	0.963	0.2	0.962	0.2
A(C2,C3,H)	110.0	110.5	0.5	110.4	0.4	110.8	0.3	110.7	0.3
A(C3,C2,H)	111.0	110.1	-0.8	110.2	-0.8	110.4	0.2	110.4	0.2
A(C3,C2,O2)	107.8	108.0	0.2	107.9	0.1	108.4	-0.4	108.3	0.4

Table 4
The charge population (in unit e) at selected atoms in ground state from NBO calculations.

Atom	DFT			DFT + PCM			
	XC1g	XC2g	Δ_{21}	XC1s	Δ_{sg}	XC2b	Δ_{sg}
(a) Ethyl acetate, Et2(1-abc)							
O1	-0.600	-0.609	-0.009	-0.650	-0.050	-0.659	-0.009
O2	-0.575	-0.580	-0.005	-0.571	0.004	-0.576	-0.005
C1	0.809	0.822	0.013	0.826	0.017	0.840	0.014
C2	-0.034	-0.039	-0.005	-0.034	0.000	-0.039	-0.005
C3	-0.590	-0.598	-0.008	-0.593	-0.003	-0.602	-0.009
C4	-0.668	-0.677	-0.009	-0.672	-0.004	-0.682	-0.010
(b) Water							
H1	0.457	0.460	0.003	0.477	0.020	0.480	0.020
O3	-0.913	-0.920	-0.007	-0.953	-0.040	-0.960	-0.040
(c) Acetic acid, Ca2(1)							
O1	-0.696	-0.703	-0.007	-0.697	-0.001	-0.704	-0.001
O3	-0.595	-0.604	-0.009	-0.643	-0.048	-0.652	-0.048
C1	0.799	0.811	0.012	0.817	0.018	0.830	0.019
C4	-0.679	-0.688	-0.009	-0.682	-0.003	-0.693	-0.005
(d) Ethanol							
H1	0.455	0.459	0.004	0.473	0.018	0.477	0.018
O2	-0.743	-0.748	-0.005	-0.776	-0.033	-0.781	-0.033
C2	-0.023	-0.027	-0.004	-0.024	-0.001	-0.029	-0.002
C3	-0.592	-0.600	-0.008	-0.596	-0.004	-0.605	-0.005

While our results show the dominance of the long-range interaction in the individual molecule, the increasing E compensates each other, hence the net energy of the product and the reactant (reaction energy) is relatively constant. The reaction energy is +0.08 eV and +0.10 eV from XC1g and XC2g calculations, respectively. The long-range correction insignificantly raises the value by 0.02 eV.

The effect of surrounding water. We also provide the results of DFT coupled with PCM in Table 3 (geometry), Table 4 (charge population), and Fig. 5 (level energy diagram). The geometry does not change (most val-

ues of err. are less than 0.5%) with respect to the calculation in the gas phase. The number of charge changes at some atoms by an order of 10^{-2} e and this is significantly changes the frontier molecular orbital wave functions as we have encountered in our previous studies [37, 38]. The perturbation from surrounding waters stabilizes the individual molecule in such a way that the net energy does not change. The reaction energy is 0.10 eV and 0.12 eV with XC1s and XC2s, respectively. The long-range correction improves the value by 0.02 eV, which is exactly the same value in the gas phase case.

Table 5
Charge population from NBO calculation at C1 and C4 of the conformers in the ground state.

Conformer	C1			C4		
	XC1g	XC2g	Δ_{21}	XC1g	XC2g	Δ_{21}
Et1(1)	0.664	0.673	0.009	0.116	0.120	0.004
Et1(2)	0.661	0.670	0.009	0.117	0.122	0.005
Et2(1-a)	0.809	0.822	0.013	-0.668	-0.677	-0.009
Et2(2-a)	0.808	0.821	0.013	-0.668	-0.677	-0.009
Et3(1-a)	0.769	0.805	0.036	-0.447	-0.458	-0.011
Et3(1-b)	0.770	0.801	0.031	-0.421	-0.436	-0.015
Et3(1-c)	0.770	0.801	0.031	-0.421	-0.436	-0.015
Et3(2-a)	0.768	0.804	0.036	-0.447	-0.459	-0.012
Et3(2-b)	0.769	0.801	0.032	-0.422	-0.437	-0.015
Et3(2-c)	0.769	0.800	0.031	-0.421	-0.435	-0.014
Et4(1-a)	0.792	0.781	-0.011	0.006	0.002	-0.004
Et4(1-b)	0.786	0.783	-0.003	0.006	-0.002	-0.004
Et4(1-c)	0.786	0.783	-0.003	0.006	0.002	-0.004
Et4(2-a)	0.791	0.781	-0.010	0.006	0.001	-0.005
Et4(2-b)	0.786	0.782	-0.004	0.006	0.002	-0.004
Et4(2-c)	0.785	0.782	-0.003	0.006	0.002	-0.004




Fig. 4. Potential energy surface of water molecule attacking the electrophilic site of ethyl acetate according to Fig. 1a scenario.

3.2. On the transition state structure

The typical optimized geometry of [Et–W] complex at the saddle point is presented in Fig. 1b. All results have one imaginary frequency as it is required for a TS structure. All imaginary frequencies belong to the normal mode involving the motion of H1 between O2 and O3. The geometry provides us some possible conformers of the [Et–W] complex. Our calculations determine more than one optimized conformer of each [Et–W] complex: two for [Et1–W] and [Et2–W] and three for [Et3–W] and [Et4–W]. Optimization calculations always end up with water in the positive z -axis in the gauche conformation and orientate CH_3 at C4 to either the (a) staggered or (d) eclipsed conformation.

The IRC calculations ensure the optimized geometries of [Et–W] are at the correct transition state according to Scheme 1. Fig. 6 shows the IRC calculation results (only two extreme cases plotted for [Et3–W] and [Et4–W]). All transition state geometries lead exactly to state 1a and 1c. Therefore, Fig. 1b is indeed the typical geometry at state 1b.

Table 6, 7, 8, and 9 display four important geometrical parameters of [Et–W] in the transition state. They are the distance of C1–O2 (R_1), O3–H1 (R_2), C1–O3 (R_3), and O2–H1 (R_4), respectively. The favorable condition for hydrolysis is the elongation of R_1 and R_2 with respect to the state 1a and a short distance of R_3 and R_4 . In case of Et2, using XC1g, R_1 and R_2 significantly elongate to 33.2% and 27.8%, respectively. R_1 and R_2 in all [Et–W] complexes are 1.73 Å and 1.19 Å on average, respectively. Since C–O in gas phase experimentally 1.48 Å [45], all [Et–W] complexes show a tendency to cleavage Et forming Ca and ethanol. Meanwhile, R_3 and R_4 determine the stability of transition state geometry: the shorter the values, the stronger the interaction

between Et and W in [Et–W] complexes, which implies more stable geometry. In that sense, the most stable conformation is gauche (in case of [Et1–W] and [Et2–W]) and eclipsed with a halogen atom at “e” position (in case of [Et3–W] and [Et2–W]).

The effect of long-range correction and surrounding water. The value of err_{21} and err_{sg} in Table 6, 7, 8, and 9 show the effect of the long-range correction and the surrounding water. Overall, the effect is more significant in [Et–W] complex (state 1b) than in individual molecules of Et and W (state 1a) and Ca and alcohol (state 1c). The significant effect of the long-range correction is on R_1 and R_3 . Both parameters involve C–O bond. When we consider the ground state analysis (Fig. 5), the results show a tendency of XC1s significantly affecting molecules with C–O bond. It explains qualitatively as to how XC1s raises E of ethyl acetate almost twice as much as the acetic acid and ethanol (see Fig. 5): ethyl acetate has two C–O bonds, acetic acid has one O–C–O, and ethanol only has one C–O.

We extend the discussion on [Et2–W] geometry in the transition state. Our results are comparable with the work of Yamabe et al. [14]. Table 10 resumes the comparison between these two works. Overall, our results are far from the results of [14]. In the water surrounding model, it is clear that PCM and Onsager’s model yields a significant difference in geometry. The difference can be understood as follows. Both PCM and Onsager’s model create a cavity to simulate the reaction field, but the method to make the cavity is completely different.

3.3. On the conformational effect

We evaluate the conformational effect on geometry based on the value of range in Table 6, 7, 8, and 9. The wider the range means the more significant the conformer variation of the complex. In our case, the overall range is in an order of 10^{-2} Å. Furthermore, we find two trends. First, the range is wider in the presence of a halogen atom (Et3 and Et4). Second, the long-range correction and the surrounding water significantly affect the ranges of R_1 and R_2 of [Et1–W] and [Et2–W]. These two trends imply that the conformer of ethyl acetate has a significant effect when it is forming a complex with water in the transition state. Accordingly, the figure 10^{-2} Å is a significant range for this study.

The difference in geometry directly affects the activation energy ($\Delta^\ddagger G^\circ$) as displayed in Table 11. We highlight three points from the table as follows.

First of all, there is a general trend of $\Delta^\ddagger G^\circ$ in the presence of the halogen atom in esters. This result is consistent with the charge population at C1 of ester in the ground state (Table 5) and R_3 (Table 6). Charge population at C1 is more negative in the presence of halogen atom (Et2 versus Et3 and Et4), so it provides more charges to form a




Fig. 5. Energy level diagram of molecules in Table 3, for (a) ethyl acetate, (b) water, (c) acetic acid, and (d) ethanol. The calculated energy by XC1g is set to be the reference.




Fig. 6. Reaction path calculations based on the optimized geometry of [Et-W] complexes for the case of (a) Et1, (b) Et2, (c) Et3, and (d) Et4 with the conformations are shown in the legend. The peak of path is the saddle point which is the transition state (1b), to the left is the initial state (1a) and to the right is the final state (1c). The energy reference is the 1a of each system, so the height corresponds to the energy barrier of each reaction with d as the difference in energy barrier between two different conformers. (Table 11 lists the complete value of energy barrier for all reactions of interest.)

covalent bond with O3 from H₂O (shorter R_3). The elongation of R_1 (Table 6) supports the C1–O3 bonding formation and directly determines the value of $\Delta^\ddagger G^\circ$. There is an exception in “d” conformation, where R_1 is shorter, making the $\Delta^\ddagger G^\circ$ higher than the other conformations. However, the overall trend is the presence of a halogen atom lowers $\Delta^\ddagger G^\circ$, which corresponds to the higher $k(T)$. This is in line with other esters [4, 5, 6] as mentioned in the introduction. Furthermore, a heavier halogen atom raises $k(T)$ which agrees with the result by

Schmeer and Sturm [12]. This part of results is important in our study because it justifies our calculation method.

Secondly, a comparison among results in the table shows the effect of the long-range correction and the surrounding water. The former’s effect alone (err.₂₁) is less than 1% on average, while the latter’s effect (err._{sg}) is more than 7% on average. Even though err.₂₁ in this work is insignificant, the exchange-correlation functional is still an important factor. As we see in Schmeer and Sturm [12], the use of Hatree-

Table 6
Distance of C1–O2 in the transition state (R_1 , in Å).

Conformer	DFT			DFT + PCM			
	XC1g	XC2g	err. ₂₁	XC1s	err. _{sg}	XC2b	err. _{sg}
(1) [Et1–W]							
(1)	1.715	1.663	–3.0	1.664	–3.0	1.615	–2.9
(2)	1.720	1.668	–3.0	1.677	–2.5	1.626	–2.5
range	0.004	0.005		0.013		0.012	
(2) [Et2–W]							
(1-a)	1.791	1.727	–3.6	1.746	–2.5	1.668	–3.4
(2-d)	1.793	1.726	–3.7	1.759	–1.9	1.680	–2.7
range	0.001	0.001		0.012		0.012	
(3) [Et3–W]							
(1-f)	1.764	1.703	–3.5	1.677	–4.9	1.619	–5.0
(1-d)	1.695	1.640	–3.2	1.654	–2.4	1.602	–2.3
(1-e)	1.712	1.654	–3.4	1.659	–3.1	1.607	–2.8
(2-f)	1.764	1.701	–3.6	1.690	–4.2	1.630	–4.1
(2-d)	1.699	1.644	–3.2	1.665	–2.0	1.612	–2.0
(2-e)	1.713	1.654	–3.4	1.671	–2.5	1.617	–2.3
range	0.069	0.063		0.035		0.028	
(4) [Et4–W]							
(1-f)	1.771	1.706	–3.7	1.692	–4.5	1.627	–4.6
(1-d)	1.687	1.636	–3.0	1.642	–2.6	1.595	–2.5
(1-e)	1.712	1.654	–3.4	1.657	–3.2	1.607	–2.8
(2-f)	1.771	1.704	–3.8	1.705	–3.7	1.639	–3.8
(2-d)	1.690	1.637	–3.1	1.653	–2.2	1.604	–2.0
(2-e)	1.717	1.659	–3.4	1.673	–2.6	1.619	–2.4
range	0.084	0.070		0.062		0.044	

Table 7
Distance of O3–H1 in the transition state (R_2 , in Å).

Conformer	DFT			DFT + PCM			
	XC1g	XC2g	err. ₂₁	XC1s	err. _{sg}	XC2b	err. _{sg}
(1) [Et1–W]							
(1)	1.236	1.232	–0.3	1.263	2.2	1.245	1.1
(2)	1.227	1.221	–0.4	1.248	1.8	1.231	0.7
range	0.009	0.010		0.015		0.014	
(2) [Et2–W]							
(1-a)	1.228	1.224	–0.4	1.268	3.2	1.257	2.6
(2-d)	1.225	1.223	–0.2	1.258	2.7	1.245	1.8
range	0.004	0.001		0.011		0.011	
(3) [Et3–W]							
(1-f)	1.210	1.203	–0.6	1.251	3.4	1.237	2.8
(1-d)	1.234	1.229	–0.4	1.252	1.5	1.235	0.5
(1-e)	1.261	1.256	–0.4	1.267	0.5	1.247	–0.7
(2-f)	1.208	1.205	–0.3	1.238	2.5	1.226	1.7
(2-d)	1.227	1.221	–0.5	1.241	1.2	1.225	0.4
(2-e)	1.256	1.250	–0.5	1.256	0.0	1.236	–1.1
range	0.053	0.052		0.029		0.021	
(3) [Et4–W]							
(1-f)	1.203	1.200	–0.2	1.238	2.9	1.231	2.6
(1-d)	1.231	1.223	–0.6	1.248	1.4	1.231	0.6
(1-e)	1.268	1.258	–0.8	1.280	1.0	1.252	–0.5
(2-f)	1.201	1.201	0.0	1.226	2.1	1.219	1.5
(2-d)	1.227	1.221	–0.5	1.239	1.0	1.224	0.2
(2-e)	1.261	1.252	–0.7	1.267	0.5	1.243	–0.7
range	0.066	0.058		0.054		0.033	

Fock method resulted in 58.74 kcal/mol and 64.47 kcal/mol of $\Delta^\ddagger G^\circ$.¹ Furthermore, the method to model the water surrounding is also important. The use of Onsager's model by Yamabe et al. [14] resulted in

¹ Schmeer and Sturm [12] used two waters and has one intermediate state and two transition states in the reaction path. The calculations were in the gas phase.

47.90 kcal/mol. The difference in $\Delta^\ddagger G^\circ$ is expected since the geometry is also different (see Table 10).

Lastly, the range of $\Delta^\ddagger G^\circ$ displays the effect of conformation. In XC1g, the range is 0.72, 1.07, 2.64, and 3.47 kcal/mol for Et1, Et2, Et3, and Et4 case, respectively. These figures give a significant ratio between the highest and the lowest $k(T)$ of each case. Since $k(T)$ is an exponential function of negative $\Delta^\ddagger G^\circ$ [see Equation (1)], 0.72 kcal/mol (in gas phase) corresponds to 3 times increment of the rate con-

Table 8
Distance of C1–O3 in the transition state (R_3 , in Å).

Conformer	DFT			DFT + PCM			
	XC1g	XC2g	err. ₂₁	XC1s	err. _{sg}	XC2b	err. _{sg}
(1) [Et1–W]							
(1)	1.780	1.731	–2.8	1.779	–0.1	1.711	–1.2
(2)	1.773	1.724	–2.8	1.771	–0.1	1.702	–1.3
range	0.007	0.008		0.008		0.009	
(2) [Et2–W]							
(1-a)	1.841	1.780	–3.3	1.859	1.0	1.776	–0.3
(2-d)	1.838	1.780	–3.2	1.859	1.1	1.775	–0.3
range	0.002	0.001		0.001		0.001	
(3) [Et3–W]							
(1-f)	1.775	1.714	–3.4	1.763	–0.7	1.692	–1.3
(1-d)	1.760	1.707	–3.0	1.756	–0.2	1.686	–1.2
(1-e)	1.829	1.771	–3.2	1.792	–2.0	1.717	–3.1
(2-f)	1.774	1.717	–3.2	1.757	–1.0	1.686	–1.8
(2-d)	1.755	1.701	–3.1	1.752	–0.2	1.681	–1.2
(2-e)	1.824	1.766	–3.2	1.789	–1.9	1.712	–3.1
range	0.074	0.070		0.040		0.035	
(4) [Et4–W]							
(1-f)	1.775	1.716	–3.3	1.759	–0.9	1.690	–1.5
(1-d)	1.749	1.695	–3.1	1.739	–0.5	1.671	–1.4
(1-e)	1.835	1.772	–3.4	1.808	–1.5	1.724	–2.7
(2-f)	1.774	1.718	–3.2	1.754	–1.2	1.687	–1.8
(2-d)	1.746	1.694	–3.0	1.736	–0.6	1.670	–1.4
(2-e)	1.832	1.773	–3.3	1.806	–1.4	1.726	–2.6
range	0.088	0.079		0.072		0.055	

Table 9
Distance of O2–H1 in the transition state (R_4 , in Å).

Conformer	DFT			DFT + PCM			
	XC1g	XC2g	Δ_{21}	XC1s	Δ_{sg}	XC2b	Δ_{sg}
(1) [Et1–W]							
(1)	1.187	1.185	–0.2	1.169	–1.6	1.181	–0.3
(2)	1.195	1.194	–0.1	1.180	–1.2	1.192	–0.1
range	0.008	0.009		0.011		0.012	
(2) [Et2–W]							
(1-a)	1.190	1.187	–0.3	1.158	–2.7	1.163	–2.0
(2-d)	1.192	1.187	–0.4	1.165	–2.3	1.171	–1.3
range	0.002	0.000		0.007		0.008	
(3) [Et3–W]							
(1-f)	1.212	1.213	0.1	1.181	–2.5	1.191	–1.8
(1-d)	1.193	1.192	–0.1	1.180	–1.1	1.193	0.0
(1-e)	1.167	1.166	–0.1	1.167	0.1	1.182	1.4
(2-f)	1.213	1.210	–0.2	1.175	–3.1	1.200	–0.9
(2-d)	1.199	1.199	0.0	1.199	0.0	1.200	0.1
(2-e)	1.170	1.170	–0.0	1.170	–0.0	1.190	1.7
range	0.046	0.048		0.032		0.018	
(4) [Et4–W]							
(1-f)	1.218	1.215	–0.2	1.191	–2.2	1.195	–1.7
(1-d)	1.197	1.199	0.2	1.186	–0.9	1.199	–0.0
(1-e)	1.160	1.163	0.2	1.156	–0.4	1.176	1.1
(2-f)	1.218	1.213	–0.5	1.200	–1.5	1.204	–0.7
(2-d)	1.200	1.201	0.1	1.193	–0.6	1.204	0.3
(2-e)	1.164	1.166	0.1	1.164	0.0	1.181	1.3
range	0.058	0.053		0.044		0.028	

Table 10
Result comparison on [Et2–W] in the transition state.

Parameter	Ref. [14]	XC1s	err.	XC2s	err.
R_1	1.790	1.746	–2.4	1.668	–6.8
R_2	1.203	1.268	5.4	1.257	4.4
R_3	1.802	1.859	3.2	1.776	–1.5
R_4	1.221	1.158	–5.2	1.163	–4.7

Here err. is the error percentage with respect to the value from Ref. [14].

Table 11

The standard Gibbs activation energy (in kcal/mol) from various [Et–W] conformers.

Conformer	DFT			DFT + PCM			
	XC1g	XC2g	err. ₂₁	XC1s	err. _{1sg}	XC2b	err. _{2sg}
(1) Et1 + W → [Et1–W] → Ca1 + ethanol							
(1)	52.93	53.29	0.7	56.77	7.3	56.78	6.5
(2)	52.21	52.44	0.4	55.67	6.6	55.81	6.4
range	0.72	0.85		1.10		0.96	
(2) Et2 + W → [Et2–W] → Ca2 + ethanol							
(1-a)	53.52	54.01	0.9	57.64	7.7	57.92	7.2
(2-d)	52.45	53.05	1.1	56.18	7.1	56.95	7.3
range	1.07	0.97		1.46		0.97	
(3) Et3 + W → [Et3–W] → Ca3 + ethanol							
(1-f)	53.42	53.89	0.9	57.93	8.4	57.72	7.1
(1-d)	54.86	54.35	–0.9	58.47	6.6	57.72	6.2
(1-e)	53.46	53.37	–0.2	57.80	8.1	57.13	7.0
(2-f)	52.80	53.02	0.4	56.90	7.8	56.87	7.3
(2-d)	53.67	53.53	–0.3	56.95	6.1	56.52	5.6
(2-e)	52.22	52.44	0.4	57.79	10.7	56.35	7.5
range	2.64	1.91		1.58		1.37	
(4) Et4 + W → [Et4–W] → Ca4 + ethanol							
(1-f)	53.10	53.57	0.9	57.57	8.4	57.28	6.9
(1-d)	55.28	54.91	–0.7	59.48	7.6	58.42	6.4
(1-e)	52.76	52.93	0.3	56.42	6.9	56.69	7.1
(2-f)	52.48	52.57	0.2	56.62	7.9	56.13	6.8
(2-d)	54.83	54.43	–0.7	58.12	6.0	57.43	5.5
(2-e)	51.81	51.96	0.3	55.83	7.8	56.19	8.1
range	3.47	2.96		3.66		2.29	

Table 12

The rate constant at 298.15 K in natural logarithmic value.

Conformer	DFT			DFT + PCM			
	XC1g	XC2g	rat. ₂₁	XC1s	rat. _{1sg}	XC2b	rat. _{2sg}
(1) Et1 + W → [Et1–W] → Ca1 + ethanol							
(1)	–59.88	–60.49	–0.61	–66.36	–6.48	–66.38	–5.89
(2)	–58.66	–59.05	–0.39	–64.50	–5.84	–64.74	–5.69
rat. _k	3.37	4.20		6.40		5.14	
(2) Et2 + W → [Et2–W] → Ca2 + ethanol							
(1-a)	–60.87	–61.70	–0.83	–67.83	–6.95	–68.30	–6.60
(2-d)	–59.07	–60.08	–1.01	–65.36	–6.30	–66.66	–6.58
rat. _k	6.09	5.05		11.75		5.14	
(3) Et3 + W → [Et3–W] → Ca3 + ethanol							
(1-f)	–60.71	–61.50	–0.79	–68.32	–7.61	–67.96	–6.46
(1-d)	–63.14	–62.27	0.86	–69.23	–6.09	–67.96	–5.69
(1-e)	–60.77	–60.62	0.15	–68.10	–7.33	–66.97	–6.35
(2-f)	–59.66	–60.03	–0.37	–66.58	–6.92	–66.53	–6.50
(2-d)	–61.13	–60.89	0.24	–66.66	–5.54	–65.94	–5.05
(2-e)	–58.68	–59.05	–0.37	–68.08	–9.40	–65.65	–6.60
rat. _k	86.13	25.12		14.15		10.09	
(3) Et4 + W → [Et4–W] → Ca4 + ethanol							
(1-f)	–60.17	–60.96	–0.79	–67.71	–7.54	–67.22	–6.26
(1-d)	–63.84	–63.22	0.62	–70.93	–7.09	–69.14	–5.92
(1-e)	–59.59	–59.88	–0.29	–65.77	–6.18	–66.22	–6.35
(2-f)	–59.12	–59.27	–0.15	–66.11	–6.99	–65.28	–6.01
(2-d)	–63.08	–62.41	0.68	–68.64	–5.55	–67.47	–5.06
(2-e)	–57.99	–58.24	–0.25	–64.77	–6.79	–65.38	–7.14
rat. _k	349.57	145.33		473.66		47.71	

stant (at 298.15 K), 1.07 kcal/mol corresponds to 6 times increment, 2.64 kcal/mol corresponds to 86 times increment, and 3.47 kcal/mol corresponds to 350 times increment. Table 12 tabulates these results and clarifies the effect of conformation. Overall, the long-range correction and the surrounding water narrow the range, but the figures are still significant. Meanwhile, the gauche conformation has the highest $k(T)$ and the halogen atom prefers in “e” position (staggered conformation, see Fig. 3).

4. Conclusion

We have reported the conformation in transition state significantly affects the activation energy of ethyl acetate neutral hydrolysis. The gauche conformers tend to have higher rate constant than the trans ones. This leads to the significant ratio between the lowest and the highest rate constant: from 3 to 350 times, or from 4 to 145 times after long-range correction (all in the gas phase). We argued that one must

take the conformers into the consideration to study the ester neutral hydrolysis. Furthermore, the halogenated ethyl acetate has higher rate constant and the halogen atom prefers in the staggered conformation. The halogenation effect from our calculations is in agreement with experimental data for other ester variant. Finally, we also presented the long-range interaction and PCM to model the surrounding water stabilize the transition state geometry but raise the activation energy from 5.5% to 8.1%.

Declarations

Author contribution statement

Febdian Rusydi: Conceived and designed the experiments; Performed the experiments; Analyzed and interpreted the data; Contributed reagents, materials, analysis tools or data; Wrote the paper.

Nufida Dwi Aisyah, Rizka Fadilla: Conceived and designed the experiments; Performed the experiments.

Hermawan Dipojono, Mudasar Mudasar, Andriwo Rusydi: Analyzed and interpreted the data.

Faozan Ahmad, Ira Puspitasari: Contributed reagents, materials, analysis tools or data.

Funding statement

This work was supported by Directorate General of Higher Education, Research, and Technology Ministry, Republic of Indonesia through National Collaboration Research Grant number 563/UN3.14/LT/2018.

Competing interest statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

Acknowledgements

Authors thank to Luslia Silfia Pulo Boli (Research Center for Quantum Engineering Design, Universitas Airlangga, Surabaya, Indonesia), Vera Khoirunisa (Engineering Physics Department, Institut Teknologi Sumatera, Lampung, Indonesia), Dr. Adhitiya Gandaryus Saputro and Dr. Muhammad Kemal Agusta (Engineering Physics Department, Institut Teknologi Bandung, Bandung, Indonesia), Prof. Azizhan Ahmad (Chemistry Department, Universiti Kebangsaan Malaysia, Malaysia), and Prof. Diño Wilson Agerico Tan and Prof. Yoshitada Morikawa (Precision Science & Technology and Applied Physics, Graduate School of Engineering, Osaka University, Japan) for their insight and valuable discussions. RNF particularly thanks to LPDP for the scholarship. All calculations using Gaussian 09 software are performed in the computer facility at the Osaka University.

References

- [1] Y.H. Hsieh, N. Weinberg, S. Wolfe, The neutral hydrolysis of methyl acetate – Part 1. Kinetic experiments, *Can. J. Chem.* 87 (2009) 539–543.
- [2] A. Skrabal, A. Zahorka, Die Wasserverseifung des Athylazetats, *Monatsh. Chem.* 53–54 (1929).
- [3] R. Wolfenden, Y. Yuan, The “neutral” hydrolysis of simple carboxylic esters in water and the rate enhancements produced by acetylcholinesterase and other carboxylic acid esterases, *J. Am. Chem. Soc.* 133 (2011) 13821–13823.
- [4] A. Moffat, H. Hunt, Solvent and chain length effects in the non-catalyzed hydrolysis of some alkyl and aryl trifluoroacetates, *J. Am. Chem. Soc.* 81 (1959) 2082–2086.
- [5] E.K. Euranto, N.J. Cleve, Kinetics of the neutral hydrolysis of chloromethyl chloroacetate, *Acta Chem. Scand.* 17 (1963) 1584–1594.
- [6] K.S. Venkatasubban, K.R. Davis, J.L. Hogg, Transition-state structure for the neutral water-catalyzed hydrolysis of ethyl trifluorothiolacetate, *J. Am. Chem. Soc.* 100 (1963) 6125–6128.
- [7] J.F. Kirsch, W.P. Jencks, Nonlinear structure-reactivity correlations. The imidazole-catalyzed hydrolysis of esters, *J. Am. Chem. Soc.* 86 (1964).
- [8] P.L. da Silva, L. Guimarães, J.R.P. Jr., Revisiting the mechanism of neutral hydrolysis of esters: water autoionization mechanisms with acid or base initiation pathways, *J. Phys. Chem. B* 117 (2013) 6487–6497.
- [9] Z. Shi, Y.H. Hsieh, N. Weinberg, S. Wolfe, The neutral hydrolysis of methyl acetate – Part 2. Is there a tetrahedral intermediate? *Can. J. Chem.* 87 (2009) 544–555.
- [10] Y.-H. Hsieh, N. Weinberg, K. Yang, C.-K. Kim, Z. Shi, S. Wolfe, Hydration of the carbonyl group – acetic acid catalysis in the co-operative mechanism, *Can. J. Chem.* 83 (2005) 769–785.
- [11] X. Chen, C.-G. Zhan, Fundamental reaction pathways and free-energy barriers for ester hydrolysis of intracellular second-messenger 3–5-cyclic nucleotide, *J. Phys. Chem. A* 108 (2004) 3789–3797.
- [12] G. Schmeer, P. Sturm, A quantum chemical approach to the water assisted neutral hydrolysis of ethyl acetate and its derivatives, *Phys. Chem. Chem. Phys.* 1 (1999) 1025–1030.
- [13] S. Wolfe, C.-K. Kim, K. Yang, N. Weinberg, Z. Shi, Hydration of the carbonyl group. A theoretical study of the cooperative mechanism, *J. Am. Chem. Soc.* 115 (1995) 4240–4260.
- [14] S. Yamabe, N. Tsuchida, Y. Hayashida, Reaction paths of the water-assisted neutral hydrolysis of ethyl acetate, *J. Phys. Chem. A* 109 (2005) 7216–7224.
- [15] C.-G. Zhan, D.W. Landry, R.L. Ornstein, Energy barriers for alkaline hydrolysis of carboxylic acid esters in aqueous solution by reaction field calculations, *J. Phys. Chem. A* 104 (2000) 7672–7678.
- [16] B. Kallies, R. Mitzner, Models of water-assisted hydrolyses of methyl formate, formamide, and urea from combined DFT-SCRF calculations, *J. Mol. Model.* 4 (1998).
- [17] R.T. Bartus, R.L. Dean, B. Beer, A.S. Lippa, The cholinergic hypothesis of geriatric memory dysfunction, *Science* 217 (1982) 408–417.
- [18] P.T. Francis, A.M. Palmer, M. Snape, G.K. Wilcock, The cholinergic hypothesis of Alzheimer – disease: a review of progress, *J. Neurol. Neurosurg. Psychiatry* 66 (1999) 137–147.
- [19] T. Svinning, H. Sorum, A reinvestigation of the crystal structure of acetylcholine bromide, *Acta Crystallogr., Sect. B* 31 (1975) 1581–1585.
- [20] M. Sax, M. Rodrigues, G. Blank, M.K. Wood, J. Pletcher, The conformation of acetylcholine and the crystal structure of 2, 2-dimethylbutyl 3, 5-dinitrobenzoate, *Acta Crystallogr., Sect. B* 32 (1976) 1953–1956.
- [21] R.W. Behling, T. Yamane, G. Navon, L.W. Jelinski, Conformation of acetylcholine bound to the nicotinic acetylcholine receptor, *Proc. Natl. Acad. Sci.* 85 (1988) 6721–6725.
- [22] K.J. Wilson, P. Derreumaux, G. Vergoten, W.L. Peticolas, Conformational studies of neuroactive ligands 2. solution-state conformations of acetylcholine, *J. Phys. Chem.* 93 (1989) 1351–1357.
- [23] B. Hernández, P. Houzé, F. Pfléger, S.G. Kruglik, M. Ghomi, Raman scattering-based multiconformational analysis for probing the structural differences between acetylcholine and acetylthiocholine, *J. Pharm. Biomed. Anal.* 138 (2017) 54–62.
- [24] H. Sorum, The crystal and molecular structure of acetyl choline bromide, *Acta Chem. Scand.* 13 (1959) 345–359.
- [25] C. Chothia, P. Pauling, Conformation of cholinergic molecules relevant to acetylcholinesterase, *Nature* 223 (1969) 919–921.
- [26] C. Chothia, P. Pauling, The conformation of cholinergic molecules at nicotinic nerve receptors, *Proc. Nat. Acad. Sci.* 65 (1970) 477–482.
- [27] T.L. Lemke, D.A. Williams, V.F. Roche, S.W. Zito (Eds.), *Foye’s Principle of Medicinal Chemistry*, 7th ed., Lippincott Williams & Wilkins, 2013.
- [28] P. Hohenberg, W. Kohn, Inhomogeneous electron gas, *Phys. Rev.* 136 (1964) B864.
- [29] W. Kohn, L.J. Sham, Self-consistent equations including exchange and correlation effects, *Phys. Rev. A* 140 (1965) A1133.
- [30] M.J. Frisch, G.W. Trucks, H.B. Schlegel, G.E. Scuseria, J.R.C.M.A. Robb, G. Scalmani, V. Barone, B. Mennucci, G.A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H.P. Hratchian, A.F. Izmaylov, J. Bloino, G. Zheng, J.L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J.A. Montgomery Jr., J.E. Peralta, F. Ogliaro, M. Bearpark, J.J. Heyd, E. Brothers, K.N. Kudin, V.N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J.C. Burant, S.S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J.M. Millam, M. Klene, J.E. Knox, J.B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R.E. Stratmann, O. Yazyev, A.J. Austin, R. Cammi, C. Pomelli, J.W. Ochterski, R.L. Martin, K. Morokuma, V.G. Zakrzewski, G.A. Voth, P. Salvador, J.J. Dannenberg, S. Dapprich, A.D. Daniels, O. Farkas, J.B. Foresman, J.V. Ortiz, J. Cioslowski, D.J. Fox, Gaussian 09, revision c. 01, 2010, “Gaussian Inc. Wallingford CT”.
- [31] A.D. Becke, Density-functional thermochemistry. iii. The role of exact exchange, *J. Chem. Phys.* 98 (1993) 5648–5652.
- [32] T. Yanaia, D.P. Tew, N.C. Handy, A new hybrid exchange–correlation functional using the coulomb-attenuating method (cam-b3lyp), *Chem. Phys. Lett.* 393 (2004) 51–57.
- [33] A.G. Saputro, F. Rusydi, M.K. Agusta, H. Kasai, H.K. Dipojono, Oxygen reduction reaction on cobalt(6) pyrrol cluster: density functional theory study, *J. Phys. Soc. Jpn.* 81 (2012) 034703.
- [34] F. Rusydi, M.K. Agusta, A.G. Saputro, H. Kasai, A first principle study on zinc porphyrin interaction with O₂ in zinc porphyrin(oxygen) complex, *J. Phys. Soc. Jpn.* 81 (2012) 124301.

- [35] F. Rusydi, M.K. Agusta, A.G. Saputro, H. Kasai, A theoretical study of ligand effects on the electronic structures of ligated zinc porphyrin using density functional theory, *J. Vac.Soc. Jpn.* 57 (2014) 102–110.
- [36] W.-F. Mark-Lee, F. Rusydi, L.J. Minggu, T. Kubo, M. Kassim, Bis(bipyridyl)-ru(ii)-benzoyl-3-(pyridine-2-yl)-1h-pyrazole as potential photosensitizer: experimental and density functional theory study, *J. Technol.* 75 (2017) 117–123.
- [37] F. Rusydi, A.G. Saputro, H. Kasai, A density-functional study on the change of q/b-band intensity ratio of zinc tetraphenylporphyrin in solvents, *J. Phys. Soc. Jpn.* 83 (2014) 084802.
- [38] F. Rusydi, G. Shukri, A.G. Saputro, H.K. Dipojono, S. Suprijadi, Dipole strength calculation based on two-level system approximation to study q/b-band intensity ratio of zntbp in solvent, *J. Phys. Soc. Jpn.* 86 (2017) 044706.
- [39] R.N. Fadilla, N.D. Aisyah, H.K. Dipojono, F. Rusydi, A theoretical study of the rearranging trans-HCOH to H₂CO via quantum tunneling with dft and wkb approximation, *Proc. Eng.* 170 (2017) 113–118.
- [40] N.D. Aisyah, R.N. Fadilla, H.K. Dipojono, F. Rusydi, A theoretical study of monodeuteriation effect on the rearrangement of trans-HCOH to H₂CO via quantum tunneling with dft and wkb approximation, *Proc. Eng.* 170 (2017) 119–123.
- [41] R.N. Fadilla, N.D. Aisyah, H.K. Dipojono, F. Rusydi, The first-principles study on the stability of trans-hcoh in various solvents, *J. Phys. Conf. Ser.* 853 (2017) 012031.
- [42] A. Fortunnelli, J. Tomasi, The implementation of density functional theory within the polarizable continuum model for solvation, *Chem. Phys. Lett.* 231 (1994) 34–39.
- [43] E.D. Glendening, A.E. Reed, J.E. Carpenter, F. Weinhold, *NBO Version 3.1*.
- [44] D.A. McQuarrie, J.D. Simon, *Physical Chemistry - A Molecular Approach*, University Science Books, 1997.
- [45] D.L. Lide, in: W.M. Haynes (Ed.), *CRC Handbook of Chemistry and Physics*, 96th ed., CRC Press/Taylor and Francis, Boca Raton FL, USA, 2015, pp. 9–48.
- [46] K. Kuchitsu (Ed.), *Structure of Free Polyatomic Molecules – Basic Data*, Springer, Heidelberg NY, USA, 1998.