

PAPER • OPEN ACCESS

Preface

To cite this article: 2020 *IOP Conf. Ser.: Mater. Sci. Eng.* **856** 011001

View the [article online](#) for updates and enhancements.

PREFACE

It is with great pleasure that I introduce the proceedings of the 2019 5th International Conference on Smart Material Research (ICSMR 2019) which was held recently in Brisbane, Australia, 13-16, November 2019. ICSMR 2019 was dedicated to technical issues related to smart material research.

The objectives of the conference were not only to provide opportunities to the international scientists to present state of the art research in smart materials but also to provide a forum to discuss ideas for future directions in the field and explore possibilities for collaborative research projects.

The conference program included keynote, oral, and poster presentations from scholars working in the areas of materials science and engineering from all over the world. It covered recent trends and progress made in the field of materials engineering and manufacturing. Professors from Australia and USA were invited to deliver four keynote lectures regarding the latest information in their respective areas of expertise.

The proceedings present a selection of high quality papers submitted to the conference by researchers from universities, research institutes, and industry. All papers were subjected to peer-review by conference committee members and international reviewers. The papers were selected based on their quality and their relevance to the conference. The proceedings present recent advances in the fields of materials science and engineering and manufacturing as well as in other related areas including materials properties, measuring methods, analytical and numerical modeling and simulations among others.

I would like to express special gratitude to members of the conference committee and organizers of the conference. I would also like to thank the reviewers for their valuable time and advice which helped in improving the quality of the papers selected for presentation at the conference and for publication in the proceedings. Finally I want to thank the authors, the members of the organizing committee, the reviewers, the chairpersons, sponsors, and all other conference participants for their support of ICSMR 2019.

Prof. Ramesh K. Agarwal
Washington University in St. Louis, USA
January 21, 2020



□ **NOTICE:** Ensuring subscriber access to content on IOPscience throughout the coronavirus outbreak - see our remote access guidelines.

Table of contents

Volume 856

2020

◀ Previous issue Next issue ▶

**2019 The 5th International Conference on Smart Materials Research 13-16
November 2019, Brisbane, Australia**

Accepted papers received: 28 April 2020

Published online: 28 May 2020

Open all abstracts

Preface

OPEN ACCESS 011001

Preface

+ Open abstract  View article  PDF

OPEN ACCESS 011002

Committee

+ Open abstract  View article  PDF

OPEN ACCESS 011003

Delegate list

+ Open abstract  View article  PDF

OPEN ACCESS 011004

Peer review statement

+ Open abstract  View article  PDF

Material Mechanics

OPEN ACCESS 012001

Punching Shear Strength of Reinforced Concrete Flat Slabs with Macro

Synthetic Fibers

Nadia Nassif, Salah Altoubat, Mohamed Maalej and Pierre Estephane

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012002

Mechanical Properties Improvement of Clays Using Silica Sand Waste and Dust Sand Foundry Waste

Anita Setyowati Srie Gunarti and Irwan Raharja

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012003

The effect of drying method on the surface structure of mesoporous sol-gel derived bioactive glass-ceramic

Neda Ghaebi Panah, Rob Atkin and Tim Sercombe

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012004

Theoretical exploration on free radical scavenging mechanism of curcumin analogues in water solvent

L S P Boli, N D Aisyah, V Khoirunisa, F Rusydi, H Rachmawati and H K Dipojono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012005

Theoretical investigation on electron transfer-based antioxidant activity of melinjo resveratrol

V Khoirunisa, R N Fadilla, L S P Boli, F Rusydi, H Rachmawati and H K Dipojono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012006

Search for the maximal threshold Co content for the oxidation behaviour point of view for 25wt.%Cr-containing TaC-strengthened cast superalloys designed for use at 1200°C on long time

P Berthod and J P Gomis

[+ Open abstract](#) [View article](#) [PDF](#)

Composite Materials

OPEN ACCESS

012007

Carbon nanotube reinforced aluminium composites fabricated by powder metallurgical and hot-rolling techniques

Hideaki Tsukamoto

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012008

Micro/nanostructured MTMOs as Electrocatalysts for Oxygen Evolution Reaction in Alkaline Medium

Charles Lois Flores and Mary Donnabelle Balela

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012009

Synthesis and antiparkinsonian activity of nanocomposite of chitosan-tripolyphosphate-*Mucuna pruriens* L extract (CS-TPP-MP)

R E Sardjono, A N Fauziyah, M D Puspitasari, I Musthapa, F Khoerunnisa, G N Azzahra, R Mamat and Erdiwansyah

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012010

Modeling and simulation of a gauge shaped beam coupled with macro fiber composite for energy harvesting application

D S Ibrahim, B B Sun, U Sharif and O A Oluseyi

[+ Open abstract](#)

[View article](#)

[PDF](#)

JOURNAL LINKS

[Journal home](#)

[Information for organizers](#)

[Information for authors](#)

[Search for published proceedings](#)

[Contact us](#)

[Reprint services from Curran Associates](#)

PAPER • OPEN ACCESS

Committee

To cite this article: 2020 *IOP Conf. Ser.: Mater. Sci. Eng.* **856** 011002

View the [article online](#) for updates and enhancements.

Conference Committee Chairs

Darren Martin, University of Queensland, Australia

Ramesh K. Agarwal, Washington University in St. Louis, USA

Program Committee Chairs

Jiazhao Wang, University of Wollongong, Australia

International Technical Committees

Agus Arsad, Universiti Teknologi Malaysia, Malaysia

Ai-Bao Chai, University of Nottingham Malaysia Campus, Malaysia

Alaa El-Din Ali El-Shafei, Mansoura University, Mansoura, Egypt

Blanka Tomková, Technical University of Liberec, Czech Republic

Carlos Rolando Rios-Soberanis, Centro de Investigación Científica de Yucatán, México

Chi-wai Kan, The Hong Kong Polytechnic University, Hong Kong

Dauda Shehu Ibrahim, Bayero University, Nigeria; Southeast University, China

Dongbin Wei, University of Technology Sydney (UTS), Australia

G. RAJARAJAN, Hindustan Institute of Technology & Science (Deemed to be University), India

Guang-Min Luo, National Kaohsiung University of Science & Technology, Taiwan

Hideaki Tsukamoto, Hosei University, Japan

Irshad Ahmad, American University of Ras Al Khaimah, UAE

Juliana Anggono, Petra Christian University, Indonesia

Lilly Mercy, Sathyabama University, India

Mochamad Chalid, Universitas Indonesia, Indonesia

Muhammad Abbas Ahmad Zaini, Universiti Teknologi Malaysia, Malaysia

Nikoloz Jalabadze, Georgian Technical University in Tbilisi, Georgia

Nilofar Asim, National University of Malaysia, Malaysia

Norhazilan Md. Noor, Universiti Teknologi Malaysia, Malaysia

P Bily, Czech Technical University in Prague, Czechia

Patricia Sanmartín Sánchez, Universidade de Santiago de Compostela, Spain

Patrice Berthod, University of Lorraine, France

Redzuan Abdullah, Universiti Teknologi Malaysia, Malaysia

Siti Rahmah Aid, Malaysia-Japan International Institute of Technology (MJIT), Universiti Teknologi Malaysia, Malaysia

Theodoros C. Rousakis, Democritus University of Thrace (D.U.Th.), Greece

Vijay Sekar K.S., SSN College of Engineering, India

Xiangjiang Liu, Zhejiang University, China

Yanisa Laoong-u-thai, Burapha University, Thailand

Yu-Cheng Chen, EPS bio technology corp., Taiwan

Zahira Yaakob, Universiti Kebangsaan Malaysia, Malaysia



PAPER • OPEN ACCESS

Theoretical exploration on free radical scavenging mechanism of curcumin analogues in water solvent

To cite this article: L S P Boli *et al* 2020 *IOP Conf. Ser.: Mater. Sci. Eng.* **856** 012004

View the [article online](#) for updates and enhancements.

Theoretical exploration on free radical scavenging mechanism of curcumin analogues in water solvent

L S P Boli^{1,2}, N D Aisyah², V Khoirunisa^{1,3}, F Rusydi^{2,4}, H Rachmawati⁵, H K Dipojono^{1*}

¹Department of Engineering Physics, Faculty of Industrial Engineering, Institut Teknologi Bandung, Bandung 40132, Indonesia

²Research Center for Quantum Engineering Design, Universitas Airlangga, Surabaya 60115, Indonesia

³Engineering Physics Program, Institut Teknologi Sumatera, Lampung Selatan 35365, Indonesia

⁴Department of Physics, Universitas Airlangga, Surabaya 60115, Indonesia

⁵School of Pharmacy, Institut Teknologi Bandung, Bandung 40132, Indonesia
Department of Physics

*dipojono@gmail.com


Abstract. One way to improve curcumin's possibility in preventing lipid peroxidation is through modification of carbon linker, β -diketone group and aromatic rings. However, there remains a need to explore the role of carbon linker on the free radical scavenging mechanisms of curcumin. This study uses density functional theory to explore two descriptors of free radical scavenging mechanisms, which are bond dissociation enthalpy (BDE) and adiabatic ionization potential (AIP) of curcumin and its analogues. Five analogues with different amount of carbon atoms in the linker in the presence / absence of β -diketone group are chosen. Our result shows that decreasing the amount of carbon atom from 7 to 5 atoms in the linker decreases the BDE at β -diketone group of the curcumin analogue. Moreover, increasing the amount of carbon atoms in the linker decreases the AIP of the analogues.

1. Introduction

Research attention on curcumin is increasing due to its antioxidant activities that offer extensive applications in the health-related fields. Curcumin ($C_{21}H_{20}O_6$) is a yellow compound isolated from the rhizome of *Curcuma longa*. In physiological conditions, curcumin exists in keto and enol tautomers [1,2]. Research about curcumin has been carried to understand antioxidant activity of curcumin which include its free radicals scavenging activity [3] and lipid peroxidation reduction [4]. This activity enables curcumin for application in treating diseases such as Alzheimer's disease, β -Thalassemia, kidney injury, skin disorders, and non-alcoholic fatty liver disease [5–9].

In performing its antioxidant activity, curcumin scavenges free radicals through different scavenging mechanisms. The mechanisms are hydrogen atom transfer (HAT), single electron transfer followed by proton transfer (SETPT), and sequential proton loss electron transfer (SPLET). The mechanisms occur depending on the environment in which curcumin exists and the radicals that react toward curcumin. For example, experiment by Jovanovic showed that in pH 3 to 7 where curcumin keto dominates, HAT mechanism rules the scavenging activity. Meanwhile, in pH above 8 where curcumin enol dominates, SET mechanism governs the activity [10]. Furthermore, theoretical calculation by Galano





Scheme 1. Structure of curcumin and its modified structures. Part **A**, **B**, and **C** denote carbon linker, β -diketone group and aromatic ring.

et al. displayed that HAT and SPLET mechanisms are dominant when curcumin reacts with methoxyl radical ($\bullet\text{OCH}_3$) and trichloromethylperoxyl radical ($\bullet\text{O}_2\text{CCl}_3$) respectively [3].

Curcumin modification is an attempt to improve the free radical scavenging activity. Geometry of curcumin contains carbon linker, β -diketone group and two aromatic rings (Scheme 1). Previous studies have reported that the modification of β -diketone group affects the scavenging mechanism. Mary et al. modified curcumin by forming a metal complex between β -diketone group and certain metal ion and found that the complex enhances HAT and SET mechanism [11]. Li et al. reported that modified curcumin with shorter carbon linker but without β -diketone group is able to scavenge 1,1-diphenyl-2-picryl-hydrazyl (DPPH) radical through the SET mechanism experimentally [12]. Considering the role of the β -diketone group and the carbon linker, further exploration of the linker effect, in the presence and absence of the group, on scavenging mechanisms is necessary.

This paper explores the role of carbon linker on the free radical scavenging mechanism of curcumin by utilizing density functional theory (DFT) calculation to calculate two thermochemical parameters of curcumin analogues. The parameters are bond dissociation enthalpy (BDE) and adiabatic ionization potential (AIP). BDE is measured as the energy releases when an X-H bond of the curcumin is breaking prior to hydrogen transfer, while AIP is measured as the energy required to remove an electron from the curcumin prior to electron transfer. Both BDE and AIP are descriptors for HAT and SET mechanisms respectively.

2. Computational Detail

This research is conducted to explore hydrogen atom transfer and electron transfer, as modelled by (1) and (2), in two stages. First is performing DFT calculation to obtain the geometry of all molecules at ground state. The molecules involved are neutral, radical, and radical ion. The neutral molecules consist of two curcumin tautomers: keto (Molecule 1, with 7 carbon atoms in the linker) and enol (Molecule 2, also with seven carbon atom in the linker), two analogues with 5 and 9 carbon atoms in the linker with β -diketone group (Molecule 3 and 4) and three analogues with 5, 7 and 9 carbon atoms in the linker without β -diketone group (Molecule 5, 6 and 7), as illustrated in scheme 1. The radicals are constructed through hydrogen abstraction at certain X-H_n site (n = 1, 2, 3, 4, and 5) of the neutral molecules. Meanwhile, the radical ions are constructed by applying +1 charge to the neutral molecules. To obtain the geometry of the molecules at ground state, geometry optimization coupled with frequency calculation are performed using M06-2X/6-311++G(d,p) for the neutral molecules and UM06-2X/6-311++G(d,p) for the radical and the radical ion. This step will produce electronic energy and thermal correction to enthalpy of each molecules. Water solvent is included to the calculation by using polarized continuum model (PCM) calculation. All calculation is performed at 298.15 K and 1 atm using Gaussian 09 software [13].

Second is calculating BDE and AIP based on the result of the DFT calculation. The BDE and AIP are calculated according to these models and are presented in (3) and (4) respectively.


Table 1. Bond dissociation enthalpy of the seven molecules in water solvent.

Molecule	Abstraction Site	BDE (kcal/mol)			Molecule	Abstraction Site	BDE (Calc., kcal/mol)
		Calc.	Ref. ^a	Ref. ^b			
1	O-H1	84.3	78.3&78.5	82.3	4	O-H1	83.0
	C-H2	97.9				C-H2	97.6
	C-H3	88.7	92.7	87.4		O-H3	104.6
2	O-H1	84.0	77.5	80.2		C-H4	97.7
	C-H2	96.2				O-H5	83.2
	O-H3	106.8			5	O-H1	84.1
	C-H4	97.8				C-H2	97.8
	O-H5	83.6	76.0		6	O-H1	83.3
3	O-H1	84.0				C-H2	97.7
	C-H2	97.7			7	O-H1	82.8
	O-H3	89.6				C-H2	96.1
	C-H4	98.2					
	O-H5	84.3					

^a Computed using B3LYP/6-311+G(2d,2p)//AM1 calculations. [14]

^b Computed using (RO) B3LYP/6-311+G (2d,2p)//AM1/AM1 calculations. [15]

For this calculation, enthalpy of hydrogen (H(H·)) in water solvent is taken from the value used by szelag et al. [16].



$$\text{BDE} = \text{H}(\text{AX}\cdot) + \text{H}(\text{H}\cdot) - \text{H}(\text{AXH}) \quad (3)$$

$$\text{AIP} = \text{H}(\text{AXH}\cdot^+) - \text{H}(\text{AXH}) \quad (4)$$

Where AXH, AX· and H· represents the 7 molecules, the radical form of the 7 molecules, and a hydrogen atom, respectively.

3. Results and Discussion

3.1. Bond dissociation enthalpy of curcumin analogues

Table 1 lists the chosen abstraction site and the results of the calculated BDE at each abstraction site. Molecule 1, 5, 6 and 7 have symmetrical shape, thus only O-H1 and C-H2 attached to one aromatic ring are chosen as the abstraction sites. C-H3 site in Molecule 1 is chosen as an extra abstraction site since it has been proposed by Jovanovic et al. as curcumin's possible abstraction site [9]. Molecule 2, 3 and 4 are not symmetrical, hence there are five abstraction sites chosen. Our results show that in molecule 1, BDE of O-H1 < C-H3 < C-H2. In Molecule 2, BDE of O-H1 ≈ O-H5 < C-H2 < C-H4 < O-H3. Comparing the BDEs in Molecule 1 and 2, it is concluded that O-H sites in aromatic ring always have the lowest BDEs and their values are similar. This is in agreement with the result reported by Barzegar et al. and Sun et al. [14,15].

Changing the amount of carbon atoms in the linker does not affect the BDE at aromatic ring but it does at the β-diketone group. Molecule 3, 2 and 4 that have 5, 7 and 9 carbon atoms in the linker respectively in the presence of the β-diketone group, show similar BDE values at O-H and C-H sites of aromatic ring. Molecule 5, 6, and 7 that have 5, 7 and 9 carbon atoms in the linker respectively in




Figure 1. Plot of the AIP of the seven molecules.

Table 2. The HOMO energy of the seven molecules.

Molecule	E_{HOMO} (eV)
1	-7.36
2	-7.05
3	-7.33
4	-7.11
5	-7.21
6	-7.08
7	-6.92

the absence of the β -diketone group also show similar BDE values at sites in aromatic ring. Regardless of the amount of carbon linker and the presence / absence of β -diketone group, the BDE at O-H site in aromatic ring remains the lowest among all abstraction sites. Different result is observed at O-H3 sites of β -diketone group in Molecule 2, 3 and 4. Molecule 3 that has 5 carbon atoms in the linker has 17.2 kcal/mol lower BDE value than Molecule 2 that has 7 carbon atoms in the linker. Meanwhile Molecule 4 that has 9 carbon atoms in the linker has only 2.2 kcal/mol lower BDE value than Molecule 2.

This results suggest two important things. First, since low BDE relates to an easy HAT, the seven molecules probably can transfer a hydrogen atom from O-H site at aromatic ring easily. Thus, it is possible for curcumin and its analogues to participate in scavenging free radicals through HAT mechanism. Furthermore, the C-H3 site in Molecule 1 also has to be considered as a possible hydrogen atom transfer site, which has been supported by Jovanovic et al. [9]. Second, changing amount of carbon atom in the linker in the presence of β -diketone group is important since the linker has shown to affect the BDE at the β -diketone group.

3.2. Adiabatic ionization potential of curcumin analogues

Figure 1 plots the calculated AIP of two groups of curcumin analogues, with and without β -diketone group. The figure shows that increasing the amount of carbon atoms from 5 to 7 to 9 reduces the AIP values in both groups with an average of 3.8 kcal/mol. Overall, molecules with β -diketone group has lower AIP values. The molecules with lower AIP values can facilitate SET, the first step of SETPT mechanism, easier [16]. Thus, our result implies that increasing of the amount of carbon linker in the presence of β -diketone group improves the possibility of the molecules to scavenge free radicals through SETPT mechanism.

Table 2 lists the results of highest occupied molecular orbital (HOMO) energy of the seven molecules. Generally, increasing the amount of carbon linker increases the HOMO energy, which could be an indication of an easy electron transfer [17]. Thus, the HOMO results confirm the AIP trend analyzed above.

4. Conclusion

This study uses two descriptors, BDE and AIP, to explore the role of carbon linker on the free radical scavenging mechanisms of curcumin. Decreasing the amount of carbon atom in the linker lowers the BDE value at β -diketone group by 17.2 kcal/mol. This implies that shorter carbon linker probably can increase the possibility of curcumin to scavenge free radical through HAT mechanism at β -diketone group. On the other hand, increasing the amount of carbon atom in the linker reduces the AIP value by 3.8 kcal/mol on average. This implies that longer carbon linker probably can increase the possibility of curcumin to involve in SETPT mechanism.

Acknowledgments

This work is supported by Directorate of Higher Education, Ministry of Research Technology and Higher Education (RISTEKDIKTI), Republic of Indonesia under grant scheme "Penelitian Dasar Unggulan Perguruan Tinggi 2019". LSPB appreciates support from Lembaga Pengelola Dana Pendidikan (LPDP) for the doctoral scholarship.

References

- [1] Tønnesen H H, Karlsen J, Mostad A, Samuelsson B, Enzell C R and Berg J-E 1982 *Acta Chem. Scand. B* **36** 475–9
- [2] Sharma R A, Gescher A J and Steward W P 2005 *Eur. J. Cancer* **41** 1955–68
- [3] Galano A, Álvarez-Diduk R, Ramírez-Silva M T, Alarcón-Ángeles G and Rojas-Hernández A 2009 *Chem. Phys.* **363** 13–23
- [4] Chaniad P, Morales N P, Rojsitthisak P and Luechapudiporn R 2018 *J. Food Biochem.* **42** 1–9
- [5] Bisceglia F, Seghetti F, Serra M, Zusso M, Gervasoni S, Verga L, Vistoli G, Lanni C, Catanzaro M, De Lorenzi E and Belluti F 2019 *ACS Chem Neurosci.* **10** 1420-1433
- [6] Nasser E, Mohammadi E, Tamaddoni A, Qujeq D, Zayeri F and Zand H 2017 *Ann. Nutr. Metab.* **71** 136–44
- [7] Abdel-Magied N and Elkady A A 2019 *Exp. Mol. Pathol.* **111** 104299
- [8] Laura V, Mattia F, Roberta G, Federico I, Emi D, Chiara T, Luca B and Elena C 2019 *Nutrients* **11** 2169
- [9] Mirhafez S R, Farimani A R, Gholami A, Hooshmand E, Tavallaie S and Nobakht M. Gh B F 2019 *Drug Metab. Pers. Ther.* **34** 20180040
- [10] Jovanovic S V, Steenken S, Boone C W and Simic M G 1999 *J. Am. Chem. Soc.* 9677–81
- [11] Mary C P V, Vijayakumar S and Shankar R 2018 *J. Mol. Graph. Model.* **79** 1–14
- [12] Li Q, Chen J, Luo S, Xu J, Huang Q and Liu T 2015 *Eur. J. Med. Chem.* **93** 461–9
- [13] Gaussian 09, Revision D.01, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2013.
- [14] Abolfazl Barzegar 2012 *Food Chem.* **135** 1369-1376
- [15] You-Min Sun, Hong-Yu Zhang, De-Zhan Chen, and Cheng-Bu Liu 2002 *Org. Lett.* **4** 2909-2911
- [16] Małgorzata Szeląg, Alicja Urbaniak, Hans A.R. Bluysen 2015 *Open Chem.* **13** 17-31
- [17] G. Mazzone et al., 2015 *Comput. Theoret. Chem.* **1077** 39-47