

Source details

Indian Journal of Forensic Medicine and Toxicology

CiteScore 2020
0.1

Scopus coverage years: from 2008 to Present
(coverage discontinued in Scopus)

Publisher: Institute of Medico-Legal Publications

SJR 2020
0.115

ISSN: 0973-9122 E-ISSN: 0973-9130

Subject area: [Social Sciences: Law](#) [Medicine: Pathology and Forensic Medicine](#)

SNIP 2020
0.243

[Pharmacology, Toxicology and Pharmaceutics: Toxicology](#) [Environmental Science: Health, Toxicology and Mutagenesis](#)

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

Improved CiteScore methodology

CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020

$$0.1 = \frac{387 \text{ Citations 2017 - 2020}}{3.509 \text{ Documents 2017 - 2020}}$$

Calculated on 05 May, 2021

CiteScoreTracker 2021

$$0.1 = \frac{410 \text{ Citations to date}}{3.815 \text{ Documents to date}}$$

Last updated on 04 July, 2021 • Updated monthly

CiteScore rank 2020

Category	Rank	Percentile
Social Sciences		
— Law	#639/722	11th
Medicine		
— Pathology and Forensic Medicine	#183/191	4th
Pharmacology, Toxicology and	#118/122	3rd

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)



① ×

Register For Free
 Reviewer Certifications
 Register for free and gain more value from your peer reviews.

reviewercredits.com OPEN

Indian Journal of Forensic Medicine and Toxicology

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
India <small>Universities and research institutions in India</small>	Environmental Science Health, Toxicology and Mutagenesis Medicine Pathology and Forensic Medicine Pharmacology, Toxicology and Pharmaceutics Toxicology Social Sciences Law	Indian Journal of Forensic Medicine and Toxicology	20

① ×

**Insights that Matter
2021**

Download The Report For Valuable Guidance That Can Inform Your Security Strategy 2021.

CrowdStrike

Open

PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	09739122, 09739130	2008-2020	Homepage How to publish in this journal editor.ijfmt@gmail.com

How to Claim a Review

Peer Review Certification

All your peer reviews will be certified by Jo and you earn credits to get benefits

reviewercredits.com

OPEN

SCOPE

"Indian Journal of Forensic Medicine & Toxicology" is a double-blind peer reviewed international journal. The frequency is quarterly. It deals with Forensic Medicine, Forensic Science, Toxicology, DNA fingerprinting, sexual medicine, environmental medicine, Forensic Pathology, legal medicine and public health laws.

 Join the conversation about this journal

 Quartiles

FIND SIMILAR JOURNALS

1
Medico-Legal Update

IND

53%
similarity

2
International Journal of
Medical Toxicology and Legal
IND





41%
similarity

3
Journal of Indian Academy of
Forensic Medicine
IND




34%
similarity

4
Journal of Punjab Academy
of Forensic Medicine and
IND

30%
similarity



● Cites / Doc. (4 years)
● Cites / Doc. (3 years)
● Cites / Doc. (2 years)



Indian Journal of Forensic Medicine and Toxicology

Q4 Health, Toxicology and Mutagenesis
... best quartile

SJR 2020 0.12

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" style="border: 1px solid #ccc; padding: 2px 5px; display: inline-block;">https://www.scimagojr.com
```






Indian Journal of Forensic Medicine & Toxicology

Website: www.ijfmt.com



Official Organ of Indian Association of Medico-Legal Experts (Regd.)

Vol. 14 No. 4 (2020): Indian Journal of Forensic Medicine & Toxicology



DOI: <https://doi.org/10.37506/ijfmt.v14i4>

Published: 2020-11-27

Articles

Age Estimation from Second & Third Molar by Modified Gleiser and Hunt Method : A Retrospective Study

Abirami Arthanari¹, Nagabhushana Doggalli², Vidhya A³, Sushma Rudraswamy⁴

1-8



Knowledge Attitude and Practice of Usage of Anti-Anxiety Drugs in Paediatric Patients among General Dental Practitioners

Ahsana A¹ , M. Dhanraj²

9-13



Effect of Silver Diamine Fluoride Treatment on the Quality of Life of Children

Ahsana. A¹ , Deepa Gurunathan²

14-18



A Pilot Study on Snake Bite Deaths in the District of Malda, West Bengal, India

Aloke Mazumder¹ , Tanushree Mondal² , Partha Pratim Mukhopadhyay³

19-25

 PDF

Profile of Dowry Death Cases Reported for Autopsy in Thoothukudi Medical College - A Retrospective Study

Seethalakshmi.M 1 , Rajalakshmi. S.2 , Rajesh.R3 , Vinoth K.V.4 ,

26-28

 PDF

Closed Mitral Valvotomy (as a Surgical Treatment Option) - Does it has any Medico Legal Bearings in the Present Scenario

Amit Chaudhary, Amit Singh2

29-32

 PDF

Chest Injury Outcomes due to Road Traffic Accidents - an Institutional Experience

Amit Chaudhary1, Amit Singh2 , Roopak Aggarwal3

33-38

 PDF

Assessment of Factors Influencing Academic Performance of Nursing Students

Amoldeep1 , Bhawna Sharma2 , Sudhir Gupta3 , Jyoti Sarin4

39-45

 PDF

A Descriptive Correlation Study on Stress and Psychological Wellbeing among Nursing Students in Selected Nursing Institutes of Ambala, Haryana

Anamika Gautam1 , Srinivasan.P2 , Gurvinder Kaur3 , Jyoti Sarin4

46-50

 PDF

Observations on Inter-Rater Agreement in Assessing Fusion Activities at Elbow, Wrist and Pelvis by Conventional and Digital X-Ray Films

Aniruddha Das1 , Srimoyee Mukherjee 2 , Arani Majumder3 , Debasis Sarkar4 , Partha Pratim Mukhopadhyay5

51-57

 PDF

Contriving an Opinion of Cause of Death in Autopsies

Anu Sasidharan1 , Nadia M. Al-Kandary2

58-62

 PDF

A Study of Patterns of Thoracic Injuries in the Victims of Vehicular Incidents Brought to the GMCH Mortuary for Medicolegal Autopsies

Anurupa Choudhury1 , Himangshu Das 2 , Ravi Kumar Sharma3

63-70

 PDF

Causative Factors of Increasing Prescription Drug Abuse based on Transcultural Nursing Theory

Ah Yusuf¹ , Muhammad Tarmizi² , Makhfudli³

972-979

 PDF

The Synergistic Effect of Ascorbic Acid and Selenium on Serum Testosterone in Restraint Stressed Rats

Alaa K. M. Al-amery¹ , Ali A. Al-Fahham²

980-985

 PDF

Comparative Test of Midazolam Hydrochloride Stability in Different Storage and Temperature Container

Anang Endaryanto¹ , Andang Miatmoko² , Mela Dwi Wulandari³ , Novi Aryanti³

986-991

 PDF

Relationship of Age, Work Period, and Education Level with Occupational Accidents in the South Borneo Coal Industry

Arahna Sevanya Tanaya¹ , Tri Martiana² , Zikri Fathur Rahman³

992-997

 PDF

Effect of SafMannan on Local Iraqi Laying Hens Performance

Yasser J. Jameel¹ , Mohannad M. Alrekabi² , and Nihad A. Ali³

998-1003

 PDF

In vitro measurement volume of the neonate thyroid gland

Atyaf Mohammed Ali¹ , Anas Hamed Musleh¹

1004-1010

 PDF

Deleted

1011-1014

Effect of Alpha Lipoic Acid on Polycystic Ovary Syndrome with Insulin Resistance

1015-1020

 PDF

Assessment of Women's Knowledge Concerning Hydatidiform Mole at Maternity Hospitals in Baghdad City

Fatima Fadel Benyian

1021-1027

 PDF

To Assess Knowledge, Attitude and Practice on the Use of Automated External Defibrillation (AED) by Emergency Medical Services (EMS) Providers in Pune, India

Parag Rishipathak¹ , Shrimathy Vijayaraghavan² , Anand Hinduja³

9297-9301

 PDF

Whither to Public Interest-The Curious Case of Compulsory Drug Licensing in Indian Pharmaceutical Industry

Rakesh Kumar Sehgal¹ , R.L. Koul²

9302-9306

 PDF

A Study on Sociodemographic Profile and Comparison of Pattern of Facial Injuries in Victims of Two Wheeler Road Traffic Accident with and without usage of Helmet in South Chennai

P. Shruthi¹ , R.Venkat²

9307-9312

 PDF

Quality of Life among Post CABG Patients

Ambina K.¹ , Shalimol U.S.² , Anjana A.P.²

9313-9320

 PDF

Pulpectomy: A Comprehensive Review

Akankhya Jena

9321-9324

 pdf

Ethical Marketing in Orthodontics: A Review

Sumita Mishra¹ , Ananya Panda² , Smruti B. Nanda³

9325-9328

 pdf

Effect of Nursing Intervention on Prevention of Exclusive Breastfeeding Discontinuation

Sahar S. Kamal¹ , Wafaa E. Ouda² , Safy S. Al-Rafay² , Fatma M. Mohamed³

9329-9334

 pdf

Deleted

211-218

Current Issue

ATOM 1.0

PDF 2.0

RSS 1.0

[Make a Submission](#)

Browse

[Open Journal Systems](#)

Information

[For Readers](#)

[For Authors](#)

[For Librarians](#)

Platform &
workflow by
OJS / PKP

Comparative Test of Midazolam Hydrochloride Stability in Different Storage and Temperature Container

Anang Endaryanto¹, Andang Miatmoko², Mela Dwi Wulandari³, Novi Aryanti³

¹Professor in Department of Pediatrics, Faculty of Medicine, Universitas Airlangga Surabaya, East Java, Indonesia, ²Lecturer in Department of Pharmaceutics, Faculty of Pharmacy, Universitas Airlangga Surabaya, East Java, Indonesia, ³Research staff in Department of Pharmacy, Dr. Soetomo Teaching Hospital, Universitas Airlangga Surabaya, East Java, Indonesia

Abstract

Background: Midazolam hydrochloride is an injection form of benzodiazepines and included in the high alert category in the Dr. Soetomo Teaching Hospital Intensive Care Unit (ICU), Surabaya, Indonesia. Nowadays, the technical preparation of midazolam drugs in the ICU room by doctors and nurses was performed by reconstituting drugs in the injection syringes to be stored for a while in the room. **Objectives:** To evaluate the preparation process for midazolam hydrochloride injection which has been analyzed in Dr. Soetomo Teaching Hospital. **Methods:** Physical, chemical, and microbiological stability of 1 mg/mL midazolam hydrochloride in aqua pro injection solvents have been evaluated at points 0, 8, 12, and 24 hours after preparation, as well as a comparison of stability in storage conditions at room temperature compared to refrigerator temperatures. **Results:** The results showed all the preparations were no foreign particles in the sample container. The midazolam did not experience precipitation in both room and refrigerator temperatures. The pH of midazolam solution during storage proved that the preparation was relatively stable (pH 4.3–4.5). There were no significant differences of midazolam levels in both storages which was in range of 95%. The Microbiology Stability Test showed negative germ growth after 24 hours incubation in both storage. **Conclusion:** This study showed no changes in physical, chemical, or microbiological stability in the midazolam injection samples up to 24 hours after the manufacturing process.

Keywords: Midazolam Hydrochloride, Injection Preparation, Aseptic Technique, Stability Test

Introduction

The stability of pharmaceutical dosage forms is strongly influenced by chemical, physical, and microbiological reactions. These reactions can lead the changes of medicinal properties during the process of shipping, storage, and use. Chemical degradation is known as the cause of a potential decrease or increase in drug toxicity. The storage condition also cause dissolution and pollination process of other medicinal ingredients, so that affect to the stability of the drug¹.

Midazolam hydrochloride is an injection form of benzodiazepines with a short reaction period. Intramuscular route is often used for midazolam administration, then continuous to iv infusion and iv slow bolus². Intramuscular midazolam administration can terminate acute seizure in children besides intranasal midazolam^{3,4}. Previous research showed the stability of midazolam is around 24 hours in normal saline 0.9% or Dextrose 5%^{5,6}. Technical midazolam preparation is usually carried out by the drug reconstitution of injection syringes preparation and drugs storage in the room. It aims to facilitate the administration in emergency condition, which needs immediate administration^{7,8}.

Corresponding Author:

Anang Endaryanto

Department of Pediatrics, Faculty of Medicine,
Universitas Airlangga Surabaya, East Java, Indonesia

Patient safety becomes priority in the management of drug use (MDU). The policy of MDU, namely standard operational procedure for mixing injectable

medicines, explains that the single component does not recommend storing reconstitution drugs in injection syringes, but in sterile vial containers covered by film paper⁹. The reconstitution drugs do not stored for more than 8 hours⁹. However, the MDU policy on injecting drug reconstitution is not supported by accurate data from preliminary research results both qualitative and quantitative aspects.

Baseline reference for the MDU policy for the reconstruction of single preparations is existed in Dr. Soetomo Teaching Hospital. This preliminary study determined the qualitative and quantitative stability of 1 mg/mL midazolam injection in aqua pro injection solvents, whether the stability of the drug was more than 8 hours or 24 hours. The stability test compared to the parameters of the storage container will be observed from the room temperature. The room temperature was 20-25 °C, compared to the refrigerator temperature (2-8 °C). The qualitative stability measured by physical, chemical, and microbiological stability. However, quantitative stability is conducted by determining levels using High-Performance Liquid Chromatography (HPLC) tools. The results of this study can be a reference data for the guidelines evaluation for injecting drug mixing policy in Dr. Soetomo Teaching Hospital. The purpose of this study determined by using qualitative and quantitative stability of midazolam reconstitution HCl 1 mg/mL with storage container parameters.

Methods

Material

Raw midazolam hydrochloride was a gift from Kalbe Farma Factory. Acetonitrile and potassium hydrogen phosphate were Wako products (Osaka, Japan). Sterile glass vials and injection syringes (Terumo®, Tokyo, Japan) were purchased from the Medical Pharmacy Service Unit, Dr. Soetomo Teaching Hospital, Surabaya, Indonesia. Other research materials were analytical grade.

Test Solution Preparation

Preparation of the test solution sample was examined by the same possible conditions that was conducted by the nurse or doctor when reconstituting midazolam hydrochloride in the ICU inpatient ward of Dr. Soetomo

Teaching Hospital. Two midazolam hydrochloride test solutions of 1 mg/mL were prepared for replication three times per test procedure. The first solution was prepared in a Terumo® injection syringe container (containing polypropylene material) and a second solution was prepared in a sterile vial storage container covered with parafilm paper. The first mixing container (5 mL injection syringe, Terumo®) was filled by midazolam hydrochloride 1 mg/mL. The concentration was obtained by mixing technically as follows: A total of 1 ampule of midazolam hydrochloride 5 mg/mL was diluted and a number of 20 mL aqua pro injection to obtain a 1 mg/mL level test solution. The solution was taken 1 mL to be stored in a 5 mL injection syringe. The second solution was prepared in a sterile vial storage container with a dilution procedure: 1 ampoule midazolam hydrochloride 5 mg/mL diluted and 20 mL aqua pro injection to obtain a 1 mg/mL level test solution. Then, 1 mL solution was stored in a sterile vial container.

This study also revealed the effect of storage temperature on the stability of midazolam hydrochloride reconstitution preparations. Three solutions test were stored at room temperature (20-25 °C) and at refrigerator temperature (3-4 °C). At the time of testing, the sample was taken by 1 mL from the storage container using aseptic techniques. The sampling point in this study was carried out at 0 o'clock (shortly after preparation), 8, 24, and 48 hours after preparation.

Preparation Stability Test

The samples were taken at 0, 8, 12, and 24 hours after preparation, then evaluated for physical, chemical, and microbiological stability. Physical stability tests was performed, including observe color's solution and it's clarity. The solution clarity was tested by placing a sample above lighting and observed by rotating the sample and flipping the container against a black or white background (Erweka Apparatus Gmens Hensen Stamm, Germany). Thus, it could clarity and the presence of foreign particles in the test sample were observed.

The sample was taken by 5 mL for physical stability checks. Microbiological stability tests were examined by the sterility requirements of intravenous preparations. Sterility tests were conducted on bacterial growth media in microbiology laboratories. Chemical stability tests was examined including pH testing and levels determination

using HPLC. PH measurement was performed because the pH can affect the preparation stability. the pH measurements of midazolam preparations after storage was conducted by using a pH meter (Eutech Instruments pH 700, Germany) at 25 °C.

Midazolam Hydrochloride Analysis Method Using HPLC

Midazolam hydrochloride levels were quantitatively measured using HPLC. The analysis conditions were as follows: the stationary phase uses an HPLC C18 column, the detector used was a photodiode array (PDA), and the analysis was analyzed at a wavelength of 220 nm, the mobile phase was acetonitrile: 0.3 M potassium dihydrogen phosphate pH 3.3 (3:7 v/v) and a flow rate of 1 mL/minute, and the injection sample volume was 10 µL

Standard solute was prepared by dissolving midazolam hydrochloride at a level of 1 mg/mL in methanol to be injected in the HPLC. Furthermore, it analyzed the peak area of the chromatogram. The results will be used to measure midazolam levels quantitatively by using a comparison of the chromatogram peak area of the sample with the standard 1 mg/mL level midazolam solution.

Statistic Analytic

For qualitative analysis, the results were compared to the initial solution visually. If there were differences, there was a qualitative change in stability. For quantitative analysis, the test solution resulted stable if midazolam hydrochloride levels in the test solution >90% initial levels. Statistical analysis used one-way analysis of variance followed by Tukey's post hoc test with a confidence level of 0.05 to determine the significance differences between all the results The test used SPSS version 23.0 (SPSS, Inc., Chicago, IL).

Results

Solution Clarity test

This study found that all preparations were clear and there were no deposits or foreign particles in the sample container (Table 1). During storage phase, the midazolam did not experience precipitation in both storage. However, there were substances in most vial samples such as fibers (unpublished data) observed in >50% of the test samples (±50 vials) which were not observed in the test samples with polypropylene syringes.

Table 1. The results of the clarity observation of the midazolam solution in Intensive Care Unit's inpatient ward Dr. Soetomo Teaching Hospital (n=3).

No	Sampling Time	pH Midazolam injection			
		Sprit		Vial	
		Room temperature	Refrigator temperature	Room temperature	Refrigator temperature
1	0th hour	*	-	*	-
2	1st hour	*	*	*	*
3	8th hour	*	*	*	*
4	12th hour	*	*	*	*
5	24th hour	*	*	*	*

Note: * = there is no sedimentation

All of samples did not experience significant pH changes, where the mean pH of the samples was around 4.3 to 4.5, at the 1st and 24th hours (Table 2). The test Thus, there was no change in the pH of the midazolam sample showed pH in the range of 4.4 as for the 0th hour. solution during storage, both room temperature and refrigerator temperature, proving that the relatively stable preparation was kept until the 24th hour.

Table 2. The results of the pH test in midazolam in the Intensive Care Unit inpatient ward of Dr. Soetomo Teaching Hospital (n=3).

No	Sampling-Time	pH Midazolam Injection		
		Sput	Vial	
1	0th hour	4.41±0.05	-	4.38±0.05
2	1st hour	4.36±0.05	4.47±0.03	4.39±0.01
3	8th hour	4.41±0.06	4.42±0.05	4.40±0.05
4	12th hour	4.32±0.05	4.49±0.08	4.45±0.06
5	24th hour	4.32±0.05	4.50±0.02	4.36±0.07
		Room temperature	Refrigerator temperature	Room temperature
				Refrigerator temperature

Microbiology Stability Test

The sterility test resulted that 6 test containers representing about 3% of the total sample and showed no microbial contamination in the vial container. In general, after incubation, the test results showed negative results, it can be declared as sterile. However, there was a sample from 3 replications showed a *Staphylococcus pseudintermedius* culture at 8 o'clock (Table 3).

Table 3. The results of the microbiological stability test of midazolam in Intensive Care Unit's inpatient ward Dr. Soetomo Teaching Hospital (n=3).

No	Sample	Microbiological Test Results at the hour		
		Storage temperature	0	1
1	Sput	Room temperature	Negative	Negative
		Refrigerator temperature	-	Negative
2	Vial	Room temperature	Negative	Negative
		Refrigerator temperature	-	Negative
			0	1
			12	24

Stability Test for Midazolam Levels

Thus, it was necessary to limit the number of midazolam injection samples to avoid incorrect technique, so that will obtain perfect sterility. Besides, there were several contaminations, such as fine fibers in most of the glass vial container samples which were not found in the injection syringe container. This was the main evaluation in the sterile vial sterilization process.

These data were expected to be a guide or reference for the preparation process of midazolam injection in hospitals, especially Dr. Soetomo Teaching Hospital, and generally other hospitals in Indonesia with proper aseptic techniques. Further studies need to concern about dose uniformity after the dilution process as important information for proper preparation of midazolam hydrochloride in the ward. This was an attempt to ensure dose uniformity between injection sample, which this study found that the coefficient of variation between replications was still too large, around 3-22%.

Conclusion

This study showed midazolam hydrochloride injection had good physical, chemical, and microbiological stability until the 24th hour after the manufacturing process. Vial and injection syringes did not show a significant effect on drug content in 24-hours storage temperature. However, midazolam injection was possible to be contaminated by microorganism because of less accurate technique during preparation.

Acknowledgement

This research was funded by the Surabaya City Ministry of Health Fund for the 2017 financial year. Thank you to the Dr. Soetomo Teaching Hospital who has facilitated the implementation of this research, especially in the Pharmacy-Hospital Installation, Dr. Microbiology Department, and *BALITBANG*, Surabaya. Furthermore, gratitude was also said to Universitas Airlangga, Faculty of Pharmacy's Testing Services Unit (ULP) for assisting in the determination of the midazolam level with HPLC, as well as all parties who helped to conduct this research well.

Ethical Clearance: This study has been approved ethically by the Medical Research Ethics Committee of Faculty of Medicine, Universitas Airlangga Surabaya.

Conflict of Interest: There is no conflict of interest

References

1. de Diego M, Godoy G, Mennickent S. Chemical stability of midazolam injection by high performance liquid chromatography. *J Sep Sci* 2007;30:1833–8. doi:10.1002/jssc.200700006.
2. Pedersen CA, Schneider PJ, Scheckelhoff DJ. ASHP national survey of pharmacy practice in hospital settings: Prescribing and transcribing—2010. *Am J Heal Pharm* 2011;68:669–88. doi:10.2146/ajhp100711.
3. Gunawan PI, Kurniawan MR, Saharso D. The efficacy and safety of intramuscular midazolam compared with rectal diazepam for anticonvulsant in children. *Pakistan Paediatr J* 2015;39:227–32.
4. Gunawan PI, Rulian F, Saharso D. Comparison of intranasal midazolam and rectal diazepam as anticonvulsant in children. *J Nepal Paediatr Soc* 2015;35:117–22. doi:10.3126/jnps.v35i2.13581.
5. Karlage K, Earhart Z, Green-Boesen K, Myrdal PB. Stability of midazolam hydrochloride injection 1-mg/ml solutions in polyvinyl chloride and polyolefin bags. *Am J Heal Pharm* 2011;68:1537–40. doi:10.2146/ajhp100413.
6. Trissel LA, American Society of Health-System Pharmacists. Handbook on injectable drugs. American Society of Health-System Pharmacists; 2013.
7. Heller M, Veach LM. Administration of Parenteral Medications. *Clin. Med. Assist.*, New York: Delmar; 2008, p. 835–82.
8. World Health Organization. WHO best practices for injections and related procedures toolkit 2010:11–2.
9. Jaruratanasirikul S, Sriwiriyan S. Stability of meropenem in normal saline solution after storage at room temperature. *Southeast Asian J Trop Med Public Health* 2003;34:627–9.
10. Inc. PC. Product Monograph Midazolam Injection USP 2017:1–40.
11. Anderson C, MacKay M. Stability of Fentanyl Citrate, Hydromorphone Hydrochloride, Ketamine Hydrochloride, Midazolam, Morphine Sulfate, and Pentobarbital Sodium in Polypropylene Syringes. *Pharmacy* 2015;3:379–85. doi:10.3390/pharmacy3040379.

By using a comparative analysis of the chromatogram peaks of the raw midazolam material. A quantitative level of the drug sample was obtained (Table 4).

Table 4. HPLC test results on drug levels of midazolam injection samples after 24 hours in the Intensive Care Unit's inpatient ward of Dr. Soetomo Teaching Hospital (n=3).

Sampling-Time	Midazolam Level (Persect)			
	Sput		Vial	
	Room temperature	Refrigator temperature	Room temperature	Refrigator temperature
24th hour	90.31±7.59	91.97±19.93	95.68±3.14	96.85±8.92

The level of midazolam at the 0 minute was lower than the data at the 24th hour, both stored at room temperature and refrigerator. After 24 hours, the levels of medicinal ingredients were still 90.31%-96.85%, both samples stored in vials and polypropylene syringes. There were no significant differences between the storage temperature in the room and the refrigerator. Moreover, midazolam concentration were in the range of 95% after 48 hours in refrigerators, either by using vial containers or injection syringes. At this point, samples at room temperature were not tested due to the technical problems. The coefficient between measurement replications was still too large (in range 3.28%-21.67%).

Discussion

This study proved that midazolam hydrochloride injection, which was prepared using aseptic technique, has good stability until the 24th hour after the manufacturing process. Both glass vial and injection syringes didn't show significant effect on drug content as well as 24-hours storage temperatures. Previous research explained that midazolam stability was influenced by pH, storage temperature, light, and storage containers⁶. This study found that the midazolam injection was still stable, both physically, chemically, and microbiologically until the 24th hour after its mixture. Recent study showed stable diluted concentration of midazolam was 0.03 mg/mL to 0.5 for 24 hours with 5% dextrose or 0.9% sodium chloride injection¹⁰. On the other hand, midazolam 5 mg/mL was stable at room temperature for 100 days in polypropylene syringes¹¹.

Treatment of samples was divided into 2 categories which used different containers, sterile glass vials and injection syringes containing polypropylene. The results showed that there was no significant difference between midazolam injection which prepared in sterile glass vial containers and injection syringes in any storage conditions. PH measurement showed unchanged results between observation times, which around 4.3-4.6 with stable physical conditions and observed the deposition of drug ingredients during storage. Moreover, more than 90% of the initial concentration of midazolam can be observed in injection samples at 24 hours after preparation, even at 48 hours in samples stored in the refrigerator. This data was in line with the previous studies which have shown that midazolam solution was stable even up to 5-weeks after manufactured and there was no midazolam adsorption process in injection syringe containers⁵. In the 0-hour sample, the results of the determination of the levels were smaller than the 24th hour which was about 81-86%. This was possible because of over filling after the dilution process of 5 mg/mL midazolam levels with aqua pro injection to a level of 1 mg/mL which results in a diluted final concentration.

In addition, sterility tests for injection samples have also been revealed. This is an important parameter for preparations of parenteral routes. The results showed that most of the samples showed sterility to germs/microbes. However, there were 1 in 3 sample replications which showed contamination by *Staphylococcus pseudintermedius* contained in the free air. This was possible because there was of working technique on a sample that less accurate due to drugs preparation.