

# The enhancement apoptosis of osteosarcoma mesenchymal stem cells co-cultivation with peripheral blood mononuclear cells sensitized by secretome and granulocyte macrophage colony-stimulating factor

*by Ferdiansyah Mahyudin*

---

**Submission date:** 09-Mar-2021 10:10AM (UTC+0800)

**Submission ID:** 1527969043

**File name:** The\_Enhancement.pdf (2.72M)

**Word count:** 4097

**Character count:** 22852

ORIGINAL ARTICLE

# The enhancement apoptosis of osteosarcoma mesenchymal stem cells co-cultivation with peripheral blood mononuclear cells sensitized by secretome and granulocyte macrophage colony-stimulating factor

Ferdiansyah Mahyudin,  
Hizbillah Yazid, Mouli Edward,  
Mohammad Hardian Basuki,  
Yunus Abdul Bari,  
Fedik Abdul Rantam<sup>1</sup>

Orthopedic and Traumatology  
Department, Faculty of Medicine,  
Dr Soetomo General Hospital,  
Airlangga University, <sup>1</sup>Stem Cell  
Research And Development Center,  
Airlangga University, Surabaya,  
Indonesia

J. Adv. Pharm. Technol. Res.

## ABSTRACT

The advanced, metastasis, and recurrent of osteosarcoma (OS) patients have a poor prognosis postaggressive surgery and chemotherapy. Peripheral blood mononuclear cells (PBMCs) as cell-based immunotherapy may successful in the OS treatment. To investigate the enhancement apoptosis of OS-mesenchymal stem cells (OS-MSCs) co-cultivated with PBMCs sensitized using the secretome and granulocyte macrophage colony-stimulating factor (GM-CSF). This true experimental study with posttest only control group design and *in vitro* study. The sample was cultured OS-MSCs which confirmed by Cluster of Differentiation-133 using immunocytochemistry (ICC) and histopathology analysis. The sample divided into six groups accordingly: OS-MSC, OS-MSC + PBMC, OS-MSC + PBMC + Secretome, OS-MSC + PBMC + GM-CSF, OS-MSC + PBMC + Secretome + GM-CSF ( $n = 5/N = 30$ ). The enhancement of OS-MSCs apoptosis was analyzed through Interleukin-2 (IL-2) level through the Enzyme-Linked Immunosorbent Assay examination, expression of Signal Transducers and Activators of Transcription (STAT)-3 and caspase-3 by ICC. One-way analysis of variance test and Tukey Honestly Significant Difference to analyze the difference between the groups ( $P < 0.05$ ). The highest of IL-2 level was found in the PBMC + Secretome + GM-CSF group. The highest expression of caspase-3 was found in OS-MSC + PBMC + Secretome + GM-CSF group with significant different between groups ( $P < 0.05$ ). There was insignificant difference of STAT-3 expression and IL-2 level between groups ( $P > 0.05$ ). The co-cultivation of OS-MSCs and PBMCs activated using secretome and GM-CSF has a great ability to enhance OS-MSCs apoptosis.

**Key words:** Granulocyte macrophage colony stimulating factor, mesenchymal stem cells, osteosarcoma, peripheral blood mononuclear cells, secretome

## Address for correspondence:

Dr. Ferdiansyah Mahyudin,  
Orthopedic and Traumatology Department, Faculty of  
Medicine, Dr. Soetomo General Hospital, Airlangga University,  
Surabaya 60132, Indonesia.  
E-mail: [ferdiansyah@fk.unair.ac.id](mailto:ferdiansyah@fk.unair.ac.id)

Submitted: 25-Apr-2020

Revised: 06-Jun-2020

Accepted: 25-Aug-2020

Published: 10-Oct-2020

8

## Access this article online

Quick Response Code:



Website:

[www.japtr.org](http://www.japtr.org)

DOI:

10.4103/japtr.JAPTR\_52\_20

## INTRODUCTION

Osteosarcoma (OS) is a malignant tumor with neoplastic cell producing bone that the most common primary sarcoma

9

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprints@medknow.com](mailto:reprints@medknow.com)

**How to cite this article:** Mahyudin F, Yazid H, Edward M, Basuki MH, Bari YA, Rantam FA. The enhancement apoptosis of osteosarcoma mesenchymal stem cells co-cultivation with peripheral blood mononuclear cells sensitized by secretome and granulocyte macrophage colony-stimulating factor. J Adv Pharm Technol Res 2020;11:213-9.



arises from the bone tissues.<sup>[1]</sup> Treatment for the newly diagnosed OS including three main modalities: presurgical chemotherapy, surgical resection, and postsurgical chemotherapy that increased the outcome of localized OS cases. However, on the advanced, metastatic, and recurrent OS cases remain to have a poor prognosis. After treated aggressively by resection and chemotherapy, the 5-year survival rate on the localized OS about 65% and <20% for metastatic cases.<sup>[2]</sup>

Recent treatment failure is related to the inability to target cancer stem cells due to capability of self-renewal, increase tumorigenicity, and chemotherapy resistance.<sup>[3,4]</sup> Cell death resistance and immune screening avoidance are the two main mechanisms of tumor growth. Reinforcing and intensifying the cancer recognition by the immune system can improve the treatment efficacy. Formerly, necrosis was considered as the only cell death mechanism that affects the immune system and stimulates inflammatory response. Otherwise, apoptosis was considered as physiologic cell death pathway without immune stimulation. It is discovered that apoptotic cells were rapidly and specifically recognized by a phagocyte. Compared to classic apoptosis or immunogenic cell death can stimulate and intensify the immune system and response.<sup>[5,6]</sup>

Apoptosis occurs through caspase-dependent and caspase-independent pathways. The caspase-dependent pathway implicates the extrinsic and intrinsic factors. Activated cytokine receptors on the membrane cancer cells will induce Janus Kinase (JAK) and Signal Transducers and Activators of Transcription (STAT-3 and STAT-5). Those activations detain Bcl-xL and activate the Cytochrome-C. Subsequently, apoptotic protease activating factor-1 (APAF-1) will be increased then activate caspase-9 and stimulate caspase-3 that lead to apoptosis of OS cells.<sup>[7,8]</sup>

Nowadays, cell and antibody-based immunotherapy have gained success in childhood cancer management. Peripheral blood mononuclear cells (PBMCs) that considered superior to other cell-based treatment sources due to isolation process is less invasive compared to bone marrow. Moreover, autologous PBMCs administration does not need long-term immunosuppressive treatment and can be performed without ethical problem.<sup>[9,10]</sup>

5 Granulocyte-macrophage colony-stimulating factor (GM-CSF) is a potent cytokine on stimulating the differentiation of myeloid cells. Furthermore, GM-CSF has a role as an additional immune stimulus to generate anti-tumoral immunity. GM-CSF also plays an important task in dendritic cells differentiation.<sup>[11,12]</sup>

The other new treatment modality is stem cell-based application, such as mesenchymal stem cells (MSCs) secretome.<sup>[13]</sup> Secretome application is also considered safer

due to containing no cellular element that makes it is less possibility of mutation and transforms into cancer-associated fibroblast within cancer microenvironment.<sup>[14,15,16]</sup>

GM-CSF, as well as secretome, will increase PBMCs immunogenicity by inducing it's containing immune cells. Thus, The aim of this study is to investigate the enhancement apoptosis of OS-MSCs cultivated with PBMCs activated by secretome and GM-CSF sensitization *in vitro*.

## 16 SUBJECT AND METHODS

### Isolation, expansion, and culture of osteosarcoma mesenchymal stem cells

OS tissues were obtained from an open biopsy of patient at Orthopaedic Department Dr. Soetomo General Hospital Surabaya. Patient was confirmed and filled the written informed consent. The tissues firstly were confirmed as OS by histopathology. The tissues were washed by phosphate-buffered saline (PBS) (Sigma Aldrich, US) containing 5% penicillin/streptomycin (Gibco, US). Tissue samples were sliced using a scissor then put into Erlenmeyer tube with 5 ml Type I collagenase 0.075% within PBS containing 2% of penicillin/streptomycin. Samples were stirred with a magnetic stirrer for 30 min on 37°C temperature. Then 5 ml of  $\alpha$ -minimal essential medium (Gibco, US) containing 20% heat-inactivated fetal bovine serum (Gibco, US) were added into the tube to neutralize the Type I collagenase activity. Samples were filtered afterward by pouring in beaker glass covered by 3 layers of sterile gauze. The supernatant of the filtration was centrifuged 2000 rpm in 5 min, disposed, and resuspension of the pellet by adding PBS. Centrifugation was repeated and disposed off again the supernatant remaining the pellet. This pellet then be resuspended within the complete medium. Those suspensions were cultivated on the culture plate and kept on the incubator 37°C, 5% CO<sub>2</sub>.<sup>[17]</sup>

### Isolation of peripheral blood mononuclear cells

The peripheral blood sample was collected from a healthy volunteer that had been screened for HIV and Hepatitis-B. The blood sample was slowly poured into conical tube 50cc and added with PBS (ratio 1:1) until homogenous suspensions then were slowly poured through the wall of conical containing Ficol 5cc and centrifuged 1600 rpm for 30 min in 26°C. After separated into 4 layers, the second layer with buffy coat form was slowly poured into the conical tube and washed using PBS 10cc. Subsequently, the conical was centrifuged 1600 rpm for 5 min. The pellet then was cultured in the medium.<sup>[17,18]</sup>

### Sensitization of peripheral blood mononuclear cells

PBMCs suspension was divided into 4 groups that would be the control group, sensitized with secretome, sensitized with GM-CSF, also the combination of secretome and



GMCSF (ratio 1:1:1). All groups then were incubated in 37°C for 2 h.

### Co-cultivation of osteosarcoma-mesenchymal stem cells with peripheral blood mononuclear cells

OS-MSCs monolayer cells were trypsinized as done until being a single cell. OS-MSCs then were put into coverslip within M24 until  $5.10^4$  cells. The cells were incubated in 37°C temperature and CO<sub>2</sub> 5% for 24 h. PBMCs sensitized with secretome, GMCSF, and those combinations then were added into each coverslip according to the classified group with ratio 1:1. All of the samples then were incubated in 37°C temperature and CO<sub>2</sub> 5% for 48 h.

### Enzyme linked immunosorbent assay, immunocytochemistry, and immunohistochemistry analysis

Enzyme Linked Immunosorbent Assay (ELISA) model was used to measure Interleukin-2 (IL-2) level was sandwich ELISA Elabscience (E-EL-H0099). Immunocytochemistry was used to evaluate Cluster of Differentiation (CD)-133 and STAT-3 expressions using a fluorescent microscope. Caspase-3 expressions were evaluated using immunohistochemistry analysis.


### Statistical analysis

One-way analysis of variance test and Tukey Honestly Significant Difference was used to evaluate differences between groups ( $P < 0.05$ ).

## RESULTS

The histopathological examination confirmed the OS chondroblastic type [Figure 1a]. Positive expression result of CD-133 was detected [Figure 1b].

The highest level IL-2 results were obtained for PBMCs that sensitized by secretome and GMCSF. Nonsensitized PBMCs as the control group had the lowest levels of IL-2 secretion [Table 1]. The PBMCs culture and results IL-2 level in PBMCs after sensitization in Figure 2a and b.



**Figure 1:** (a) Histopathological examination showed chondroblastic type osteosarcoma tissue (b) Cluster of Differentiation-133 marker results show positive immunocytofluorescence with magnification  $\times 200$

Microscopic evaluation of cultivation from between groups can be seen in Figure 3a OS-MSCs cultivation; 3b OS-MSCs and pure PBMCs cultivation; 3c co-cultivation of OS-MSCs and PBMCs sensitized with secretome; 3d OS-MSCs co-cultivated with PBMCs activated by GMCSF; 3e Co-cultivation of OS-MSCs and PBMCs activated by GMCSF and secretome. The greatest IL-2 level was found in OS-MSCs + PBMC + Secretome group [Table 2]. There was no significant different of IL-2 level between groups [Figure 3f].

The STAT-3 expression as shown in Figure 4a OS-MSCs Cultivation; 4b OS-MSCs and PBMCs Cultivation sensitized with secretome; 4c OS-MSCs cultivation and PBMCs sensitized with GMCSF; 4d The OS-MSCs co-cultivated with PBMCs activated by GMCSF and secretome; The greatest expression of STAT-3 was found in OS + PBMC + GMCSF group [Table 3]. There was no significant different of STAT-3 expression between groups [Figure 4e].

The caspase-3 expression as shown in Figure 5a OS-MSCs only; 5b OS-MSCs co-cultivated with PBMCs sensitized by secretome; 5c OS-MSCs cultivated with PBMCs sensitized by GMCSF; 5d OS-MSCs co-cultivated with PBMCs activated by GMCSF and secretome. The greatest expression of Caspase-3 was found in OS-MSCs + PBMC + Secretome + GMCSF

**Table 1: The interleukin-2 level in peripheral blood mononuclear cells sensitized by secretome, granulocyte macrophage colony stimulating factor and combination**

Group	Mean $\pm$ SD	ANOVA
PBMCs	0.06145 $\pm$ 0.00247	0.011*
PBMCs + secretome	0.06545 $\pm$ 0.00106	
PBMCs + GMCSF	0.06260 $\pm$ 0.00127	
PBMCs + secretome + GMCSF	0.07495 $\pm$ 0.00318	


\*Significant at  $P < 0.05$ . GMCSF: Granulocyte macrophage colony stimulating factor, PBMCs: Peripheral blood mononuclear cells, SD: Standard deviation

**Table 2: Interleukin-2 level in osteosarcoma -mesenchymal stem cells co-cultivated with peripheral blood mononuclear cells sensitized by secretome, Granulocyte macrophage colony stimulating factor and combination**

Group	Mean $\pm$ SD	ANOVA
OS-MSCs	0.0622 $\pm$ 0.00335	0.083
OS-MSCs + PBMCs	0.06670 $\pm$ 0.00945	
OS-MSCs + PBMCs + secretome	0.08150 $\pm$ 0.1127	
OS-MSCs + PBMCs + GMCSF	0.06976 $\pm$ 0.00567	
OS-MSCs + PBMCs + secretome + GMCSF	0.07083 $\pm$ 0.00318	

OS-MSCs: Osteosarcoma -mesenchymal stem cells, GMCSF: Granulocyte macrophage colony stimulating factor, PBMCs: Peripheral blood mononuclear cells, SD: Standard deviation





**Figure 2:** (a) Cultured peripheral blood mononuclear cells in each group with  $\times 200$  magnification (b) the Interleukin-2 level of peripheral blood mononuclear cells sensitized by secretome, granulocyte macrophage colony stimulating factor, and combination

group [Table 4]. There was significant different of Caspase-3 expression between groups [Figure 5e].

## DISCUSSION

In this study, isolation of OS-MSCs was obtained from an open biopsy sample of an OS patient that have been histopathologically confirmed through CD-113 marker.<sup>[19-21]</sup> Most PBMC's is found as naïve cells which possess no function as the effector.<sup>[9,22]</sup> Sensitized PBMC's with secretome and GMCSF before co-cultivation with OS-MSCs hopefully able to increase the potential of PBMCs. PBMCs sensitized with secretome and GMCSF showed a higher amount of secreted IL-2 compared to all the groups. The amount of IL-2 level reflects the capability of PBMCs as immunotherapy agent. IL-2 may act directly as a cytokine which trigger the apoptotic process while also activate Natural killer-Cell which is a crucial cellular immunity in againts cancerous cells.<sup>[23]</sup>

**Table 3: Signal transducers and activators of transcription-3 expression in osteosarcoma -mesenchymal stem cells co-cultivated with peripheral blood mononuclear cells sensitized by secretome, granulocyte macrophage colony stimulating factor and combination**

Group	Mean $\pm$ SD	ANOVA
OS-MSCs	16.230 $\pm$ 2.420	0.978
OS-MSCs+PBMCs+secretome	17.243 $\pm$ 4.785	
OS-MSCs+PBMCs+GMCSF	17.580 $\pm$ 5.495	
OS-MSCs + PBMCs + secretome + GMCSF	17.566 $\pm$ 4.231	

OS-MSCs: Osteosarcoma -mesenchymal stem cells, GMCSF: Granulocyte macrophage colony stimulating factor, PBMCs: Peripheral blood mononuclear cells, SD: Standard deviation


**Table 4: Caspase-3 expression in Osteosarcoma -mesenchymal stem cells co-cultivated with peripheral blood mononuclear cells sensitized by secretome, Granulocyte macrophage colony stimulating factor and combination**

Group	Mean $\pm$ SD	ANOVA
OS-MSCs	8.250 $\pm$ 1.892	0.001*
OS-MSCs + PBMCs	8.500 $\pm$ 2.081	
OS-MSCs + PBMCs + secretome	12.750 $\pm$ 2.986	
OS-MSCs + PBMCs + GMCSF	15.000 $\pm$ 4.966	
OS-MSCs + PBMCs + secretome + GMCSF	20.500 $\pm$ 3.316	

\*Significant at  $P < 0.05$ . OS-MSCs: Osteosarcoma -mesenchymal stem cells, GMCSF: Granulocyte macrophage colony stimulating factor, PBMCs: Peripheral blood mononuclear cells, SD: Standard deviation

The expression of STAT-3 and the level of IL-2 showed insignificant result. However, caspase-3 expression in this study showed significant results regarding the superiority of apoptotic potential in OS-MSCs cultivated with PBMCs activated using GMCSF and secretome. Caspase-3 expression is heavily associated with the process of cell apoptosis through the dependent caspase pathway given its role as an effector component in the final stages of apoptosis.<sup>[7]</sup> PBMCs that were sensitized with secretome and GMCSF have the highest apoptotic potential in co-cultivation with OS-MSCs; and PBMCs that are sensitized with GMCSF alone can be an alternative to be the choice of immune-based therapy regimens for OS.

The GMCSF has a role in the differentiation, maturation, and activation of the PBMCs component in anti-cancer management.<sup>[24]</sup> In addition, secretome can modulate lymphocyte proliferation and resemble the role of MSCs in the treatment of several diseases.<sup>[25,26]</sup> However, if used separately, PBMCs sensitization with GMCSF has statistically better results than sensitization with secretome



**Figure 3:** Microscopic evaluation of cultivation (a) osteosarcoma-Mesenchymal stem cells only (b) osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cells (c) osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cell sensitized with secretome (d) osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cell sensitized with granulocyte macrophage colony stimulating factor (e) osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cell activated using granulocyte macrophage colony stimulating factor and secretome (f) Interleukin-2 levels in osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cells sensitized by secretome, granulocyte macrophage colony stimulating factor, and combination

in caspase-3 expression as the originator of apoptosis OS-MSCs.


This study investigates OS-MSCs apoptosis by the evaluation of intrinsic pathway components including IL-2, STAT-3, and caspase-3 through a process in the mitochondria.<sup>[27,28]</sup> The process was initiated by IL-2 in cytokine receptors that activate JAK and STAT-3 and STAT-5. This activation will inhibit the function of Bcl-xL and stimulate Cytochrome-C activation that increase in APAF-1 triggers caspase-9

activation and caspase-3 stimulation resulted in OS stem cell apoptosis.<sup>[29]</sup> The ability to enhance the apoptosis of OS-MSCs through co-cultivation with PBMCs activated using secretome and GMCSF can be the basis of therapy with less immunological impact compared to chemotherapy.

## CONCLUSION

OS-MSCs co-cultivated with PBMCs activated using by secretome and GMCSF has the greater ability enhancing






**Figure 4:** The positive expression number of Signal Transducers and Activators of Transcription-3 (a) osteosarcoma-Mesenchymal stem cells only, (b) osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cells sensitized by secretome (c) osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cells sensitized by granulocyte macrophage colony stimulating factor (d) osteosarcoma-Mesenchymal stem cells cultivated with Peripheral blood mononuclear cells activated using granulocyte macrophage colony stimulating factor and secretome (e) Signal Transducers and Activators of Transcription-3 expressions in osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cells sensitized by secretome, granulocyte macrophage colony stimulating factor, and combination

OS-MSCs apoptosis. The further investigation before clinical trial on human is necessary.

**Financial support and sponsorship**  
Nil.



**Figure 5:** The positive expression of caspase-3 in (a) osteosarcoma-Mesenchymal stem cells only, (b) osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cells sensitized by secretome (c) osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cells sensitized by granulocyte macrophage colony stimulating factor (d) osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cells activated using granulocyte macrophage colony stimulating factor and secretome (e) The caspase-3 expression in osteosarcoma-Mesenchymal stem cells co-cultivated with Peripheral blood mononuclear cells sensitized by secretome, granulocyte macrophage colony stimulating factor, and combination

**Conflicts of interest**

There are no conflicts of interest.

**REFERENCES**

1. Abarrategi A, Tornin J, Martinez-Cruzado L, Hamilton A, Martinez-Campos E, Rodrigo JP, *et al.* Osteosarcoma: Cells-of-origin, cancer stem cells, and targeted therapies. *Stem Cells Int* 2016;2016:3631764.
2. Wang Z, Li B, Ren Y, Ye Z. T-cell-based immunotherapy for osteosarcoma: Challenges and opportunities.

- Front Immunol 2016;7:353.
3. Durfee RA, Mohammed M, Luu HH. Review of osteosarcoma and current management. *Rheumatol Ther* 2016;3:221-43.
  4. Hattinger CM, Fanelli M, Tavanti E, Vella S, Ferrari S, Picci P, *et al.* Advances in emerging drugs for osteosarcoma. *Expert Opin Emerg Drugs* 2015;20:495-514.
  5. Montico B, Nigro A, Casolaro V, Dal Col J. Immunogenic apoptosis as a novel tool for anticancer vaccine development. *Int J Mol Sci* 2018;19:594.
  6. Van Acker HH, Capsomidis A, Smits EL, Van Tendeloo VF. CD56 in the immune system: More than a marker for cytotoxicity? *Front Immunol* 2017;8:892.
  7. Nagata S. Apoptosis and clearance of apoptotic cells. *Annu Rev Immunol* 2018;36:489-517.
  8. Fortin A, Cregan SP, MacLaurin JG, Kushwaha N, Hickman ES, Thompson CS, *et al.* APAF1 is a key transcriptional target for p53 in the regulation of neuronal cell death. *J Cell Biol* 2001;155:207-16.
  9. Zhang M, Huang B. The multi-differentiation potential of peripheral blood mononuclear cells. *Stem Cell Res Ther* 2012;3:48.
  10. Koek WN, van der Eerden BC, Alves RD, van Driel M, Schreuders-Koedam M, Zillikens MC, *et al.* Osteoclastogenic capacity of peripheral blood mononuclear cells is not different between women with and without osteoporosis. *Bone* 2017;95:108-14.
  11. Yan WL, Shen KY, Tien CY, Chen YA, Liu SJ. Recent progress in GM-CSF-based cancer immunotherapy. *Immunotherapy* 2017;9:347-60.
  12. Shiomi A, Usui T. Pivotal Roles of GMCSF in Autoimmunity and Inflammation. *Mediators of Inflammation*; 2015. p. 2015.
  13. Vieira de Castro J, Gomes ED, Granja S, Anjo SI, Baltazar F, Manadas B, *et al.* Impact of mesenchymal stem cells' secretome on glioblastoma pathophysiology. *J Transl Med* 2017;15:200.
  14. Purnamawati P, Pawitan JA, Rachman A, Wanandi SI. Effects of umbilical cord-and adipose-derived stem cell secretomes on ALDH1A3 expression and autocrine TGF- $\beta$ 1 signaling in human breast cancer stem cells. *F1000 Research* 2018;7:249.
  15. Vizoso FJ, Eiro N, Cid S, Schneider J, Perez-Fernandez R. Mesenchymal stem cell secretome: Toward Cell-free therapeutic strategies in regenerative medicine. *Int J Mol Sci* 2017;18:E1852.
  16. Hartwig S, Raschke S, Knebel B, Scheler M, Irmiler M, Passlack W, *et al.* Secretome profiling of primary human skeletal muscle cells. *Biochim Biophys Acta* 2014;1844:1011-7.
  17. Rantam FA, Ferdiansyah, Nasronudin, Purwati. Stem cell exploration. Surabaya: Airlangga University Press; 2009. p. 70-5.
  18. Riedhammer C, Halbritter D, Weissert R. Peripheral blood mononuclear cells: isolation, freezing, thawing, and culture. *Methods Mol Biol* 2016;1304:53-61.
  19. Brown HK, Tellez-Gabriel M, Heymann D. Cancer stem cells in osteosarcoma. *Cancer Lett* 2017;386:189-95.
  20. Isakoff MS, Bielack SS, Meltzer P, Gorlick R. Osteosarcoma: Current treatment and a collaborative pathway to success. *J Clin Oncol* 2015;33:3029-35.
  21. Li J, Zhong XY, Li ZY, Cai JF, Zou L, Li JM, *et al.* CD133 expression in osteosarcoma and derivation of CD133<sup>+</sup> cells. *Mol Med Rep* 2013;7:577-84.
  22. Kleiveland CR. Peripheral blood mononuclear cells. In: Verhoeckx K, Cotter P, Lopez-Exposito I, Kleiveland C, Lea T, Mackie A, *et al.* editors. *The Impact of Food Bioactives on Health*. Cham(CH): Springer; 2015. p. 161-7.
  23. Guma SR, Lee DA, Ling Y, Gordon N, Kleinerman ES. Aerosol interleukin-2 induces natural killer cell proliferation in the lung and combination therapy improves the survival of mice with osteosarcoma lung metastasis. *Pediatr Blood Cancer* 2014;61:1362-8.
  24. Kaufman HL, Ruby CE, Hughes T, Slingluff CL Jr. Current status of granulocyte-macrophage colony-stimulating factor in the immunotherapy of melanoma. *J Immunother Cancer* 2014;2:11.
  25. Chen J, Sun MX, Hua YQ, Cai ZD. Prognostic significance of serum lactate dehydrogenase level in osteosarcoma: A meta-analysis. *J Cancer Res Clin Oncol* 2014;140:1205-10.
  26. Makridakis M, Vlahou A. Secretome proteomics for discovery of cancer biomarkers. *J Proteomics* 2010;73:2291-305.
  27. Levy DE, Lee CK. What does STAT3 Do? *J Clin Invest* 2002;109:1143-8.
  28. Youle RJ, Strasser A. The BCL-2 protein family: Opposing activities that mediate cell death. *Nat Rev Mol Cell Biol* 2008;9:47-59.
  29. Khan S, Ahmad K, Alshammari EM, Adnan M, Baig MH, Lohani M, *et al.* Implication of caspase-3 as a common therapeutic target for multineurodegenerative disorders and its inhibition using nonpeptidyl natural compounds. *Biomed Res Int* 2015;2015:379817.



# The enhancement apoptosis of osteosarcoma mesenchymal stem cells co-cultivation with peripheral blood mononuclear cells sensitized by secretome and granulocyte macrophage colony-stimulating factor

## ORIGINALITY REPORT

12%

SIMILARITY INDEX

10%

INTERNET SOURCES

9%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://eprints.manipal.edu">eprints.manipal.edu</a> Internet Source	2%
2	Neal J. Meropol. "Coincident Kaposi sarcoma and T-cell lymphoma in a patient with the Wiskott-Aldrich syndrome", American Journal of Hematology, 06/1992 Publication	2%
3	<a href="http://eprints.iums.ac.ir">eprints.iums.ac.ir</a> Internet Source	1%
4	Swagat Mohapatra, Akshaya Kumar Kundu, Smruti Ranjan Mishra, Shantibhusan Senapati, Tushar Jyotiranjana, Gopalaxmi Panda. "HSF1 and GM-CSF expression, its association with cardiac health, and assessment of organ function during heat stress in crossbred Jersey cattle", Research in Veterinary Science, 2021 Publication	1%

5	<a href="https://open.library.ubc.ca">open.library.ubc.ca</a> Internet Source	1 %
6	<a href="http://www.igi-global.com">www.igi-global.com</a> Internet Source	1 %
7	<a href="http://biosignaling.biomedcentral.com">biosignaling.biomedcentral.com</a> Internet Source	1 %
8	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	1 %
9	Lukas Widhiyanto, Dwikora Novembri Utomo, Adrianto Prasetyo Perbowo, Kukuh Dwiputra Hernugrahanto. "Macroscopic and histologic evaluation of cartilage regeneration treated using xenogenic biodegradable porous sponge cartilage scaffold composite supplemented with allogenic adipose derived mesenchymal stem cells (ASCs) and secretome: An in vivo experimental study", <i>Journal of Biomaterials Applications</i> , 2020 Publication	<1 %
10	<a href="http://www.e-journal.unair.ac.id">www.e-journal.unair.ac.id</a> Internet Source	<1 %
11	<a href="http://dspace.nitrkl.ac.in">dspace.nitrkl.ac.in</a> Internet Source	<1 %
12	Soo Hyun Park, Kyoung Woo Kim, Yeoun Sook Chun, Jae Chan Kim. "Human mesenchymal stem cells differentiate into keratocyte-like	<1 %



cells in keratocyte-conditioned medium",  
Experimental Eye Research, 2012

Publication

13

[Www.dovepress.com](http://www.dovepress.com)

Internet Source

<1 %

14

[www.aginganddisease.org](http://www.aginganddisease.org)

Internet Source

<1 %

15

[www.scirp.org](http://www.scirp.org)

Internet Source

<1 %

16

Mansoureh Mirabdollahi, Shaghayegh  
Haghjooy Javanmard, Hojjat Sadeghi-Aliabadi.  
"In Vitro Assessment of Cytokine Expression  
Profile of MCF-7 Cells inResponse to hWJ-  
MSCs Secretome", Advanced Pharmaceutical  
Bulletin, 2019

Publication

<1 %

17

[agritrop.cirad.fr](http://agritrop.cirad.fr)

Internet Source

<1 %

Exclude quotes

Off

Exclude matches

< 10 words

Exclude bibliography

On

# The enhancement apoptosis of osteosarcoma mesenchymal stem cells co-cultivation with peripheral blood mononuclear cells sensitized by secretome and granulocyte macrophage colony-stimulating factor

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---