

Prevalence, intensity and
histopathology of
Zeylanicobdella arugamensis
infestation on groupers reared
on different aquaculture
systems

by Gunanti Mahasri

Submission date: 24-Feb-2021 06:35PM (UTC+0800)

Submission ID: 1516931490

File name: 5._JFD-2020_Sudah_Published.pdf (1.36M)

Word count: 7708

Character count: 41694



Prevalence, intensity and histopathology of *Zeylanicobdella arugamensis* infestation on groupers reared on different aquaculture systems

Gunanti Mahasri¹ | Umi Hafidloh² | Funky P. Pratama² | Dicky Rahmawan² | Sri Subekti³ | Putri D. Wulansari¹ | Muhamad Amin¹

¹Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, Indonesia

²Aquaculture Study Program, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, Indonesia

³Department of Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, Indonesia

Correspondence

Muhamad Amin, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyosari, Surabaya 60113, Indonesia.
Email: muhamad.amin@fpk.unair.ac.id

Abstract

The marine leech, *Zeylanicobdella arugamensis*, is a major threat to aquaculture in grouper-producing countries including Indonesia. This study aimed at investigating prevalence, intensity and histopathology of the ectoparasite in humpback and hybrid groupers cultured in different rearing systems. A total of 260 groupers (60 humpback groupers and 200 hybrid groupers) were used for samples. The marine leech was observed on skin, fins, gills and mouth, followed by histopathological assay on the skin tissue. The results showed that prevalence of the leech in both groupers was higher when they were cultured in the floating net cages compared with the hatchery, $p < .05$. Furthermore, humpback grouper had a higher prevalence than hybrid grouper when they were cultured in a similar system, $p < .05$. Meanwhile, there was no significant difference in intensity between the two groupers, $p > .05$. Within the hybrid groupers, the highest prevalence was obtained from hybrid groupers reared in the earthen ponds. Histopathological studies showed that the infected groupers exhibited inflammation, congestion and erosion of the epidermis layer. Hybrid grouper had more severe histopathological lesions in the skin tissues. These results suggested that species and type of aquaculture system had significantly determined the prevalence, intensity and severity of lesion in *Z. arugamensis* infestation.

KEYWORDS

fish disease, grouper, histopathology, intensity, prevalence

1 | INTRODUCTION

Indonesia is the third largest grouper-producing country in the world after China and Taiwan (Rimmer & Glamuzina, 2019; Session, 2017). The total Indonesian production of groupers in 2015 was recorded at 8,972 tons, 15,089 tons in 2016 and increased significantly to 605,132 tons in 2018 (BKIPM, 2018). The most common species of groupers which are cultured by Indonesian farmers are humpback grouper (*Cromileptes altivelis*) and hybrid "Cantang" grouper (*Epinephelus* sp.). Among the cultured species, humpback grouper

is the most expensive, ~IDR 450,000/kg or ~US\$ 45, much higher than the hybrid grouper, ~IDR 85,000/kg or ~US\$ 8. However, even though cheaper, the hybrid grouper is cultured by many fish farmers due to faster growth, required 8 months to reach marketable size (~400–500 g) compared with humpback grouper which needs at least 20 months (De, Ghaffar, & Das, 2014). However, the development of Indonesian grouper aquaculture has nowadays been hampered by diseases including *Zeylanicobdella arugamensis* infestation.

Zeylanicobdella arugamensis is a marine leech which belongs to the phylum Annelida and Family of Piscicolidae (De Silva, 1963).

The pathogenicity of *Z. arugamensis* has been previously reported in several studies. Sommerville (1998), for instance, reported that a severe infestation intensity of ectoparasites in grouper resulted in acute death. Similarly, a study by Kua (2008) confirmed that fish infested by the ectoparasite died within 3 days following infestation due to secondary infections with pathogenic bacteria such as *Vibrio alginolyticus*. According to Cruz-Lacierda, Toledo, Tan-Fermin, and Bureson (2000), the parasite has been commonly found in Indonesia. However, the first case of *Z. arugamensis* infestation was reported in detail in 2008 by Slamet et al. (2008) in several species including brown-marbled grouper (*Epinephelus fuscoguttatus*), green grouper (*Epinephelus suillus*), humpback grouper (*C. altivelis*), coral grouper (*Epinephelus corallicola*), seabass (*Lates calcarifer*) and snapper (*Lates argentinaculatus*). The next study was reported by Mahardika, Mastuti, Sudewi, and Zafran (2018) infecting hybrid grouper (*Epinephelus fuscoguttatus* × *Epinephelus lanceolatus*) in Gondol, Bali Island (Mahasri, Wulansari, & Imani, 2019), in Situbondo, East Java, Indonesia. The above studies did not provide data on differences in pathogenicity in the different species of groupers, nor did they examine the influence of rearing conditions on the prevalence and intensity of infestation. Thus, more studies that cover broader topics are still required.

This present study was conducted to investigate the prevalence and infestation intensity of *Z. arugamensis* on the two most common groupers species cultured in Indonesia, hybrid grouper and humpback grouper. In addition, the effect of different aquaculture systems (hatchery, floating net cage, concrete pond and earthen pond) was also studied. Furthermore, histopathology of *Z. arugamensis* in skin tissues of infested groupers was further analysed.

2 | MATERIALS AND METHODS

2.1 | Study area

The present study was carried out at the brackish water Aquaculture Development Center (BBAP), Situbondo, East Java, Indonesia (Figure 1). Four aquaculture systems were chosen for collecting samples which were hatchery, floating net cage, concrete pond and earthen pond. Prevalence and intensity were examined *in situ* while the characterization and histopathology assay were carried out at the Laboratory of Anatomy and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga Surabaya, East Java, Indonesia, from September to December 2019.




FIGURE 1 Sampling locations: (a) Earthen pond, (b) concrete pond, (c) hatchery and (d) floating net cage in the Brackishwater Aquaculture Development Center, Situbondo, East Java, Indonesia

2.2 | Fish samples

A total of 260 groupers consisting of 60 humpback groupers and 200 hybrid groupers (*Epinephelus* sp.) with 17–23 g wet weight or 10–13 cm length were collected from four locations at Brackishwater Aquaculture Development Center Situbondo, East Java, Indonesia (Table 1). Each fish was collected randomly using a scope net and placed on a bucket containing sea water and brought to the laboratory for examination.

2.3 | Examination of marine leech

The examination of marine leech on the fish samples was performed by a native method according to Noga (2010). In brief, marine leech was examined on skin, fins, gills and mouth directly and followed by scraping by gently scraping along the side of the body, fins with a scalpel, while gills were clipped. The result of scraping was placed on an object glass and added a drop of physiological buffer solution (PBS). The scraping results were observed using a binocular microscope with a magnification of 40× and 100×. Marine leeches found in this stage were collected, counted and preserved in 5% glycerine alcohol until further uses. Prevalence and intensity of the marine leech on groupers were calculated according to Mahasri, Koesdarto, et al. (2019) with the following formula:

$$\text{Prevalence} = \frac{\sum \text{infected fish}}{\sum \text{fish samples}} \times 100\%$$

$$\text{Intensity} = \frac{\sum \text{marine leeches}}{\sum \text{infected fish}}$$

In addition, the present study determined the level of infestation of the marine parasite in each fish according to a category made by Williams and Bunkley-Williams (1996) (Table 2). The level of infestation was defined as the average number of marine leeches found in every fish sample.

TABLE 1 Fish size (total wet weight and total length) and total samples collected from four different aquaculture systems

Species	Rearing system	Fish size		Number of samples (fish)
		WW ± SD(g)	TL ± SD (cm)	
Humpback grouper	Hatchery	17.64 ± 3.49 ^a	11.63 ± 0.85 ^{A,B}	30
	Floating net cages	17.84 ± 2.66 ^a	11.29 ± 0.77 ^A	30
Hybrid grouper (Cantang)	Hatchery	17.29 ± 2.59 ^a	11.09 ± 0.67 ^A	50
	Floating net cages	17.34 ± 1.64 ^a	11.38 ± 0.43 ^A	50
	Concrete pond	21.86 ± 3.64 ^b	12.33 ± 0.41 ^B	50
	Earthen pond	18.54 ± 1.94 ^{a,b}	11.41 ± 0.63 ^A	50

Abbreviations: SD, standard deviation; TL, total length; WW, wet weight.

Different superscripts indicate that the values are significantly different at $p < .05$.

2.4 | Staining procedure and identification

The marine leeches were stained using Semichon's acetic carmine according to the protocol of Ngamniyom, Manaboon, and Panyarachun (2012). Briefly, the marine leech in 5% glycerine alcohol was washed with piperazine (PZ) solution. Then, the leech was tied and flattened between two slides, followed by immersion in 70% alcohol for 2 min. After that, the leech was immersed in a carmine solution for 8 hr. The next step was transferring the leech into the acidic alcohol solution (70% alcohol + HCl) for 2 min and moved into an alkaline alcohol solution (alcohol + NaHCO₃) for 20 min; thereafter, gradual dehydration was performed by putting in serial alcohol solutions: 70% alcohol for 5 min, 85% alcohol for 5 min and 95% alcohol for 5 min. Then, the leech was mounted in Hung's I solution for 20 min and placed on a clean glass object and drops of Hung's II solution and covered by a glass cover. The leech was afterwards observed under a binocular microscope equipped with a Lucida camera for identification purposes.

The identification of marine leech was performed based on its morphology and morphometric measurement results which were later compared with the parasitic identification key book and journal by Kabata (1985).

2.5 | Histopathology

The histopathological assay was carried out according to a protocol of Muntiha (2001), with slight modifications. In brief, 5 infected fish from each grouper and rearing system were killed and tissue specimens of skin were cut into small pieces of 3 mm thickness and fixed in 10% buffered neutral formalin (BNF) for 24 hr. The fixed tissues were rinsed in tap water, dehydrated through a graded series of ethanol, infiltrated with xylene and then embedded in paraffin wax. Five-micron-thick sections were cut in a rotary microtome from the tissue blocks and mounted on glass slides. The sections were deparaffinized in xylene, rehydrated through decreasing concentrations of ethanol, stained with haematoxylin and eosin (HE) and examined

TABLE 2 Level of infestation

Level of infestation (parasites)	Categories
<1	Very light
1-5	Light
6-10	Moderate
>10-50	Medium
51-100	Heavy
>100-999	Very heavy
>1,000	Superinfection

Note: Sumber: Williams and Bunkley-Williams (1996)

under a light microscope. Histopathological changes including types of lesions such as erosion of epidermis layer, inflammation and congestion were examined from the fish samples, and then, the severity of lesion was determined as light, medium and heavy. The severity of lesion was considered "light" when the type of lesion was found in less than 30% of infected fish, moderate 30%–70% of infected fish and heavy when the type of lesion was found in more than 70% of occurrence.

2.6 | Water quality measurements

Water quality measurements were carried out every week for three weeks at the hatchery, floating cage net, concrete pond and earthen ponds. The water quality parameters were measured using a thermometer for temperature, DO meter (Lutron DO-55509) to measure dissolved oxygen, a pH meter (pHep, HANNA) to measure pH, a digital refractometer (Benchtop) to measure salinity and an ammonia test kit (Redwood Aquatic, API) to measure ammonia level.

2.7 | Data analysis

Data on the prevalence and infestation intensity of *Z. arugamensis* were compared between species and aquaculture systems using multivariate analysis of variance (MANOVA) with statistical software (SPSS version 22) after three assumptions (normal distribution, equal variance and independent) were fulfilled. However,

histopathological data were analysed descriptively by comparing the type and severity of lesions on the infected skin sections as well as the percentage of fish affected by similar types of lesions.

3 | RESULTS

3.1 | Prevalence and intensity

The present study consisted of two studies which were (a) a comparison of prevalence and intensity of marine leech in two-grouper species (humpback grouper and hybrid grouper) which were cultured in similar rearing system (hatchery and floating net cages), and (b) a comparison of prevalence and intensity of marine leech within hybrid grouper cultured in four different rearing systems (hatchery, floating net cage, concrete pond and earthen pond).

The first result showed that the prevalence of marine leech in the humpback grouper was significantly higher than that in the hybrid grouper when they were cultured in similar types of aquaculture rearing systems (floating net cage or hatchery) ($F = 5.75$, $df 1,14$, $p = .03$). Cultured in hatcheries, the mean prevalence of the marine leech was 16.67% for the humpback grouper and zero prevalence for hybrid grouper. While cultured in floating net cages, the prevalence was 53.33% for humpback grouper and 10% for the hybrid grouper (Table 3). Furthermore, the prevalences of the marine leech were significantly higher when both fish were cultured in the floating net cages than when the fish were cultured in the hatchery ($t = 2.851$, $df 10$, $p = .017$). The second comparison indicated that hybrid groupers reared in the earthen pond had the highest prevalence followed by those hybrid groupers reared in concrete ponds and floating net cage, and the lowest was in the hatchery ($F = 346.41$, $df 3,8$, $p < .01$). As presented in Table 3, the prevalence was 100% in the earthen ponds, 31.11% in the concrete ponds, 10% in the floating net cages and 0% in the hatchery. The result also indicated that the marine leeches were mostly attached to pectoral, ventral, anal and caudal fins in both farms (Figure 2). Large numbers of the leeches were also found in the skin folds behind the lower jaw, under the operculum, eyes and inside the mouth.

A similar pattern was obtained in the infestation intensity. As presented in Table 3, the infestation intensity was not significantly different between the two-grouper species when they were cultured in

Fish species	Aquaculture systems	Prevalence (%)	Intensity (leeches/fish)
Humpback grouper	Hatchery	16.67 ± 5.77 ^{a,b}	1.67 ± 0.29 ^B
	Floating net cage	53.33 ± 25.82 ^{a,b}	5.87 ± 4.37 ^B
Hybrid grouper (Cantang)	Hatchery	0.00 ± 0.00 ^a	0.00 ± 0.00 ^A
	Floating net cage	10.00 ± 3.33 ^a	4.83 ± 1.61 ^B
	Concrete pond	31.11 ± 7.69 ^{a,b}	3.79 ± 0.51 ^B
	Earthen pond	100.00 ± 0.00 ^c	14.05 ± 1.71 ^C

Note: (a, b, c), (A,B,C) Superscript in common represents that the values are no significantly different, $p > .05$.

TABLE 3 Prevalence and intensity of marine leech, *Zeylanicobdella arugamensis*, on humpback grouper and hybrid grouper reared in the hatchery, floating net cages, concrete ponds and earthen ponds

FIGURE 2 Marine leech, *Zeylanicobdella arugamensis*, attached to caudal fin (a), gill (b, c) and pectoral fin (d)



a similar type of rearing system ($F = 3.11, df 3, 11, p = .71$). However, infestation intensity was significantly higher in those hybrid groupers when the culture is in the earthen pond compared with the hatchery, floating net cage or concrete ponds ($F = 73.98, df 3, 8, p < .01$). The highest infestation rate was 14, 4, 5 and 0 for the earthen pond, concrete pond, floating net cage and hatchery, respectively. For hump-back grouper, the values ranged from 1 to 2 (hatchery) and 3 to 11 leeches/fish in the floating net cages. While hybrid grouper ranged from 5 to 6 (floating net cage), 1 to 6 (concrete ponds) and 3 to 25 in earthen ponds.

The morphological characteristics of the marine leech viewed under a light microscope with 400x magnification were as follows:

- Colour: appeared to dark-brown colour
- Body form: elongated and cylindrical body narrowing at both ends with oral and caudal suckers, lines with lighter colours on the dorsal and dark black on the ventral.
- Body size: Total length ranged from ~6.6 mm including both anterior and posterior suckers, the area and body width 3.6 mm, the oral sucker up to 1.9 mm in diameter and having a pair of eyes located on the dorsal surface the anterior or oral sucker. While posterior sucker was generally wider than the oral sucker, up to 2.85 mm in diameter

Based on these morphological characteristics, the marine leech was identified as *Z. arugamensis*. The morphological appearance of *Z. arugamensis* live and preserved specimens isolated from the present study is presented in Figure 3.

The infected fish were also showed several abnormalities including in behaviour or anatomical features. The infected fish for instance were swimming slowly, mostly found on the near of aeration, less appetite and often rubbing the body on the surface of pond walls or the net in the floating net cage. The fish skin also appeared to be darker especially around the attachment area of the *Zeylanicobdella*. Other results showed that the attachment and feeding sites of the *Z. arugamensis* exhibited, haemorrhages, frayed fins and swelling around both anterior and posterior sucker areas on the host's skin (Figure 4). Lesions characterized by the presence of white ulcers on the skin surface were observed on the attachment site of *Zeylanicobdella*.

3.2 | Histopathology

There were several histopathological changes in the skin tissues of infected fish including erosion of the epidermal layer, inflammation and congestion. Among the lesion types, the epidermis layer was the worse type being considered as heavy due to more than 70% occurrence (Table 4). Upon excision, skin tissues of infected fish showed that the infected fish was partial loss of the epithelium and disintegrate due to the ectoparasite attachment. While the two other lesion types (inflammation and congestion) were considered moderate and light due to the occurrence of <70% and 30%, respectively. The inflammation or inflammatory cells in the skin tissue due to ectoparasitic infestation were indicated by dilation of local blood vessels resulting in excessive local blood flow marked by leucocytes or white blood cells. Furthermore, the other




FIGURE 3 Morphology of *Zeylanicobdella arugamensis* viewed under a light microscope with 40x magnification after staining with carmine. Description: (1) anterior sucker, (2) posterior sucker. Anterior and posterior parts viewed under a binocular microscope with 100x magnification. (a) Anterior part. Anterior sucker (As), eyespot (Es), suboesophageal ganglionic mass (Sgm), proboscyst (Pb), testis (T), ductus jaculatorius (De), testicular ganglion (Tg) and ovary (Ov). (b) Posterior part. Pulsatile vesicle (Pv), posterior crop caecum (Pcc) and posterior sucker (Ps)

abnormality on the skin section of infested fish was congestion, indicated by the increasing number of eosinophil granule cells due to blood comes out of blood vessels. The histopathological data also showed that the level of skin tissue damages in infested fish showed that the infestation of *Z. arugamensis* had a more severe

histopathological impact on epidermis erosion and congestion on the hybrid grouper compared with the humpback grouper. In addition, the most severe effects of the *Z. arugamensis* infestation were observed from skin tissue of fish reared in the earthen and concrete ponds (Table 4).




FIGURE 4 Frayed fins, haemorrhages and swollen areas of fish skin infected by *Z. arugamensis*. (a) haemorrhages of fin base; (b) pale colour and haemorrhages; (c) swelling and reddish; and (d) swelling and pale colour on fish skin

TABLE 4 Type and severity of lesions on skin histology section of humpback and hybrid groupers infected by *Z. arugamensis*

Species	Rearing system	Type of lesions	Severity
Humpback grouper	Hatchery	Epidermis erosion	Moderate
		Inflammation	Light
		Congestion	Light
	Floating net cages	Epidermis erosion,	Moderate
		Inflammation	Light
		Congestion	Light
Hybrid grouper (Cantang)	Hatchery	-	24
	Floating net cages	Epidermis erosion	Heavy
		Inflammation	Light
		Congestion	Moderate
	Concrete pond	Epidermis erosion	Heavy
		Inflammation	Light
		Congestion	Moderate
	Earthen pond	Epidermis erosion	Heavy
		Inflammation	Light
		Congestion	Moderate

Note: Five fish samples were taken from each fish and rearing groups. Light: <30% occurrence in fish samples, moderate: 30%–70% occurrence and heavy: >70% occurrence.

TABLE 5 Temperature, salinity, pH dissolved oxygen (DO) and ammonia levels in rearing water of groupers reared in the hatchery, concrete ponds, earthen pond and floating net cages

Parameters	Humpback grouper		Hybrid grouper			
	Hatchery	Floating net cage	Hatchery	Floating net cage	Concrete pond	Earthen pond
Temperature (°C)	27.07 ± 3.72 ^a	32.00 ± 1.00 ^a	27.67 ± 3.06 ^a	29.93 ± 1.20 ^a	28.67 ± 0.58 ^a	29.90 ± 1.15 ^a
Salinity (ppt)	31.33 ± 1.53 ^A	30.67 ± 0.58 ^A	32.33 ± 2.52 ^A	33.67 ± 1.53 ^A	32.00 ± 2.65 ^A	31.67 ± 2.52 ^A
pH	7.43 ± 0.40 ^x	8.00 ± 0.20 ^x	7.57 ± 0.51 ^x	7.97 ± 0.15 ^x	7.53 ± 0.45 ^x	7.07 ± 0.12 ^x
DO (mg/L)	5.65 ± 1.55 ^X	4.73 ± 0.31 ^X	5.62 ± 0.46 ^X	4.59 ± 1.47 ^X	3.83 ± 0.46 ^X	5.13 ± 1.94 ^X
Ammonia (mg/L)	2.75 ± 1.39 ^z	1.50 ± 0.50 ^z	2.58 ± 1.66 ^z	0.58 ± 0.14 ^z	2.00 ± 1.32 ^z	0.77 ± 0.12 ^z

Note: Values are mean with three measurement replicates. Common superscripts indicate that the mean values are not significantly different, $p > .05$.

3.3 | Water quality

Water quality measurements showed that temperature, dissolved oxygen (DO), pH, level of ammonia and salinity were not significantly different among the rearing waters, $p_{\text{values}} > .05$. In general, the measurement had 27°C–32°C water temperature, 30–33 ppt salinity, 7–8 pH, 3–6 mg/L DO, 0.5–3 mg/L ammonia (Table 5).

4 | DISCUSSIONS

The present study reported prevalence, intensity and histopathology of *Z. arugamensis* infestation in humpback and hybrid groupers cultured in a hatchery, floating net cages, concrete ponds and earthen ponds in Situbondo, East Java, Indonesia. Based on its morphological characteristics, the marine leech was identified as *Z. arugamensis* (Cruz-Lacierda et al., 2000; Mahardika, Mastuti, Sudewi, et al., 2018;

Murwantoko, Negoro, Isnansetyo, & Zafran, 2018; Nagasawa & Uyeno, 2009; Palm, Yulianto, Theisen, Rueckert, & Kleinertz, 2015). In Indonesia, the similar marine leech had been identified using a molecular method based on nucleotides sequence of mitochondrial cytochrome oxidase subunit I (COI) and confirmed that the leech as *Z. arugamensis* (Murwantoko et al., 2018). Based on these identification results, we could also confirm that the marine leech infecting groupers in the present study was *Z. arugamensis*. The marine ectoparasite was mostly found attached to the body surface, eyes, mouth, ventral fins and dorsal fins. These results were very similar with previous studies in the same species (Azmay, Taruna, Taha, & Arai, 2020; Jamaris, Roza, & Mahardika, 2019; Mahardika, Mastuti, Sudewi, et al., 2018; Murwantoko et al., 2018) or different fish species (Azuar, Raza'i, & Miranti, 2019; Nagasawa, Shimadzu, & Ikeda, 2012; Nagasawa & Uyeno, 2009; Ravi & Shariman Yahaya, 2017). Accordingly, a study by Cruz-Lacierda et al. (2000) also showed that *Z. arugamensis* mostly found attached on fins (dorsal,

pectoral, ventral and caudal), gills, mouth and ventral sites. These results suggest that the marine leech infested their host in very similar sites.

The prevalence level of *Z. arugamensis* in both humpback grouper and hybrid grouper was significantly lower when they were cultured in the hatchery than in the floating net cages. The reason could be the environmental conditions such as temperature, dissolved oxygen, organic matter and salinity in the hatchery were more controllable compared with the floating net cage (Sanders, Fryer, Leith, & Moore, 1972). In addition, high fluctuation of these water quality parameters in the floating net cage could trigger stress conditions and later influenced the immune response of the fish towards parasites (Seng, 2002). Furthermore, the lower prevalence in the hatchery may due to filtration and other treatment of water supply which reduces potential disease including the marine leech or its eggs from external water (Sanders et al., 1972). Other study results indicated that groupers reared in earthen ponds had significantly higher prevalence and infestation intensity compared with groupers reared in the floating net cages. Some possible reasons are higher organic content, high turbidity and water stagnation in the ponds, and these speculative reasons have been confirmed in several studies. Cruz-Lacierta et al. (2000), for example, indicated that a high infestation of *Z. arugamensis* was highly supported by high turbidity of rearing water due to suspended material. Besides, quiet water conditions are known to have an impact on the life cycle of *Z. arugamensis* (Kua, Azmi, & Hamid, 2010). While the infestation intensity obtained in the present study was considered low in all species and rearing system, except in hybrid grouper reared in the earthen pond. According to Williams and Bunkley-Williams (1996), infestation intensity which was >11 individuals/fish is considered as a medium (Table 6). The prevalence levels of *Z. arugamensis* obtained in the present study were slightly lower compared with previous studies from other countries, including 17% on adult and 83% on juvenile fish (Cruz-Lacierta et al., 2000), 100% with an intensity of 4 leeches/fish in Malaysia (Kua, Choong, & Leaw, 2014) and 100% in Brunei Darussalam (Azmeiy et al., 2020). However, the present study results also showed a higher prevalence than previously reported from the hybrid grouper from seven farms in Indonesia which were 59% (Murwantoko et al., 2018).

Ectoparasite infestation has been confirmed to cause stress for fish and has been associated with decreased food intake and reduced growth (Ooué, Terui, Urabe, & Nakamura, 2017; Stauffer et al., 2017) due to their ability to take nutrient from their host. Additionally, the attachment of *Z. arugamensis* can cause skin tissue damages. The damage intensity obtained in the present study was based on scoring value according to Pantung, Helander, Helander, and Cheevaporn (2008). According to Kua (2008) skin damages in infected fish was not solely caused by leech infestation, but could also due to the secondary infection by bacteria or fungi. This was in agreement with Rach, Gaikowski, and Ramsay (2000) who stated that parasites often induce tissue or organ damage in fish due to their presence, attachment and feeding. Furthermore, the parasite induced tissue damage and increased the fish's susceptibility to secondary fungal or bacterial infections, and erosions at the attachment site may create possible routes for the secondary pathogens to enter the body.

The present study compared histopathological changes including types and severity of lesion in skin tissue of humpback and hybrid groupers infested by *Z. arugamensis* reared at different aquaculture system. The result showed that erosion of epidermis layer in the infected fish skin was observed to be the most severe lesion followed by congestion and inflammation. The type and severity of lesions were compared among the fish groups; the result showed that hybrid grouper in general had more severe lesion than humpback groupers (Table 4). The histopathological lesions reported in fish at leech attachment sites caused severe erosion and cause mucus cell loss on the epidermis layer. Similarly, Hirazawa, Ishizuka, and Hagiwara (2016) previously reported the same abnormalities of fish skin tissue due to parasite infestation. A similar result was also documented by Elgendy et al. (2018) in tilapia skin due to the infestation of ectoparasite. While inflammation level of *Z. arugamensis* on both grouper species was not significantly different, a histological section grouper has been reported to show a mild inflammatory response in the epidermis layer due to a parasite infestation by Rizgalla et al. (2016). Furthermore, according to Rohman (2009), the inflammation began with tissue damage and causes rupture of mast cells (cells containing granules) followed by the release of inflammatory mediators, as well as vasodilation

TABLE 6 Prevalence and infestation intensity of *Z. arugamensis* in humpback and hybrid groupers reared in a hatchery, floating net cages and ponds

Fish species	Aquaculture systems	Prevalence (%)	Prevalence category	Intensity (leeches/fish)	Intensity category
Humpback grouper	Hatchery	16.67 ± 5.77 ^{a,b}	Occasionally	1.67 ± 0.29 ^B	Low
	Floating net cage	53.33 ± 25.82 ^{a,b}	Frequently	5.87 ± 4.37 ^B	Low
Hybrid grouper (Cantang)	Hatchery	0.00 ± 0.00 ^c	Almost never	0.00 ± 0.00 ^A	Normal
	Floating net cage	10.00 ± 3.33 ^c	Occasionally	4.83 ± 1.61 ^B	Low
	Concrete pond	31.11 ± 7.69 ^{a,b}	Common	3.79 ± 0.51 ^B	Low
	Earthen pond	100.00 ± 0.00 ^c	Always	14.05 ± 1.71 ^C	Medium
Reference			Williams & Bunkley-Williams (1996)		

7

Note: Different superscripts indicate that the values are significantly different at $p < .05$.

which then causes leucocyte cell migration. In addition, another damage of skin tissue due to the ectoparasite infestation was in the form of congestion, the occurrence of blood damages caused by circulatory disorders which can result in a lack of oxygen and nutrients (Nabib & Maidie, 1987). Congestion is generally preceded by a swelling of the cell where the cell will enlarge so that the blood flow is disrupted; this causes blood damages in several places. According to Flik and Wiegertjes (2005), congestion is an abundance of blood in the area of certain blood vessels. Congestion is generally followed by an increase in the number of eosinophil granule cells. Ressang (1984) added that congestion is the blocking of blood caused by circulatory disorders which can result in a lack of oxygen and nutrients. The cell swelling might be caused by an increase in cell permeability, where cells are unable to maintain ionic homeostasis and fluid resulting in the transfer of extracellular fluid into the cell (Rizgalla et al., 2016).

Many papers have described a strong correlation between the quality parameters of rearing water and parasite infestation in aquaculture industries including temperature (Jansen & Bakke, 1991; Karvonen et al., 2013; Möller, 1978; Scott & Nokes, 2009; Strepparava et al., 2018), salinity (Cheung, Nigrelli, & Ruggieri, 1979; Ernst, Whittington, Corneillie, & Talbot, 2005; Möller, 1978; Rogowski & Stockwell, 2006; Soleng & Bakke, 1997), dissolved oxygen (Mikheev, Pasternak, Valtonen, & Taskinen, 2014; Molnar, 1994; Yoshinaga, 2001), pH and ammonia (Banu, Khan, & Thulin, 2004). However, to the authors' knowledge, publications concerning the correlation of *Z. arugamensis* infestation on groupers are still very limited, only a correlation with temperature (Kua et al., 2010, 2014; Mastuti & Zarman, 2018). Thus, many studies are still required to describe deeper on *Z. arugamensis* infestation on groupers.

15 The five parameters of water quality in the sampling locations fell within the desirable and acceptable limits for the culture of groupers except for temperature and ammonia (Table 7). The water temperature in hybrid grouper fluctuated within a minimum of 25°C and a maximum of 27°C, which is slightly lower than the optimal temperature for the fish, 28°C–32°C (Mursitorini & Ramdhani, 2013; SNI, 2011). At the temperature of 28°C–32°C, *Z. arugamensis* took 2 weeks to grow to a mature adult and start producing cocoons (Kua et al., 2010). Furthermore, Kua et al. (2014) reported that the best temperature for the cocoons to hatch was 27°C, and having the highest survival rate in juvenile and adult stages was recorded at 25°C. This factor might explain why the prevalence and intensity of *Z. arugamensis* were higher in hybrid

grouper cultured in the hatchery and earthen ponds, with a water temperature of 25°C–27°C.

While ammonia level obtained in the present study ranged from 0.5 to 5 mg/L, and according to (SNI, 2011), the value was higher than tolerance levels for grouper, 0.01 mg/L. These water conditions might trigger the emergence of ectoparasite helminth (Fauzi, Mokoginta, & Yaniharto, 2008). To the authors' knowledge, there were no papers that documented the effect of ammonia concentration in rearing water on the prevalence of infestation of *Z. arugamensis*. However, the high level of ammonia could suppress the immune system of groupers. A study by Schuwerack, Lewis, Hoole, and Morley (2002) confirmed that several types of leucocytes were decreased when the fish was exposed to ammonia at a concentration of 0.5 mg/L. Accordingly, Ojwala, Otachi, and Kitaka (2018) concluded that the exposure of fish to a high level of ammonia (>0.05) induced cellular and immunological changes in fish which later highly determined their vulnerability to parasite infestation. Thus, better water quality management is highly recommended for decreasing parasite infestation in grouper cultures.

5 | CONCLUSIONS

The present study demonstrated the infestation of humpback and hybrid groupers cultured in floating net cages, concrete pond, earthen pond and hatchery in Situbondo, East Java, Indonesia. The prevalence and infestation intensity of *Z. arugamensis* on humpback grouper and hybrid grouper were higher when they were cultured in floating net cages than in hatcheries. Another result showed also that hybrid groupers reared in the earthen pond had the highest prevalence and intensity of *Z. arugamensis* compared with those groupers reared in the hatchery, floating net cage or concrete ponds. Most of the marine leech was abundantly distributed in fins, gills and mouth. The marine leech was mostly found attached to fins, gills and mouth. These results may suggest hybrid grouper was more vulnerable to the marine leech infestation, especially when they were cultured in earthen ponds.

ACKNOWLEDGEMENTS

We wish to express our thanks to Brackishwater Aquaculture Development Centre, Situbondo, East Java, Indonesia, for kindly providing fish samples and amenities for this experiment.

10 **34** CONFLICT OF INTEREST

The authors declare that there was no conflict of interest.

8
TABLE 7 Temperature, salinity, pH dissolved oxygen (DO) and ammonia levels in rearing water of groupers reared in the hatchery, ponds and floating net cages

References	Temperature (°C)	Salinity (ppt)	pH	DO (mg/L)	Ammonia (mg/L)
SNI (2011)	28–32	28–33	7.5–8.5	≥5	≤0.01
Mursitorini and Ramdhani (2013)	27–31	≥5	7.8–8.2	30–33	≤1.0

Abbreviations: CP, concrete ponds; EP, earthen pond; FNC, floating net cage; HbG, hybrid grouper; Hc, hatchery; HG, humpback grouper.

DATA AVAILABILITY STATEMENT

36 data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Muhamad Amin  <https://orcid.org/0000-0002-7633-8388>

REFERENCES

- Azmezy, S., Taruna, M., Taha, H., & Arai, T. (2020). Prevalence and infestation intensity of a piscicolid leech, *Zeylanicobdella arugamensis* on cultured hybrid grouper in Brunei Darussalam. *Veterinary Parasitology: Regional Studies and Reports*, 20, 100398. <https://doi.org/10.1016/j.vprsr.2020.100398>
- Azuar, A., Raza'i, T. S., & Miranti, S. (2019). Identifikasi Prevalensi dan Intensitas Ektoparasit pada Ikan Bawal Bintang (*Trachinotus blochii*) di Lokasi Budidaya Kota Tanjungpinang. *Intek Akuakultur*, 3(1), 66–79. <https://doi.org/10.31629/intek.v3i1.1003>
- Banu, A., Khan, M., & Thulin, M. (2004). Water quality, stocking density and parasites of freshwater fish in four selected areas of Bangladesh. *Pakistan Journal of Biological Sciences*, 7(3), 436–440. <https://doi.org/10.3923/pjbs.2004.436.440>
- BKIPM K. (2018). *BKIPM Statistik: Volume Ekspor Produk Perikanan Hidup Ikan Kerapu Tahun 2018*. Retrieved from www.bkipm.kkp.go.id, 11 November 2018.
- Cheung, P., Nigrelli, R., & Ruggieri, G. (1979). Studies on cryptocaryoniasis in marine fish: Effect of temperature and salinity on the reproductive cycle of *Cryptocaryon irritans* Brown, 1951. *Journal of Fish Diseases*, 2(2), 93–97. <https://doi.org/10.1111/j.1365-2761.1979.tb00146.x>
- Cruz-Lacierta, E. R., Toledo, J. D., Tan-Fermin, J. D., & Burreson, E. M. (2000). Marine leech (*Zeylanicobdella arugamensis*) infestation in cultured orange-spotted grouper, *Epinephelus coioides*. *Aquaculture*, 185(3), 191–196. [https://doi.org/10.1016/S0044-8486\(99\)00356-7](https://doi.org/10.1016/S0044-8486(99)00356-7)
- De, M., Ghaffar, M. A., & Das, S. K. (2014). *Temperature effect on gastric emptying time of hybrid grouper (Epinephelus spp.)*. Paper presented at the AIP Conference proceedings.
- De Silva, P. (1963). *Zeylanicobdella arugamensis* gen. nov. and sp. nov. from Arugam Kalapu, Eastern Province, Ceylon. *Spolia Zeylanica*, 30(1), 47–53.
- Elgendy, M. Y., Hassan, A. M., Zaher, M. F. A., Abbas, H. H., Soliman, W.-E.-D., & Bayoumy, E. M. (2018). *Nerocila bivittata* massive infestations in *Tilapia zillii* with emphasis on hematological and histopathological changes. *Asian Journal of Scientific Research*, 11, 134–144. <https://doi.org/10.3923/ajsr.2018.134.144>
- Ernst, I., Whittington, I. D., Corneillie, S., & Talbot, C. (2005). Effects of temperature, salinity, desiccation and chemical treatments on egg embryonation and hatching success of *Benedenia seriola* (Monogenea: Capsalidae), a parasite of farmed *Seriola* spp. *Journal of Fish Diseases*, 28(3), 157–164. <https://doi.org/10.1111/j.1365-2761.2004.00605.x>
- Fauzi, I. A., Mokoginta, I., & Yaniharto, D. (2008). Pemeliharaan ikan kerapu bebek (*Cromileptes altivelis*) yang diberi pakan pelet dan ikan rucah di keramba jaring apung. *Jurnal Akuakultur Indonesia*, 7(1), 65–70.
- Flik, G., & Wiegertjes, G. (2005). *Host-parasite interactions*. New York (NY): Taylor & Francis.
- Hirazawa, N., Ishizuka, R., & Hagiwara, H. (2016). The effects of *Neobenedenia girellae* (Monogenea) infection on host amberjack *Seriola dumerilii* (Carangidae): Hematological and histopathological analyses. *Aquaculture*, 461, 32–39. <https://doi.org/10.1016/j.aquaculture.2016.04.007>
- Jamaris, Z., Roza, D., & Mahardika, K. (2019). Prevalensi ektoparasit pada ikan budidaya di keramba jaring apung di teluk kaping, buleleng, bali. *JFMR (Journal of Fisheries and Marine Research)*, 3(1), 32–40. <https://doi.org/10.21776/ub.jfmr.2019.003.01.4>
- Jansen, P., & Bakke, T. (1991). Temperature-dependent reproduction and survival of *Gyrodactylus salaris* Malmberg, 1957 (Platyhelminthes: Monogenea) on Atlantic salmon (*Salmo salar* L.). *Parasitology*, 102(1), 105–112. <https://doi.org/10.1017/S0031182000060406>
- Kabata, Z. (1985). *Parasites and diseases of fish cultured in the tropics*. London and Philadelphia, PA: Taylor & Francis Ltd.
- Karvonen, A., Kristjánsson, B. K., Skúlason, S., Lanki, M., Rellstab, C., & Jokela, J. (2013). Water temperature, not fish morph, determines parasite infections of sympatric Icelandic threespine sticklebacks (*Gasterosteus aculeatus*). *Ecology and Evolution*, 3(6), 1507–1517. <https://doi.org/10.1002/ece3.568>
- Kua, B. (2008). What is the risk of leech infestation in cultured marine seabass? *FRI Newslet.: Publ. Fish. Res. Inst., Dep. Fish. Malaysia*, 12(1), 22.
- Kua, B. C., Azmi, M. A., & Hamid, N. K. A. (2010). Life cycle of the marine leech (*Zeylanicobdella arugamensis*) isolated from sea bass (*Lates calcarifer*) under laboratory conditions. *Aquaculture*, 302(3), 153–157. <https://doi.org/10.1016/j.aquaculture.2010.02.029>
- Kua, B., Choong, F., & Leaw, Y. (2014). Effect of salinity and temperature on marine leech, *Zeylanicobdella arugamensis* (De Silva) under laboratory conditions. *Journal of Fish Diseases*, 37(3), 201–207. <https://doi.org/10.1111/jfd.12087>
- Mahardika, K., Mastuti, I., Sudewi, S., & Zafran, Z. (2018). Identification and life cycle of marine leech isolated from cultured hybrid grouper in the northern Bali waters of Indonesia. *Indonesian Aquaculture Journal*, 13(1), 41–49. <https://doi.org/10.15578/iaj.13.1.2018.41-49>
- Mahardika, K., Mastuti, I., & Zarman, M. R. (2018). Respon Lintah Laut (*Zeylanicobdella arugamensis*) terhadap salinitas berbeda secara laboratorium. *JFMR-Journal of Fisheries and Marine Research*, 2(3), 208–214. <https://doi.org/10.21776/ub.jfmr.2018.002.03.9>
- Mahasri, G., Koesdarto, S., Sari, D. P. W., Santanumurti, M. B., Kandi, I. W. K., Fitri, S. F., & Amin, M. (2019). Prevalence and intensity of *Trypanosoma* sp. in wild swamp eels (*Synbranchus bengalensis*) marketed in Surabaya, Indonesia. *Biodiversitas Journal of Biological Diversity*, 20(11), <https://doi.org/10.13057/biodiv/d201119>
- Mahasri, G., Wulansari, P. D., & Imani, I. H. (2019). Intensitas Cacing Ektoparasit Ikan Kerapu Tikus *Cromileptes altivelis* pada Keramba Jaring Apung di Perairan Situbondo Jawa Timur. *Jurnal Kelautan Tropis*, 22(2), 135–140. <https://doi.org/10.14710/jkt.v22i2.5295>
- Mikheev, V. N., Pasternak, A. F., Valtonen, E. T., & Taskinen, J. (2014). Increased ventilation by fish leads to a higher risk of parasitism. *Parasites and Vectors*, 7(1), 281. <https://doi.org/10.1186/1756-3305-7-281>
- Möller, H. (1978). The effects of salinity and temperature on the development and survival of fish parasites. *Journal of Fish Biology*, 12(4), 311–323. <https://doi.org/10.1111/j.1095-8649.1978.tb04176.x>
- Molnar, K. (1994). Effect of decreased water oxygen content on common carp fry with *Dactylogyrus vastator* (Monogenea) infection of varying severity. *Diseases of Aquatic Organisms*, 20, 153–157. <https://doi.org/10.3354/dao020153>
- Muntiha, M. (2001). Teknik pembuatan preparat histopatologi dari jaringan hewan dengan pewarnaan hematoksilin dan eosin (H&E). *Temu Teknis Fungsional Non Peneliti*, 1001, 156–163.
- Mursitorini, E., & Ramdhani, P. (2013). Penyakit Ikan Kerapu. *Loka Pemeriksaan Penyakit dan Lingkungan* [Press release].
- Murwantoko, M., Negoro, S. L. C., Isnansetyo, A., & Zafran, Z. (2018). Identification of marine leech and assessment of its prevalence and intensity on cultured hybrid groupers (*Epinephelus* sp.). *Biodiversitas Journal of Biological Diversity*, 19(5), 1798–1804. <https://doi.org/10.13057/biodiv/d190529>
- Nabib, R., & Maidie, M. (1987). *Patologi Khusus Veteriner. Cetakan ke-3*. Bogor: Bagian Patologi, Fakultas Kedokteran Hewan, Institut Pertanian Bogor.
- Nagasawa, K., Shimadzu, N., & Ikeda, Y. (2012). Four new host records for the fish leech *Zeylanicobdella arugamensis* (Hirudinida: Piscicolidae), with an updated host list (1963–2012). *Biogeography*, 14, 143–146.

- Nagasawa, K., & Uyeno, D. (2009). *Zeylanicobdella arugamensis* (Hirudinida, Piscicolidae), a leech infesting brackish-water fishes, new to Japan. *Biogeography*, 11, 125–130.
- Ngamniyom, A., Manaboon, M., & Panyarachun, B. (2012). Thai Medaka, *Oryzias minutillus* Smith, 1945 (Beloniformes: Adrianichthyidae): A new host species of *Clinostomum complanatum* metacercariae (Digenea: Clinostomatidae) and the surface topography by using SEM. *Ching Mai Journal of Science*, 39(3), 540–544.
- Noga, E. J. (2010). *Fish disease: Diagnosis and treatment*, Iowa: John Wiley & Sons.
- Ojwala, R. A., Otachi, E. O., & Kitaka, N. K. (2018). Effect of water quality on the parasite assemblages infecting Nile tilapia in selected fish farms in Nakuru County, Kenya. *Parasitology Research*, 117(11), 3459–3471. <https://doi.org/10.1007/s00436-018-6042-0>
- Ooue, K., Terui, A., Urabe, H., & Nakamura, F. (2017). A delayed effect of the aquatic parasite *Margaritifera laevis* on the growth of the salmonid host fish *Oncorhynchus masou masou*. *Limnology*, 18(3), 345–351. <https://doi.org/10.1007/s10201-017-0514-2>
- Palm, H. W., Yulianto, I., Theisen, S., Rueckert, S., & Kleinertz, S. (2015). *Epinephelus fuscoguttatus* mariculture in Indonesia: Implications from fish parasite infections. *Regional Studies in Marine Science*, 2, 54–70. <https://doi.org/10.1016/j.rsma.2015.07.003>
- Pantung, N., Helander, K. G., Helander, H. F., & Cheevaporn, V. (2008). Histopathological alterations of hybrid walking catfish (*Clarias macrocephalus* x *Clarias gariepinus*) in acute and subacute cadmium exposure. *Environment Asia*, 1, 22–27.
- Rach, J. J., Gaikowski, M. P., & Ramsay, R. T. (2000). Efficacy of hydrogen peroxide to control parasitic infestations on hatchery-reared fish. *Journal of Aquatic Animal Health*, 12(4), 267–273. [https://doi.org/10.1577/1548-8667\(2000\)012<0267:EOHPTC>2.0.CO;2](https://doi.org/10.1577/1548-8667(2000)012<0267:EOHPTC>2.0.CO;2)
- Ravi, R., & Shariman Yahaya, Z. (2017). *Zeylanicobdella arugamensis*, the marine leech from cultured crimson snapper (*Lutjanus erythropterus*), Jerejak Island, Penang, Malaysia. *Asian Pacific Journal of Tropical Biomedicine*, 7(5), 473–477. <https://doi.org/10.1016/j.apjtb.2017.01.018>
- Ressang, A. (1984). *Patologi Khusus Veteriner: Edisi Kedua* (pp. 567–574). Departemen Urusan Research Nasional Republik Indonesia.
- Rimmer, M. A., & Glamuzina, B. (2019). A review of grouper (Family Serranidae: Subfamily Epinephelinae) aquaculture from a sustainability science perspective. *Reviews in Aquaculture*, 11(1), 58–87. <https://doi.org/10.1111/raq.12226>
- Rizgalla, J., Bron, J., Shinn, A., Herath, T., Paladini, G., & Ferguson, H. (2016). Ulcerative dermatitis in wild dusky grouper *Epinephelus marginatus* (Lowe) from Libyan waters. *Journal of Fish Diseases*, 39(12), 1457–1466. <https://doi.org/10.1111/jfd.12485>
- Rogowski, D. L., & Stockwell, C. A. (2006). Parasites and salinity: Costly tradeoffs in a threatened species. *Oecologia*, 146(4), 615–622. <https://doi.org/10.1007/s00442-005-0218-x>
- Rohman, A. (2009). *Kromatografi untuk analisis obat* (p. 1). Yogyakarta: Graha Ilmu.
- Sanders, J., Fryer, J., Leith, D., & Moore, K. (1972). Control of the infectious protozoan *Ceratomyxa shasta* by treating hatchery water supplies. *The Progressive Fish-Culturist*, 34(1), 13–17. [https://doi.org/10.1577/1548-8640\(1972\)34\[13:COTIPC\]2.0.CO;2](https://doi.org/10.1577/1548-8640(1972)34[13:COTIPC]2.0.CO;2)
- Schuwertack, P. M. M., Lewis, J. W., Hoole, D., & Morley, N. J. (2002). Ammonia-induced cellular and immunological changes in juvenile *Cyprinus carpio* infected with the blood fluke *Sanguinicola inermis*. *Parasitology*, 122(3), 339–345. <https://doi.org/10.1017/S0031182001007260>
- Scott, M. E., & Nokes, D. J. (2009). Temperature-dependent reproduction and survival of *Gyrodactylus bullatarudis* (Monogenea) on guppies (*Poecilia reticulata*). *Parasitology*, 89(2), 221–228. <https://doi.org/10.1017/S003118200001256>
- Seng, L. T. (2002). Practical approaches to health management for cage cultured marine fishes. *Aquaculture Asia*, 7(3), 42–48.
- Session, S.-S. (2017). Asia-Pacific Fishery Commission. *Aquaculture*, 2014, 20,000.
- Slamet, B., Tridjoko, T., Agus, P., Setiadharna, T., Giri, N., & Suwiry, K. (2008). Inventarisasi dan Pengendalian Penyakit Parasit pada Induk Ikan Laut di Bak Pemeliharaan. *Jurnal Perikanan Universitas Gadjah Mada*, 10(2), 276–281.
- SNL. (2011). 6487.3. *Ikan krapu bebek (Cromileptes altivelis, Valenciennes)-Bagian 3: Produksi benih* (Vol. 6487.3:2011). Standar Nasional Indonesia.
- Soleng, A., & Bakke, T. A. (1997). Salinity tolerance of *Gyrodactylus salaris* (Platyhelminthes, Monogenea): Laboratory studies. *Canadian Journal of Fisheries and Aquatic Sciences*, 54(8), 1837–1864. <https://doi.org/10.1139/f97-089>
- Stauffer, J., Bruneaux, M., Panda, B., Visse, M., Vasemägi, A., & Ilmonen, P. (2017). Telomere length and antioxidant defense associate with parasite-induced retarded growth in wild brown trout. *Oecologia*, 185(3), 365–374. <https://doi.org/10.1007/s00442-017-3953-x>
- Strepparava, N., Segner, H., Ros, A., Hartikainen, H., Schmidt-Posthaus, H., & Wahli, T. (2018). Temperature-related parasite infection dynamics: The case of proliferative kidney disease of brown trout. *Parasitology*, 145(3), 281–291. <https://doi.org/10.1017/S0031182017001482>
- Sommerville, C. (1998). Parasites of farmed fish. In K. D. Black & A. D. Pickering (Eds.), *Biology of farmed fish* (pp. 146–179). Sheffield, UK: Sheffield Academic Press.
- Williams, E. H., & Bunkley-Williams, L. (1996). *Parasites of offshore big game fishes of Puerto Rico and the western Atlantic*. University of Puerto Rico.
- Yoshinaga, T. (2001). Effects of high temperature and dissolved oxygen concentration on the development of *Cryptocaryon irritans* (Ciliophora) with a comment on the autumn outbreaks of cryptocaryoniasis. *Fish Pathology*, 36(4), 231–235. <https://doi.org/10.3147/jsfp.36.231>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Mahasri G, Hafidloh U, Pratama FP, et al. Prevalence, intensity and histopathology of *Zeylanicobdella arugamensis* infestation on groupers reared on different aquaculture systems. *J Fish Dis*. 2020;00:1–11. <https://doi.org/10.1111/jfd.13219>

Prevalence, intensity and histopathology of *Zeylanicobdella arugamensis* infestation on groupers reared on different aquaculture systems

ORIGINALITY REPORT

10%

SIMILARITY INDEX

7%

INTERNET SOURCES

7%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1 www.infona.pl 2%

Internet Source
- 2 Cruz-Lacierda, E.R.. "Marine leech (*Zeylanicobdella arugamensis*) infestation in cultured orange-spotted grouper, *Epinephelus coioides*", *Aquaculture*, 20000525 1%

Publication
- 3 www.jeeng.net 1%

Internet Source
- 4 Jeff Rach, Mark Gaikowski, Robert Ramsay. "Efficacy of Hydrogen Peroxide to Control Parasitic Infestations on Hatchery-Reared Fish", *Journal of Aquatic Animal Health*, 2000 1%

Publication
- 5 Rachid Boukedjouta, Tobia Pretto, Miriam Abbadi, Lorena Biasini, Anna Toffan, Karim Mezali. " Viral encephalopathy and retinopathy is endemic in wild groupers <1%

(genus spp.) of the Algerian coast ", Journal of Fish Diseases, 2020

Publication

6

Rajiv Ravi, Zary Shariman Yahaya. "Zeylanicobdella arugamensis , the marine leech from cultured crimson snapper (Lutjanus erythropterus), Jerejak Island, Penang, Malaysia", Asian Pacific Journal of Tropical Biomedicine, 2017

Publication

<1 %

7

www.tandfonline.com

Internet Source

<1 %

8

Pingky D. Fitria, Muhamad Amin, Widya P. Lokapirnasari, Mirni Lamid. "Supplementation of fermented coffee-peel flour to increase high-density lipoprotein (HDL) cholesterol, docosahexaenoic acids (DHA) and eicosapentaenoic acids (EPA) deposition in tilapia fillet", Biocatalysis and Agricultural Biotechnology, 2020

Publication

<1 %

9

epdf.pub

Internet Source

<1 %

10

paperity.org

Internet Source

<1 %

11

S H Liyana, L A Sari, N N Dewi, E D Masithah, A M Sahidu, K T Pursetyo. "Distribution

<1 %

patterns and the biomass of bivalves at Segoro Tambak estuary, Sedati, Sidoarjo, East Java", IOP Conference Series: Earth and Environmental Science, 2019

Publication

12

www.banglajol.info

Internet Source

<1 %

13

Hongyu He, Cheng Xu, Ziqin Zhao, Xinyu Qin, Hongmei Xu, Hongwei Zhang. "Low expression of SLC22A18 predicts poor survival outcome in patients with breast cancer after surgery", Cancer Epidemiology, 2011

Publication

<1 %

14

Submitted to Universitas Airlangga

Student Paper

<1 %

15

link.springer.com

Internet Source

<1 %

16

ir.lib.hiroshima-u.ac.jp

Internet Source

<1 %

17

Ekasari, Julie, Muhammad Hanif Azhar, Enang H. Surawidjaja, Sri Nuryati, Peter De Schryver, and Peter Bossier. "Immune response and disease resistance of shrimp fed biofloc grown on different carbon sources", Fish & Shellfish Immunology, 2014.

Publication

<1 %

18

N D Rahayu, L Sulmartiwi, G Mahasri, Muntalim, B Angwarmas, G D Pamenang. " Inventory of ectoparasite helminth on the Hybrid Grouper () from traditional ponds in the Kampung Kerapu Lamongan East Java Indonesia ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

19

P Lestari, D Suprpto, G Mahasri. " Isolation and identification of bacteria in gastrointestinal of eel () that has potential as probiotic ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

20

S O Wijaya, S Subekti, Kismiyati. " The prevalence of benedeniasis in humpback grouper () in floating net cages in Situbondo Regency, East Java, Indonesia ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

21

Saima Siddique, Zarrien Ayub. " Population dynamics and reproduction of (Holothuroidea: Echinodermata) in coastal waters of Pakistan, North Arabian Sea ", Journal of the Marine Biological Association of the United Kingdom, 2015

Publication

<1 %

22	enaca.org Internet Source	<1 %
23	eprints.kaums.ac.ir Internet Source	<1 %
24	id.scribd.com Internet Source	<1 %
25	scitepress.org Internet Source	<1 %
26	www.ansinet.org Internet Source	<1 %
27	www.yumpu.com Internet Source	<1 %
28	H.W. Palm, I. Yulianto, S. Theisen, S. Rueckert, S. Kleinertz. "Epinephelus fuscoguttatus mariculture in Indonesia: Implications from fish parasite infections", Regional Studies in Marine Science, 2015 Publication	<1 %
29	Ketut Mahardika, Indah Mastuti, Sudewi Sudewi, Zafran Zafran. "IDENTIFICATION AND LIFE CYCLE OF MARINE LEECH ISOLATED FROM CULTURED HYBRID GROUPER IN THE NORTHERN BALI WATERS OF INDONESIA", Indonesian Aquaculture Journal, 2018 Publication	<1 %

- | | | |
|----|--|------|
| 30 | Muhammad Dawood Shah, Kazuki Tani, Balu Alagar Venmathi Maran, Yoong Soon Yong et al. "High-resolution chemical profiling and antiparasitic potential of the tropical shrub <i>Dillenia suffruticosa</i> ", Fisheries Science, 2020
Publication | <1 % |
| 31 | theses.dur.ac.uk
Internet Source | <1 % |
| 32 | hdl.handle.net
Internet Source | <1 % |
| 33 | onlinelibrary.wiley.com
Internet Source | <1 % |
| 34 | www.hindawi.com
Internet Source | <1 % |
| 35 | www.slideshare.net
Internet Source | <1 % |
| 36 | Marcos Tavares - Dias. "Toxicity, physiological, histopathological and antiparasitic effects of the formalin, a chemotherapeutic of fish aquaculture", Aquaculture Research, 2021
Publication | <1 % |
| 37 | Muhamad Amin, Mark B Adams, Christopher M Burke, Christopher JS Bolch. "Isolation and screening of lactic acid bacteria associated with the gastrointestinal tracts of abalone at | <1 % |

various life stages for probiotic candidates", Aquaculture Reports, 2020

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Prevalence, intensity and histopathology of Zeylanicobdella arugamensis infestation on groupers reared on different aquaculture systems

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11
