

Oxygen Dissolved Nanobubble Technology Improved the Quality of Pacific White Shrimp Cultivation

by Gunanti Mahasri

Submission date: 22-Feb-2022 05:23PM (UTC+0800)

Submission ID: 1768235319

File name: C6..pdf (349.89K)

Word count: 1665

Character count: 8770

Linko, K. and Paloheimo, M. (1989) Monitoring of the inspired and end-tidal oxygen, carbon dioxide and nitrous oxide concentrations: clinical applications during anaesthesia and recovery. *J. Clin. Monit. Comput.*, **5**: 149-156.

Ramakrishnan, K.S. (2012) Evaluation of buprenorphine and butorphanol as preanaesthetic in midazolam-ketamine-isoflurane anaesthesia for ovariohysterectomy in cats. M.V.Sc. thesis submitted to The Tamil Nadu Veterinary and Animal Sciences University, Chennai, Tamil Nadu.

Senthilkumar, S. (2013) Cardiopulmonary and stress response to low flow isoflurane anaesthesia with opioid analgesics in cattle. Ph.D. thesis submitted to Tamil Nadu Veterinary and Animal Sciences University, Chennai, Tamilnadu.

Thangadurai, R. (2012) General anaesthesia with intermittent positive pressure ventilation for thorcotomy in cattle with

traumatic pericarditis. Ph.D. thesis submitted to The Tamil Nadu Veterinary and Animal Sciences University, Chennai, Tamil Nadu.

Tranquilli, W.J., Thurmon, J.C. and Grimm, K.A. (2007) Lumb and Jones' Veterinary Anesthesia and Analgesia. 4th edn., Blackwell Publishers, Iowa, USA. pp. 513.

Turner, P., Kerr, C.L., Healy, A. J. and Taylor, M. (2006) Effect of meloxicam and butorphanol on minimum alveolar concentration of isoflurane in rabbits. *Am. J. Vet. Res.*, **67**: 770-774.

Yamashita, K., Okano, N., Yamashita, M., Umar, M.A., Kushiro, T. and Muir, W.W. (2008) Effects of carprofen and meloxicam with or without butorphanol on minimum alveolar concentration of sevoflurane in dogs. *J. Vet. Med. Sci.*, **70**: 29-35.

Indian Vet. J., May 2019, 96 (05) : 37 - 39

Oxygen Dissolved Nanobubble Technology Improved the Quality of Pacific White Shrimp Cultivation

Gunanti Mahasri¹, Ade Irmalia Harifa and Sudarno

Department of Fish Health Management, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia.

(Received : October, 2018 358/18 Accepted : December, 2018)

Abstract

This study aims to determine the effect of nanobubble on dissolved oxygen to reduce Presumptive Vibrio Count (PVC) in Pacific White Shrimp (*Litopenaeus vannamei*) aquaculture system. The results of the analysis showed that the use of nanobubble had a significant effect ($p < 0.05$) in dissolved oxygen and reduce PVC while it had no significant difference in PVC on aquaculture media. The use of nanobubble technology provided optimal dissolved oxygen for Pacific White Shrimp aquaculture and lower PVC value of *Vibrio* sp. in the hepatopancreas compared to aerators treatment.

Key words: Nanobubble, Dissolved Oxygen, Quality, Pacific White Shrimp

The high loss due to *Vibrio* sp. bacte-

rial infection is one of the challenges on Pacific White Shrimp. One of the technologies developed to improve the quality of aquaculture is nanobubble since it can be made available for a long time so as to maintain the dissolved oxygen in the waters remains stable and related to the respiratory and metabolic systems (Mahasri et al., 2018; Takahashi and Chiba, 2007). The purpose of this study was to determine its effect on dissolved oxygen (DO) and Presumptive Vibrio Count (PVC) on Pacific White Shrimp.

Materials and Methods

This study used 240 of Pacific White Shrimp (7-10 cm, 4-6 g) and 30 of each treatment. The treatments were A1 (nanobubble), A2 (aerator), B1 (0 days), B2 (10 days), B3 (20 days) and B4 (30 days). The dissolved oxygen was measured by DO meter tool. PVC was a calculation method to determine the number of *Vibrio* sp. bacteria

¹Corresponding author : Email : mahasritot@gmail.com

Table I. DO on the aquaculture media of Pacific White Shrimp (Meant ± SD)

Treatments	Duration (days)	Dissolved Oxygen ± SD
Nanobubble	0	11.06 ^c ± 0.72
	10	9.63 ^d ± 0.24
	20	9.04 ^{cd} ± 0.21
	40	9.41 ^c ± 0.36
Aerator	0	3.4 ^a ± 0.30
	10	5.41 ^b ± 0.22
	20	4.85 ^b ± 0.42
	40	5.06 ^b ± 0.26

Means bearing different superscripts in a column differs significantly ($p < 0.05$)

Table II. PVC of *Vibrio* sp. bacteria on the aquaculture media of Pacific White Shrimp (Meant ± SD)

Treatments	Duration (days)	PVC ± SD
Nanobubble	0	2.65 ^c ± 0.04
	10	1.75 ^{abc} ± 0.13
	20	1.2 ^a ± 0.31
	40	2.13 ^{bc} ± 0.08
Aerator	0	3.4 ^a ± 0.04
	10	5.41 ^b ± 0.08
	20	4.85 ^b ± 0.22
	40	5.06 ^b ± 0.11

Means bearing different superscripts in a column differs significantly ($p < 0.05$)

in the sample. PVC *Vibrio* sp. measurement was conducted in TCBS media growth and calculated with SNI01-2332.3-2006 method (Letchumanan *et al.*, 2005). The targets were hepatopancreas and water media. They were taken based on a previous study (Kusdarwati *et al.*, loc cit). Data analysis of DO and PVC used ANOVA (Analysis of Variance) and Duncan Multiple Range Test (Kusriningrum, 2008).

Results and Discussion

The results of DO measurement in the aquaculture media of Pacific White Shrimp could be seen in Table I. It showed significant differences between nanobubble with aerators treatments ($p < 0.05$).

The nanobubble application in the Pacific White Shrimp showed higher and more stable DO compare with aerator. Nanobubble had a bubble size that contained oxygen gas in the water which was < 200 nm (Ebina *et al.*, 2013). It was due to its float rate at a level of 0.361 feet per second or 3610 times faster than

microbubble that floated at rate of 0.0001 feet per second (Takahashi and Chiba, loc cit).

The result of *Vibrio* sp. PVC on aquaculture media is showed in Table II. It indicated a nonsignificant difference between nanobubble treatment with aerator ($p > 0.05$).

The result of *Vibrio* sp. PVC hepatopancreas is showed in Table III. It indicated a significant difference between nanobubble treatment with aerator ($p > 0.05$). The application of nanobubble on PVC of *Vibrio* sp in aquaculture media showed nonsignificant different than aerator. The bubbles from aerator and nanobubble would capture solids and pollutants suspended in the water and lift them to the surface. They were able to penetrate small cavities in contaminants so that they could wrap and lift the pollutants to the surface. It would maintain the water quality and organic matters in water (Maimunah and Kilawati, 2014). The pollutant or organic matter that accumulated in water was decomposed and decrease the ammonia level which posed a

Table III. PVC of *Vibrio* sp. bacteria on the hepatopancreas of Pacific White Shrimp (Meant ± SD)

Treatments	Duration (days)	PVC ± SD
Nanobubble	0	3.65 ^{dc} ± 0.04
	10	3.25 ^{cd} ± 0.01
	20	2.65 ^{ab} ± 0.04
	40	2.53 ^a ± 0.03
Aerator	0	3.55 ^{dc} ± 0.03
	10	3.08 ^c ± 0.05
	20	2.9 ^{bc} ± 0.02
	40	3.7 ^e ± 0.03

Means bearing different superscripts in a column differs significantly ($p < 0.05$)

threat to aquatic organisms (Mukti *et al.*, 2014). Poor water quality could cause stress on shrimp and disrupted its growth and body's resistance to infectious disease (Diansari *et al.*, 2013).

The application of nanobubble on PVC of *Vibrio* sp in hepatopancreas showed a significantly different than aerator. The increased oxygen in the medium increased the metabolism of Pacific White Shrimp so it could accelerate the deleterious effects of the pathogen in its body (Barbieri *et al.*, 2016). Characteristics of *Vibrio* sp. was an opportunistic pathogen, it could develop as pathogen if its environmental conditions and host deteriorate (Utami, 2016). The nanobubble application could improve the quality of the aquaculture system by maintaining a balance between the environment, host (shrimp) and pathogen. Prevention was the most effective action compared to treatment since it was carried out before the outbreak occurred and less cost (Mahasri, 2016).

Summary

The application of nanobubble has improved the quality of aquaculture by maintaining a balance between the environment, host and the pathogen.

References

Barbieri, E., de Medeiros, A.M.Z. and Henriques, M.B. (2016) Oxygen consumption and ammonia excretion of juvenile pink shrimp (*Farfantepenaeus paulensis*) in culture: temperature effects. *Marine and freshwater behaviour and physiology*, **49**(1): 19-25.

Diansari, R. V. R., Arini, E., and Elfitasari, T. (2013) Pengaruh kepadatan yang berbeda terhadap kelulushidupan dan pertumbuhan ikan nila (*Oreochromis niloticus*) pada sistem resirkulasi dengan filter zeolit. *J. of Aqua. Manage. and Tech.*,

2(3): 37-45.

Ebina, K., Shi, K., Hirao, M., Hashimoto, J., Kawato, Y., Kaneshiro, S., Morimoto, T., Koizumi, K. and Yoshikawa, H., (2013) Oxygen and air nanobubble water solution promote the growth of plants, fishes, and mice. *PLoS One*, **8**(6): p.e65339.

Kusdarwati, R., Sudarno, Rozi and Nindarwi, D.D. (2016) Petunjuk Praktikum Analisis Penyakit Ikan I. Fakultas Perikanan dan Kelautan Universitas Airlangga. pp. 8-49.

Kusriningrum, R.S. (2008) Perancangan percobaan. *Universitas Airlangga. Surabaya*. pp.43-51.

Letchumanan, V., Yin, W. F., Lee, L. H., and Chan, K. G. (2015) Prevalence and antimicrobial susceptibility of *Vibrio* parahaemolyticus isolated from retail shrimps in Malaysia. *Frontiers in Microb.*, **6**: 33.

Mahasri, G. (2016) Pengembangan teknologi nanobubble dan sistem budidaya terpadu pada nila dan ikan lele sebagai upaya pengendalian penyakit dan peningkatan produksi perikanan. Laporan Akhir Intensif Riset Sinas RT 10.04. Lemabaga Penelitian dan Inovasi Universitas Airlangga Surabaya, p 59.

Mahasri, G., Saskia, A., Apandi, P.S., Dewi, N.N. and Usuman, N.M. (2018) Development of an aquaculture system using nanobubble technology for the optimization of dissolved oxygen in culture media for Nile tilapia (*Oreochromis niloticus*). *IOP Conference Series: Earth and Environmental Science*, **137**(1): p. 012046. IOP Publishing.

Maimunah, Y., and Kilawati, Y. (2015) Kualitas Lingkungan Tambak Insentif Litopenaeus Vannamei Dalam Kaitannya Dengan Prevalensi Penyakit White Spot Syndrome Virus. *Res. J. of Life Sci.*, **2**(1): 50-59.

Mukti, A. T., M. Arief and W. H. Satyantini. (2014) Dasar-dasar Akuakultur. Program Studi Budidaya Perairan, Fakultas Perikanan dan Kelautan, Universitas Airlangga, pp. 27-52

Takahashi, M., and Chiba, K. M. (2007) Oxygen Nanobubble Water and Method of Producing the Same. *US20070286795*.

Utami, W. (2016) Pengaruh Salinitas Terhadap Efek Infeksi *Vibrio* Harveyi Pada Udang Vaname (*Litopenaeus Vannamei*). *J. of Aqua. Manag. and Tech.*, **5**(1): 82-90.

Oxygen Dissolved Nanobubble Technology Improved the Quality of Pacific White Shrimp Cultivation

ORIGINALITY REPORT

11%

SIMILARITY INDEX

6%

INTERNET SOURCES

9%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Y R Alissianto, Z A Sandriani, B S Rahardja, Agustono, Rozi. " The effect of amino acid lysine and methionine addition on feed toward the growth and retention on mud crab () ", IOP Conference Series: Earth and Environmental Science, 2018
Publication 2%
- 2 www.lib.unair.ac.id
Internet Source 2%
- 3 Makesh, M.. "Development of monoclonal antibodies against VP28 of WSSV and its application to detect WSSV using immunocomb", Aquaculture, 20061116
Publication 1%
- 4 Submitted to Universitas Airlangga
Student Paper 1%
- 5 scientific.cloud-journals.com
Internet Source 1%
- 6 www.science.gov
Internet Source 1%

7 S P Febri, A Fikri, Hanisah, S Nazlia, Zuraidah, A Putriningtias, T M Faisal, Murdhiani, Muliari. "Application of virgin coconut oil in feed in efforts to increase growth and survival rate of red tilapia (Oreochromis sp.)", IOP Conference Series: Earth and Environmental Science, 2021
Publication

1 %

8 Nutrition & Food Science, Volume 46, Issue 3 (2016)
Publication

1 %

9 S.V. Rama Rao, M.V.L.N. Raju, A.K. Panda, M.R. Reddy. "Sunflower seed meal as a substitute for soybean meal in commercial broiler chicken diets", British Poultry Science, 2007
Publication

1 %

10 Dwi Febrianti, Munti Yuhana, Widanarni .. "Dietary Synbiotic Microcapsule Influence the Immune Responses, Growth Performance and Microbial Populations to White Spot Syndrome Virus in Pacific White Shrimp (Litopenaeus vannamei)", Journal of Fisheries and Aquatic Science, 2015
Publication

1 %

Exclude bibliography On

Oxygen Dissolved Nanobubble Technology Improved the Quality of Pacific White Shrimp Cultivation

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
